

## DOMAIN DEFORMATIONS AND EIGENVALUES OF THE DIRICHLET LAPLACIAN IN A RIEMANNIAN MANIFOLD

AHMAD EL SOUFI AND SAÏD ILIAS

ABSTRACT. For any bounded regular domain  $\Omega$  of a real analytic Riemannian manifold  $M$  we denote by  $\lambda_k(\Omega)$  the  $k$ -th eigenvalue of the Dirichlet Laplacian of  $\Omega$ . In this paper, we consider  $\lambda_k$  as a functional on the set of domains of fixed volume in  $M$ . We introduce and investigate a natural notion of critical domain for this functional. In particular, we obtain necessary and sufficient conditions for a domain to be critical, locally minimizing or locally maximizing for  $\lambda_k$ . These results rely on Hadamard type variational formulae that we establish in this general setting.

As an application, we obtain a characterization of critical domains of the trace of the heat kernel under Dirichlet boundary conditions.

### 1. Introduction

Isoperimetric eigenvalue problems constitute one of the main topics in spectral geometry and shape optimization. Given a Riemannian manifold  $M$ , a natural integer  $k$  and a positive constant  $V$ , the problem is to optimize the  $k$ -th eigenvalue of the Dirichlet Laplacian, considered as a functional on the set of all bounded domains of volume  $V$  of  $M$ .

The first result in this subject is the famous Faber-Krahn Theorem [14], [20], originally conjectured by Rayleigh, stating that Euclidean balls minimize the first eigenvalue of the Dirichlet Laplacian among all domains of given volume. Extensions of this classical result to higher order eigenvalues, combinations of eigenvalues, as well as domains of other Riemannian manifolds or subjected to other types of constraints, have been obtained during the last decades, and a very rich literature is devoted to this subject (see, for instance, [2], [3], [4], [5], [8], [9], [10], [11], [18], [25], [26], [30], [35], [36], [37] and the references therein).

A fundamental tool in the proof of many results concerning the first Dirichlet eigenvalue is the following variation formula, known as Hadamard's formula

---

Received July 18, 2005; received in final form May 4, 2007.

2000 *Mathematics Subject Classification.* 49R50, 35P99, 58J50, 58J32.

©2007 University of Illinois

(see [17], [15], [32], [33]):

$$\frac{d}{d\varepsilon}\lambda_1(\Omega_\varepsilon)|_{\varepsilon=0} = - \int_{\partial\Omega_0} v \left( \frac{\partial\phi}{\partial\nu} \right)^2 d\sigma,$$

where  $\lambda_1(\Omega_\varepsilon)$  stands for the first Dirichlet eigenvalue of the domain  $\Omega_\varepsilon$ ,  $\frac{\partial\phi}{\partial\nu}$  denotes the normal derivative of the first normalized eigenfunction  $\phi$  of the Dirichlet Laplacian on  $\Omega_0$  and  $v$  is the normal displacement of the boundary induced by the deformation. This formula shows that a necessary and sufficient condition for a domain  $\Omega \subset \mathbb{R}^n$  to be critical for the Dirichlet first eigenvalue functional under fixed volume variations is that its first Dirichlet eigenfunctions are solutions of the following overdetermined problem:

$$\begin{cases} \Delta\phi = \lambda_1(\Omega)\phi \text{ in } \Omega, \\ \phi = 0 \text{ on } \partial\Omega, \\ \left| \frac{\partial\phi}{\partial\nu} \right| = c \text{ on } \partial\Omega, \end{cases}$$

for some constant  $c$ . Since the first Dirichlet eigenfunction does not change sign in  $\Omega$ , it follows from the well known symmetry result of Serrin [34] that  $\phi$  is radial and  $\Omega$  is a round ball. Therefore, Euclidean balls are the only critical domains of the Dirichlet first eigenvalue functional under fixed volume deformations.

Notice that Hadamard's formula remains valid for any higher order eigenvalue  $\lambda_k$  as long as  $\lambda_k(\Omega)$  is simple. However, when  $\lambda_k(\Omega)$  is degenerate, a differentiability problem arises. Our first aim in this paper (see Section 3) is to overcome this problem and introduce a natural and simple notion of critical domain. Indeed, using perturbation theory of unbounded self-adjoint operators in Hilbert spaces, we will see that, for any deformation  $\Omega_\varepsilon$ , analytic in  $\varepsilon$ , of a domain  $\Omega$  of a real analytic Riemannian manifold  $M$ , and any natural integer  $k$ , the function  $\varepsilon \mapsto \lambda_k(\Omega_\varepsilon)$  admits left-sided and right-sided derivatives at  $\varepsilon = 0$ . Of course, when  $\Omega$  is a local extremum of  $\lambda_k$ , these derivatives have opposite signs. This suggests to define critical domains of  $\lambda_k$  to be the domains  $\Omega$  such that, for any analytic volume-preserving deformation  $\Omega_\varepsilon$  of  $\Omega$ , the right-sided and left-sided derivatives of  $\lambda_k(\Omega_\varepsilon)$  at  $\varepsilon = 0$  have opposite signs. That is,

$$\frac{d}{d\varepsilon}\lambda_k(\Omega_\varepsilon)|_{\varepsilon=0^+} \times \frac{d}{d\varepsilon}\lambda_k(\Omega_\varepsilon)|_{\varepsilon=0^-} \leq 0,$$

which means that  $\lambda_k(\Omega_\varepsilon) \leq \lambda_k(\Omega) + o(\varepsilon)$  or  $\lambda_k(\Omega_\varepsilon) \geq \lambda_k(\Omega) + o(\varepsilon)$  as  $\varepsilon \rightarrow 0$ .

After giving, in Section 2, a general Hadamard type variation formula, we derive in Section 3 necessary and sufficient conditions for a domain  $\Omega$  of the Riemannian manifold  $M$  to be critical for the  $k$ -th Dirichlet eigenvalue functional under volume-preserving domain deformations. For instance, we show (Theorem 3.3) that if  $\Omega$  is a critical domain of the  $k$ -th Dirichlet eigenvalue under volume-preserving domain deformations, then there exists a family of

eigenfunctions  $\phi_1, \dots, \phi_m$  satisfying the following system:

$$(1) \quad \begin{cases} \Delta\phi_i = \lambda_k(\Omega) \phi_i \text{ in } \Omega, \text{ for all } i \leq m, \\ \phi_i = 0 \text{ on } \partial\Omega, \text{ for all } i \leq m, \\ \sum_{i=1}^m \left(\frac{\partial\phi_i}{\partial\nu}\right)^2 = 1 \text{ on } \partial\Omega. \end{cases}$$

Moreover, this necessary condition is also sufficient when either  $\lambda_k(\Omega) > \lambda_{k-1}(\Omega)$  or  $\lambda_k(\Omega) < \lambda_{k+1}(\Omega)$ , which means that  $\lambda_k(\Omega)$  corresponds to the first one or the last one in a cluster of equal eigenvalues. On the other hand, we prove that if  $\lambda_k(\Omega) > \lambda_{k-1}(\Omega)$  (resp.  $\lambda_k(\Omega) < \lambda_{k+1}(\Omega)$ ) and if  $\Omega \subset M$  is a local minimizer (resp. maximizer) of the  $k$ -th Dirichlet eigenvalue functional under volume-preserving domain deformations, then  $\lambda_k(\Omega)$  is simple and the absolute value of the normal derivative of its corresponding eigenfunction is constant along the boundary  $\partial\Omega$  (Theorem 3.1).

The final section deals with the trace of the heat kernel under Dirichlet boundary conditions, defined for a domain  $\Omega \subset M$  by

$$Y_\Omega(t) = \int_\Omega H(t, x, x) v_g = \sum_{k \geq 1} e^{-\lambda_k(\Omega)t},$$

where  $H$  is the fundamental solution of the heat equation in  $\Omega$  under Dirichlet boundary conditions. Luttinger [23] proved an isoperimetric Faber-Krahn-like result for  $Y(t)$  considered as a functional on the set of bounded Euclidean domains; that is, for any bounded domain  $\Omega \subset \mathbb{R}^n$  and any  $t > 0$ , one has  $Y_\Omega(t) \leq Y_{\Omega^*}(t)$ , where  $\Omega^*$  is an Euclidean ball whose volume is equal to that of  $\Omega$ .

For any smooth deformation  $\Omega_\varepsilon$  of  $\Omega$ , the corresponding heat trace function  $Y_\varepsilon(t)$  is always differentiable w.r.t.  $\varepsilon$  and the domain  $\Omega$  will be called critical for the trace of the Dirichlet heat kernel at time  $t$  if, for any volume-preserving deformation  $\Omega_\varepsilon$  of  $\Omega$ , we have

$$\frac{d}{d\varepsilon} Y_\varepsilon(t) \Big|_{\varepsilon=0} = 0.$$

After giving the first variation formula for this functional (Theorem 4.1), we show that a necessary and sufficient condition for a domain  $\Omega$  to be critical for the trace of the Dirichlet heat kernel at time  $t$  is that the Laplacian of the function  $x \mapsto H(t, x, x)$  must be constant along the boundary  $\partial\Omega$  (Corollary 4.1).

Using the Minakshisundaram-Pleijel asymptotic expansion of  $Y(t)$ , one can derive necessary conditions for a domain to be critical for the trace of the Dirichlet heat kernel at every time  $t > 0$ . For instance, we show that the boundary of such a domain necessarily has constant mean curvature (Theorem 4.2).

Thanks to Alexandrov type results (see [1], [24]), one deduces that when the ambient space  $M$  is Euclidean, hyperbolic, or a standard hemisphere, then

geodesic balls are the only critical domains of the trace of the Dirichlet heat kernel at every time  $t > 0$  (Corollary 4.3).

**2. Hadamard type variation formulae**

Let  $\Omega$  be a regular bounded domain of a Riemannian oriented manifold  $(M, g)$ . We will denote by  $\bar{g}$  the metric induced by  $g$  on the boundary  $\partial\Omega$  of  $\Omega$ . Let us start with the following general formula:

PROPOSITION 2.1. *Let  $(g_\varepsilon)$  be a differentiable variation of the metric  $g$ . Let  $\phi_\varepsilon \in C^\infty(\Omega)$  be a differentiable family of functions and  $\Lambda_\varepsilon$  a differentiable family of real numbers such that, for all  $\varepsilon$ ,  $\|\phi_\varepsilon\|_{L^2(\Omega, g_\varepsilon)} = 1$  and*

$$\begin{cases} \Delta_{g_\varepsilon} \phi_\varepsilon = \Lambda_\varepsilon \phi_\varepsilon & \text{in } \Omega, \\ \phi_\varepsilon = 0 & \text{on } \partial\Omega. \end{cases}$$

Then,

$$\begin{aligned} \frac{d}{d\varepsilon} \Lambda_\varepsilon|_{\varepsilon=0} &= \int_\Omega \phi_0 \Delta' \phi_0 v_g \\ &= - \int_\Omega \langle d\phi_0 \otimes d\phi_0 + \frac{1}{4} \Delta \phi_0^2 g, h \rangle v_g, \end{aligned}$$

where  $h := \frac{d}{d\varepsilon} g_\varepsilon|_{\varepsilon=0}$ ,  $\Delta' := \frac{d}{d\varepsilon} \Delta_{g_\varepsilon}|_{\varepsilon=0}$  and  $\langle \cdot, \cdot \rangle$  is the inner product induced by  $g$  on the space of covariant tensors.

*Proof.* For simplicity, let us introduce the following notations:

$$\lambda := \Lambda_0, \quad \phi := \phi_0, \quad \phi' := \frac{d}{d\varepsilon} \phi_\varepsilon|_{\varepsilon=0}, \quad \Lambda' := \frac{d}{d\varepsilon} \Lambda_\varepsilon|_{\varepsilon=0}.$$

Differentiating the two sides of the equality  $\Delta_{g_\varepsilon} \phi_\varepsilon = \Lambda_\varepsilon \phi_\varepsilon$  we obtain

$$\Delta' \phi + \Delta \phi' = \Lambda' \phi + \Lambda \phi'.$$

After multiplication by  $\phi$  and integration we get

$$\int_\Omega \phi \Delta' \phi v_g + \int_\Omega \phi \Delta \phi' v_g = \Lambda' + \lambda \int_\Omega \phi \phi' v_g.$$

Integration by parts gives

$$\int_\Omega \phi \Delta \phi' v_g = \lambda \int_\Omega \phi \phi' v_g + \int_{\partial\Omega} \left( \frac{\partial \phi}{\partial \nu} \phi' - \phi \frac{\partial \phi'}{\partial \nu} \right) v_{\bar{g}}.$$

Thus,

$$\Lambda' = \int_\Omega \phi \Delta' \phi v_g + \int_{\partial\Omega} \left( \phi' \frac{\partial \phi}{\partial \nu} - \phi \frac{\partial \phi'}{\partial \nu} \right) v_{\bar{g}}.$$

It is clear that the boundary integral in this last equation vanishes (since  $\phi_\varepsilon = 0$  on  $\partial\Omega$ ). In conclusion, we have

$$(2) \quad \Lambda' = \int_\Omega \phi \Delta' \phi v_g.$$

Now,  $\Delta'$  is given by (see [4])

$$(3) \quad \Delta' \phi = \langle D^2 \phi, h \rangle - \langle d\phi, \delta h + \frac{1}{2} d\tilde{h} \rangle,$$

where  $\tilde{h}$  is the trace of  $h$  w.r.t.  $g$  (that is,  $\tilde{h} = \langle g, h \rangle$ ). Integration by parts yields

$$(4) \quad \begin{aligned} \int_{\Omega} \phi \langle d\phi, \delta h \rangle v_g &= \frac{1}{2} \int_{\Omega} \langle D^2 \phi^2, h \rangle v_g \\ &= \int_{\Omega} \langle d\phi \otimes d\phi + \phi D^2 \phi, h \rangle v_g \end{aligned}$$

and

$$(5) \quad \int_{\Omega} \phi \langle d\phi, d\tilde{h} \rangle v_g = \frac{1}{2} \int_{\Omega} \tilde{h} \Delta \phi^2 v_g.$$

Combining (2), (3), (4) and (5) we obtain

$$\Lambda' = - \int_{\Omega} \langle d\phi \otimes d\phi + \frac{1}{4} \Delta \phi^2 g, h \rangle v_g,$$

which completes the proof of the proposition. □

In the particular case of domain deformations, Proposition 2.1 gives rise to the following variation formulae.

**COROLLARY 2.1.** *Let  $\Omega_\varepsilon = f_\varepsilon(\Omega)$  be a deformation of  $\Omega$ . Let  $\phi_\varepsilon \in C^\infty(\Omega_\varepsilon)$  and  $\Lambda_\varepsilon \in \mathbf{R}$  be two differentiable curves such that, for all  $\varepsilon$ ,  $\|\phi_\varepsilon\|_{L^2(\Omega_\varepsilon, g)} = 1$  and*

$$\begin{cases} \Delta \phi_\varepsilon = \Lambda_\varepsilon \phi_\varepsilon & \text{in } \Omega_\varepsilon, \\ \phi_\varepsilon = 0 & \text{on } \partial\Omega_\varepsilon. \end{cases}$$

Then,

$$\frac{d}{d\varepsilon} \Lambda_\varepsilon \Big|_{\varepsilon=0} = - \int_{\partial\Omega} v \left( \frac{\partial\phi}{\partial\nu} \right)^2 v_{\bar{g}},$$

where  $\phi = \phi_0$  and  $v = g \left( \frac{d}{d\varepsilon} f_\varepsilon \Big|_{\varepsilon=0}, \nu \right)$  is the normal component of the variation vector field of the deformation  $\Omega_\varepsilon$ .

*Proof.* Let us apply Proposition 2.1 with  $g_\varepsilon = f_\varepsilon^* g$  and  $\bar{\phi}_\varepsilon = \phi_\varepsilon \circ f_\varepsilon$ . Indeed, one can easily check that  $\|\bar{\phi}_\varepsilon\|_{L^2(\Omega, g_\varepsilon)} = 1$ ,  $\Delta_{g_\varepsilon} \bar{\phi}_\varepsilon = \Lambda_\varepsilon \bar{\phi}_\varepsilon$  in  $\Omega$  and  $\bar{\phi}_\varepsilon = 0$  on  $\partial\Omega$ . Hence,

$$(6) \quad \frac{d}{d\varepsilon} \Lambda_\varepsilon \Big|_{\varepsilon=0} = - \int_{\Omega} \langle d\phi \otimes d\phi + \frac{1}{4} \Delta \phi^2 g, h \rangle v_g$$

with  $\phi := \phi_0 = \bar{\phi}_0$  and  $h = \frac{d}{d\varepsilon} f_\varepsilon^* g \Big|_{\varepsilon=0} = \mathcal{L}_V g$ , where  $\mathcal{L}_V g$  is the Lie derivative of  $g$  w.r.t. the vector field  $V = \frac{d}{d\varepsilon} f_\varepsilon \Big|_{\varepsilon=0}$ .

Expressing  $\mathcal{L}_V g$  in terms of the covariant derivative  $\nabla V$  of  $V$  and integrating by parts, we obtain

$$\begin{aligned} \int_{\Omega} \langle d\phi \otimes d\phi, \mathcal{L}_V g \rangle v_g &= \int_{\Omega} \mathcal{L}_V g(\nabla\phi, \nabla\phi) v_g = 2 \int_{\Omega} \langle \nabla_{\nabla\phi} V, \nabla\phi \rangle v_g \\ &= \int_{\Omega} \operatorname{div}(\langle V, \nabla\phi \rangle \nabla\phi) v_g + 2 \int_{\Omega} \langle V, \nabla\phi \rangle \Delta\phi v_g - 2 \int_{\Omega} D^2\phi(V, \nabla\phi) v_g \\ &= 2 \int_{\partial\Omega} \langle V, \nabla\phi \rangle \frac{\partial\phi}{\partial\nu} v_{\bar{g}} + \lambda \int_{\Omega} \langle V, \nabla\phi^2 \rangle v_g - 2 \int_{\Omega} D^2\phi(V, \nabla\phi) v_g, \end{aligned}$$

with  $\lambda := \Lambda_0$ , and

$$\begin{aligned} \frac{1}{4} \int_{\Omega} \Delta\phi^2 \langle g, \mathcal{L}_V g \rangle v_g &= \frac{1}{2} \int_{\Omega} \Delta\phi^2 \operatorname{div} V v_g \\ &= \lambda \int_{\Omega} \phi^2 \operatorname{div} V v_g - \int_{\Omega} |\nabla\phi|^2 \operatorname{div} V v_g \\ &= \int_{\Omega} (-\lambda \langle V, \nabla\phi^2 \rangle + 2D^2\phi(V, \nabla\phi)) v_g \\ &\quad + \int_{\partial\Omega} (\lambda\phi^2 - |\nabla\phi|^2) \langle V, \nu \rangle v_{\bar{g}}. \end{aligned}$$

Substituting this in (6), we get

$$\frac{d}{d\varepsilon} \Lambda_{\varepsilon} |_{\varepsilon=0} = \int_{\partial\Omega} \left\{ -2\langle V, \nabla\phi \rangle \frac{\partial\phi}{\partial\nu} + \langle V, \nu \rangle |\nabla\phi|^2 - \lambda \langle V, \nu \rangle \phi^2 \right\} v_{\bar{g}}.$$

Since  $\phi$  is identically zero on the boundary, we have at any point of  $\partial\Omega$ ,  $\nabla\phi = \frac{\partial\phi}{\partial\nu} \nu$ . In particular,  $|\nabla\phi|^2 = \left(\frac{\partial\phi}{\partial\nu}\right)^2$  and

$$\langle V, \nabla\phi \rangle = \langle V, \nu \rangle \frac{\partial\phi}{\partial\nu} = v \frac{\partial\phi}{\partial\nu}.$$

Thus,

$$\frac{d}{d\varepsilon} \Lambda_{\varepsilon} |_{\varepsilon=0} = - \int_{\partial\Omega} v \left(\frac{\partial\phi}{\partial\nu}\right)^2 v_{\bar{g}}. \quad \square$$

### 3. Critical domains

Throughout this section, the ambient Riemannian manifold  $(M, g)$  is assumed to be real analytic.

**3.1. Preliminary results and definitions.** Let  $\Omega$  be a regular bounded domain of a Riemannian manifold  $(M, g)$ . An analytic deformation  $(\Omega_{\varepsilon})$  of  $\Omega$  is given by an analytic 1-parameter family of diffeomorphisms  $f_{\varepsilon} : \Omega \rightarrow \Omega_{\varepsilon}$  such that  $f_{\varepsilon}(\partial\Omega) = \partial\Omega_{\varepsilon}$  and  $f_0 = \operatorname{Id}$ . Such a deformation is called volume-preserving if the Riemannian volume of  $\Omega_{\varepsilon}$  w.r.t. the metric  $g$  does not depend on  $\varepsilon$ .

The spectrum of the Dirichlet Laplacian  $\Delta_g$  on  $\Omega_\varepsilon$  will be denoted

$$\text{Sp}_D(\Delta_g, \Omega_\varepsilon) = \{ \lambda_{1,\varepsilon} < \lambda_{2,\varepsilon} \leq \dots \leq \lambda_{k,\varepsilon} \uparrow + \infty \}.$$

The functions  $\varepsilon \mapsto \lambda_{k,\varepsilon}$  are continuous, but not differentiable in general, except for  $\lambda_{1,\varepsilon}$ , which is always differentiable since it is simple. Nevertheless, as we will see below, the general perturbation theory of unbounded self-adjoint operators enables us to show that the function  $\lambda_{k,\varepsilon}$  admits right-sided and left-sided derivatives at  $\varepsilon = 0$ . In the sequel, a family of functions  $\phi_\varepsilon \in C^\infty(\Omega_\varepsilon)$  will be called differentiable (resp. analytic) w.r.t.  $\varepsilon$ , if this property holds for  $\phi_\varepsilon \circ f_\varepsilon \in C^\infty(\Omega)$ .

LEMMA 3.1. *Let  $\lambda \in \text{Sp}_D(\Delta_g, \Omega)$  be an eigenvalue of multiplicity  $p$  of the Dirichlet Laplacian in  $\Omega$ . For any analytic deformation  $\Omega_\varepsilon$  of  $\Omega$  there exist  $p$  families  $(\Lambda_{i,\varepsilon})_{i \leq p}$  of real numbers and  $p$  families  $(\phi_{i,\varepsilon})_{i \leq p} \subset C^\infty(\Omega_\varepsilon)$  of functions, depending analytically on  $\varepsilon$  and satisfying, for all  $\varepsilon \in (-\varepsilon_0, \varepsilon_0)$  and for all  $i \in \{1, \dots, p\}$ :*

- (a)  $\Lambda_{i,0} = \lambda$ .
- (b) The family  $\{\phi_{1,\varepsilon}, \dots, \phi_{p,\varepsilon}\}$  is orthonormal in  $L^2(\Omega_\varepsilon, g)$ .
- (c) We have  $\begin{cases} \Delta \phi_{i,\varepsilon} = \Lambda_{i,\varepsilon} \phi_{i,\varepsilon} \text{ in } \Omega_\varepsilon, \\ \phi_{i,\varepsilon} = 0 \text{ on } \partial\Omega_\varepsilon. \end{cases}$

The proof is based on the perturbation theory of unbounded self-adjoint operators in Hilbert spaces. Results concerning the differentiability of eigenvalues and eigenvectors have been first obtained by Rellich [31] and later by Kato [19] in the analytic case. Many results were also obtained under weaker differentiability conditions (see, for instance, [21], [22] for recent contributions to this subject). However, even a smooth curve  $\varepsilon \mapsto P_\varepsilon$  of self-adjoint operators may lead to noncontinuous eigenvectors w.r.t.  $\varepsilon$  (see Rellich's example [19, Chap. II, Example 5.3]). Since we need to differentiate eigenvectors w.r.t.  $\varepsilon$ , we imposed analyticity assumptions in order to obtain analytic curves of operators.

*Proof of Lemma 3.1.* In order to work in the framework of perturbation theory, we first need to modify our operators so that they all have the same domain. Indeed, for any  $\varepsilon$  we set  $g_\varepsilon = f_\varepsilon^* g$  and denote by  $\Delta_\varepsilon$  the Laplace operator of  $(\Omega, g_\varepsilon)$ . Clearly, we have

$$\text{Sp}_D(\Delta_g, \Omega_\varepsilon) = \text{Sp}_D(\Delta_\varepsilon, \Omega).$$

Notice that since  $f_\varepsilon$  depends analytically on  $\varepsilon$  and  $g$  is real analytic, the curves  $\varepsilon \mapsto g_\varepsilon$  and, hence,  $\varepsilon \mapsto \Delta_\varepsilon$ , are analytic w.r.t.  $\varepsilon$ .

The operator  $\Delta_\varepsilon$  is symmetric w.r.t. the inner product in  $L^2(\Omega, g_\varepsilon)$ , but not necessarily w.r.t. the inner product in  $L^2(\Omega, g)$ . Therefore, we need to introduce a conjugation as follows. Let  $U_\varepsilon : L^2(\Omega, g) \rightarrow L^2(\Omega, g_\varepsilon)$  be the

unitary isomorphism given by

$$U_\varepsilon : v \mapsto \left( \frac{|g|}{|g_\varepsilon|} \right)^{1/4} v,$$

where  $|g| = \det(g_{ij})$  is the determinant of the matrix  $(g_{ij})$  of the components of  $g$  in a local coordinate system. We define the operator  $P_\varepsilon$  to be

$$P_\varepsilon = U_\varepsilon^{-1} \circ \Delta_\varepsilon \circ U_\varepsilon.$$

Therefore, we have  $\text{Sp}_D(P_\varepsilon, \Omega) = \text{Sp}_D(\Delta_\varepsilon, \Omega)$  and, if  $v_\varepsilon \in C^\infty(\Omega)$  is an eigenfunction of  $P_\varepsilon$ , then  $\phi_\varepsilon = U_\varepsilon(v_\varepsilon) \circ f_\varepsilon^{-1} \in C^\infty(\Omega_\varepsilon)$  is an eigenfunction of  $\Delta_g$  with the same eigenvalue. Again, since for all  $\varepsilon$ ,  $(M, g_\varepsilon)$  is real analytic, the curves  $\varepsilon \mapsto U_\varepsilon$  and  $\varepsilon \mapsto P_\varepsilon$  are analytic. The result of the lemma then follows from the Rellich-Kato theory applied to  $\varepsilon \mapsto P_\varepsilon$ .  $\square$

Now, let us fix a positive integer  $k$  and let  $\Lambda_{1,\varepsilon}, \dots, \Lambda_{p,\varepsilon}$  be the family of eigenvalues associated with  $\lambda_k$  by Lemma 3.1. Using the continuity of  $\lambda_{k,\varepsilon}$  and the analyticity of  $\Lambda_{i,\varepsilon}$  w.r.t.  $\varepsilon$ , we can easily see that there exist two integers  $i \leq p$  and  $j \leq p$  such that

$$\lambda_{k,\varepsilon} = \begin{cases} \Lambda_{i,\varepsilon} & \text{if } \varepsilon \leq 0, \\ \Lambda_{j,\varepsilon} & \text{if } \varepsilon \geq 0. \end{cases}$$

Hence,  $\lambda_{k,\varepsilon}$  admits left-sided and right-sided derivatives with

$$\frac{d}{d\varepsilon} \lambda_{k,\varepsilon} \Big|_{\varepsilon=0^+} = \frac{d}{d\varepsilon} \Lambda_{j,\varepsilon} \Big|_{\varepsilon=0}$$

and

$$\frac{d}{d\varepsilon} \lambda_{k,\varepsilon} \Big|_{\varepsilon=0^-} = \frac{d}{d\varepsilon} \Lambda_{i,\varepsilon} \Big|_{\varepsilon=0}.$$

DEFINITION 3.1. The domain  $\Omega$  is said to be “critical” for the  $k$ -th eigenvalue of the Dirichlet problem if, for any analytic volume-preserving deformation  $\Omega_\varepsilon$  of  $\Omega$ , the right-sided and left-sided derivatives of  $\lambda_{k,\varepsilon}$  at  $\varepsilon = 0$  have opposite signs, that is,

$$\frac{d}{d\varepsilon} \lambda_{k,\varepsilon} \Big|_{\varepsilon=0^+} \times \frac{d}{d\varepsilon} \lambda_{k,\varepsilon} \Big|_{\varepsilon=0^-} \leq 0.$$

It is easy to see that

$$\frac{d}{d\varepsilon} \lambda_{k,\varepsilon} \Big|_{\varepsilon=0^+} \leq 0 \leq \frac{d}{d\varepsilon} \lambda_{k,\varepsilon} \Big|_{\varepsilon=0^-} \iff \lambda_{k,\varepsilon} \leq \lambda_{k,0} + o(\varepsilon)$$

and

$$\frac{d}{d\varepsilon} \lambda_{k,\varepsilon} \Big|_{\varepsilon=0^-} \leq 0 \leq \frac{d}{d\varepsilon} \lambda_{k,\varepsilon} \Big|_{\varepsilon=0^+} \iff \lambda_{k,\varepsilon} \geq \lambda_{k,0} + o(\varepsilon).$$



Therefore, the domain  $\Omega$  is critical for the  $k$ -th eigenvalue of the Dirichlet problem if and only if one of the following inequalities holds:

$$\begin{aligned} \lambda_{k,\varepsilon} &\leq \lambda_{k,0} + o(\varepsilon), \\ \lambda_{k,\varepsilon} &\geq \lambda_{k,0} + o(\varepsilon). \end{aligned}$$

REMARK 3.1. Suppose that for an integer  $k$  we have  $\lambda_k < \lambda_{k+1}$ . Then, for sufficiently small  $\varepsilon$ , we will have

$$\lambda_{k,\varepsilon} = \max_{i \leq p} \Lambda_{i,\varepsilon},$$

where  $\Lambda_{1,\varepsilon}, \dots, \Lambda_{p,\varepsilon}$  are the eigenvalues associated to  $\lambda_k$  by Lemma 3.1 (indeed,  $\Lambda_{i,0} = \lambda_k < \lambda_{k+1}$  for any  $1 \leq i \leq p$ ). Hence,

$$\frac{d}{d\varepsilon} \lambda_{k,\varepsilon}|_{\varepsilon=0^-} \leq \frac{d}{d\varepsilon} \lambda_{k,\varepsilon}|_{\varepsilon=0^+}.$$

In particular,  $\Omega$  is critical for the functional  $\Omega \mapsto \lambda_k(\Omega)$  if and only if

$$\frac{d}{d\varepsilon} \lambda_{k,\varepsilon}|_{\varepsilon=0^-} \leq 0 \leq \frac{d}{d\varepsilon} \lambda_{k,\varepsilon}|_{\varepsilon=0^+}$$

(or, equivalently,  $\lambda_{k,\varepsilon} \leq \lambda_{k,0} + o(\varepsilon)$ ).

Similarly, if  $\lambda_{k-1} < \lambda_k$ , then, for sufficiently small  $\varepsilon$ ,

$$\lambda_{k,\varepsilon} = \min_{i \leq p} \Lambda_{i,\varepsilon}$$

and

$$\frac{d}{d\varepsilon} \lambda_{k,\varepsilon}|_{\varepsilon=0^+} \leq \frac{d}{d\varepsilon} \lambda_{k,\varepsilon}|_{\varepsilon=0^-}.$$

LEMMA 3.2. Let  $\lambda \in \text{Sp}_p(\Delta_g, \Omega)$  be an eigenvalue of multiplicity  $p$  of the Dirichlet Laplacian in  $\Omega$  and let us denote by  $E_\lambda$  the corresponding eigenspace. Let  $\Omega_\varepsilon = f_\varepsilon(\Omega)$  be an analytic deformation of  $\Omega$  and let  $(\Lambda_{i,\varepsilon})_{i \leq p}$  and  $(\phi_{i,\varepsilon})_{i \leq p} \subset C^\infty(\Omega_\varepsilon)$  be as in Lemma 3.1. Then  $\Lambda'_1 := \frac{d}{d\varepsilon} \Lambda_{1,\varepsilon}|_{\varepsilon=0}, \dots, \Lambda'_p := \frac{d}{d\varepsilon} \Lambda_{p,\varepsilon}|_{\varepsilon=0}$  are the eigenvalues of the quadratic form  $q_v$  defined on the space  $E_\lambda \subset L^2(\Omega, g)$  by

$$q_v(\phi) = - \int_{\partial\Omega} v \left( \frac{\partial\phi}{\partial\nu} \right)^2 v_{\bar{g}},$$

where  $v = g \left( \frac{d}{d\varepsilon} f_\varepsilon|_{\varepsilon=0}, \nu \right)$ . Moreover, the  $L^2$ -orthonormal basis  $\phi_{1,0}, \dots, \phi_{p,0}$  diagonalizes  $q_v$  on  $E_\lambda$ .

*Proof.* For simplicity, we set

$$g_\varepsilon := f_\varepsilon^* g, \quad \Delta' := \frac{d}{d\varepsilon} \Delta_{g_\varepsilon}|_{\varepsilon=0}, \quad \Lambda_i := \Lambda_{i,0}, \quad \phi_i := \phi_{i,0}, \quad \Lambda'_i := \frac{d}{d\varepsilon} \Lambda_{i,\varepsilon}|_{\varepsilon=0}.$$

From  $\Delta_{g_\varepsilon}(\phi_{i,\varepsilon}) = \Lambda_{i,\varepsilon}(\phi_{i,\varepsilon})$  we deduce

$$\Delta' \phi_i + \Delta \phi'_i = \Lambda'_i \phi_i + \Lambda_i \phi'_i.$$

We multiply by  $\phi_j$  and integrate to get

$$\int_{\Omega} \phi_j \Delta' \phi_i v_g + \int_{\Omega} \phi_j \Delta \phi'_i v_g = \Lambda'_i \int_{\Omega} \phi_i \phi_j v_g + \lambda \int_{\Omega} \phi_i \phi'_j v_g.$$

Integration by parts gives (since  $\phi_j = \phi'_i = 0$  on  $\partial\Omega$ )

$$\int_{\Omega} \phi_j \Delta \phi'_i v_g = \lambda \int_{\Omega} \phi_i \phi'_j v_g.$$

Therefore

$$\int_{\Omega} \phi_j \Delta' \phi_i v_g = \Lambda'_i \int_{\Omega} \phi_i \phi_j v_g.$$

It follows that the  $L^2$ -orthonormal basis  $\phi_1, \dots, \phi_p$  diagonalizes the quadratic form  $\phi \rightarrow \int_{\Omega} \phi \Delta' \phi v_g$  on  $E_{\lambda}$ , the corresponding eigenvalues being  $\Lambda'_1, \dots, \Lambda'_p$ . As we have seen in the proof of Corollary 2.1, this last quadratic form coincides with  $q_{\nu}$  on  $E_{\lambda}$ .  $\square$

Any volume-preserving deformation  $\Omega_{\varepsilon} = f_{\varepsilon}(\Omega)$  induces a function  $v := g(\frac{d}{d\varepsilon} f_{\varepsilon}|_{\varepsilon=0}, \nu)$  on  $\partial\Omega$  satisfying  $\int_{\partial\Omega} v v_{\bar{g}} = 0$  (indeed, this last integral is, up to a constant, equal to  $\frac{d}{d\varepsilon} \text{vol}(\Omega_{\varepsilon})|_{\varepsilon=0}$ ). In the sequel, we will denote by  $\mathcal{A}_0(\partial\Omega)$  the set of regular functions on  $\partial\Omega$  such that  $\int_{\partial\Omega} v v_{\bar{g}} = 0$ . The following elementary lemma will be useful in the proof of our main results.

**LEMMA 3.3.** *Let  $v \in \mathcal{A}_0(\partial\Omega)$ . Then there exists an analytic volume-preserving deformation  $\Omega_{\varepsilon} = f_{\varepsilon}(\Omega)$  so that  $v = g(\frac{d}{d\varepsilon} f_{\varepsilon}|_{\varepsilon=0}, \nu)$ .*

*Proof.* Let  $U \subset M$  be an open neighborhood of  $\bar{\Omega}$  and let  $\tilde{v}$  and  $\tilde{\nu}$  be smooth extensions to  $U$  of  $v$  and  $\nu$ , respectively. For  $\varepsilon$  sufficiently small, the map  $\varphi_{\varepsilon}(x) = \exp_x \varepsilon \tilde{v}(x) \tilde{\nu}(x)$  is a diffeomorphism from  $\Omega$  to  $\varphi_{\varepsilon}(\Omega)$ . Moreover, since  $(M, g)$  is real analytic, the curve  $\varepsilon \rightarrow \varphi_{\varepsilon}$  is analytic w.r.t.  $\varepsilon$ . The deformation  $\varphi_{\varepsilon}(\Omega)$  is not necessarily volume-preserving. However, let  $X$  be any analytic vector field on  $U$  such that  $\int_{\Omega} \text{div } X v_g \neq 0$  and denote by  $(\gamma_t)_t$  the associated 1-parameter local group of diffeomorphisms. The function  $(t, \varepsilon) \mapsto F(t, \varepsilon) = \text{vol}(\gamma_t \circ \varphi_{\varepsilon}(\Omega))$  satisfies  $\frac{\partial}{\partial t} F(0, 0) = \int_{\Omega} \text{div } X v_g \neq 0$ . Applying the implicit function theorem in the analytic setting, we get the existence of a function  $t(\varepsilon)$  depending analytically on  $\varepsilon \in (-\eta, \eta)$ , for some  $\eta > 0$  sufficiently small, such that  $F(t(\varepsilon), \varepsilon) = F(0, 0)$ , for all  $\varepsilon \in (-\eta, \eta)$ . The deformation  $f_{\varepsilon} = \gamma_{t(\varepsilon)} \circ \varphi_{\varepsilon}$  is clearly analytic and volume-preserving. Moreover, one has

$$t'(0) = -\frac{\frac{d}{d\varepsilon} \text{vol}(\varphi_{\varepsilon}(\Omega))|_{\varepsilon=0}}{\frac{d}{dt} \text{vol}(\gamma_t(\Omega))|_{t=0}} = -\frac{\int_{\Omega} \text{div } \tilde{v} \tilde{\nu} v_g}{\int_{\Omega} \text{div } X v_g} = -\frac{\int_{\partial\Omega} v v_{\bar{g}}}{\int_{\partial\Omega} \langle X, \nu \rangle v_{\bar{g}}} = 0.$$

Therefore, for all  $x \in \partial\Omega$ ,

$$\frac{d}{d\varepsilon} f_{\varepsilon}(x)|_{\varepsilon=0} = t'(0)X(x) + \frac{d\varphi_{\varepsilon}(x)}{d\varepsilon}|_{\varepsilon=0} = v(x)\nu(x). \quad \square$$

**3.2. Critical domains for the  $k$ -th eigenvalue of the Dirichlet**

**Laplacian.** In the sequel, we will denote by  $\lambda_k$  the  $k$ -th eigenvalue of the Dirichlet problem in  $\Omega$  and by  $E_k$  the corresponding eigenspace.

In the following results, a special role is played by the eigenvalues  $\lambda_k$  satisfying  $\lambda_k > \lambda_{k-1}$  or  $\lambda_k < \lambda_{k+1}$ . This means that the index  $k$  is the lowest or the highest one among all indices corresponding to the same eigenvalue. Let us start with the following necessary condition that must be satisfied by any locally minimizing or locally maximizing domain. Here, a local minimizer (resp. maximizer) for the  $k$ -th eigenvalue of the Dirichlet Laplacian is a domain  $\Omega$  such that, for any volume-preserving deformation  $\Omega_\varepsilon$ , the function  $\varepsilon \mapsto \lambda_{k,\varepsilon}$  admits a local minimum (resp. maximum) at  $\varepsilon = 0$ .

**THEOREM 3.1.** *Let  $k$  be a natural integer such that  $\lambda_k > \lambda_{k-1}$  (resp.  $\lambda_k < \lambda_{k+1}$ ) and assume that  $\Omega$  is a local minimizer (resp. local maximizer) for the  $k$ -th eigenvalue of the Dirichlet Laplacian. Then  $\lambda_k$  is simple and the absolute value of the normal derivative of its corresponding eigenfunction is constant on  $\partial\Omega$ . That is, there exists a unique (up to sign) function  $\phi$  satisfying*

$$\begin{cases} \Delta\phi = \lambda_k\phi \text{ in } \Omega, \\ \phi = 0 \text{ on } \partial\Omega, \\ \left| \frac{\partial\phi}{\partial\nu} \right| = 1 \text{ on } \partial\Omega. \end{cases}$$

*Proof.* Suppose that  $\lambda_k > \lambda_{k-1}$  and let  $\Omega_\varepsilon = f_\varepsilon(\Omega)$  be a volume preserving analytic deformation of  $\Omega$ . Let  $(\Lambda_{i,\varepsilon})_{i \leq p}$  and  $(\phi_{i,\varepsilon})_{i \leq p}$  be families of eigenvalues and eigenfunctions associated to  $\lambda_k$  according to Lemma 3.1. Since  $\Lambda_{i,0} = \lambda_k > \lambda_{k-1}$ , we have, for sufficiently small  $\varepsilon$ , for continuity reasons,

$$\Lambda_{i,\varepsilon} > \lambda_{k-1,\varepsilon}.$$

Hence,

$$\Lambda_{i,\varepsilon} \geq \lambda_{k,\varepsilon}.$$

As the function  $\varepsilon \mapsto \lambda_{k,\varepsilon}$  admits a local minimum at  $\varepsilon = 0$  with  $\Lambda_{i,0} = \lambda_{k,0} = \lambda_k$ , it follows that the differentiable function  $\varepsilon \mapsto \Lambda_{i,\varepsilon}$  achieves a local minimum at  $\varepsilon = 0$  and that  $\frac{d}{d\varepsilon} \Lambda_{i,\varepsilon} \Big|_{\varepsilon=0} = 0$ . Applying Lemma 3.2, we deduce that the quadratic form  $q_v$  is identically zero on the eigenspace  $E_k$ , where  $v = g(\frac{d}{d\varepsilon} f_\varepsilon \Big|_{\varepsilon=0}, \nu)$ . The volume-preserving deformation being arbitrary, it follows that the form  $q_v$  vanishes on  $E_k$  for any  $v \in \mathcal{A}_0(\partial\Omega)$  (Lemma 3.3).

Therefore, for all  $\phi \in E_k$  and for all  $v \in \mathcal{A}_0(\partial\Omega)$ , we have  $\int_{\partial\Omega} v \left( \frac{\partial\phi}{\partial\nu} \right)^2 v_{\bar{g}} = 0$ , which implies that  $\frac{\partial\phi}{\partial\nu}$  is locally constant on  $\partial\Omega$  for any  $\phi \in E_k$ . Now, if  $\phi_1$  and  $\phi_2$  are two eigenfunctions in  $E_k$ , one can find a linear combination  $\phi = \alpha\phi_1 + \beta\phi_2$  so that  $\frac{\partial\phi}{\partial\nu}$  vanishes on at least one connected component of  $\partial\Omega$ . We apply the Holmgren uniqueness theorem (see, for instance, [27,

Theorem 2, p. 42], and recall that  $(M, g)$  is assumed to be real analytic) to deduce that  $\phi$  is identically zero in  $\Omega$  and that  $\lambda_k$  is simple.

To finish the proof, we must show that, for all  $\phi \in E_k$ ,  $|\frac{\partial\phi}{\partial\nu}|$  takes the same constant value on all the components of  $\partial\Omega$ . Indeed, let  $\Sigma_1$  and  $\Sigma_2$  be two distinct connected components of  $\partial\Omega$  and let  $v \in \mathcal{A}_0(\partial\Omega)$  be the function given by  $v = \text{vol}(\Sigma_2)$  on  $\Sigma_1$ ,  $v = -\text{vol}(\Sigma_1)$  on  $\Sigma_2$  and  $v = 0$  on the other components. Then the condition  $\int_{\partial\Omega} v \left(\frac{\partial\phi}{\partial\nu}\right)^2 v_{\bar{g}} = 0$  implies that  $\left(\frac{\partial\phi}{\partial\nu}\right)^2 \Big|_{\Sigma_1} = \left(\frac{\partial\phi}{\partial\nu}\right)^2 \Big|_{\Sigma_2}$ .

Of course, the same arguments work in the case  $\lambda_k < \lambda_{k+1}$ . □

The criticality of the domain  $\Omega$  for the  $k$ -th eigenvalue of the Dirichlet Laplacian is closely related to the definiteness of the quadratic forms  $q_v$  introduced in Lemma 3.2 above, on the eigenspace  $E_k$ . Indeed, we have the following theorem:

**THEOREM 3.2.** *Let  $k$  be any natural integer.*

- (1) *If  $\Omega$  is a critical domain for the  $k$ -th eigenvalue of the Dirichlet Laplacian, then, for all  $v \in \mathcal{A}_0(\partial\Omega)$ , the quadratic form  $q_v(\phi) = -\int_{\partial\Omega} v \left(\frac{\partial\phi}{\partial\nu}\right)^2 v_{\bar{g}}$  is not definite on  $E_k$ .*
- (2) *Assume that  $\lambda_k > \lambda_{k-1}$  or  $\lambda_k < \lambda_{k+1}$ , and that, for all  $v \in \mathcal{A}_0(\partial\Omega)$ , the quadratic form  $q_v(\phi) = -\int_{\partial\Omega} v \left(\frac{\partial\phi}{\partial\nu}\right)^2 v_{\bar{g}}$  is not definite on  $E_k$ . Then  $\Omega$  is a critical domain for the  $k$ -th eigenvalue of the Dirichlet Laplacian.*

*Proof.* (1) Consider a function  $v \in \mathcal{A}_0(\partial\Omega)$  and let  $\Omega_\varepsilon = f_\varepsilon(\Omega)$  be an analytic volume-preserving deformation of  $\Omega$  so that  $v := g(\frac{d}{d\varepsilon} f_\varepsilon|_{\varepsilon=0}, \nu)$  (Lemma 3.3). Let  $(\Lambda_{i,\varepsilon})_{i \leq p}$  and  $(\phi_{i,\varepsilon})_{i \leq p}$  be families of eigenvalues and eigenfunctions associated to  $\lambda_k$  according to Lemma 3.1. As we have seen above, there exist two integers  $i \leq p$  and  $j \leq p$  so that  $\frac{d}{d\varepsilon} \lambda_{k,\varepsilon}|_{\varepsilon=0^-} = \frac{d}{d\varepsilon} \Lambda_{i,\varepsilon}|_{\varepsilon=0}$  and  $\frac{d}{d\varepsilon} \lambda_{k,\varepsilon}|_{\varepsilon=0^+} = \frac{d}{d\varepsilon} \Lambda_{j,\varepsilon}|_{\varepsilon=0}$ . The criticality of  $\Omega$  then implies that  $\frac{d}{d\varepsilon} \Lambda_{i,\varepsilon}|_{\varepsilon=0} \times \frac{d}{d\varepsilon} \Lambda_{j,\varepsilon}|_{\varepsilon=0} \leq 0$ . Applying Lemma 3.2, we deduce that the quadratic form  $q_v$  admits both nonnegative and nonpositive eigenvalues on  $E_k$ , which proves assertion (1).

(2) Assume that  $\lambda_k > \lambda_{k-1}$  and let  $\Omega_\varepsilon = f_\varepsilon(\Omega)$  be a volume-preserving deformation of  $\Omega$ . Let  $(\Lambda_{i,\varepsilon})_{i \leq p}$  and  $(\phi_{i,\varepsilon})_{i \leq p}$  be families of eigenvalues and eigenfunctions associated to  $\lambda_k$  according to Lemma 3.1. As we have seen in Remark 3.1, we have, for sufficiently small  $\varepsilon$ ,  $\lambda_{k,\varepsilon} = \min_{i \leq p} \Lambda_{i,\varepsilon}$ . Hence,

$$\frac{d}{d\varepsilon} \lambda_{k,\varepsilon} \Big|_{\varepsilon=0^+} = \min_{i \leq p} \frac{d}{d\varepsilon} \Lambda_{i,\varepsilon} \Big|_{\varepsilon=0}$$

and

$$\frac{d}{d\varepsilon} \lambda_{k,\varepsilon} \Big|_{\varepsilon=0^-} = \max_{i \leq p} \frac{d}{d\varepsilon} \Lambda_{i,\varepsilon} \Big|_{\varepsilon=0}.$$

Now, the nondefiniteness of  $q_v$  on  $E_k$  means that its smallest eigenvalue is nonpositive and its largest one is nonnegative. According to Lemma 3.2, this implies that

$$\frac{d}{d\varepsilon} \lambda_{k,\varepsilon} \Big|_{\varepsilon=0^+} = \min_{i \leq p} \frac{d}{d\varepsilon} \Lambda_{i,\varepsilon} \Big|_{\varepsilon=0} \leq 0$$

and

$$\frac{d}{d\varepsilon} \lambda_{k,\varepsilon} \Big|_{\varepsilon=0^-} = \max_{i \leq p} \frac{d}{d\varepsilon} \Lambda_{i,\varepsilon} \Big|_{\varepsilon=0} \geq 0,$$

which implies the criticality of the domain  $\Omega$ .

The case  $\lambda_k < \lambda_{k+1}$  can be handled similarly. □

The indefiniteness of  $q_v$  for any  $v \in \mathcal{A}_0(\partial\Omega)$  can be interpreted intrinsically in the following manner:

LEMMA 3.4. *Let  $k$  be a natural integer. The following two conditions are equivalent:*

- (i) *For all  $v \in \mathcal{A}_0(\partial\Omega)$ , the quadratic form  $q_v$  is not definite on  $E_k$ .*
- (ii) *There exists a finite family of eigenfunctions  $(\phi_i)_{i \leq m} \subset E_k$  satisfying*

$$\sum_{i=1}^m \left( \frac{\partial \phi_i}{\partial \nu} \right)^2 = 1 \text{ on } \partial\Omega.$$

*Proof.* To see that (ii) implies (i), it suffices to notice that, for any  $v \in \mathcal{A}_0(\partial\Omega)$

$$\sum_{i \leq m} q_v(\phi_i) = - \sum_{i \leq m} \int_{\partial\Omega} v \left( \frac{\partial \phi_i}{\partial \nu} \right)^2 v_{\bar{g}} = - \int_{\partial\Omega} v v_{\bar{g}} = 0.$$

Therefore,  $q_v$  is not definite on  $E_k$ .

The proof of “(i) implies (ii)” uses arguments similar to those used in the case of closed manifolds by Nadirashvili [25] and the authors [12]. Let  $K$  be the convex hull of  $\{(\frac{\partial \phi}{\partial \nu})^2, \phi \in E_k\}$  in  $C^\infty(\partial\Omega)$ . Then we need to show that the constant function 1 belongs to  $K$ .

Let us suppose, to the contrary, that  $1 \notin K$ . Then, from the Hahn-Banach theorem (applied to the finite dimensional vector space spanned by  $K$  and 1 and endowed with the  $L^2(\partial\Omega, \bar{g})$  inner product), there exists a function  $v \in C^\infty(\partial\Omega)$  such that  $\int_{\partial\Omega} v v_{\bar{g}} > 0$  and, for all  $\phi \in E_k$ ,

$$\int_{\partial\Omega} v \left( \frac{\partial \phi}{\partial \nu} \right)^2 v_{\bar{g}} \leq 0.$$

Hence, the zero mean value function

$$v_o = v - \frac{1}{\text{vol}(\partial\Omega)} \int_{\partial\Omega} v v_{\bar{g}}$$

satisfies, for all  $\phi \in E_k$ ,

$$\begin{aligned} q_{v_o}(\phi) &= - \int_{\partial\Omega} v_o \left( \frac{\partial\phi}{\partial\nu} \right)^2 v_{\bar{g}} \\ &= - \int_{\partial\Omega} v \left( \frac{\partial\phi}{\partial\nu} \right)^2 v_{\bar{g}} + \frac{1}{\text{vol}(\partial\Omega)} \int_{\partial\Omega} v v_{\bar{g}} \int_{\partial\Omega} \left( \frac{\partial\phi}{\partial\nu} \right)^2 v_{\bar{g}} \\ &\geq \frac{1}{\text{vol}(\partial\Omega)} \int_{\partial\Omega} v v_{\bar{g}} \int_{\partial\Omega} \left( \frac{\partial\phi}{\partial\nu} \right)^2 v_{\bar{g}}, \end{aligned}$$

with  $\int_{\partial\Omega} \left( \frac{\partial\phi}{\partial\nu} \right)^2 v_{\bar{g}} > 0$  for any nontrivial Dirichlet eigenfunction  $\phi$  (due to the Holmgren uniqueness theorem). In conclusion, the function  $v_o \in \mathcal{A}_0(\partial\Omega)$  is such that the quadratic form  $q_{v_o}$  is positive definite on  $E_k$ , which contradicts condition (i).  $\square$

A consequence of this lemma and Theorem 3.2 is the following result:

**THEOREM 3.3.** *Let  $k$  be any natural integer.*

- (1) *If  $\Omega$  is a critical domain for the  $k$ -th eigenvalue of the Dirichlet Laplacian, then there exists a finite family of eigenfunctions  $(\phi_i)_{i \leq m} \subset E_k$  satisfying  $\sum_{i=1}^m \left( \frac{\partial\phi_i}{\partial\nu} \right)^2 = 1$  on  $\partial\Omega$ , that is, the functions  $(\phi_i)_{i \leq m}$  are solutions of the following system:*

$$\begin{cases} \Delta\phi_i = \lambda_k\phi_i \text{ in } \Omega, & \text{for all } i \leq m, \\ \phi_i = 0 \text{ on } \partial\Omega, & \text{for all } i \leq m, \\ \sum_{i=1}^m \left( \frac{\partial\phi_i}{\partial\nu} \right)^2 = 1 & \text{on } \partial\Omega. \end{cases}$$

- (2) *Assume that  $\lambda_k > \lambda_{k-1}$  or  $\lambda_k < \lambda_{k+1}$  and that there exists a finite family of eigenfunctions  $(\phi_i)_{i \leq m} \subset E_k$  such that  $\sum_{i=1}^m \left( \frac{\partial\phi_i}{\partial\nu} \right)^2$  is constant on  $\partial\Omega$ . Then the domain  $\Omega$  is critical for the  $k$ -th eigenvalue of the Dirichlet Laplacian.*

**COROLLARY 3.1.** *Assume that  $\lambda_k$  is simple. The domain  $\Omega$  is critical for the  $k$ -th eigenvalue of the Dirichlet Laplacian if and only if the following overdetermined Pompeiu-Schiffer type system admits a solution:*

$$\begin{cases} \Delta\phi = \lambda_k\phi \text{ in } \Omega, \\ \phi = 0 \text{ on } \partial\Omega, \\ \left| \frac{\partial\phi}{\partial\nu} \right| = 1 \text{ on } \partial\Omega. \end{cases}$$

**3.3. Nonexistence of critical domains under metric variations.** In this subsection, we point out the inconsistency of the notion of critical domains w.r.t. metric variations under the Dirichlet boundary condition. Indeed, if  $g_\varepsilon$  is an analytic variation of the metric  $g$ , then we can associate to each eigenvalue  $\lambda_k$  of the Dirichlet problem in  $\Omega$  analytic families  $(\Lambda_{i,\varepsilon})_{i \leq p} \subset \mathbb{R}$  and  $(\phi_{i,\varepsilon})_{i \leq p} \subset C^\infty(\Omega)$  (where  $p$  is the multiplicity of  $\lambda_k$ ) satisfying, for sufficiently small  $\varepsilon$ :

- (1)  $(\phi_{i,\varepsilon})_{i \leq p}$  is  $L^2(\Omega, g_\varepsilon)$  orthonormal.
- (2) For all  $i \in \{1, \dots, p\}$ ,  $\Lambda_{i,0} = \lambda_k$ .
- (3) For all  $i \leq p$ ,  $\begin{cases} \Delta_{g_\varepsilon} \phi_{i,\varepsilon} = \Lambda_{i,\varepsilon} \phi_{i,\varepsilon} \text{ in } \Omega, \\ \phi_{i,\varepsilon} = 0 \text{ on } \partial\Omega. \end{cases}$

Therefore  $\lambda_{k,\varepsilon}$  admits left-sided and right-sided derivatives at  $\varepsilon = 0$ , and we can mimic Definition 3.1 to introduce the notion of critical domain for the  $k$ -th eigenvalue of the Dirichlet problem w.r.t. volume-preserving variations of the metric. Using Proposition 2.1 and arguments similar to those used above (see also [12], [25]), we can show that, if the domain  $(\Omega, g)$  is critical for the  $k$ -th eigenvalue of the Dirichlet problem, then there exists a family of eigenfunctions  $\phi_1, \dots, \phi_m \in E_k$  satisfying

$$(7) \quad \sum_{i=1}^m d\phi_i \otimes d\phi_i = g.$$

Now, if we consider only volume-preserving conformal variations  $g_\varepsilon$  of  $g$  (that is  $g_\varepsilon = \alpha_\varepsilon g$  with  $\int_\Omega \alpha_\varepsilon^{n/2} v_g = \text{vol}(\Omega, g)$ ), then the necessary condition (7) for  $(\Omega, g)$  to be critical w.r.t. such variations becomes  $\sum_{i=1}^m \phi_i^2 = 1$  in  $\Omega$ . As the eigenfunctions of the Dirichlet Laplacian vanish on the boundary  $\partial\Omega$ , this last condition can never be fulfilled by functions of  $E_k$ . Thus, we have the following result:

**PROPOSITION 3.1.** *There is no critical domain  $(\Omega, g)$  for the  $k$ -th eigenvalue of the Dirichlet Laplacian under conformal volume-preserving variations of the metric  $g$ .*

### 4. Applications to the trace of the heat kernel

This section deals with critical domains of the trace of the heat kernel under the Dirichlet boundary condition.

Recall that the Dirichlet heat kernel  $H$  of  $(\Omega, g)$  is defined to be the solution of the following parabolic problem:

$$\begin{cases} (\frac{\partial}{\partial t} - \Delta_y)H(t, x, y) = 0, \\ H(0, x, y) = \delta_x, \\ \text{for all } y \in \partial\Omega, H(t, x, y) = 0, \end{cases}$$

Its trace is the function

$$Y(t) = \int_{\Omega} H(t, x, x) v_g.$$

The relationship between this kernel and the spectrum of the Dirichlet Laplacian is given by

$$H(t, x, y) = \sum_{k \geq 1} e^{-\lambda_k t} \phi_k(x) \phi_k(y),$$

where  $(\phi)_{k \geq 1}$  is an  $L^2(\Omega, g)$ -orthonormal family of eigenfunctions satisfying

$$\begin{cases} \Delta \phi_k = \lambda_k \phi_k & \text{in } \Omega, \\ \phi_k = 0 & \text{on } \partial\Omega. \end{cases}$$

Therefore

$$(8) \quad Y(t) = \sum_{k \geq 1} e^{-\lambda_k t}.$$

Let  $\Omega_\varepsilon$  be a smooth deformation of  $\Omega$  and let  $Y_\varepsilon(t) = \sum_{k \geq 1} e^{-\lambda_{k,\varepsilon} t}$  be the corresponding heat trace function. Unlike the eigenvalues, the function  $Y_\varepsilon(t)$  is always differentiable in  $\varepsilon$  and *the domain  $\Omega$  will be called critical for the trace of the Dirichlet heat kernel at time  $t$  if, for any volume-preserving deformation  $\Omega_\varepsilon$  of  $\Omega$ , we have*

$$\frac{d}{d\varepsilon} Y_\varepsilon(t) \Big|_{\varepsilon=0} = 0.$$

From the results of Section 3 above, one can deduce the variation formula for the heat trace. For this, we need to introduce the mixed second derivative  $d_S H(t)|_x$  of  $H$  at the point  $x$ , defined as the smooth 2-tensor given by

$$d_S H(t)|_x(X, X) = \frac{\partial^2}{\partial \alpha \partial \beta} H(t, c(\alpha), c(\beta)) \Big|_{\alpha=\beta=0},$$

where  $c$  is a curve in  $\Omega$  such that  $c(0) = x$  and  $\dot{c}(0) = X$ . It is easy to check that

$$d_S H(t) = \sum_{k \geq 1} e^{-\lambda_k t} d\phi_k \otimes d\phi_k.$$

**THEOREM 4.1.** *Let  $\Omega_\varepsilon = f_\varepsilon(\Omega)$  be a volume-preserving deformation of  $\Omega$ . We have, for all  $t > 0$ ,*

$$\frac{d}{d\varepsilon} Y_\varepsilon(t) \Big|_{\varepsilon=0} = -t \int_{\partial\Omega} v d_S H(t)(\nu, \nu) v_{\bar{g}} = \frac{t}{2} \int_{\partial\Omega} v \Delta H(t, x, x) v_{\bar{g}},$$

where  $v = g(\frac{d}{d\varepsilon} f_\varepsilon \Big|_{\varepsilon=0}, \nu)$ .



*Proof.* The formula of Theorem 4.1 can be derived from the first variation formula of the heat kernel given in the paper of Ray and Singer [28, Proposition 6.1]. However, at least in the case where the ambient manifold is real analytic, it can also be obtained as an immediate consequence of Hadamard’s type formula of Section 2, thanks to the relation (8) above. Indeed, in this manner we obtain, for all  $t > 0$ ,

$$\frac{d}{d\varepsilon} Y_\varepsilon(t) \Big|_{\varepsilon=0} = -t \sum_{k \geq 1} e^{-\lambda_k t} \int_{\partial\Omega} v \left( \frac{\partial\phi_k}{\partial\nu} \right)^2 v_{\bar{g}},$$

where  $(\lambda_k, \phi_k)$  are as above. To get the desired formula for  $Y_\varepsilon(t)$  it suffices to notice that

$$d_S H(t)(\nu, \nu) = \sum_{k \geq 1} e^{-\lambda_k t} d\phi_k \otimes d\phi_k(\nu, \nu) = \sum_{k \geq 1} e^{-\lambda_k t} \left( \frac{\partial\phi_k}{\partial\nu} \right)^2. \quad \square$$

An immediate consequence is the following result:

**COROLLARY 4.1.** *The following conditions are equivalent:*

- (i) *The domain  $\Omega$  is critical for the trace of the Dirichlet heat kernel at the time  $t$  under volume-preserving domain deformations.*
- (ii)  *$\Delta H(t, x, x)$  is constant on the boundary  $\partial\Omega$ .*
- (iii) *For any positive integer  $k$  and any  $L^2(\Omega, g)$ -orthonormal basis  $\phi_1, \dots, \phi_p$  of the eigenspace  $E_k$  of  $\lambda_k$ ,  $\sum_{i \leq p} \left( \frac{\partial\phi_i}{\partial\nu} \right)^2$  is constant on  $\partial\Omega$ .*

Recall that if  $\rho$  is an isometry of  $(\Omega, g)$ , then, for all  $x \in \Omega$  and for all  $t > 0$ ,  $H(t, \rho(x), \rho(x)) = H(t, x, x)$ . In particular, if  $\Omega$  is a ball of  $\mathbb{R}^n$  endowed with a rotationally symmetric Riemannian metric  $g$  given in polar coordinates by  $g = a^2(r)dr^2 + b^2(r)d\sigma^2$ , where  $d\sigma^2$  is the standard metric of the unit sphere  $\mathbb{S}^{n-1}$ , then  $H(t, x, x)$  is radial (that is, depends only on the parameter  $r$ ). Therefore, the function  $\Delta H(t, x, x)$  is also radial and hence constant on the boundary of the ball.

**COROLLARY 4.2.** *Let  $g$  be a rotationally symmetric Riemannian metric on  $\mathbb{R}^n$ . The geodesic balls centered at the origin are critical domains for the trace of the Dirichlet heat kernel under volume-preserving domain deformations.*

In particular, geodesic balls of Riemannian space forms are critical for the trace of the Dirichlet heat kernel under volume-preserving domain deformations.

The Minakshisundaram-Pleijel asymptotic expansion of the trace of the heat kernel also provides information about the geometric properties of extremal or critical domains. Indeed, it is well known that there exists a sequence

$(a_i)_{i \in \mathbb{N}}$  of real numbers such that for sufficiently small  $t > 0$ , we have

$$Y(t) = (4\pi t)^{-n/2} \sum_{k \geq 0} a_k t^{k/2}$$

with (see, for instance, [6], [7])

$$\begin{aligned} a_0 &= \text{vol}(\Omega, g), \\ a_1 &= -\frac{\sqrt{\pi}}{2} \text{vol}(\partial\Omega, \bar{g}), \\ a_2 &= \frac{1}{6} \left\{ \int_{\Omega} \text{scal}_g v_g + 2 \int_{\partial\Omega} \text{tr } A v_{\bar{g}} \right\}, \\ a_3 &= \frac{\sqrt{\pi}}{192} \left\{ \int_{\partial\Omega} (-16 \text{scal}_g - 7(\text{tr } A)^2 + 10|A|^2 + 8\rho_g(\nu, \nu)) v_{\bar{g}} \right\}, \end{aligned}$$

where  $\text{scal}_g$  and  $\rho_g$  are, respectively, the scalar and Ricci curvatures of  $(\Omega, g)$ ,  $A$  is the shape operator of the boundary  $\partial\Omega$  (i.e., for all  $X \in T\partial\Omega$ ,  $A(X) = D_X\nu$ ) and  $\text{tr } A$  is the trace of  $A$  (i.e.,  $(n-1)$ -times the mean curvature of  $\partial\Omega$ ).

An immediate consequence of these formulae is the following: Suppose that for any domain  $\Omega'$  having the same volume as  $\Omega$  we have  $Y_{\Omega'}(t) \leq Y_{\Omega}(t)$ , for all  $t > 0$ . Then  $\text{vol } \partial\Omega' \geq \text{vol } \partial\Omega$ . Consequently, we have the following result:

**PROPOSITION 4.1.** *If the domain  $\Omega$  maximizes  $Y$  at every time  $t > 0$  among all domains of the same volume, then  $\Omega$  is a solution of the isoperimetric problem in  $(M, g)$ , that is, for all  $\Omega' \subset M$  such that  $\text{vol } \Omega = \text{vol } \Omega'$  we have  $\text{vol } \partial\Omega' \geq \text{vol } \partial\Omega$ .*

Another consequence of the Minakshisundaram-Pleijel asymptotic expansion is the following result:

**THEOREM 4.2.** *If the domain  $\Omega$  is a critical domain of the trace of the Dirichlet heat kernel at every time  $t > 0$ , then  $\partial\Omega$  has constant mean curvature. If in addition the Ricci curvature (resp. the sectional curvature) of the ambient space  $(M, g)$  is constant in a neighborhood of  $\Omega$ , then  $\text{tr}(A^2)$  (resp.  $\text{tr}(A^3)$ ) is constant on  $\partial\Omega$ .*

*Proof.* Let  $\Omega_{\varepsilon} = f_{\varepsilon}(\Omega)$  be a volume-preserving variation of  $\Omega$  and let us denote for any  $\varepsilon$  by  $(a_{i,\varepsilon})_{i \geq 0}$  the coefficients of the asymptotic expansions of  $Y_{\varepsilon}(t)$ . Since  $\frac{d}{d\varepsilon} Y_{\varepsilon}(t)|_{\varepsilon=0} = 0$ , we have for any  $i \geq 0$ ,  $\frac{d}{d\varepsilon} a_{i,\varepsilon}|_{\varepsilon=0} = 0$  (see, for instance, [16] for an analytic justification for this last assertion). In particular,  $\frac{d}{d\varepsilon} \text{vol}(\partial\Omega_{\varepsilon})|_{\varepsilon=0} = 0$  for any volume-preserving variation of  $\Omega$ . This property is known to be equivalent to the fact that the mean curvature of  $\partial\Omega$  is constant (see, for instance, [29]).

Now, let us suppose that the Ricci curvature of  $(M, g)$  is constant in a neighborhood of  $\Omega$ . Then for any small  $\varepsilon$  we have

$$\begin{aligned} a_{2,\varepsilon} &= \frac{1}{6} \left\{ \text{scal}_g \text{vol}(\Omega_\varepsilon) + 2 \int_{\partial\Omega_\varepsilon} (\text{tr } A_\varepsilon) v_{\bar{g}} \right\} \\ &= \frac{1}{6} \left\{ \text{scal}_g \text{vol}(\Omega) + 2 \int_{\partial\Omega_\varepsilon} (\text{tr } A_\varepsilon) v_{\bar{g}} \right\}. \end{aligned}$$

Hence, we have (see, for instance, [29])

$$\begin{aligned} \frac{d}{d\varepsilon} \int_{\partial\Omega_\varepsilon} (\text{tr } A_\varepsilon) v_{\bar{g}} \Big|_{\varepsilon=0} &= \int_{\partial\Omega} (\Delta_{\bar{g}} v - \rho(\nu, \nu)v - (\text{tr } A^2)v) v_{\bar{g}} \\ &\quad + \frac{1}{2} \int_{\partial\Omega} \text{tr } A (\text{div}_{\bar{g}} V^T + v \text{tr } A) v_{\bar{g}}, \end{aligned}$$

where  $V = \frac{df_\varepsilon}{d\varepsilon} \Big|_{\varepsilon=0} = v\nu + V^T$  on the boundary  $\partial\Omega$ . Since  $\int_{\partial\Omega} v v_{\bar{g}} = 0$  and  $\text{tr } A$  and  $\rho(\nu, \nu)$  are constant on  $\partial\Omega$ , we have

$$\frac{d}{d\varepsilon} a_{2,\varepsilon} \Big|_{\varepsilon=0} = \frac{1}{3} \int_{\partial\Omega} (\text{tr } A^2)v v_{\bar{g}} = 0.$$

It follows that  $\text{tr } A^2$  is constant on  $\partial\Omega$ .

As before, we have

$$\frac{d}{d\varepsilon} a_{3,\varepsilon} \Big|_{\varepsilon=0} = \frac{\sqrt{\pi}}{192} \left( -7 \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \int_{\partial\Omega_\varepsilon} (\text{tr } A_\varepsilon)^2 v_{\bar{g}} + 10 \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \int_{\partial\Omega_\varepsilon} \text{tr } A_\varepsilon^2 v_{\bar{g}} \right),$$

but

$$\begin{aligned} \frac{d}{d\varepsilon} \int_{\partial\Omega_\varepsilon} (\text{tr } A_\varepsilon)^2 v_{\bar{g}} \Big|_{\varepsilon=0} &= 2 \int_{\partial\Omega} \text{tr } A (\Delta_{\bar{g}} v - \rho(\nu, \nu)v - (\text{tr } A^2)v) v_{\bar{g}} \\ &\quad + \frac{1}{2} \int_{\partial\Omega} (\text{tr } A)^2 (\text{div}_{\bar{g}} V^T + v \text{tr } A) v_{\bar{g}} \\ &= 0, \end{aligned}$$

since  $\text{tr } A$ ,  $\text{tr } A^2$  and  $\rho(\nu, \nu)$  are constants. Thus,

$$\frac{d}{d\varepsilon} a_{3,\varepsilon} \Big|_{\varepsilon=0} = \frac{10\sqrt{\pi}}{192} \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \int_{\partial\Omega_\varepsilon} \text{tr } A_\varepsilon^2 v_{\bar{g}}.$$

After some straightforward but long computations we obtain, using the fact that the sectional curvature is constant in a neighborhood of  $\Omega$  and that  $\text{tr } A$  and  $\text{tr } A^2$  are constant,

$$\frac{d}{d\varepsilon} a_{3,\varepsilon} \Big|_{\varepsilon=0} = c \int_{\partial\Omega} \text{tr } A^3 v v_{\bar{g}} = 0,$$

where  $c$  is a constant. This proves that  $\text{tr } A^3$  is constant. □

Alexandrov's Theorem [1] shows that in the Euclidean space the geodesic spheres are the only embedded compact hypersurfaces of constant mean curvature. This theorem was extended to hypersurfaces of the hyperbolic space and the standard hemisphere (see [24]). Since the boundary of a critical domain of the trace of the heat kernel is an embedded hypersurface of constant mean curvature, we have the following corollary:

**COROLLARY 4.3.** *Let  $(M, g)$  be one of the following spaces:*

- *The Euclidean space.*
- *The hyperbolic space.*
- *The standard hemisphere.*

*Then a domain  $\Omega$  of  $(M, g)$  is critical for the trace of the Dirichlet heat kernel if and only if  $\Omega$  is a geodesic ball.*

**Acknowledgments.** The authors would like to thank Professors Bernard Helffer and Peter Gilkey for valuable discussions. They also thank the referee for pointing out some mistakes in the first version of the paper and for valuable comments.

#### REFERENCES

- [1] A. D. Aleksandrov, *Uniqueness theorems for surfaces in the large. I*, Vestnik Leningrad. Univ. **11** (1956), 5–17. MR 0086338 (19,167c)
- [2] M. S. Ashbaugh, *Isoperimetric and universal inequalities for eigenvalues*, Spectral theory and geometry (Edinburgh, 1998), London Math. Soc. Lecture Note Ser., vol. 273, Cambridge Univ. Press, Cambridge, 1999, pp. 95–139. MR 1736867 (2001a:35131)
- [3] M. S. Ashbaugh and R. D. Benguria, *A sharp bound for the ratio of the first two eigenvalues of Dirichlet Laplacians and extensions*, Ann. of Math. (2) **135** (1992), 601–628. MR 1166646 (93d:35105)
- [4] M. Berger, *Sur les premières valeurs propres des variétés riemanniennes*, Compositio Math. **26** (1973), 129–149. MR 0316913 (47 #5461)
- [5] P. Bérard, G. Besson, and S. Gallot, *Sur une inégalité isopérimétrique qui généralise celle de Paul Lévy-Gromov*, Invent. Math. **80** (1985), 295–308. MR 788412 (86j:58017)
- [6] N. Blažić, N. Bokan, and P. Gilkey, *The spectral geometry of the Laplacian and the conformal Laplacian for manifolds with boundary*, Global differential geometry and global analysis (Berlin, 1990), Lecture Notes in Math., vol. 1481, Springer, Berlin, 1991, pp. 5–17. MR 1178513 (94a:58200)
- [7] T. P. Branson and P. B. Gilkey, *The asymptotics of the Laplacian on a manifold with boundary*, Comm. Partial Differential Equations **15** (1990), 245–272. MR 1032631 (90m:58201)
- [8] P. Bérard and D. Meyer, *Inégalités isopérimétriques et applications*, Ann. Sci. École Norm. Sup. (4) **15** (1982), 513–541. MR 690651 (84h:58147)
- [9] S. Y. Cheng, *Eigenvalue comparison theorems and its geometric applications*, Math. Z. **143** (1975), 289–297. MR 0378001 (51 #14170)
- [10] J. F. Escobar, *Uniqueness theorems on conformal deformation of metrics, Sobolev inequalities, and an eigenvalue estimate*, Comm. Pure Appl. Math. **43** (1990), 857–883. MR 1072395 (92f:58038)
- [11] A. El Soufi and S. Ilias, *Immersiones minimales, première valeur propre du laplacien et volume conforme*, Math. Ann. **275** (1986), 257–267. MR 854009 (87j:53088)

- [12] ———, *Riemannian manifolds admitting isometric immersions by their first eigenfunctions*, Pacific J. Math. **195** (2000), 91–99. MR 1781616 (2001f:53128)
- [13] ———, *Critical metrics of the trace of the heat kernel on a compact manifold*, J. Math. Pures Appl. (9) **81** (2002), 1053–1070. MR 1946915 (2004b:58035)
- [14] G. Faber, *Beweiss, dass unter allen homogenen Membranen von gleicher Fläche und gleicher Spannung die kreisförmige den tiefsten Grundton gibt*, Sitzungsber. Bayer Akad. Wiss., Math.-Phys. Abt., München, (1923), 169–172.
- [15] P. R. Garabedian and M. Schiffer, *Convexity of domain functionals*, J. Analyse Math. **2** (1953), 281–368. MR 0060117 (15,627a)
- [16] P. B. Gilkey and L. Smith, *The eta invariant for a class of elliptic boundary value problems*, Comm. Pure Appl. Math. **36** (1983), 85–131. MR 680084 (84h:58141)
- [17] J. Hadamard, *Mémoire sur le problème d'analyse relatif à l'équilibre des plaques élastiques encastrées*, Oeuvres, vol. 2, 1968, pp. 515–631.
- [18] A. Henrot, *Minimization problems for eigenvalues of the Laplacian*, J. Evol. Equ. **3** (2003), 443–461. MR 2019029 (2005a:49078)
- [19] T. Kato, *Perturbation theory for linear operators*, second ed., Grundlehren der Mathematischen Wissenschaften, Vol. 132, Springer-Verlag, Berlin, 1976. MR 0407617 (53 #11389)
- [20] E. Krahn, *Über eine von Rayleigh formulierte Minimaleigenschaft des Kreises*, Math. Ann. **94** (1924), 97–100.
- [21] A. Kriegl and P. W. Michor, *The convenient setting of global analysis*, Mathematical Surveys and Monographs, vol. 53, American Mathematical Society, Providence, RI, 1997. MR 1471480 (98i:58015)
- [22] ———, *Differentiable perturbation of unbounded operators*, Math. Ann. **327** (2003), 191–201. MR 2006008 (2005e:47030)
- [23] J. M. Luttinger, *Generalized isoperimetric inequalities*, J. Mathematical Phys. **14** (1973), 586–593. MR 0337197 (49 #1969)
- [24] S. Montiel and A. Ros, *Compact hypersurfaces: the Alexandrov theorem for higher order mean curvatures*, Differential geometry, Pitman Monogr. Surveys Pure Appl. Math., vol. 52, Longman Sci. Tech., Harlow, 1991, pp. 279–296. MR 1173047 (93h:53062)
- [25] N. Nadirashvili, *Berger's isoperimetric problem and minimal immersions of surfaces*, Geom. Funct. Anal. **6** (1996), 877–897. MR 1415764 (98f:53061)
- [26] B. Osgood, R. Phillips, and P. Sarnak, *Extremals of determinants of Laplacians*, J. Funct. Anal. **80** (1988), 148–211. MR 960228 (90d:58159)
- [27] J. Rauch, *Partial differential equations*, Graduate Texts in Mathematics, vol. 128, Springer-Verlag, New York, 1991. MR 1223093 (94e:35002)
- [28] D. B. Ray and I. M. Singer, *R-torsion and the Laplacian on Riemannian manifolds*, Advances in Math. **7** (1971), 145–210. MR 0295381 (45 #4447)
- [29] R. C. Reilly, *Variational properties of functions of the mean curvatures for hypersurfaces in space forms*, J. Differential Geometry **8** (1973), 465–477. MR 0341351 (49 #6102)
- [30] ———, *Applications of the Hessian operator in a Riemannian manifold*, Indiana Univ. Math. J. **26** (1977), 459–472. MR 0474149 (57 #13799)
- [31] F. Rellich, *Perturbation theory of eigenvalue problems*, Gordon and Breach Science Publishers, New York, 1969. MR 0240668 (39 #2014)
- [32] M. Schiffer, *Partial differential equations of elliptic type*, Symposium on partial differential equations, Univ. of California, Berkeley, 1957, pp. 97–149.
- [33] ———, *Hadamard's formula and variation of domain-functions*, Amer. J. Math. **68** (1946), 417–448. MR 0018750 (8,325a)

- [34] J. Serrin, *A symmetry problem in potential theory*, Arch. Rational Mech. Anal. **43** (1971), 304–318. MR 0333220 (48 #11545)
- [35] N. Shimakura, *La première valeur propre du laplacien pour le problème de Dirichlet*, J. Math. Pures Appl. (9) **62** (1983), 129–152. MR 713393 (85e:35091)
- [36] G. Szegö, *Inequalities for certain eigenvalues of a membrane of given area*, J. Rational Mech. Anal. **3** (1954), 343–356. MR 0061749 (15,877c)
- [37] H. F. Weinberger, *An isoperimetric inequality for the  $N$ -dimensional free membrane problem*, J. Rational Mech. Anal. **5** (1956), 633–636. MR 0079286 (18,63c)

AHMAD EL SOUFI, UNIVERSITÉ DE TOURS, LABORATOIRE DE MATHÉMATIQUES ET PHYSIQUE THÉORIQUE, UMR-CNRS 6083, PARC DE GRANDMONT, 37200 TOURS, FRANCE  
*E-mail address:* `elsoufi@univ-tours.fr`

SAÏD ILIAS, UNIVERSITÉ DE TOURS, LABORATOIRE DE MATHÉMATIQUES ET PHYSIQUE THÉORIQUE, UMR-CNRS 6083, PARC DE GRANDMONT, 37200 TOURS, FRANCE  
*E-mail address:* `ilias@univ-tours.fr`