

Dose-Response Association of CD8⁺ Tumor-Infiltrating Lymphocytes and Survival Time in High-Grade Serous Ovarian Cancer

Ovarian Tumor Tissue Analysis (OTTA) Consortium

[+ Supplemental content](#)

IMPORTANCE Cytotoxic CD8⁺ tumor-infiltrating lymphocytes (TILs) participate in immune control of epithelial ovarian cancer; however, little is known about prognostic patterns of CD8⁺ TILs by histotype and in relation to other clinical factors.

OBJECTIVE To define the prognostic role of CD8⁺ TILs in epithelial ovarian cancer.

DESIGN, SETTING, AND PARTICIPANTS This was a multicenter observational, prospective survival cohort study of the Ovarian Tumor Tissue Analysis Consortium. More than 5500 patients, including 3196 with high-grade serous ovarian carcinomas (HGSOCs), were followed prospectively for over 24 650 person-years.

EXPOSURES Following immunohistochemical analysis, CD8⁺ TILs were identified within the epithelial components of tumor islets. Patients were grouped based on the estimated number of CD8⁺ TILs per high-powered field: negative (none), low (1-2), moderate (3-19), and high (≥ 20). CD8⁺ TILs in a subset of patients were also assessed in a quantitative, uncategorized manner, and the functional form of associations with survival was assessed using penalized B-splines.

MAIN OUTCOMES AND MEASURES Overall survival time.

RESULTS The final sample included 5577 women; mean age at diagnosis was 58.4 years (median, 58.2 years). Among the 5 major invasive histotypes, HGSOCs showed the most infiltration. CD8⁺ TILs in HGSOCs were significantly associated with longer overall survival; median survival was 2.8 years for patients with no CD8⁺ TILs and 3.0 years, 3.8 years, and 5.1 years for patients with low, moderate, or high levels of CD8⁺ TILs, respectively (P value for trend = 4.2×10^{-16}). A survival benefit was also observed among women with endometrioid and mucinous carcinomas, but not for those with the other histotypes. Among HGSOCs, CD8⁺ TILs were favorable regardless of extent of residual disease following cytoreduction, known standard treatment, and germline *BRCA1* pathogenic mutation, but were not prognostic for *BRCA2* mutation carriers. Evaluation of uncategorized CD8⁺ TIL counts showed a near-log-linear functional form.

CONCLUSIONS AND RELEVANCE This study demonstrates the histotype-specific nature of immune infiltration and provides definitive evidence for a dose-response relationship between CD8⁺ TILs and HGSOC survival. That the extent of infiltration is prognostic, not merely its presence or absence, suggests that understanding factors that drive infiltration will be the key to unraveling outcome heterogeneity in this cancer.

JAMA Oncol. 2017;3(12):e173290. doi:10.1001/jamaoncol.2017.3290
Published online October 12, 2017.

The Authors/Members of the Ovarian Tumor Tissue Analysis Consortium (OTTA) are listed at the end of this article.

Corresponding Author: Ellen L. Goode, PhD, MPH, Department of Health Sciences Research, Mayo Clinic, 200 First St SW, Rochester, MN 55905 (egoode@mayo.edu).

Epithelial ovarian cancer (OC) is the most lethal gynecologic cancer and is responsible for approximately 14 000 deaths annually in the United States.¹ While initial remission is often achieved, most patients relapse and die from their disease. Immune checkpoint inhibitors have demonstrated clinical activity in a small subset of patients with OC.^{2,3} Understanding the endogenous immune response to OC—including the frequency of CD8⁺ tumor-infiltrating lymphocytes (TILs) and their impact on prognosis—has biological and clinical relevance.

Earlier studies⁴⁻⁸ demonstrated that OC prognosis is associated with TILs at the time of primary cytoreductive surgery. CD8⁺ T cells are stimulated by peptides from degraded proteins bound to human leukocyte antigen class I molecules.⁹ This can trigger CD8⁺ T cells to kill tumor cells and secrete proinflammatory cytokines. While the presence of CD8⁺ TILs within the epithelial component of OCs has been associated with favorable prognosis,^{2,6-8,10-12} most prior analyses used simple dichotomous classification of CD8⁺ TILs and neglected to specify the inclusion or exclusion of stromal tissue. Prior analyses have been inadequately powered to evaluate histotype-specific survival associations. This is critical, as the invasive histotypes (high-grade serous OC [HGSOC], the most common and most lethal¹³; endometrioid OC [ENOC]¹⁴; clear cell OC [CCOC]^{14,15}; mucinous OC [MOC]¹⁶; and low-grade serous OC [LGSOC]¹⁷⁻²⁰) represent distinct biological processes, with distinct proposed cells of origin, clinical courses, and responses to chemotherapy.²¹⁻²³

We conducted a large-scale assessment of intraepithelial CD8⁺ TILs in more than 5000 prospectively followed patients with OC. Our goals were to clarify the associations and evaluate the functional form of CD8⁺ TILs with overall survival in HGSOCs, and to explore association of CD8⁺ TIL levels with overall survival in patients with other histotypes.

Methods

Study Design and Participants

We assembled a prospective cohort of 7377 women with a primary diagnosis of epithelial ovarian, peritoneal, or fallopian tube cancer, with a final sample size of 5577. Patients were followed from enrollment in an institutional review board-approved protocol until death from any cause (see eTable 1 in Supplement 1).²⁴⁻²⁶ We requested and received institutional review board/ethics board approval from more than 20 institutions participating in the Ovarian Tumor Tissue Analysis Consortium. None of the studies provided compensation for participants (Supplement 2). Tumor specimens were obtained at initial debulking surgery, formalin fixed, paraffin embedded, and arrayed on tissue microarrays (TMAs). Clinical covariates and vital status underwent standardized quality control measures. We excluded 288 patients owing to loss to follow-up, 11 with missing age at diagnosis, 65 with non-epithelial disease, and 1436 owing to inadequate quality or amount of arrayed tumor tissue. The final sample size of 5577 included 5078 women with tumors of the 5 major invasive histotypes (HGSOC, ENOC, CCOC, MOC, and LGSOC) (see

Key Points

Question To what extent are CD8⁺ tumor-infiltrating lymphocytes (TILs) prognostic in epithelial ovarian cancer?

Findings This cohort study of more than 24 000 person-years of follow-up on more than 5500 patients shows improved survival with increasing CD8⁺ TIL counts in high-grade serous, endometrioid, and mucinous ovarian cancers (*P* value for trends were 4.2×10^{-16} ; .008, and .04, respectively). Among high-grade serous ovarian cancers, this nearly log-linear relationship was present regardless of extent of residual disease following cytoreduction, receipt of standard treatment, and germline *BRCA1* mutation.

Meaning CD8⁺ TILs are a key prognostic factor in certain ovarian cancer histotypes and warrant additional study in the context of immunotherapy.

eTable 2 in Supplement 1). The median time from diagnosis to enrollment was zero days (interquartile range, 0-63 days); however, 38% of patients were enrolled more than 1 month after diagnosis. Because some HGSOC may be mistaken as ENOC,²⁷ we used WT1 and TP53 immunohistochemical staining from 17 studies to reclassify 82 ENOC cases as HGSOC; overall survival of these reclassified patients was consistent with HGSOC (see eFigure 1 in Supplement 1).

Immunohistochemical Analysis and Scoring

For most patients (4669 [84%]), staining was performed at the Mayo Clinic using the Leica Bond RX stainer; however, for patients enrolled at the Study of Epidemiology and Risk Factors and Cancer Heredity (SEA) and Mayo Clinic Ovarian Cancer Study (MAY1) study sites (476 [8%] and 432 [9%], respectively), previously stained slides were used. Immunohistochemical methods are provided in the eMethods in Supplement 1. Scoring was conducted at the University of Calgary; each core was screened for a hotspot of CD8⁺ TILs using a Nikon eclipse 80i microscope at original magnification $\times 200$. Within each hotspot, 1 high-power field at original magnification $\times 400$ with a 0.55-mm field diameter was evaluated, ensuring comparable area despite different core sizes across studies. Only CD8⁺ TILs within the epithelial component of the tumor (tumor islets) were considered, and CD8⁺ cells in the stroma or abutting tumor cells were disregarded (as seen, eg, in eFigure 2 in Supplement 1 classified as negative). A 4-point ordinal score was defined a priori based on CD8⁺ TIL counts per high-powered field: negative (none), low (1-2 TILs), moderate (3-19 TILs), and high (≥ 20 TILs), similar to the validated method of Zhang and colleagues,⁴ except that we decreased the low to moderate cutoff from 6 TILs to 3 TILs. We did this to increase ease and consistency of scoring, as a cutoff of 3 or more TILs is routinely used in colorectal carcinoma reporting to assess Lynch syndrome.²⁸ Multiple cores from 156 cases were evaluated blindly by 2 gynecologic pathologists (W.C. and M.K.), and a weighted κ statistic was estimated. Differences in interpretation were discussed at a multiheaded microscope, and the 2 pathologists scored 24% and 76% of the remaining cohort, respectively. The TMAs included an average of 2.4 cores per patient; for cases with more than 1 scored core, the maximum score was used, consistent with the scoring of hotspot regions.

Statistical Analysis

χ^2 Tests compared CD8⁺ TIL categories across clinical factors. Kaplan-Meier curves visually compared survival across categories. Cox proportional hazards regression estimated hazard ratios (HRs) and 95% CIs. Primary analyses were based on tests for trend, modeling the ordered CD8⁺ TIL categories as a 1 *df* linear term. Regression models included age at diagnosis (continuous), stage (I/II, III/IV, unknown), and study site as covariates; we also ran sensitivity analyses adjusting for extent of residual disease and postsurgical treatment. Separate analyses were conducted by histotype and among histopathological groupings (eg, combining LGSOC with their suspected precursor, serous borderline tumors), and by relevant clinical factors. This report meets reporting recommendations for tumor marker prognostic studies (REMARK)²⁹; additional statistical methods are provided in Supplement 1.

CD8⁺ TIL Cut Point Analysis

Because categorical CD8⁺ TIL cut points may artificially restrict variability in the data and can be somewhat arbitrary, the pathologist (M.K.) rescored all cores from a subset of 2175 patients (1449 with HGSOC), recording CD8⁺ TIL count as a numeric marker. Each core was given a value of 0 to 20 or greater, using a threshold of 20 for counts that exceeded that number. As before, the maximum score was used for cases with more than 1 scored core. Among patients with HGSOC, we compared survival distributions of those with rescored levels to those without using Cox proportional hazards regression. Among HGSOC cases with rescored CD8⁺ TIL levels, we ran 5 additional sets of Cox regression analyses. We first categorized the levels using our original thresholds (0, 1-2, 3-19, and ≥ 20 CD8⁺ TILs) to confirm that our original results using all HGSOC cases did not differ from the subset who were rescored. Second, we categorized the levels using the thresholds of Zhang and colleagues⁴ to determine the robustness of our original results to these cut points. Third, we assessed the functional form of the association between CD8⁺ TIL levels and survival using penalized B-splines.³⁰ Fourth, we fitted the numerically valued CD8⁺ TIL levels as a 1 *df* linear term. Finally, we carried out a formal cut point analysis similar to that described by Budczies and colleagues.³¹ Briefly, this approach examines all possible contiguous dichotomizations of TIL levels (ie, 0 vs ≥ 1 , 0-1 vs ≥ 2 , 0-2 vs ≥ 3) using Cox proportional hazards regression to identify the threshold that best discriminates survivors from nonsurvivors based on evidence of association.

Results

Distribution of CD8⁺ TILs by Histotype

The final sample included 5577 women; mean age at diagnosis was 58.4 years (median, 58.2 years). Epithelial CD8⁺ TILs were assessed using a 4-tiered scoring system (interobserver agreement, 81.8%; weighted κ , 0.846; 95% CI, 0.804-0.888). We observed intratumoral heterogeneity in CD8⁺ TILs across cores per patient (intraclass correlation coefficient, 0.56; 95% CI, 0.54-0.57). CD8⁺ TILs varied across the major invasive OC histotypes (HGSOC, ENOC, CCOC, MOC, and LGSOC; $\chi^2 P = 2.8 \times 10^{-103}$; see eFigure 3 in Supplement 1). Most HGSOC

cases (83%) had evidence of CD8⁺ TILs, with a lower proportion seen in LGSOC and ENOC cases (73% and 72%, respectively) and CCOC and MOC cases (52% and 51%, respectively). Most borderline tumors showed evidence of CD8⁺ TILs (serous, 84%; mucinous, 70%; see eTable 3 in Supplement 1).

Associations of CD8⁺ TILs With Overall Survival by Histotype

We observed a striking association for longer survival time with increasing levels of CD8⁺ TILs in HGSOC cases (*P* value for trend adjusted for study, age, and stage = 4.2×10^{-16}) (Table; Figure). Median survival was 2.8 years for women negative for CD8⁺ TILs, and 3.0 years, 3.8 years, and 5.1 years for low, moderate, or high levels, respectively. At the extremes, women with high levels of CD8⁺ TILs (≥ 20 per field) had a 43% reduced risk of death compared with women with no evidence of CD8⁺ TILs (HR, 0.57; 95% CI, 0.49-0.65) (Table). Associations were similar after adjustment for residual disease (see eTable 4 in Supplement 1).

Increasing levels of CD8⁺ TILs were also associated with longer survival time among women with ENOC (*P* value for trend = .008) (Table; Figure). This association was also apparent in separate analyses of grade 1 ENOC and grades 2 and 3 ENOC, although these were limited in sample size (see eTable 5 in Supplement 1). While there was a statistically significant dose-response similar to HGSOC, it is noteworthy that ENOCs with moderate levels (3-19 per field) showed the greatest improvement in survival time compared with women with ENOC and no detectable CD8⁺ TILs (HR, 0.50; 95% CI, 0.34-0.74).

A similar association was observed for women with MOC (*P* = .04) (Table; Figure), although, as the histotype with the lowest overall levels of CD8⁺ TILs, only 13 women (4%) had high TIL levels. Kaplan-Meier plots indicate a dose-response relationship, at least for negative to moderate levels (Figure). In contrast, CCOCs and LGSOCs showed no apparent association between CD8⁺ TILs and survival time (Table; see eFigure 4 in Supplement 1). Because LGSOC is the rarest of the invasive histotypes, the null association in this group should be interpreted with caution. As some prior studies combined LGSOC and HGSOC, we also analyzed invasive serous cases as a group, including those with missing grade. We found that the striking HGSOC results were attenuated (see eTable 5 in Supplement 1), suggesting that the relevance of CD8⁺ TILs among serous cases may be limited to HGSOC and confirming that immunohistochemistry-aided histotype classification is a critical first step to improving the classification of OC cases.^{27,32} No other patterns were observed in analyses of histopathological groups (see eTable 5 in Supplement 1).

Among the 5 major invasive histotypes, time to disease progression was known for 52% of cases (*n* = 2681). Progression-free survival results were remarkably similar to overall survival results (see eTable 6 in Supplement 1).

Associations of CD8⁺ TILs With Clinical Features in HGSOC

The extent of residual disease following primary cytoreductive surgery was available for 2173 HGSOC cases. Our results showed that a greater proportion of tumors without macroscopic residual disease had high CD8⁺ TIL levels than those with macroscopic disease (26% vs 20%; *P* = .006; see eFigure 3 in Supplement 1). Increasing CD8⁺ TILs were associated with

Table. Multivariable-Adjusted Association of CD8⁺ Tumor-Infiltrating Lymphocytes (TIL) and Overall Survival Among 5078 Patients With the 5 Most Common Invasive Epithelial Ovarian Cancer Histotype^a

Histotype	CD8 ⁺ TILs	No.	Person-years	Events, %	HR (95% CI)	P Value for Trend	P Value for 3 df Test
High-grade serous	Negative	546	1709.75	76.2	1 [Reference]	4.2×10^{-16}	7.2×10^{-15}
	Low	546	1908.39	72.3	0.86 (0.75-0.99)		
	Moderate	1394	5264.82	69.0	0.77 (0.69-0.87)		
	High	710	3110.87	56.5	0.57 (0.49-0.65)		
Endometrioid	Negative	206	1118.53	33.5	1 [Reference]	.008	.006
	Low	130	675.44	34.6	0.80 (0.54-1.18)		
	Moderate	283	1844.53	18.0	0.50 (0.34-0.74)		
	High	110	657.59	22.7	0.76 (0.47-1.23)		
Clear cell	Negative	309	1640.28	41.1	1 [Reference]	.50	.52
	Low	141	658.25	45.4	1.16 (0.84-1.60)		
	Moderate	118	618.79	41.5	0.88 (0.62-1.24)		
	High	80	412.44	40.0	0.92 (0.61-1.39)		
Mucinous	Negative	168	750.75	44.0	1 [Reference]	.04	.16
	Low	77	375.26	31.2	0.91 (0.55-1.51)		
	Moderate	85	470.62	27.1	0.56 (0.34-0.93)		
	High	13	72.35	23.1	0.79 (0.23-2.68)		
Low-grade serous	Negative	43	198.06	48.8	1 [Reference]	.91	>.99
	Low	44	184.89	65.9	0.94 (0.50-1.74)		
	Moderate	63	272.13	47.6	0.98 (0.52-1.83)		
	High	12	49.53	41.7	0.92 (0.33-2.59)		

Abbreviation: TIL, tumor-infiltrating lymphocytes.

^a Adjusted for study, age (continuous), and stage (I/II, III/IV, unknown); levels based on counts of TIL per high-powered field (HPF): negative, no CD8⁺ TILs; low, 1-2 CD8⁺ TILs; moderate, 3-19 CD8⁺ TILs; high, 20 or more CD8⁺ TILs per

HPF; adjusted for study, age (continuous), and stage (I/II, III/IV, unknown); HR, hazard ratio; P value trend, from a 1 df trend test; P value for 3 df test, from an unordered test comparing risk across the 4 CD8⁺ TIL groups.

improved survival in a dose-response manner in both surgical outcome groups, indicating that immune response improves prognosis regardless of the remaining residual disease after surgery (see eTable 7 in Supplement 1).

Our study included 133 *BRCA1* and 66 *BRCA2* mutation carriers and 844 tested noncarriers. The extent of CD8⁺ TILs differed by mutation status ($P = .02$), as 29% of *BRCA1* mutation carriers had high TIL counts, yet only 18% of noncarriers and 15% of *BRCA2* mutation carriers did (see eFigure 3 in Supplement 1). The survival benefit associated with CD8⁺ TILs was also found to differ by mutation status (P value for interaction = .006). Increased CD8⁺ TILs were associated with favorable survival among cases without mutations ($P = 5.1 \times 10^{-7}$) and among cases with a *BRCA1* mutation ($P = .003$) (see eTable 7 in Supplement 1). Among *BRCA2* mutation carriers, there was no evidence of association between CD8⁺ TILs and survival ($P = .62$).

Treatment details were documented for 501 patients with HGSOE who received standard first-line chemotherapy, including 295 who received the standard dose (carboplatin area under the curve, 5 or 6, and paclitaxel, 135 mg/m² or 175 mg/m²). Association with CD8⁺ TIL level and overall survival was also observed within this group (P value for trend = .003) (see eTable 7 in Supplement 1).

Among HGSOE cases, CD8⁺ TIL level was associated with earlier stage ($P = 4.3 \times 10^{-4}$) and younger age at diagnosis ($P = 1.6 \times 10^{-4}$). In stratified analyses CD8⁺ TIL level was consistently prognostic in stage and age subgroups (see eTable 5 in Supplement 1). We also observed that patients born more


recently showed higher levels ($n = 2734$, $P = .001$); we also adjusted all analyses for year of birth, and results were similar. CD8⁺ TIL level was not associated with year of diagnosis ($P = .71$), self-reported racial group ($P = .74$), or pretreatment or posttreatment level of protein CA 125 ($P = .42$ and $.89$, respectively).

Analysis of CD8⁺ TIL Cut Points in HGSOE

Of the 3196 HGSOE cases, 1449 (45%) were rescored using a numeric count. There were no differences in survival between women who were rescored and those who were not ($P = .12$; κ values comparing original values to rescored values, 0.91 (95% CI, 0.89-0.92). eTable 8 in Supplement 1 displays associations of categorized CD8⁺ TIL levels and survival in women with rescored tumors. After adjustment for age and stage, strong dose-response associations were observed using both the original threshold values (0, 1-2, 3-19, and ≥ 20 CD8⁺ TILs) and those used by Zhang and colleagues⁴ (0, 1-5, 6-19, ≥ 20 CD8⁺ TILs) ($P < 10^{-5}$ for each). As before, associations were slightly attenuated but remained significant after adjustment for extent of residual disease and postsurgical treatment ($P < 10^{-4}$ for each).

Assessment of the functional form of the association between numeric CD8⁺ TIL levels and survival using penalized B-splines, after adjustment for age and stage, is shown in eFigure 5 in Supplement 1. We observed a strong negative association with survival, indicating that increasing CD8⁺ TIL levels are progressively protective across this spectrum of values. The results of fitting CD8⁺ TIL levels as a 1 df linear term are also

Figure. Kaplan-Meier Overall Survival Plots by CD8⁺ Tumor-Infiltrating Lymphocyte (TIL) Levels for the High-Grade Serous, Endometrioid, and Mucinous Ovarian Cancer



Negative, no CD8⁺ TILs; low, 1-2 CD8⁺ TILs; moderate, 3-19 CD8⁺ TILs; high, 20 or more CD8⁺ TILs per high-powered field. The numbers just below the x-axis represent the number of women at risk in 2-year time intervals. Number at risk on date of diagnosis may be smaller than number at risk later owing to left truncation of follow-up resulting from delayed study enrollment.

shown in eFigure 5 in Supplement 1 and track very closely to those using penalized B-splines, indicating that the association between CD8⁺ TIL levels and survival in women with HGSOc is virtually log-linear in nature.

Results of a formal cut point analysis examining all possible sets of contiguous dichotomizations of TIL levels can be found in eTable 9 in Supplement 1. The best discrimination of survivors from nonsurvivors occurred when comparing those with 0 to 13 TILs with those with 14 or more (HR, 0.75; 95% CI, 0.65-0.86; $P = 1.5 \times 10^{-5}$). However, each of the 19 dichotomizations yielded highly significant results (P values for all comparisons, $\leq 1.1 \times 10^{-3}$), with HRs consistently ranging from 0.75 to 0.83, again indicating that greater TIL levels are protective across the entire spectrum of values examined.

Discussion

To our knowledge, our study is the largest report on intraepithelial CD8⁺ TILs in OC to date and shows a robust dose-dependent increase in survival for increasing TIL levels in women with HGSOc. Analyses on a subset of individuals using numeric TIL counts confirmed a progressively protective, nearly log-linear survival effect as CD8⁺ TILs counts increased from 0 to 20 or more per high-powered field, suggesting that the quantity of CD8⁺ TILs, not merely their presence, is informative and that the most immune-rich HGSOcs are the most likely to have improved clinical outcome. This effect was not modified or confounded by the extent of residual disease after cytoreductive surgery. Because there are fewer than a handful of other validated prognostic biomarkers for HGSOc (eg, *BRCA1* and *BRCA2* status³³ and PR expression²⁶), these results may provide increased prognostic prediction.

In addition, to our knowledge, this is the first CD8⁺ TIL study in histotypes other than HGSOc; we revealed a significant reduction in risk of death for patients diagnosed as having ENOC and MOC. In ENOC, patients with moderate CD8⁺ TIL levels had the most favorable survival, with no additional benefit observed beyond this threshold. Because prior reports suggest that ENOCs with high CD8⁺ TIL are more commonly mismatch repair deficient,³⁴ we speculate that, similar to endometrial cancers,^{35,36} ENOC with high CD8⁺ TIL levels may be associated with an intermediate outcome owing to the association with mismatch repair deficiency. No survival associations were seen in CCOC.

Other investigations have noted higher response rates to immune checkpoint blockade among patients with a higher burden of neoantigens,^{37,38} suggesting that increased neoantigens increases the likelihood that T lymphocytes recognize tumor as foreign and mount an immune response. It has also been demonstrated that *BRCA1*-mutated HGSOc tumors have a higher average neoantigen number than nonmutated tumors.³⁹ In this study, patients with HGSOc with germline *BRCA1* mutations demonstrated higher levels of CD8⁺ TILs than patients with *BRCA2* mutations or those tested negative for mutation. While neoantigen load may explain higher levels of CD8⁺ TILs in *BRCA1*-mutated tumors, and their association with better outcome, it does not explain the better outcome of *BRCA2*-mutated tumors.^{40,41}

Limitations

Given its robust prognostic ability, relative ease of testing, and low interobserver variability (percent agreement = 81.8%,

weighted $\kappa = 0.846$), quantitation of CD8⁺ TILs should be considered for clinical evaluation as suggested for other cancers.⁴²⁻⁴⁴ Unfortunately, as expected, we found intratumoral heterogeneity in CD8⁺ TILs across cores per patient (intraclass correlation coefficient = 0.56; 95% CI, 0.54-0.57). To account for this, we use the maximum score, which is akin to the hotspot assessment of proliferation in other cancers, and is more feasible for surgical specimens with many tumor-containing slides. We also propose that, similar to the system for patients with breast cancer, a practical and robust scoring system should be developed.⁴³ Additional issues requiring large-scale study that were not evaluated here include utility of image analysis; evaluation of stromal CD8⁺ TILs; consistency across multiple tumor sites per patient; impact of neoadjuvant chemotherapy⁴⁵⁻⁴⁷; relationships among CD8⁺ TIL levels, HGSOC molecular subtypes,⁴⁸⁻⁵⁰ common genetic variation,⁵¹ and epidemiologic risk factors⁵²; and evaluation of other lymphocyte subsets, such as CD4⁺ TILs, CD20⁺ TILs (B cells), tertiary lymphoid structures, and plasma cells.^{10,12-14,53} Clinically, it will be important to test whether CD8⁺ TILs predict response to certain therapies including standard

chemotherapy and immune therapy, as, for example, CD8⁺ TILs predict chemoresponse in subtypes of breast cancer.⁵⁴ It will also be critical to study whether the immune response of CD8⁺ TILs can be activated by checkpoint blockade.

Conclusions

These large-scale analyses show that CD8⁺ TILs vary by histotype with HGSOC tumors having the highest levels and a strong association with survival, regardless of extent of residual disease or first-line chemotherapy treatment. Penalized B-splines revealed that this association was nearly log-linear in nature, indicating that progressively greater TIL counts yield progressively better prognoses for HGSOC tumors. We showed for the first time that CD8⁺ TILs in HGSOC cases with germline *BRCA2* mutations may not be associated with survival. Finally, we found that ENOC and MOC tumors show trends associating CD8⁺ TILs with survival time and that CCOC do not show these trends. A clinically applicable scoring system for CD8⁺ TILs should be developed to incorporate into clinical trials.

ARTICLE INFORMATION

Accepted for Publication: August 5, 2017.

Published Online: October 12, 2017.

doi:10.1001/jamaoncol.2017.3290

The Authors/Members of the Ovarian Tumor Tissue Analysis (OTTA) Consortium:

Ellen L. Goode, PhD, MPH; Matthew S. Block, MD, PhD; Kimberly R. Kalli, PhD; Robert A. Vierkant, MS; Wenqian Chen, MD; Zachary C. Fogarty, MS; Aleksandra Gentry-Maharaj, PhD; Aleksandra Toloczko, BS; Alexander Hein, MD; Alicia L. Bouligny; Allan Jensen, PhD; Ana Osorio, PhD; Andreas D. Hartkopf, MD; Andy Ryan, PhD; Anita Chudecka-Glaz, MD, PhD; Anthony M. Magliocco, MD; Arndt Hartmann, MD, PhD; Audrey Y. Jung, PhD; Bo Gao, BMed, PhD; Brenda Y. Hernandez, PhD, MPH; Brooke L. Fridley, PhD; Bryan M. McCauley, BS; Catherine J. Kennedy, BSc (Hons); Chen Wang, PhD; Chloe Karpinskyj, BS; Christiani B. de Sousa, MD, PhD; Daniel G. Tiezzi, MD, PhD; David L. Wachter, MD; Esther Herpel, MD, PhD; Florin Andrei Taran, MD; Francesmary Modugno, PhD; Gregg Nelson, MD, PhD; Jan Lubiński, MD, PhD; Janusz Menkiszak, MD, PhD; Jennifer Alsop, BA; Jenny Lester, MPH, CCRP; Jesús García-Donas, MD; Jill Nation, MD; Jillian Hung, MHS; José Palacios, MD, PhD; Joseph H. Rothstein, MS; Joseph L. Kelley, MD; Jurandy M. de Andrade, MD, PhD; Luis Robles-Díaz, MD; Maria P. Intermaggio, MS; Martin Widschwendter, MD; Matthias W. Beckmann, MD, PhD; Matthias Ruebner, PhD; Mercedes Jimenez-Linan, MD; Naveena Singh, MD; Oleg Oszurek, PhD; Paul R. Harnett, MBBS, PhD; Peter F. Rimbau, MD; Peter Sinn, MD, PhD; Philipp Wagner, MD; Prafull Ghatage, MD; Raghwa Sharma, MBBS, FRCPA; Robert P. Edwards, MD; Roberta B. Ness, MD, MPH; Sandra Orsulic, PhD; Sara Y. Brucker, MD, PhD; Sharon E. Johnatty, PhD; Teri A. Longacre, MD; Ursula Eilber, PhD; Valerie McGuire, PhD; Weiva Sieh, PhD; Yanina Natanzon, PhD, MS; Zheng Li, MD; Alice S. Whittemore, PhD; Anna deFazio, PhD; Annette Staebler, MD; Beth Y. Karlan, MD; Blake Gilks, MD; David D. Bowtell, PhD; Estrid Høgdall, PhD; Francisco J. Candido dos Reis, MD,

PhD; Helen Steed, MD; Ian G. Campbell, PhD; Jacek Gronwald, MD, PhD; Javier Benítez, PhD; Jennifer M. Koziak, BSc; Jenny Chang-Claude, PhD; Kirsten B. Moysich, PhD; Linda E. Kelemen, ScD; Linda S. Cook, PhD; Marc T. Goodman, PhD; María José García, PhD; Peter A. Fasching, MD; Stefan Kommos, MD; Suha Deen, MChB; Susanne K. Kjaer, MD; Usha Menon, MD; James D. Brenton, MD, PhD; Paul D. P. Pharoah, MD, PhD; Georgia Chenevix-Trench, PhD; David G. Huntsman, MD; Stacey J. Winham, PhD; Martin Köbel, MD; Susan J. Ramus, PhD.

Affiliations of the Authors/Members of the Ovarian Tumor Tissue Analysis (OTTA)

Consortium: Department of Health Sciences Research, Mayo Clinic, Rochester, Minnesota (Goode, Vierkant, Fogarty, Bouligny, McCauley, Wang, Natanzon, Li, Winham); Department of Medical Oncology, Mayo Clinic, Rochester, Minnesota (Block, Kalli); Department of Pathology and Laboratory Medicine, University of Calgary, Calgary, Alberta, Canada (Chen, Rimbau, Köbel); Gynaecological Cancer Research Centre, Department of Women's Cancer, Institute for Women's Health, University College London, London, England (Gentry-Maharaj, Ryan, Karpinskyj, Widschwendter, Menon); International Hereditary Cancer Center, Department of Genetics and Pathology, Pomeranian Medical University, Szczecin, Poland (Toloczko, Lubiński, Oszurek, Gronwald); Department of Gynecology and Obstetrics, University Hospital Erlangen, Friedrich-Alexander-University, Erlangen-Nuremberg Comprehensive Cancer Center, Erlangen EMN, Germany (Hein, Beckmann, Ruebner, Fasching); Department of Health Sciences, Spelman College, Atlanta, Georgia (Bouligny); Virus, Lifestyle and Genes, Danish Cancer Society Research Center, Copenhagen, Denmark (Jensen, Høgdall, Kjaer); Human Genetics Group, Spanish National Cancer Research Centre (CNIO), Madrid, Spain (Osorio, Benítez, García); Centre for Biomedical Network Research on Rare Diseases (CIBERER), Instituto de Salud Carlos III, Madrid, Spain (Osorio, Benítez,

García); Tübingen University Hospital, Department of Women's Health, Tübingen, Germany (Hartkopf, Taran, Wagner, Brucker, Kommos); Department of Gynecological Surgery and Gynecological Oncology of Adults and Adolescents, Pomeranian Medical University, Szczecin, Poland (Chudecka-Glaz, Menkiszak); Department of Anatomic Pathology, Moffitt Cancer Center, Tampa, Florida (Magliocco); Institute of Pathology, University Hospital Erlangen, Friedrich-Alexander University Erlangen Nuremberg, Comprehensive Cancer Center Erlangen-EMN, Erlangen, Germany (Hartmann, Wachter); Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany (Jung, Eilber, Chang-Claude); Centre for Cancer Research, The Westmead Institute for Medical Research, The University of Sydney, Sydney, New South Wales, Australia (Gao, Kennedy, Hung, Harnett, deFazio); The Crown Princess Mary Cancer Centre, Westmead Hospital, The University of Sydney, Australia (Gao, Harnett); Cancer Center, University of Hawaii, Honolulu (Hernandez); Department of Biostatistics and Bioinformatics, Division of Population Sciences, Moffitt Cancer Center, Tampa, Florida (Fridley); Department of Gynaecological Oncology, Westmead Hospital, Sydney, New South Wales, Australia (Kennedy, Hung, deFazio); Department of Gynecology and Obstetrics, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, Brazil (de Sousa, Tiezzi, de Andrade, Candido dos Reis); Tissue Bank of the National Center for Tumor Diseases (NCT) Heidelberg, Germany and Institute of Pathology, University of Heidelberg, Heidelberg, Germany (Herpel); Department of Obstetrics, Gynecology and Reproductive Sciences, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania (Modugno, Kelley, Edwards); Department of Epidemiology, University of Pittsburgh Graduate School of Public Health, Pittsburgh, Pennsylvania (Modugno); Women's Cancer Research Program, Magee-Women's Research Institute and University of Pittsburgh Cancer Institute, Pittsburgh, Pennsylvania (Modugno); Department of Oncology, Division of

Gynecologic Oncology, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada (Nelson, Nation, Ghatage); Department of Oncology, University of Cambridge, Strangeways Research Laboratory, Cambridge, England. (Alsop, Pharoah); Women's Cancer Program, Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, California (Lester, Orsulic, Karlan); Medical Oncology Service, HM Hospitales-Centro Integral Oncológico HM Clara Campal, Madrid, Spain (García-Donas); Department of Pathology, Hospital Universitario Ramón y Cajal, IRYCIS, Universidad de Alcalá, Madrid, Spain (Palacios); Department of Population Health Science and Policy and Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, New York (Rothstein, Sieh); Familial Cancer Unit and Medical Oncology Department, Hospital Universitario 12 de Octubre, Madrid, Spain (Robles-Díaz); School of Women's and Children's Health, University of New South Wales, Sydney, Australia (Intermaggio, Ramus); Department of Histopathology, Addenbrookes Hospital, Cambridge, England (Jimenez-Linan); Department of Pathology, Barts Health National Health Service Trust, London, England (Singh); Department of Pathology, Catholic University of Health and Allied Sciences-Bugando, Mwanza, Tanzania (Rambau); Department of Pathology, Institute of Pathology, Heidelberg University Hospital, Germany (Sinn); Pathology West ICPMR Westmead, Westmead Hospital, The University of Sydney, Sydney, Australia (Sharma); University of Western Sydney at Westmead Hospital, Westmead, New South Wales, Australia (Sharma); University of Texas School of Public Health, Houston, (Ness); Department of Genetics, QIMR Berghofer Medical Research Institute, Brisbane, Australia (Johnatty, Chenevix-Trench); Department of Pathology, Stanford University, Stanford, California (Longacre); Department of Health Research and Policy, Stanford University School of Medicine, Stanford, California (McGuire); Department of Gynecologic Oncology, The Third Affiliated Hospital of Kunming Medical University (Yunnan Tumor Hospital), Kunming, China (Li); Department of Health Research and Policy and Department of Biomedical Data Science, Stanford University School of Medicine, Stanford, California (Whittemore); Tübingen University Hospital, Institute of Pathology, Tübingen, Germany (Staebler); Genetic Pathology Evaluation Centre, Vancouver General Hospital and University of British Columbia, Vancouver, British Columbia, Canada (Gilks); Cancer Genomics Program, Research Department, Peter MacCallum Cancer Centre, East Melbourne, Victoria, Australia (Bowtell); Sir Peter MacCallum Department of Oncology, University of Melbourne, Parkville, Victoria, Australia (Bowtell); The Garvan Institute, Sydney, New South Wales, Australia (Bowtell, Ramus); Molecular Unit, Department of Pathology, Herlev Hospital, University of Copenhagen, Copenhagen, Denmark (Høgdaal); Department of Obstetrics and Gynecology, Division of Gynecologic Oncology, Royal Alexandra Hospital, Edmonton, Alberta, Canada (Steed); Cancer Genetics Laboratory, Research Division, Peter MacCallum Cancer Centre, Melbourne, Australia (Campbell); Sir Peter MacCallum Department of Oncology, University of Melbourne, Australia (Campbell); Department of Pathology, University of Melbourne, Melbourne, Victoria, Australia (Campbell); Alberta Health Services-Cancer Care,

Calgary, Alberta, Canada (Koziak); University Cancer Center Hamburg, University Medical Center Hamburg-Eppendorf, Hamburg, Germany (Chang-Claude); Department of Cancer Prevention and Control, Roswell Park Cancer Institute, Buffalo, New York (Moysich); Department of Public Health Sciences, Medical University of South Carolina and Hollings Cancer Center, Charleston (Kelemen); Division of Epidemiology and Biostatistics, University of New Mexico, Albuquerque (Cook); Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, California (Goodman); David Geffen School of Medicine, Department of Medicine Division of Hematology and Oncology, University of California at Los Angeles (Fasching); Department of Histopathology, Queen's Medical Centre, Nottingham University Hospitals NHS Trust, Nottingham, England (Deen); Department of Gynecology, Rigshospitalet, University of Copenhagen, Copenhagen, Denmark (Kjaer); Department of Oncology, University of Cambridge, Hutchison/MRC Research Centre, Cambridge, England (Brenton); Cancer Research UK Cambridge Institute, University of Cambridge, Li Ka Shing Centre, Robinson Way, Cambridge, England (Brenton); Cambridge Experimental Cancer Medicine Centre, Cambridge, England (Brenton); Department of Public Health and Primary Care, University of Cambridge, Strangeways Research Laboratory, Worts Causeway, Cambridge, England (Pharoah); Department of Pathology and Laboratory Medicine, University of British Columbia, Vancouver, British Columbia, Canada (Huntsman); Centre for Translational and Applied Genomics, British Columbia Cancer Agency, Vancouver, British Columbia, Canada (Huntsman).

Author Contributions: Dr Goode had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. Mr Vierkant and Dr Winham conducted and were responsible for the data analysis.

Study concept and design: Goode, Block, Vierkant, Hartmann, Nation, Jimenez-Linan, Ness, Longacre, Bowtell, Moysich, Deen, Brenton, Pharoah, Köbel, Ramus.

Acquisition, analysis, or interpretation of data: Goode, Block, Kalli, Vierkant, Chen, Fogarty, Gentry-Maharaj, Tołoczko, Hein, Bouligny, Jensen, Osorio, Hartkopf, Ryan, Chudecka-Glaz, Magliocco, Hartmann, Jung, Gao, Hernandez, Fridley, McCauley, Kennedy, Wang, Karpinskyj, de Sousa, Tiezzi, Wachter, Herpel, Taran, Modugno, Nelson, Lubiński, Menkiszak, Alsop, Lester, García-Donas, Hung, Palacios, Rothstein, Kelley, de Andrade, Robles-Díaz, Intermaggio, Widschwendter, Beckmann, Ruebner, Singh, Oszurek, Harnett, Rambau, Sinn, Wagner, Ghatage, Sharma, Edwards, Orsulic, Brucker, Johnatty, Eilber, McGuire, Sieh, Natanzon, Li, Whittemore, deFazio, Staebler, Karlan, Gilks, Bowtell, Høgdaal, Candido dos Reis, Steed, Campbell, Gronwald, Benítez, Koziak, Chang-Claude, Moysich, Kelemen, Cook, Goodman, García, Fasching, Kommoss, Kjaer, Menon, Pharoah, Chenevix-Trench, Huntsman, Winham, Köbel, Ramus.

Drafting of the manuscript: Goode, Block, Vierkant, Fogarty, Jimenez-Linan, Rambau, Sharma, Brucker, Li, Bowtell, Köbel, Ramus; OTTA.

Critical revision of the manuscript for important intellectual content: Goode, Block, Kalli, Vierkant, Chen, Gentry-Maharaj, Tołoczko, Hein, Bouligny,

Jensen, Osorio, Hartkopf, Ryan, Chudecka-Glaz, Magliocco, Hartmann, Jung, Gao, Hernandez, Fridley, McCauley, Kennedy, Wang, Karpinskyj, de Sousa, Tiezzi, Wachter, Herpel, Taran, Modugno, Nelson, Lubiński, Menkiszak, Alsop, Lester, García-Donas, Nation, Hung, Palacios, Rothstein, Kelley, de Andrade, Robles-Díaz, Intermaggio, Widschwendter, Beckmann, Ruebner, Singh, Oszurek, Harnett, Sinn, Wagner, Ghatage, Sharma, Edwards, Ness, Orsulic, Brucker, Johnatty, Longacre, Eilber, McGuire, Sieh, Natanzon, Whittemore, deFazio, Staebler, Karlan, Gilks, Høgdaal, Candido dos Reis, Steed, Campbell, Gronwald, Benítez, Koziak, Chang-Claude, Moysich, Kelemen, Cook, Goodman, García, Fasching, Kommoss, Deen, Kjaer, Menon, Brenton, Pharoah, Chenevix-Trench, Huntsman, Winham, Köbel, Ramus.

Statistical analysis: Vierkant, Chen, Fogarty, Gao, Fridley, McCauley, Wang, Winham.

Obtained funding: Goode, Modugno, Edwards, Ness, Moysich, Kelemen, Cook, Goodman, Kjaer, Chenevix-Trench.

Administrative, technical, or material support:

Goode, Block, Kalli, Gentry-Maharaj, Tołoczko, Hein, Bouligny, Jensen, Osorio, Hartkopf, Chudecka-Glaz, Magliocco, Hartmann, Jung, Hernandez, Kennedy, Karpinskyj, de Sousa, Tiezzi, Wachter, Herpel, Taran, Lubiński, Alsop, Lester, Nation, Rothstein, de Andrade, Intermaggio, Widschwendter, Beckmann, Ruebner, Jimenez-Linan, Singh, Harnett, Wagner, Ness, Orsulic, Johnatty, Eilber, McGuire, Sieh, Natanzon, deFazio, Staebler, Karlan, Gilks, Bowtell, Høgdaal, Candido dos Reis, Campbell, Benítez, Koziak, Chang-Claude, Moysich, Kelemen, Cook, García, Fasching, Kommoss, Deen, Menon, Huntsman, Ramus.

Study supervision: Goode, Vierkant, Nelson, Menkiszak, García-Donas, Ghatage, Sharma, Edwards, Ness, Moysich, Brenton, Huntsman.

Conflict of Interest Disclosures: Dr Bowtell receives funding from Astra Zeneca and Genentech Roche for the conduct of research studies and clinical trials unrelated to the work described in this article. Dr Huntsman is a founder and Chief Medical Officer of Contextual Genomics, a somatic mutation testing laboratory; the company's work and interests do not overlap with the subject of and methodologies used in this article. No other disclosures are reported.

Funding/Support: Funding was provided by Canadian Institutes for Health Research (MOP-86727); Brazilian National Council for Scientific and Technological Development, grant No. 478416/2009-1; Calgary Laboratory Services Internal Research Competition RS10-533; German Federal Ministry of Education and Research of Germany (01 GB 9401); German Cancer Research Center (DKFZ); US National Cancer Institute K07-CA80668, P50-CA159981, RO1CA095023; National Institutes of Health/National Center for Research Resources/General Clinical Research Center grant MO1-RR000056; US Army Medical Research and Materiel Command DAMD17-02-1-0669; NIH (R01-CA122443, P50-CA136393, P30-CA15083, U01-CA71966, U01-CA69417, R01-CA16056, K07-CA143047); Cancer Research UK (C490/A10119, C490/A10123, C490/A16561); UK National Institute for Health Research Biomedical Research Centres at the University of Cambridge and at University College Hospital, "Women's Health Theme"; NIH SFB 685; the Eve Appeal; the Oak Foundation; and

Deutsche Forschungsgemeinschaft. The Australian Ovarian Cancer Study Group was supported by the US Army Medical Research and Materiel Command under DAMD17-01-1-0729; The Cancer Council Victoria; Queensland Cancer Fund; The Cancer Council, New South Wales; the Tom Baker Cancer Centre Translational Laboratories; The Cancer Council, South Australia; The Cancer Foundation of Western Australia; The Cancer Council Tasmania; and the National Health and Medical Research Council of Australia (NHMRC; ID400413, ID400281). The AOCs gratefully acknowledges additional support from the Peter MacCallum Cancer Foundation and Ovarian Cancer Australia (OCA). The Gynaecological Oncology Biobank at Westmead, a member of the Australasian Biospecimen Network-Oncology group, was funded by the National Health and Medical Research Council Enabling grants ID 310670 and ID 628903 and the Cancer Institute NSW grants 12/RIG/1-17 and 15/RIG/1-16. Funding for MALOVA was provided by research grant R01-CA61107 from the National Cancer Institute, Bethesda, Maryland; research grant 94 222 52 from the Danish Cancer Society, Copenhagen, Denmark; and the Mermaid I project. Drs DeFazio and Harnett are funded by Cancer Institute NSW grant 15/TRC/1-01. Dr Karlan is funded by the American Cancer Society Early Detection Professorship (SIOP-06-258-01-COUN) and the National Center for Advancing Translational Sciences (NCATS), grant UL1TRO00124.

Role of the Funder/Sponsor: The funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

REFERENCES

- Siegel RL, Miller KD, Jemal A. Cancer Statistics, 2017. *CA Cancer J Clin*. 2017;67(1):7-30.
- Hamanishi J, Mandai M, Ikeda T, et al. Safety and antitumor activity of anti-PD-1 antibody, nivolumab, in patients with platinum-resistant ovarian cancer. *J Clin Oncol*. 2015;33(34):4015-4022.
- Brahmer JR, Tlykodi SS, Chow LQ, et al. Safety and activity of anti-PD-L1 antibody in patients with advanced cancer. *N Engl J Med*. 2012;366(26):2455-2465.
- Zhang L, Conejo-Garcia JR, Katsaros D, et al. Intratumoral T cells, recurrence, and survival in epithelial ovarian cancer. *N Engl J Med*. 2003;348(3):203-213.
- Tomsová M, Melichar B, Sedláková I, Steiner I. Prognostic significance of CD3⁺ tumor-infiltrating lymphocytes in ovarian carcinoma. *Gynecol Oncol*. 2008;108(2):415-420.
- Han LY, Fletcher MS, Urbauer DL, et al. HLA class I antigen processing machinery component expression and intratumoral T-cell infiltrate as independent prognostic markers in ovarian carcinoma. *Clin Cancer Res*. 2008;14(11):3372-3379.
- Clarke B, Tinker AV, Lee CH, et al. Intraepithelial T cells and prognosis in ovarian carcinoma: novel associations with stage, tumor type, and BRCA1 loss. *Mod Pathol*. 2009;22(3):393-402.
- Hwang WT, Adams SF, Tahirovic E, Hagemann IS, Coukos G. Prognostic significance of tumor-infiltrating T cells in ovarian cancer: a meta-analysis. *Gynecol Oncol*. 2012;124(2):192-198.
- Blum JS, Wearsch PA, Cresswell P. Pathways of antigen processing. *Annu Rev Immunol*. 2013;31:443-473.
- Sato E, Olson SH, Ahn J, et al. Intraepithelial CD8⁺ tumor-infiltrating lymphocytes and a high CD8⁺/regulatory T cell ratio are associated with favorable prognosis in ovarian cancer. *Proc Natl Acad Sci U S A*. 2005;102(51):18538-18543.
- Callahan MJ, Nagymanyoki Z, Bonome T, et al. Increased HLA-DMB expression in the tumor epithelium is associated with increased CTL infiltration and improved prognosis in advanced-stage serous ovarian cancer. *Clin Cancer Res*. 2008;14(23):7667-7673.
- Preston CC, Maurer MJ, Oberg AL, et al. The ratios of CD8⁺ T cells to CD4⁺CD25⁺ FOXP3⁺ and FOXP3⁻ T cells correlate with poor clinical outcome in human serous ovarian cancer. *PLoS One*. 2013;8(11):e80063.
- Kurman RJ. Origin and molecular pathogenesis of ovarian high-grade serous carcinoma. *Ann Oncol*. 2013;24(suppl 10):x16-x21.
- Worley MJ, Welch WR, Berkowitz RS, Ng SW. Endometriosis-associated ovarian cancer: a review of pathogenesis. *Int J Mol Sci*. 2013;14(3):5367-5379.
- Mabuchi S, Sugiyama T, Kimura T. Clear cell carcinoma of the ovary: molecular insights and future therapeutic perspectives. *J Gynecol Oncol*. 2016;27(3):e31.
- Kelemen LE, Köbel M. Mucinous carcinomas of the ovary and colorectum: different organ, same dilemma. *Lancet Oncol*. 2011;12(11):1071-1080.
- Emmanuel C, Chiew YE, George J, et al; Australian Ovarian Cancer Study (AOCs). Genomic classification of serous ovarian cancer with adjacent borderline differentiates RAS pathway and TP53-mutant tumors and identifies NRAS as an oncogenic driver. *Clin Cancer Res*. 2014;20(24):6618-6630.
- Oswald AJ, Gourley C. Low-grade epithelial ovarian cancer: a number of distinct clinical entities? *Curr Opin Oncol*. 2015;27(5):412-419.
- Grabowski JP, Harter P, Heitz F, et al. Operability and chemotherapy responsiveness in advanced low-grade serous ovarian cancer: an analysis of the AGO Study Group metadatabase. *Gynecol Oncol*. 2016;140(3):457-462.
- Fader AN, Java J, Ueda S, et al; Gynecologic Oncology Group (GOG). Survival in women with grade 1 serous ovarian carcinoma. *Obstet Gynecol*. 2013;122(2, pt 1):225-232.
- Köbel M, Bak J, Bertelsen BI, et al. Ovarian carcinoma histotype determination is highly reproducible, and is improved through the use of immunohistochemistry. *Histopathology*. 2014;64(7):1004-1013.
- Ramalingam P. Morphologic, immunophenotypic, and molecular features of epithelial ovarian cancer. *Oncology (Williston Park)*. 2016;30(2):166-176.
- Köbel M, Kalloger SE, Boyd N, et al. Ovarian carcinoma subtypes are different diseases: implications for biomarker studies. *PLoS Med*. 2008;5(12):e232.
- Köbel M, Madore J, Ramus SJ, et al; AOCs Study Group. Evidence for a time-dependent association between FOLR1 expression and survival from ovarian carcinoma: implications for clinical testing: an Ovarian Tumour Tissue Analysis consortium study. *Br J Cancer*. 2014;111(12):2297-2307.
- Köbel M, Kalloger SE, Lee S, et al; Ovarian Tumour Tissue Analysis consortium. Biomarker-based ovarian carcinoma typing: a histologic investigation in the ovarian tumor tissue analysis consortium. *Cancer Epidemiol Biomarkers Prev*. 2013;22(10):1677-1686.
- Sieh W, Köbel M, Longacre TA, et al. Hormone-receptor expression and ovarian cancer survival: an Ovarian Tumour Tissue Analysis consortium study. *Lancet Oncol*. 2013;14(9):853-862.
- Köbel M, Rahimi K, Rambau PF, et al. An immunohistochemical algorithm for ovarian carcinoma typing. *Int J Gynecol Pathol*. 2016;35(5):430-441.
- College of American Pathologists. Protocol for the examination of specimens from patients with primary carcinoma of the colon and rectum. 2016; <http://www.cap.org/ShowProperty?nodePath=/UCMCon/Contribution%20Folders/WebContent/pdf/cp-colon-16protocol-3400.pdf>. Accessed August 22, 2017.
- McShane LM, Altman DG, Sauerbrei W, Taube SE, Gion M, Clark GM; Statistics Subcommittee of the NCI-EORTC Working Group on Cancer Diagnostics. Reporting recommendations for tumour MARKer prognostic studies (REMARK). *Br J Cancer*. 2005;93(4):387-391.
- Eilers PHC, Marx BD. Flexible smoothing with B-splines and penalties. *Stat Sci*. 1996;11(2):89-102.
- Budczies J, Klauschen F, Sinn BV, et al. Cutoff Finder: a comprehensive and straightforward web application enabling rapid biomarker cutoff optimization. *PLoS One*. 2012;7(12):e51862.
- Kar SP, Berchuck A, Gayther SA, et al. Common genetic variation and susceptibility to ovarian cancer: current insights and future directions [published online June 14, 2017]. *Cancer Epidemiol Biomarkers Prev*. 2017;cebp.0315.2017. doi:10.1158/1055-9965.EPI-17-0315
- Candido-dos-Reis FJ, Song H, Goode EL, et al; for EMBRACE; kConFab Investigators; Australian Ovarian Cancer Study Group. Germline mutation in BRCA1 or BRCA2 and ten-year survival for women diagnosed with epithelial ovarian cancer. *Clin Cancer Res*. 2015;21(3):652-657.
- Rambau PF, Duggan MA, Ghatage P, et al. Significant frequency of MSH2/MSH6 abnormality in ovarian endometrioid carcinoma supports histotype-specific Lynch syndrome screening in ovarian carcinomas. *Histopathology*. 2016;69(2):288-297.
- McMeekin DS, Trichtler DL, Cohn DE, et al. Clinicopathologic significance of mismatch repair defects in endometrial cancer: an NRG Oncology/Gynecologic Oncology Group study. *J Clin Oncol*. 2016;34(25):3062-3068.
- Kandoth C, Schultz N, Cherniack AD, et al; Cancer Genome Atlas Research Network. Integrated genomic characterization of endometrial carcinoma. *Nature*. 2013;497(7447):67-73.
- Rizvi NA, Hellmann MD, Snyder A, et al. Cancer immunology. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science*. 2015;348(6230):124-128.
- McGranahan N, Furness AJ, Rosenthal R, et al. Clonal neoantigens elicit T cell immunoreactivity

and sensitivity to immune checkpoint blockade. *Science*. 2016;351(6280):1463-1469.

39. Strickland KC, Howitt BE, Shukla SA, et al. Association and prognostic significance of BRCA1/2-mutation status with neoantigen load, number of tumor-infiltrating lymphocytes and expression of PD-1/PD-L1 in high grade serous ovarian cancer. *Oncotarget*. 2016;7(12):13587-13598.
40. Bolton KL, Chenevix-Trench G, Goh C, et al; EMBRACE; kConFab Investigators; Cancer Genome Atlas Research Network. Association between BRCA1 and BRCA2 mutations and survival in women with invasive epithelial ovarian cancer. *JAMA*. 2012;307(4):382-390.
41. Sun C, Li N, Ding D, et al. The role of BRCA status on the prognosis of patients with epithelial ovarian cancer: a systematic review of the literature with a meta-analysis. *PLoS One*. 2014;9(5):e95285.
42. Galon J, Pagès F, Marincola FM, et al. Cancer classification using the Immunoscore: a worldwide task force. *J Transl Med*. 2012;10:205.
43. Denkert C, Wienert S, Poterier A, et al. Standardized evaluation of tumor-infiltrating lymphocytes in breast cancer: results of the ring studies of the international immuno-oncology biomarker working group. *Mod Pathol*. 2016;29(10):1155-1164.
44. Hida AI, Ohi Y. Evaluation of tumor-infiltrating lymphocytes in breast cancer: proposal of a simpler method. *Ann Oncol*. 2015;26(11):2351.
45. Mesnage SJL, Auguste A, Genestie C, et al. Neoadjuvant chemotherapy (NACT) increases immune infiltration and programmed death-ligand 1 (PD-L1) expression in epithelial ovarian cancer (EOC). *Ann Oncol*. 2017;28(3):651-657.
46. Vergote I, Tropé CG, Amant F, et al; European Organization for Research and Treatment of Cancer-Gynaecological Cancer Group; NCIC Clinical Trials Group. Neoadjuvant chemotherapy or primary surgery in stage IIIC or IV ovarian cancer. *N Engl J Med*. 2010;363(10):943-953.
47. Melamed A, Hinchcliff EM, Clemmer JT, et al. Trends in the use of neoadjuvant chemotherapy for advanced ovarian cancer in the United States. *Gynecol Oncol*. 2016;143(2):236-240.
48. Verhaak RG, Tamayo P, Yang JY, et al; Cancer Genome Atlas Research Network. Prognostically relevant gene signatures of high-grade serous ovarian carcinoma. *J Clin Invest*. 2013;123(1):517-525.
49. Tothill RW, Tinker AV, George J, et al; Australian Ovarian Cancer Study Group. Novel molecular subtypes of serous and endometrioid ovarian cancer linked to clinical outcome. *Clin Cancer Res*. 2008;14(16):5198-5208.
50. Konecny GE, Wang C, Hamidi H, et al. Prognostic and therapeutic relevance of molecular subtypes in high-grade serous ovarian cancer. *J Natl Cancer Inst*. 2014;106(10):dju249.
51. Phelan CM, Kuchenbaecker KB, Tyrer JP, et al; AOCs study group; EMBRACE Study; GEMO Study Collaborators; HEBON Study; KConFab Investigators; OPAL study group. Identification of 12 new susceptibility loci for different histotypes of epithelial ovarian cancer. *Nat Genet*. 2017;49(5):680-691.
52. Cannioto RA, Trabert B, Poole EM, Schildkraut JM. Ovarian cancer epidemiology in the era of collaborative team science. *Cancer Causes Control*. 2017;28(5):487-495.
53. Kroeger DR, Milne K, Nelson BH. Tumor-infiltrating plasma cells are associated with tertiary lymphoid structures, cytolytic T-cell responses, and superior prognosis in ovarian cancer. *Clin Cancer Res*. 2016;22(12):3005-3015.
54. Loi S, Sirtaine N, Piette F, et al. Prognostic and predictive value of tumor-infiltrating lymphocytes in a phase III randomized adjuvant breast cancer trial in node-positive breast cancer comparing the addition of docetaxel to doxorubicin with doxorubicin-based chemotherapy: BIG 02-98. *J Clin Oncol*. 2013;31(7):860-867.