

## ORIGINAL ARTICLE

# Down under the tunic: bacterial biodiversity hotspots and widespread ammonia-oxidizing archaea in coral reef ascidians

Patrick M Erwin<sup>1,2</sup>, Mari Carmen Pineda<sup>1</sup>, Nicole Webster<sup>3</sup>, Xavier Turon<sup>4</sup> and Susanna López-Legentil<sup>1,2</sup>

<sup>1</sup>Departament de Biologia Animal and Institut de Recerca de la Biodiversitat (IRBio), Universitat de Barcelona (UB), Barcelona, Spain; <sup>2</sup>Center for Marine Science, University of North Carolina Wilmington, Wilmington, NC, USA; <sup>3</sup>Australian Institute of Marine Science, PMB 3, Townsville MC, Queensland, Australia and <sup>4</sup>Center for Advanced Studies of Blanes (CEAB-CSIC), Blanes, Girona, Spain

**Ascidians are ecologically important components of marine ecosystems yet the ascidian microbiota remains largely unexplored beyond a few model species. We used 16S rRNA gene tag pyrosequencing to provide a comprehensive characterization of microbial symbionts in the tunic of 42 Great Barrier Reef ascidian samples representing 25 species. Results revealed high bacterial biodiversity (3217 unique operational taxonomic units (OTU<sub>0.03</sub>) from 19 described and 14 candidate phyla) and the widespread occurrence of ammonia-oxidizing *Thaumarchaeota* in coral reef ascidians (24 of 25 host species). The ascidian microbiota was clearly differentiated from seawater microbial communities and included symbiont lineages shared with other invertebrate hosts as well as unique, ascidian-specific phylotypes. Several rare seawater microbes were markedly enriched (200–700 fold) in the ascidian tunic, suggesting that the rare biosphere of seawater may act as a conduit for horizontal symbiont transfer. However, most OTUs (71%) were rare and specific to single hosts and a significant correlation between host relatedness and symbiont community similarity was detected, indicating a high degree of host-specificity and potential role of vertical transmission in structuring these communities. We hypothesize that the complex ascidian microbiota revealed herein is maintained by the dynamic microenvironments within the ascidian tunic, offering optimal conditions for different metabolic pathways such as ample chemical substrate (ammonia-rich host waste) and physical habitat (high oxygen, low irradiance) for nitrification. Thus, ascidian hosts provide unique and fertile niches for diverse microorganisms and may represent an important and previously unrecognized habitat for nitrite/nitrate regeneration in coral reef ecosystems.**

*The ISME Journal* (2014) 8, 575–588; doi:10.1038/ismej.2013.188; published online 24 October 2013

**Subject Category:** Microbe-microbe and microbe-host interactions

**Keywords:** Sea-squirt; pyrosequencing; holobiont; thaumarchaeota; coral reef; microbiota

## Introduction

Symbiotic microbial communities are a common feature of marine invertebrates and include diverse lineages of bacteria, archaea, fungi, microalgae and viruses (Rowan, 1998; Taylor *et al.*, 2007). Prokaryotic symbionts are a particularly rich component of invertebrate microbiota and encompass nearly all major branches of bacterial and archaeal life. Many of these symbiont lineages are primarily host-associated (i.e., obligate symbionts) and represent novel microbial taxa from species level (e.g.,

*Synechococcus spongiarum* in sponges, Usher *et al.*, 2004) to phylum level, (e.g., *Poribacteria*, Fieseler *et al.*, 2004) while others exist in both free-living and host-associated states, (i.e., facultative symbionts) though generally enriched in the invertebrate microhabitat and rare in seawater communities (Sunagawa *et al.*, 2010). The phylogenetic diversity of symbiotic microbes is associated with a diversity of metabolic pathways in the carbon, (Wilkinson, 1983) nitrogen (Hoffmann *et al.*, 2009) and sulfur cycles (Hoffmann *et al.*, 2005), spurred by the utilization of host waste products (e.g., ammonia), the presence of dimethylsulfoniopropionate (DMSP, Raina *et al.*, 2010) and physico-chemical conditions of the host microenvironment (e.g., oxygen gradients; Hoffmann *et al.*, 2008; Kühl *et al.*, 2012). The structural and functional diversity of symbiotic microbial communities indicate that invertebrate hosts provide fertile microbial niches

Correspondence: S López-Legentil, Departament de Biologia Animal and Institut de Recerca de la Biodiversitat (IRBio), Universitat de Barcelona (UB), Diagonal Avenue 643, Barcelona 08028, Spain. E-mail: lopezlegentils@uncw.edu

Received 30 May 2013; revised 11 September 2013; accepted 20 September 2013; published online 24 October 2013

that contribute to prokaryotic biodiversity and nutrient cycling in coastal marine ecosystems.

Invertebrate-microbe symbioses also play critical roles in host ecological success through the provision of supplemental nutrition and production of defensive secondary metabolites. For example, sponges, corals and ascidians are able to supplement their heterotrophic filter-feeding activities with fixed carbon sourced from photosynthetic symbionts (Muscatine and Porter, 1977; Pardy and Lewin, 1981; Freeman and Thacker, 2011), utilizing autotrophic symbiont metabolism to enhance their growth rates in nutrient-limited environments. Sponge symbionts are also responsible for the synthesis of vitamin B1, which animals need to obtain from their diet (Fan *et al.*, 2012), while the cyanobacteria in the genus *Prochloron* appear to provide UV-absorbing molecules to their ascidian hosts (Hirose *et al.*, 2004). Further, symbiont biosynthesis of secondary metabolites contributes to the chemical defenses of marine invertebrates (Schmidt *et al.*, 2005; Freeman *et al.*, 2012), a key strategy for sessile organisms to deter predation, avoid surface fouling and compete for substrate (Armstrong *et al.*, 2001; Pawlik, 2011). In addition to their roles in host biology and ecology, many of these unique and structurally diverse secondary metabolites have pharmaceutical applications and substantial importance for biotechnology and drug discovery (Paul and Ritson-Williams, 2008; Erwin *et al.*, 2010).

Ascidians (Class Ascidiacea) are sessile, filter-feeding invertebrates that inhabit diverse benthic ecosystems in tropical, temperate and polar marine environments. As a basal lineage in the phylum Chordata, ascidians occupy a key stage in deuterostome evolution (Delsuc *et al.*, 2006). Ascidians are also a prolific source of novel marine natural products (Erwin *et al.*, 2010) and the involvement of microbial symbionts in bioactive compound production (Schmidt and Donia, 2010) has prompted recent studies of the ascidian microbiota (Donia *et al.*, 2011; Kwan *et al.*, 2012). Historically, most studies of microbial symbionts in ascidians have focused on cyanobacteria, in particular the genera *Prochloron* and *Synechocystis*. These symbionts associate with colonial ascidians on the colony surface, inside the common cloacal cavities or as endosymbionts in the tunic, a polysaccharide envelope surrounding the zooids (Cox *et al.*, 1985; Cox, 1986; Hernández-Mariné *et al.*, 1990; Hirose *et al.*, 1996, 2006a, b, 2012; Turon *et al.*, 2005; Martínez-García *et al.*, 2007). Even when inhabiting the colonial tunic, the symbionts are mostly extracellular, with only a few instances of intracellular associations (Hirose *et al.*, 1996; Moss *et al.*, 2003; Kojima and Hirose, 2010). However, few studies to date have employed the molecular approaches required to accurately assess microbial biodiversity in ascidians (Martínez-García *et al.*, 2007, 2008, 2011; Münchhoff *et al.*, 2007;

Tait *et al.*, 2007; López-Legentil *et al.*, 2011; Behrendt *et al.*, 2012; Erwin *et al.*, 2013). For example, DNA sequence analysis and fluorescence *in situ* hybridization techniques only recently revealed the first archaeal symbionts in the ascidian tunic, indicating that *Thaumarchaeota* may be involved in nitrification inside host tissues (Martínez-García *et al.*, 2008).

A growing body of literature suggests that ascidian-associated microbes may play a critical role in the metabolic needs of their host, (Hirose and Maruyama, 2004; Martínez-García *et al.*, 2008; Kühl *et al.*, 2012), yet the microbial communities inhabiting most ascidian species remain unknown. The advent of high-throughput, next-generation DNA sequencing platforms offers new opportunities for in-depth microbial diversity evaluation across large sample sets. Deep sequencing of microbial communities from soils, seawater and sponges has revealed diversity estimates over an order of magnitude higher than that recovered by traditional sequencing techniques (Huber *et al.*, 2007; Roesch *et al.*, 2007; Webster *et al.*, 2010), including the detection of bacterial phyla not represented in first-generation sequencing datasets (e.g., Webster and Taylor, 2012). Similarly, the recent application of next generation sequencing to the ascidian microbiota has revealed a high diversity of symbiotic microbes and uncovered new ascidian-associated microbial lineages in the colonial host *Lissoclinum patella* (Behrendt *et al.*, 2012) and solitary host *Styela plicata* (Erwin *et al.*, 2013), highlighting the depth of microbial biodiversity and unknown facultative and obligate symbiotic microbes awaiting discovery within ascidian hosts.

In this study, we used 16S rRNA gene tag pyrosequencing to investigate the diversity, structure and specificity of microbial communities inhabiting the tunic of 42 samples of Great Barrier Reef (GBR) ascidians (representing 25 species, 7 families and 3 orders) in order to provide the most comprehensive characterization of the ascidian microbiome to date. The diversity and composition of ascidian-associated microbial communities were compared to free-living communities in ambient seawater and among ascidian host species, including intraspecific variability among replicates for 10 ascidian species. In addition, the spatial localization of symbionts within the ascidian tunic was visualized by electron microscopy, and the genetic identity of ascidian hosts was established by analysis of mitochondrial (cytochrome oxidase subunit I) and ribosomal (18S rRNA) gene sequences. This comprehensive assessment of microbial diversity in GBR ascidians will provide the basis for future research within the fields of symbiosis, drug discovery and ascidian holobiont resilience to environmental change or anthropogenic disturbance. Exploration of ascidian microbiomes may also highlight a hidden reservoir for primary productivity and nitrogen metabolism and enable

more reliable predictions of biogeochemical cycling in coral reef environments.

## Material and methods

### Sample collection

Ascidian ( $n = 42$ ) and seawater ( $n = 3$ ) samples were collected by SCUBA between 2–14 m depth from several localities within the Great Barrier Reef, North Queensland, Australia (Supplementary Table S1). Ascidian samples were processed for: (1) taxonomic analyses, by preservation in 4% formaldehyde, (2) molecular analyses, by immediate submersion in liquid nitrogen and storage at  $-80^{\circ}\text{C}$  and (3) electron microscopy analyses, by preservation in 2.5% glutaraldehyde using filtered seawater as buffer. Seawater samples (2l) were transported to the laboratory, concentrated on 0.2  $\mu\text{m}$  sterivex filters (Durapore; Millipore, North Ryde, New South Wales, Australia) with a peristaltic pump and aseptically frozen at  $-80^{\circ}\text{C}$ .

### DNA extraction

Frozen ascidian tissues (approximately 0.5 g per sample) were thawed, dissected under a stereomicroscope into inner tunic and zooid fractions and aseptically transferred to 1.5 ml tubes using sterile scalpels and tweezers. Inner tunic (i.e., beneath the surface layer) was chosen to avoid epibionts and ambient seawater microbes. These tunic samples were processed for microbial analysis, while zooids were processed for barcoding each ascidian specimen. DNA extraction was conducted separately for inner tunic and zooid tissue fractions with the Power Plant DNA Isolation Kit (MoBio Laboratories, Carlsbad, CA, USA) following the manufacturer's protocol. DNA extraction from concentrated seawater samples (filters) was performed by the addition of 1.8 ml lysis buffer (40 mM EDTA, 50 mM Tris and 0.75 M sucrose) and 200  $\mu\text{l}$  of Lysozyme (10 mg/ml), incubation at  $37^{\circ}\text{C}$  for 45 min, the addition of 40  $\mu\text{l}$  of Proteinase K (10  $\mu\text{g}$  of Proteinase K in 1 ml of 10% SDS) and incubation at  $55^{\circ}\text{C}$  for 1 h. Lysates were transferred to sterile tubes and DNA was extracted using standard phenol:chloroform procedures and resuspended in 20  $\mu\text{l}$  of distilled water. All PCR products were visualized on 1% agarose gels to assess amplification specificity and initial product quantity.

### Identification and barcoding of host ascidians

Ascidian samples were assigned to the lowest taxonomic group possible based on morphological examination (Supplementary Text S1). Genetic identification was also performed using the mitochondrial gene cytochrome oxidase subunit I (COI) and 18S rRNA gene sequences. Both gene regions are commonly used to determine species boundaries and diversity among ascidian taxa (Tarjuelo *et al.*,

2004; López-Legentil and Turon, 2005; Yokobori *et al.*, 2006; Pérez-Portela *et al.*, 2009) and COI is the metazoan standard for the Barcode of Life Project ([www.barcodeoflife.org](http://www.barcodeoflife.org)).

DNA extractions from zooid tissue were used as templates for PCR amplification of a 519 – 621 bp fragment of the COI gene. Total PCR reaction volume was 50  $\mu\text{l}$ , including 10  $\mu\text{l}$  of  $5 \times$  Buffer, 0.4  $\mu\text{l}$  of bovine serum albumin (BSA; 10 mg/ml), 0.25  $\mu\text{l}$  of My Taq DNA Polymerase (Bioline, London, United Kingdom), 2  $\mu\text{l}$  of each primer (10  $\mu\text{M}$ ) and 1  $\mu\text{l}$  of template DNA. Two sets of primer pairs were used for COI amplification, the 'universal' primers LCO1490 and HCO2198 (Folmer *et al.*, 1994) and the ascidian-specific primers Tun\_forward and Tun\_Reverse2 (Stefaniak *et al.*, 2009). PCR conditions for amplification with universal primers were: an initial denaturing step of  $94^{\circ}\text{C}$  for 2 min; 30 cycles of  $94^{\circ}\text{C}$  for 45 s,  $50^{\circ}\text{C}$  for 45 s and  $72^{\circ}\text{C}$  for 50 s; and a final elongation step at  $72^{\circ}\text{C}$  for 5 min. PCR conditions for amplification with ascidian-specific primers were: an initial denaturing step of  $94^{\circ}\text{C}$  for 1 min; 60 cycles of  $94^{\circ}\text{C}$  for 10 s,  $50^{\circ}\text{C}$  for 30 s and  $72^{\circ}\text{C}$  for 50 s; and a final elongation step at  $72^{\circ}\text{C}$  for 10 min. PCR products were purified and bi-directionally sequenced at Macrogen, Inc. (Seoul, South Korea). Quality-checked sequences are archived in GenBank under accession numbers KC017426 to KC017444. Additional genetic identification and phylogenetic analyses of host ascidians were performed with 18S rRNA gene sequences recovered from the non-target, eukaryotic data component of the pyrosequencing run (Supplementary Text S2, Figure S4).

### 16S rRNA gene tag pyrosequencing

DNA extractions from inner tunic tissue and seawater samples were used as templates for PCR amplification of a ca. 466 bp fragment of the 16S rRNA gene encompassing the V6 – V8 regions using the primer set pyro926F (5'-AAACTYAAAKGAA TTGRCGG-3') and pyro1392R (5'-ACGGGCGGTG TGTRC-3') complemented with adaptors B and A, respectively (Roche, Basel, Switzerland), as detailed previously (Erwin *et al.*, 2013). Multiplex identifier (MID) barcodes unique to each sample were attached to reverse primers (Supplementary Table S2). PCR products were sent to Macrogen, Inc. for purification and further processing. Amplicon library was constructed using 5  $\mu\text{g}$  of DNA from each sample (ascidian and seawater), resulting in a final concentration of 700 513 297 molecules/ $\mu\text{l}$ . Massively parallel 16S rRNA gene tag pyrosequencing was performed using the Roche 454 GS-FLX Titanium system, and the resulting data were deposited as flowgrams (sff file) in the Sequence Read Archive (SRA) of the National Center for Biotechnology Information under the accession number SRA056317.

Sequence data were processed with stringent filtering and screening criteria to minimize the occurrence of spurious sequences and overestimation of microbial diversity (Huse *et al.*, 2010; Schloss *et al.*, 2011), using the mothur software package (Schloss *et al.*, 2009), as detailed previously (Erwin *et al.*, 2013). Briefly, adaptor, MID and primer sequences were removed from raw sequences and the dataset de-noised (removal of reads with ambiguous base calls, long homopolymers and barcode or primer mismatches) and quality filtered (removal of short sequences and low quality reads). Non-target sequences (e.g. eukaryotic 18S rRNA, mitochondria, chloroplast) were removed using Metaxa v1.1, (Bengtsson *et al.*, 2011) resulting in a dataset consisting solely of archaeal and bacterial 16S rRNA gene sequences. These sequences were aligned to the Greengenes database, trimmed to an overlapping alignment space (449 bp) and putatively chimeric sequences were removed (UCHime; Edgar *et al.*, 2011).

#### Data analysis

High quality sequences ( $n = 94\,637$ ) were assigned to taxonomic groups based on the improved Greengenes taxonomy template (McDonald *et al.*, 2012) with *Thaumarchaeota* elevated to the rank of phylum (Brochier-Armanet *et al.*, 2008, Spang *et al.*, 2010), grouped into OTU<sub>0.03</sub> based on 97% sequence similarity and the average neighbor clustering algorithm, and the taxonomic assignment of each OTU<sub>0.03</sub> was constructed by majority consensus (Schloss and Westcott, 2011).

Sampling coverage and expected total OTU diversity were calculated using rarefaction analysis and the bootstrap estimator (Smith and Van Belle, 1984) at six different OTU definitions corresponding approximately to the species (OTU<sub>0.03</sub>), genus (OTU<sub>0.05</sub>), family (OTU<sub>0.10</sub>), order (OTU<sub>0.15</sub>), class (OTU<sub>0.20</sub>) and phylum (OTU<sub>0.25</sub>) levels (97%, 95%, 90%, 85%, 80% and 75% similarity, respectively). All subsequent analyses were based on OTUs at 97% sequence identity (OTU<sub>0.03</sub>). Sub-sampling of sequence pools from samples with greater than 2000 reads were performed in the mothur software package to standardize sampling effort and determine its effect on diversity estimates. Host-specificity of the ascidian microbiota was assessed by partitioning OTUs into core (present in >70% of host species), variable (present in at least two host species) and specific (present in a single host species) groups (*sensu* Schmitt *et al.*, 2012). To broaden the analysis of the specificity of the ascidian microbiota, abundant ascidian-associated OTUs (i.e., those represented by >100 total sequence reads) were compared to sequences in the GenBank database using a nucleotide-nucleotide BLAST search (Altschul *et al.*, 1990). To compare microbial community similarity across hosts, Bray-Curtis similarity matrices were constructed using

square root transformations of relative OTU abundance per host and visualized in cluster plots using Primer v6 (Plymouth Marine Laboratory, United Kingdom). Finally, Mantel tests were conducted to test for correlations between host relatedness (18S rRNA sequence similarity) and symbiont similarity (Bray-Curtis similarity) using the ade4 package for R (Dray and Dufour, 2007).

#### Transmission electron microscopy

Bacterial cells in the tunic of the representative ascidian species *Phallusia julinea*, *Polycarpa aurata*, *Pycnoclavella* sp., *Clavelina meridionalis*, *Lissoclinum badium* and *Synoicum castellatum* were visualized by transmission electron microscopy. Resin blocks and semi-thin and ultra-thin sections were prepared at the Microscopy Unit of the Scientific and Technical Services of the University of Barcelona as described in López-Legentil *et al.* (2011). Transmission electron microscopy observations were conducted on a JEOL JEM-1010 (Tokyo, Japan) electron microscope coupled with an Orius CDD camera (Gatan, Germany).


## Results

#### Diversity and phylogeny of ascidian hosts

The 42 host ascidians examined for microbial symbionts were classified in 25 species from 7 families and all 3 recognized orders in the class Ascidiacea, with 18 species belonging to the Aplousobranchia, the largest ascidian order in terms of species and family richness (Shenkar and Swalla, 2011). Analyses of 18S rRNA gene sequences (23 of the 25 host species) and COI sequences (19 of 25 host species) confirmed morphological identifications and provide molecular datasets to facilitate additional research on the ascidian microbiota. All reference works used to identify each specimen and pertinent taxonomic remarks are provided (Supplementary Text S1), including underwater images (Supplementary Figures S1, S2 and S3) and a phylogenetic analysis using 18S rRNA sequences (Supplementary Text S2, Figure S4).

#### Richness and diversity of the ascidian microbiota

Collective analysis of 16S rRNA sequence reads derived from ascidian hosts ( $n = 67\,826$ ) revealed a remarkable richness and diversity of microbial communities associated with GBR ascidians. A total of 3321 unique microbial OTU<sub>0.03</sub> represented the combined GBR ascidian microbiome and corresponded to 19 described bacterial phyla, 14 candidate bacterial phyla and 3 described archaeal phyla (Figure 1). This increases the taxonomic diversity known to inhabit ascidians by 14 microbial phyla. Coverage estimates of total diversity sampled were high across all taxonomic levels, ranging from 82 (OTU<sub>0.03</sub>) to 85% (OTU<sub>0.25</sub>). Rarefaction analysis



**Figure 1** Taxonomic diversity of the ascidian microbiota. (a) Phylum level distribution of the 3321 microbial OTU<sub>0.03</sub> recovered from 42 GBR ascidian hosts, depicting common phyla (in color, >1% OTU<sub>0.03</sub> diversity), rare phyla (in gray, <1%; SBR1093, *Lentisphaerae*, *Chlamydiae*, *Tenericutes*, TM7, WS3, *Spirochaetes*, *Nitrospirae*, OP3, TM6, *Crenarchaeota*, *Chlorobi*, OP11, *Thermi*, *Armatimonadetes*, *Fusobacteria*, NKB19, *Caldithrix*, OP8, PAUC34f, BRC1, *Elusimicrobia*, GN04, KSB1 and SM2F11) and bacterial OTUs unclassified at the phylum level (in black). (b) Class level distribution of proteobacterial OTUs.

revealed that observed OTU diversity was approaching expected OTU diversity at higher level taxonomic rankings (e.g., phylum and class; Supplementary Figure S5) while additional sampling would continue to uncover new microbial OTUs at lower taxonomic levels (e.g., genus and species; Supplementary Figure S6) due to a rich rare component of the microbiota (1817 singletons).

Analyses of individual hosts and ascidian species revealed up to 486 microbial OTU<sub>0.03</sub> per individual and 697 unique OTU<sub>0.03</sub> per species (Tables 1 and 2), with many ascidians hosting more diverse microbial communities than those recovered from ambient seawater in terms of observed and expected (Chao1) OTU richness and common diversity indices (Shannon, Simpson Inverse; Supplementary Table S3). 16S rRNA sequence reads derived from seawater ( $n = 26\,811$ ) grouped into 385 unique OTU<sub>0.03</sub> (129–284 per replicate). While high variability in sampling effort (sequence reads per sample) can obscure direct comparisons among host species and between ascidians and seawater, over 25% ( $n = 11$ ) of the sampled ascidians exhibited higher microbial OTU<sub>0.03</sub> diversity than the most well-sampled seawater replicate, despite lower sampling effort (7 500–13 500 fewer sequence reads; Table 1). Further, this trend was maintained after sub-sampling of sequence pools to standardize sampling efforts across ascidian and seawater sources (Supplementary Figure S6).

#### Composition of the ascidian microbiota

Microbial communities in GBR ascidians were composed of diverse bacterial phyla and archaeal lineages (Figure 1, Supplementary Table S4). Bacterial OTUs dominated the ascidian microbiota, accounting for 97% ( $n = 3217$ ) of OTU<sub>0.03</sub> diversity and 82% of all sequence reads ( $n = 55\,698$ ). The most dominant bacterial phylum was

*Proteobacteria*, representing over one-third (38%) of OTU<sub>0.03</sub> diversity ( $n = 1251$ ) and the only phylum detected in all examined ascidians. *Proteobacteria* accounted for over half of all sequence reads in 12 ascidian individuals and over 90% of sequences from *Aplidium protectans*, *Lissoclinum* cf. *capsulatum* and *Didemnum granulatum* (Figure 2). Within the *Proteobacteria*, the classes *Alphaproteobacteria* and *Gammaproteobacteria* were most prevalent (517 OTUs and 397 OTUs, respectively), followed by *Deltaproteobacteria* and *Betaproteobacteria* (125 OTUs and 6 OTUs, respectively). Representatives from the phyla *Bacteroidetes* and *Planctomycetes* were also common, each accounting for over 15% of OTU<sub>0.03</sub> diversity ( $n = 496$  and 486, respectively, Figure 1) and detected in the majority (>88%) of ascidian hosts (Figure 2, Supplementary Table S4).

*Cyanobacteria* was the fourth most diverse phyla associated with ascidians (172 OTUs, 5% of OTU<sub>0.03</sub> diversity) and included the genus *Prochloron*, present only in *Lissoclinum patella* (OTU0810), and 4 OTUs that were closely related (95–98% sequence identity) to the recently described *Candidatus* ‘*Acaryochloris bahamiensis*’ (López-Legentil et al., 2011). Most notably, two *Acaryochloris* OTUs (OTU0125, 0126) were common in all 3 individuals of the host *Eudistoma amplum* (0.7 to 8.9% relative abundance). An additional 5 described phyla were common in ascidians, including *Chloroflexi* (103 OTU<sub>0.03</sub>), *Acidobacteria* (87), *Actinobacteria* (62), *Verrucomicrobia* (51) and *Firmicutes* (45), each accounting for 1 to 3% of OTU<sub>0.03</sub> diversity and detected in at least half of the ascidian hosts examined. The remaining 24 described and candidate phyla present in the ascidian microbiota were rare overall (each <1% of total OTU<sub>0.03</sub> diversity) and within each host ascidian (<2% of sequence reads; Figure 2, Supplementary Table S4), with the exception of *Spirochaetes* in *Polycarpa aurata* (18% relative abundance) and SBR1093 in *Eudistoma amplum* (11%).

Archaeal OTUs accounted for 18% ( $n = 12\,128$ ) of sequence reads but only 3% ( $n = 104$ ) of the OTU<sub>0.03</sub> richness in the ascidian microbiota. *Thaumarchaeota* were particularly abundant ( $n = 11\,993$ ; 53 OTUs) and common (present in 93% of host individuals), with most archaeal sequence reads (98%) matching to the ammonia-oxidizing genera *Nitrosopumilus* ( $n = 11\,630$ ; 36 OTUs) and *Cenarchaeum* ( $n = 261$ ; 5 OTUs). In fact, the most common OTU<sub>0.03</sub> in the ascidian microbiota (OTU0001, *Nitrosopumilus* sp.) was present in 37 of the 42 host individuals (22 of 25 host species) at relative abundances up to 95% (*Lissoclinum badium*), while extremely rare in ambient seawater (0–0.04%). In addition, a common archaeal symbiont in *Leptoclinides madara* (OTU0025, 17–28% relative abundance) was classified to the genus *Cenarchaeum* and closely matched (98% sequence identity) an uncultivated archaeon reported in the marine

**Table 1** Taxonomic classification of ascidian hosts and sequence data summary for ascidian and seawater samples. Total values in bold refer to summed reads and unique OTUs

Species	Order	Family	Total		Archaea		Bacteria	
			Reads	OTU <sub>0.03</sub>	Reads	OTU <sub>0.03</sub>	Reads	OTU <sub>0.03</sub>
<i>Clavelina arafurensis</i>	Aplousobranchia	Clavelinidae	490	190	57	4	433	186
<i>Clavelina meridionalis</i>			249	103	15	8	234	95
<i>Clavelina meridionalis</i>			1207	333	44	10	1163	323
<i>Clavelina meridionalis</i>			1023	411	38	11	985	400
<i>Pycnoclavella</i> sp.			1449	313	93	11	1356	302
<i>Pycnoclavella</i> sp.			116	47	3	3	113	44
<i>Pycnoclavella diminuta</i>			2040	384	294	9	1746	375
<i>Pycnoclavella diminuta</i>			1188	301	434	9	754	292
<i>Pycnoclavella diminuta</i>			347	167	66	6	281	161
<i>Didemnum</i> cf. <i>albopunctatum</i>		Didemnidae	3654	154	906	12	2748	142
<i>Didemnum</i> cf. <i>granulatum</i>			386	22	11	4	375	18
<i>Didemnum multispirale</i>			3035	102	10	2	3025	100
<i>Didemnum multispirale</i>			2799	142	21	3	2778	139
<i>Didemnum multispirale</i>			2979	209	25	6	2954	203
<i>Didemnum</i> sp.1			6905	486	255	6	6650	480
<i>Didemnum</i> sp.2			2684	448	762	12	1922	436
<i>Leptoclinides madara</i>			979	74	165	2	814	72
<i>Leptoclinides madara</i>			281	18	79	2	202	16
<i>Lissoclinum badium</i>			3224	27	3055	4	169	23
<i>Lissoclinum badium</i>			4670	29	4464	4	206	25
<i>Lissoclinum</i> cf. <i>capsulatum</i>			598	36	2	1	596	35
<i>Lissoclinum patella</i>			2489	86	1	1	2488	85
<i>Eudistoma amplum</i>		Polycitoridae	517	164	177	16	340	148
<i>Eudistoma amplum</i>			444	175	89	13	355	162
<i>Eudistoma amplum</i>			825	286	112	11	713	275
<i>Polycitor giganteus</i>			1602	95	6	3	1596	92
<i>Aplidium protectans</i>		Polyclinidae	4272	129	30	3	4242	126
<i>Aplidium</i> sp.			1968	176	64	7	1904	169
<i>Synoicum castellatum</i>			3846	382	4	2	3842	380
<i>Synoicum castellatum</i>			6447	344	60	3	6387	341
<i>Synoicum castellatum</i>			120	46	27	4	93	42
<i>Phallusia arabica</i>	Phlebobranchia	Asciidiidae	105	23	2	1	103	22
<i>Phallusia arabica</i>			338	53	39	8	299	45
<i>Phallusia arabica</i>			54	17	2	2	52	15
<i>Phallusia julinea</i>			562	97	55	4	507	93
<i>Phallusia philippinensis</i>			28	8	12	1	16	7
<i>Ecteinascidia diaphanis</i>		Perophoridae	1168	344	17	4	1151	340
<i>Perophora</i> aff. <i>modificata</i>			1541	189	184	9	1357	180
<i>Polycarpa argentata</i>	Stolidobranchia	Styelidae	561	68	446	7	115	61
<i>Polycarpa aurata</i>			449	18	0	0	449	18
<i>Polycarpa aurata</i>			159	23	2	2	157	21
<i>Polycarpa aurata</i>			28	8	0	0	28	8
<b>Ascidian Microbiota Total =</b>			<b>67 826</b>	<b>3321</b>	<b>12 128</b>	<b>104</b>	<b>55 698</b>	<b>3217</b>
Filtered Seawater	n.a.	n.a.	9 573	221	289	24	9 284	197
Filtered Seawater	n.a.	n.a.	14 441	248	134	21	14 307	227
Filtered Seawater	n.a.	n.a.	2 797	129	3	3	2 794	126
<b>Ambient Seawater Total =</b>			<b>26 811</b>	<b>385</b>	<b>426</b>	<b>26</b>	<b>26 385</b>	<b>359</b>
<b>Grand Total =</b>			<b>94 637</b>	<b>3604</b>	<b>12 554</b>	<b>124</b>	<b>82 083</b>	<b>3480</b>

**Table 2** Intra-specific variation in the ascidian microbiota highlighting the shared components (i.e., present in all host individuals) of each species' microbiota

Species	Species Cluster	No. Samples	Total Sequences	Total OTU <sub>0.03</sub>	Shared Sequences (%)	Shared OTU <sub>0.03</sub> (%)
<i>Clavelina meridionalis</i>	Y	3	2479	697	1338 (54)	26 (4)
<i>Pycnoclavella</i> sp.	N	2	1565	341	1077 (69)	19 (6)
<i>Pycnoclavella diminuta</i>	N	3	3575	673	1731 (48)	35 (5)
<i>Didemnum multispirale</i>	Y	3	8813	367	6192 (70)	24 (6)
<i>Leptoclinides madara</i>	Y	2	1260	81	1116 (89)	11 (14)
<i>Lissoclinum badium</i>	Y	2	7894	41	7848 (99)	15 (37)
<i>Eudistoma amplum</i>	Y	3	1786	491	809 (45)	31 (6)
<i>Synoicum castellatum</i>	N	3	10 413	620	5237 (50)	17 (3)
<i>Phallusia arabica</i>	N	3	497	82	104 (21)	2 (2)
<i>Polycarpa aurata</i>	N	3	636	39	514 (81)	3 (8)


Species cluster refers to Figure 2.

sponge *Axinella verrucosa* (GenBank accession number AF420237).


#### Specificity of the ascidian microbiota

Comparison of the rich ascidian microbiota with ambient seawater microbes revealed low overlap

between free-living and host-associated microbial communities. A total of 283 OTUs were present in the seawater communities and absent from the ascidian microbiota, while 102 OTUs were present in both ascidian and seawater samples, representing only 3% of total OTU<sub>0.03</sub> diversity in the ascidian



**Figure 2** Microbial community similarity and composition in 42 samples of GBR ascidians. Dendrogram (*left*) based on Bray-Curtis (BC) similarity of microbial communities in ascidian hosts. Ordinal classifications of ascidians hosts are shown as circles, Aplousobranchia (*white*), Phlebobranchia (*gray*) and Stolidobranchia (*black*) and zooid organization as triangles, colonial (*white*) and solitary (*black*). Bar charts (*right*) show the relative abundance of microbial phyla in each host ascidian, with host species names listed on the right. **Bold** names indicate species with replicate samples.



**Figure 3** Relative abundance of seawater microbes in the ascidian microbiota. (a) Rank-abundance plots showing the relative abundance of 102 microbial OTUs present in both seawater (*black line*) and ascidian hosts (*gray bars*). Asterisks denote OTUs > 200 times more abundant in ascidian hosts than seawater. (b) Classification and relative abundance of 5 rare seawater biosphere OTUs among ascidian hosts.

microbiota. Further, over one-third ( $n = 40$ ) of these shared microbial OTUs exhibited greater than an order of magnitude difference in relative abundance in seawater and ascidians assemblages, including 5

OTUs that were 200x to 700x more abundant in host ascidians (Figure 3). For example, OTU0001 (*Nitrosopumilus* sp.) accounted for 17% of sequence reads from the ascidian microbiota. The remaining 4

OTUs were specific to particular host families (e.g., OTU0301 in Didemnidae), species (e.g., OTU1798 in 3 individuals of *Clavelina meridionalis*) or individuals (e.g., OTU0225 in 1 of 3 *Didemnum multi-spirale* individuals) and rare or absent in most ascidian hosts (Figure 3).

Additional analysis of abundant components of the ascidian microbiota revealed symbiont overlap between ascidians and other invertebrate hosts, as well as a unique component of the ascidian microbiota (Table 3). A total of 56 microbial OTUs accounted for 78% of sequences obtained from ascidian hosts. Over two-thirds of these OTUs ( $n = 38$ ) matched closely ( $>97\%$  sequence identity)

to previously characterized sequences (Table 3), most commonly derived from seawater ( $n = 14$ ), corals ( $n = 9$ ), sponges ( $n = 6$ ) and sediment ( $n = 3$ ). In some cases, OTUs that were widespread among ascidians hosts and in the rare biosphere of seawater matched closely to other invertebrate-associated sequences. For example, OTU0264 (*Bacteroidetes*, *Flavobacteriaceae*) was present in 24 ascidian individuals, was rare in seawater ( $<0.05\%$  relative abundance) and matched identically to coral-derived sequences from Caribbean (*Montastraea faveolata*) and Indo-Pacific (*Montipora aequituberculata*) stony corals and an Indo-Pacific soft coral (*Sinularia* sp.). The remaining 18 OTUs exhibited

**Table 3** Abundant OTUs in the ascidian microbiota, showing their representation in ascidian (ASC) and seawater (SW) datasets, number of host species, closest known relative and taxonomic classification

OTU	Reads (ASC)	Hosts (ASC)	Reads (SW)	BLAST Match Source (Identity, Acc. No.)	Phylum	Lowest Taxonomic Rank
0001	11338	39	9	Sponge (98.3, AF420237)	Thaumarchaeota	<i>G. Nitrosopumilus</i>
0140	9981	42	11105	Seawater (100, GU119217)	Cyanobacteria	<i>G. Prochlorococcus</i>
0287	4964	11	0	Bivalve (92.9, EU857739)	Unclassified	<i>K. Bacteria</i>
0364	3669	13	13	Sponge (100, HQ241801)	$\gamma$ -proteobacteria	<i>G. Coxiella</i>
0188	2836	30	33	Seawater (100, HQ338142)	$\alpha$ -proteobacteria	<i>F. Rhodobacteraceae</i>
0292	1836	34	30	Seawater (100, GU119442)	Cyanobacteria	<i>G. Prochlorococcus</i>
0189	1790	28	13	Seawater (100, JF514245)	$\alpha$ -proteobacteria	<i>G. Mesorhizobium</i>
0225	1633	25	1	Seawater (100, JF769651)	$\alpha$ -proteobacteria	<i>O. Rhizobiales</i>
0301	1432	10	1	Ascidian (100, DQ860066)	$\alpha$ -proteobacteria	<i>O. Rhizobiales</i>
1128	1346	3	0	Seafloor Lava (93.2, EU491218)	Proteobacteria	<i>P. Proteobacteria</i>
1129	985	2	0	Sediment (88.7, GU046335)	Unclassified	<i>K. Bacteria</i>
0851	858	5	0	Sponge (94.1, EU883386)	$\alpha$ -proteobacteria	<i>O. Rhodospirillales</i>
0310	779	32	10685	Seawater (100, JN547429)	Cyanobacteria	<i>G. Prochlorococcus</i>
1063	671	3	0	Soil (97.9, JQ059148)	$\alpha$ -proteobacteria	<i>F. Rhodospirillaceae</i>
1798	567	5	1	Seawater (95.8, HQ715140)	$\alpha$ -proteobacteria	<i>O. Rhizobiales</i>
1379	379	5	0	Soil (90.4, GQ127925)	Unclassified	<i>K. Bacteria</i>
3180	354	1	0	Sediment (95.4, AB374687)	Bacteroidetes	<i>F. Flammeovirgaceae</i>
1101	336	16	118	Seawater (100, AB540006)	Bacteroidetes	<i>F. Flavobacteriaceae</i>
0355	333	25	0	Coral (100, FJ809316)	SBR1093	<i>C. VHS-B5-50</i>
0862	329	16	0	Sponge (100, EU335078)	Chloroflexi	<i>C. Anaerolineae</i>
0931	327	13	0	Algae (96.7, HM474939)	Chloroflexi	<i>C. Anaerolineae</i>
0164	326	30	3	Seawater (100, GU119490)	Planctomycetes	<i>O. Pirellulales</i>
1032	326	3	0	Sediment (96.6, JQ989595)	$\alpha$ -proteobacteria	<i>O. Rhizobiales</i>
0866	324	22	0	Coral (100, DQ416621)	Bacteroidetes	<i>F. Flavobacteriaceae</i>
0293	261	11	0	Sponge (100, FJ625530)	Planctomycetes	<i>O. Pirellulales</i>
2687	260	2	0	Biofilm (94.6, FJ901434)	Cyanobacteria	<i>F. Phormidiaceae</i>
0296	246	2	0	Sediment (96.7, JN977252)	$\gamma$ -proteobacteria	<i>C. <math>\gamma</math>-proteobacteria</i>
0273	245	15	0	Coral (99.6, JQ347330)	Cyanobacteria	<i>F. Pseudanabaenaceae</i>
0025	241	2	0	Sponge (98.3, AF420237)	Thaumarchaeota	<i>G. Cenarchaeum</i>
0875	211	3	0	Coral (97.1, FJ425620)	Bacteroidetes	<i>F. Flammeovirgaceae</i>
0003	208	19	0	Cyanobacteria (100, JX197041)	Thaumarchaeota	<i>G. Nitrosopumilus</i>
0335	206	26	59	Seawater (100, EU592360)	$\alpha$ -proteobacteria	<i>F. Rhodobacteraceae</i>
0300	202	4	0	Sponge (100, JN128259)	$\gamma$ -proteobacteria	<i>G. Microbulbifer</i>
0344	198	9	0	Sponge (100, DQ097259)	$\alpha$ -proteobacteria	<i>G. Pseudovibrio</i>
2656	193	3	0	Diatom Bloom (94.4, EU734047)	$\beta$ -proteobacteria	<i>C. <math>\beta</math>-proteobacteria</i>
0318	186	20	0	Coral (100, FJ489710)	SBR1093	<i>C. EC214</i>
2229	183	3	0	Seawater (98.3, HM798908)	$\alpha$ -proteobacteria	<i>F. Rhodospirillaceae</i>
0187	179	17	2	Seawater (100, HM103531)	$\alpha$ -proteobacteria	<i>F. Rhodobacteraceae</i>
0161	165	19	0	Sediment (100, GQ249478)	$\gamma$ -proteobacteria	<i>F. Chromatiaceae</i>
0306	157	16	0	Coral (100, FJ203575)	$\alpha$ -proteobacteria	<i>F. Hyphomicrobiaceae</i>
0850	153	3	0	Biofilm (98.7, DQ167245)	$\alpha$ -proteobacteria	<i>G. Kiloniella</i>
2389	152	3	0	Coral (95.8, EF206859)	$\gamma$ -proteobacteria	<i>C. <math>\gamma</math>-proteobacteria</i>
0133	147	20	0	Coral (100, GU118991)	Bacteroidetes	<i>F. Flammeovirgaceae</i>
0264	147	24	13	Coral (100, FJ809398)	Bacteroidetes	<i>F. Flavobacteriaceae</i>
0294	145	3	0	Algae (99.6, GU451475)	$\alpha$ -proteobacteria	<i>G. Pseudovibrio</i>
1065	143	4	0	Coral (93.2, GU118840)	$\alpha$ -proteobacteria	<i>O. Rhodospirillales</i>
0186	138	17	0	Sediment (99.6, FJ358900)	Bacteroidetes	<i>F. Flammeovirgaceae</i>
0307	137	13	0	Algae (99.6, HM474882)	$\alpha$ -proteobacteria	<i>F. Rhodospirillaceae</i>
2811	137	2	0	Seawater (94.5, EF572701)	Bacteroidetes	<i>F. Flavobacteriaceae</i>
0297	132	12	0	Coral (99.6, FJ203345)	Planctomycetes	<i>O. Pirellulales</i>
0939	130	13	0	Sediment (100, DQ256661)	Cyanobacteria	<i>G. Leptolyngbya</i>
2749	124	1	0	Sediment (96.2, EU287328)	$\alpha$ -proteobacteria	<i>O. Rhizobiales</i>
0686	121	7	0	Bivalve (92.5, EU857738)	Unclassified	<i>K. Bacteria</i>
2875	117	1	0	Seawater (98.3, JN216763)	$\alpha$ -proteobacteria	<i>C. <math>\alpha</math>-proteobacteria</i>
1132	107	1	0	Mammal Gut (89.2, EU459272)	Unclassified	<i>K. Bacteria</i>
0172	101	4	0	Seawater (99.2, GQ349494)	$\delta$ -proteobacteria	<i>G. Nitrospina</i>



greater divergence from both free-living and host-associated microbes, including 11 OTUs that exhibited <95% sequence identity to known microbial sequences (Table 3).

#### Core, variable and specific microbial OTUs

Comparison of the microbial communities among ascidian hosts revealed a high degree of host specificity in the ascidian microbiota and the presence of a small number of very abundant and widespread microbial OTUs. No universal symbiont OTUs (i.e., present in all host species) were detected and core OTUs (present in >70% of host species) were represented by 7 OTUs at high relative abundance, accounting for 40% of all sequence reads. These OTUs corresponded to 2 *Prochlorococcus* sp. (*Cyanobacteria*; OTU0140, OTU0310) that were also common in seawater communities (41 and 40% relative abundance, respectively), as well as *Nitrosopumilus* sp. (*Thaumarchaeota*; OTU0001), *Prochlorococcus* sp. (*Cyanobacteria*; OTU0292), Rhodobacteraceae sp. (*Alphaproteobacteria*; OTU0188), *Pirellulales* sp. (*Planctomycetes*; OTU0164) and an OTU from the candidate phylum SBR1093 (OTU0355) that were rare (0.01–0.12% relative abundance) or absent in seawater samples. Variable OTUs (present in at least 2 host species) were represented by 950 OTUs and accounted for 49% of sequence reads, while specific OTUs (present in a single host species) were represented by 2364 OTUs and accounted for 11% of sequence reads.

Community-level analysis of tunic-associated microbes among ascidian species revealed a significant correlation between host relatedness (18S rRNA sequence similarity) and symbiont community similarity (Mantel test,  $r=0.37$ ,  $P<0.001$ ). This relationship was maintained when replicate samples were removed ( $r=0.28$ ,  $P<0.001$ ) and when using sub-sampled sequence pools to standardize sampling effort ( $r=0.50$ ,  $P<0.001$ ), indicating that high symbiont similarity among individuals of the same species and sampling artifacts were not the sole drivers of the observed correlation. Indeed, while symbiont communities were consistent across replicate individuals for 5 colonial ascidian species, other host species exhibited high intra-specific variability among replicates, including two solitary and three colonial species (Table 2). The lowest intra-specific diversity in symbiont structure was seen in *Lissoclinum badium*, where shared symbionts accounted for 37% of OTU<sub>0.03</sub> diversity and 99% of sequence reads. The highest intra-specific diversity was seen in *Phallusia arabica*, where shared symbionts only accounted for 2% of OTU<sub>0.03</sub> diversity and 21% of sequence reads (Table 2). Symbiont communities did not strictly cluster by higher-level host taxonomy (order to genus-level) or lifestyle (solitary or colonial; Figure 2), likely obscured by the observed variability in symbiont specificity among hosts.

#### Bacterial ultrastructure in the ascidian tunic

Transmission electron microscopy examination of the solitary ascidians *Phallusia julinea* and *Polycarpa aurata* revealed randomly distributed and extremely rare bacterial cells in the inner tunic of these two species. All bacterial morphotypes observed in *P. julinea* were ovoid to rod-shaped cells (ca. 0.4  $\mu\text{m} \times 2 \mu\text{m}$ ; Supplementary Figure S7A), while ovoid cells (ca. 0.12  $\mu\text{m}$ ), cyanobacteria (ca. 0.15  $\mu\text{m}$ , with ca. 5 thylakoids evenly spaced along the periphery of the cell), and a spiral bacterium (Supplementary Figure S7B) were observed in *P. aurata*. Colonial ascidians were characterized by a higher number of bacteria in their tunic. *Pycnoclavella* sp. featured groups of 2 to 5 cyanobacteria encased in a network of fibers (Supplementary Figure S7C). Both clavelinids (*Pycnoclavella* sp. and *C. meridionalis*) contained ovoid-shaped bacteria often surrounded by irregular inclusions spread throughout the tunic (Supplementary Figure S7D). In *Lissoclinum badium* and *Synoicum castellatum*, all bacterial cells were ovoid or rod-shaped (ca. 0.5  $\mu\text{m} \times 2 \mu\text{m}$ , and ca. 0.3  $\mu\text{m} \times 1 \mu\text{m}$ , respectively) and observed either in isolation or forming small groups of 2–6 bacteria in close proximity to ascidian cells (Supplementary Figure S7E and S7F, respectively).

## Discussion

#### Bacterial biodiversity hotspots in the ascidian tunic

In this study, we provide the most comprehensive characterization of the ascidian microbiota to date and reveal exceptional bacterial biodiversity inhabiting the tunic of GBR ascidians. Encompassing 3321 unique OTU<sub>0.03</sub> from 19 described bacterial phyla, 14 candidate bacterial phyla and 3 described archaeal phyla, the ascidian microbiota exhibited comparable diversity to the rich microbiota associated with marine sponges (Schmitt *et al.*, 2012) and corals (Sunagawa *et al.*, 2010) and indicates that the ascidian tunic represents a previously unrecognized hotspot for marine microbial diversity. Visualization of microbial cells by transmission electron microscopy confirmed the presence of microbes in the ascidian tunic and was consistent with results from 16S rRNA gene tag pyrosequencing, for example, the prevalence of cyanobacterial OTUs (>50% of sequence reads) and cyanobacterial cells encased in a fiber network in *Pycnoclavella* sp. and the detection of a *Spirochaetes* OTU (18% relative abundance) and a bacterium with spiral morphology in *Polycarpa aurata*.

Phylum-level composition of the ascidian microbiota retrieved herein was similar to what has been described for other ascidian species and was comprised of mostly *Proteobacteria*, *Bacteroidetes* and *Planctomycetes* (Martínez-García *et al.*, 2007; Tait *et al.*, 2007; Behrendt *et al.*, 2012). Moreover, as found for other tropical ascidians (e.g., Behrendt

*et al.*, 2012), *Cyanobacteria* were particularly abundant in most of our ascidian samples. In addition, the ascidian microbiota demonstrated some overlap with other host-associated microbial communities yet clear distinction from ambient planktonic communities in coral reef seawater, except for the widespread presence of *Cyanobacteria* from the *Prochlorococcus* genus. Consistently, previous studies have noted multiple shared symbiont lineages among microbiota of sponges and corals (Taylor *et al.*, 2007; Simister *et al.*, 2012), indicating microbial lineages adapted to host-associated lifestyles may disperse among disparate host organisms. However, the ascidian microbiota also maintained distinguishing characteristics in comparison to other host-associated communities. For example, the phylum *Planctomycetes* exhibited high diversity in ascidian hosts, whereas members of this phylum are typically rare in microbiota of sponge (Schmitt *et al.*, 2012; Webster and Taylor, 2012) and coral hosts (Sunagawa *et al.*, 2010; Barott *et al.*, 2011). Further, 11 of the 56 most common OTUs in the ascidian microbiota exhibited high sequence divergence (>5%) from any previously described marine microbe. The unique niches inside invertebrate tissues are becoming recognized hotspots for microbial biodiversity and our results suggest that ascidian tunics offer a similarly fertile habitat for marine microorganisms.

#### *Rare seawater microbes enriched in the ascidian tunic*

The vast majority of OTUs in the ascidian microbiota were not present in planktonic communities. However, a cautionary note is necessary here as seawater samples were collected in only one of our sampling sites and at one given time (October 2011), while ascidian samples were collected from different locations (separated by less than 120 km) and times (May through November 2011; Supplementary Table S1). Microbes in seawater are known to vary seasonally, occur in patches or be stratified according to their microenvironmental requirements or to microscale turbulences (e.g., Giovannoni and Stingl, 2005). Accordingly, the low number of shared OTUs (3%) between the seawater and the ascidian samples may be partly due to an insufficient sampling of the surrounding seawater.

Nevertheless, we found that several microbes from the rare biosphere of seawater exhibited high relative abundance in ascidian-associated communities. Five microbial OTUs exhibited 200 to 700 times higher relative abundance in the ascidian tunic than in the plankton, suggesting the selective enrichment of rare seawater microbes in ascidian hosts as observed for the microbiota in marine sponges (Webster *et al.*, 2010; Taylor *et al.*, 2013) and reef-building corals (Sunagawa *et al.*, 2010). Notably, 3 of the 5 OTUs enriched in the ascidian microbiota were classified to the order Rhizobiales, a lineage of *Alphaproteobacteria* well known for

their nitrogen-fixation capacity and mutualistic relationships with terrestrial plants (Lodwig *et al.*, 2003) and more recently documented as dominant nitrogen-fixing symbionts in the coral microbiome (Lema *et al.*, 2012). In this study, a total of 176 OTUs affiliated with Rhizobiales were present in the ascidian microbiota and detected in all 25 ascidian host species prompting further study of nitrogen-fixing bacteria in the ascidian microbiota and their potential contribution to nitrogen cycles in the ascidian holobiont. These results also indicate the potential for horizontal symbiont transfer among hosts with the rare biosphere of seawater acting as a conduit among host habitats.

#### *Host specificity of the ascidian microbiota*

The vast majority of symbiont OTUs (71%) were present in a single host species and absent in seawater, indicating a high degree of host specificity in the microbiota of coral reef ascidians. Indeed, no universal symbionts (i.e., present in all ascidian hosts) occurred and only 7 core OTUs (of 3321 total OTUs) were detected. While few 16S rRNA gene sequence datasets from ascidians are available for comparative analyses, several OTUs exhibited specific associations with particular host taxa across a broad geographic range. For example, OTU3073 from *Ecteinascidia diaphanis* matched to the candidate genus *Endoecteinascidia*, a distinct lineage of *Gammaproteobacteria* described solely from ascidians in the genus *Ecteinascidia*, including *E. turbinata* from the Mediterranean (Moss *et al.*, 2003) and Caribbean (Pérez-Matos *et al.*, 2007). The detection of this candidate genus from a GBR ascidian expands the known geographic range of this symbiont taxon and further supports its specificity to the host genus *Ecteinascidia*. In addition, this symbiont lineage is particularly notable for its putative role in secondary metabolite synthesis within the animal cell, including the production of the anticancer agent ET-743 (Rath *et al.*, 2011), which may constitute a key functional aspect of ascidian-bacterial symbioses (Kwan *et al.*, 2012).

Even among replicate individuals of the same ascidian species, some intra-specific variability was observed. Consistent microbial community structure was observed in 5 of the 10 ascidian species where multiple individuals were analyzed, while the remaining half exhibited greater similarity to the microbiota of unrelated species than to conspecific hosts, suggesting a non-obligate symbiosis. These results suggest different factors structuring the symbiont communities in different ascidian species, with more homogenous communities potentially maintained in some hosts by vertical symbiont transmission or specific functional requirements and more heterogeneous communities in other hosts determined by more stochastic or dynamic factors. This observation is in agreement with mounting evidence suggesting that colonial ascidians, such as

the Didemnidae, establish stable symbiotic microbial associations that are vertically transmitted (Kott, 1980, 1982, 2001; Hirose, 2000; Schuett *et al.*, 2005; Hirose *et al.*, 2006a, b; Hirose and Hirose, 2007; Bright and Bulgheresi, 2010; López-Legentil *et al.*, 2011; Kojima and Hirose, 2012), while others, such as solitary ascidians may selectively acquire symbionts from the surrounding seawater (Erwin *et al.*, 2013).

#### *Widespread ammonia-oxidizing archaea (AOA) in the ascidian microbiota*

Nitrification is a key process in the global nitrogen cycle that results in the conversion of ammonia to nitrite (ammonia-oxidation) and nitrite to nitrate (nitrite-oxidation), a two-step process mediated solely by prokaryotic organisms (Ward *et al.*, 2007). The archaeal component of the ascidian microbiota was notably comprised of lineages with known ammonia-oxidization capabilities. In particular, sequences affiliated with the genus *Nitrosopumilus* dominated the archaeal communities in GBR ascidians and several *Nitrosopumilus* OTUs exhibited a widespread distribution among hosts and high relative abundance within hosts. In coral reef waters, observations of high nitrite/nitrate concentrations compared to adjacent, open water habitats have long suggested active nitrification among reef-associated microbes (Webb *et al.*, 1975). More recent studies have reported that host-associated microbes in sponges and corals contributed to nitrification in these reef habitats to a larger extent than reported for free-living communities in sediments and seawater (Diaz and Ward, 1997; Southwell *et al.*, 2008). The finding herein of widespread ammonia-oxidizing archaea in coral reef ascidians suggests an additional and potentially important source of nitrification in reef habitats.

In fact, the most dominant of all OTUs in the ascidian microbiota (17% of total reads) was classified in the genus *Nitrosopumilus* and matched nearly identically (>99% sequence identity) to a symbiotic ammonia-oxidizing archaea (AOA) previously described in the Mediterranean ascidian *Cystodytes dellechiajei*, where active nitrification was detected in the tunic layer (Martínez-García *et al.*, 2008). Another OTU recovered from two individuals of the ascidian *Leptoclinides madara* at high relative abundance (17–28%) was classified in the genus *Cenarchaeum*, a candidate taxon erected for the sponge-associated symbiont *Cenarchaeum symbiosum* (Preston *et al.*, 1996) whose genome includes homologues of genes associated with chemolithotrophic ammonia oxidation (Hallam *et al.*, 2006). Finally, some ascidians (e.g., *Lissoclinum badium*) hosted *Nitrospina* symbionts, a genus of *Deltaproteobacteria* whose members are capable of nitrite-oxidation, in addition to dominant AOA lineages, suggesting that the complete nitrification

process may occur in the ascidian tunic of at least some species.

Ammonia is the primary form of nitrogenous waste produced by ascidians (Goodbody, 1974) and may be recycled *via* uptake or oxidation by resident microbes. For example, the widespread AOA reported herein may utilize the ammonia-rich waste products of their host ascidians as substrate for nitrification reactions. Indeed, nitrifying microbes require not only a reduced form of inorganic nitrogen, but also high oxygen and low irradiance levels, as marine AOA are particularly susceptible to photoinhibition at higher irradiance levels (Merbt *et al.*, 2012). Thus, the ascidian tunic habitat not only satisfies the ammonia and oxygen requirements of AOA (Kühl *et al.*, 2012), but may also shelter these populations from the high irradiance levels characteristic of shallow water reefs (e.g., Vermeij and Bak, 2002) and represent important habitats for nitrite/nitrate regeneration in coral reef environments. Further, the dynamic chemical landscapes in and around ascidians (Behrendt *et al.*, 2012, Kühl *et al.*, 2012) may offer periodic windows of optimal conditions for additional metabolic pathways and maintain the complex microbiota observed in ascidian tunics.

While the taxonomic scope of the ascidian species examined herein was broad, the geographic scope was restricted to shallow water habitats of the GBR. Yet even within this single biome, our results show a remarkably rich and diverse microbial community associated with coral reef ascidians. Given the broad distribution of ascidians in the marine environment, (Lambert, 2005) expanded efforts to document the diversity of the ascidian microbiota will continue to clarify the role of ascidians as habitats for novel microbial communities and their importance for microbial-mediated processes in marine biogeochemical cycles.

#### Conflict of Interest

The authors declare no conflict of interest.

#### Acknowledgements

This research was funded by the Marie Curie International Reintegration Grant FP7-PEOPLE-2010-RG 277038 (within the 7th European Community Framework Program), the Spanish Government projects CTM2010-17755 and CTM2010-22218 and the Catalan Government grant 2009 SGR-484 for Consolidated Research Groups. NSW was funded through an Australian Research Council Future Fellowship (FT1200100480).

#### References

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. (1990). Basic local alignment search tool. *J Mol Biol* **215**: 403–410.

- Armstrong E, Yan L, Boyd KG, Wright PC, Burgess JG. (2001). The symbiotic role of marine microbes on living surfaces. *Hydrobiologia* **461**: 37–40.
- Barott KL, Rodriguez-Brito B, Janoušek J, Marhaver KL, Smith JE, Keeling P *et al.* (2011). Microbial diversity associated with four functional groups of benthic reef algae and the reef-building coral *Montastraea annularis*. *Environ Microb* **13**: 1192–1204.
- Behrendt L, Larkum Anthony WD, Trampe E, Norman A, Sørensen SJ, Kühl M. (2012). Microbial diversity of biofilm communities in microniches associated with the didemnid ascidian *Lissoclinum patella*. *ISME J* **6**: 1222–1237.
- Bengtsson J, Eriksson KM, Hartmann M, Wang Z, Shenoy BD, Grelet G-A *et al.* (2011). Metaxa: a software tool for automated detection and discrimination among ribosomal small subunit (12S/16S/18S) sequences of archaea, bacteria, eukaryotes, mitochondria, and chloroplasts in metagenomes and environmental sequencing datasets. *Anton Leeuw* **100**: 471–475.
- Bright M, Bulgheresi S. (2010). A complex journey: transmission of microbial symbionts. *Nat Rev Microbiol* **8**: 218–230.
- Brochier-Armanet C, Boussau B, Gribaldo S, Forterre P. (2008). Mesophilic Crenarchaeota: proposal for a third archaeal phylum, the Thaumarchaeota. *Nat Rev Microbiol* **6**: 245–252.
- Cox G. (1986). Comparison of *Prochloron* from different hosts: I. Structural and ultrastructural characteristics. *New Phytol* **104**: 429–445.
- Cox GC, Hiller RG, Larkum AWD. (1985). An unusual cyanophyte, containing phycourobilin and symbiotic with ascidians and sponges. *Mar Biol* **89**: 149–163.
- Diaz MC, Ward BB. (1997). Sponge-mediated nitrification in tropical benthic communities. *Mar Ecol Prog Ser* **156**: 97–107.
- Donia MS, Fricke WF, Partensky F, Cox J, Elshahawi SI, White JR *et al.* (2011). Complex microbiome underlying secondary and primary metabolism in the tunicate-*Prochloron* symbiosis. *Proc Natl Acad Sci USA* **108**: E1423–E1432.
- Dray S, Dufour A. (2007). The ade4 package: Implementing the duality diagram for ecologists. *J Stat Softw* **22**: 1–20.
- Delsuc F, Brinkmann H, Chourrout D, Philippe H. (2006). Tunicates and not cephalochordates are the closest living relatives of vertebrates. *Nature* **439**: 965–968.
- Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. (2011). UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* **27**: 2194–2200.
- Erwin PM, López-Legentil S, Schuhmann PW. (2010). The pharmaceutical value of marine biodiversity for anti-cancer drug discovery. *Ecol Econ* **70**: 445–451.
- Erwin PM, Pineda MC, Webster N, Turon X, López-Legentil S. (2013). Small core communities and high variability in bacteria associated with the introduced ascidian *Styela plicata*. *Symbiosis* **59**: 35–46.
- Fan L, Reynolds D, Liu M, Stark M, Kjelleberg S, Webster NS *et al.* (2012). Functional equivalence and evolutionary convergence in complex communities of microbial sponge symbionts. *Proc Natl Acad Sci USA* **109**: E1878–E1887.
- Fieseler L, Horn M, Wagner M, Hentschel U. (2004). Discovery of the novel candidate phylum “Poribacteria” in marine sponges. *Appl Environ Microbiol* **70**: 3724–3732.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. (1994). DNA primers for amplification of mitochondrial cytochrome *c* oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotech* **3**: 294–299.
- Freeman CJ, Thacker RW. (2011). Complex interactions between marine sponges and their symbiotic microbial communities. *Limnol Oceanogr* **56**: 1577–1586.
- Freeman MF, Gurgui C, Helf MJ, Morinaka BI, Uria AR, Oldham NJ *et al.* (2012). Metagenome mining reveals polytheonamides as posttranslationally modified ribosomal peptides. *Science* **338**: 387–390.
- Giovannoni SJ, Stingl U. (2005). Molecular diversity and ecology microbial plankton. *Nature* **437**: 343–348.
- Goodbody I. (1974). The physiology of ascidians. *Adv Mar Biol* **12**: 121–149.
- Hallam SJ, Konstantinidis KT, Putnam N, Schleper C, Watanabe Y, Sugahara J *et al.* (2006). Genomic analysis of the uncultivated marine crenarchaeote *Cenarchaeum symbiosum*. *Proc Natl Acad Sci USA* **103**: 18296–18301.
- Hernández-Mariné M, Turon X, Catalan J. (1990). A marine *Synechocystis* (Chroococcales, Cyanophyta) epizooic on didemnid ascidians from the Mediterranean Sea. *Phycologia* **29**: 275–284.
- Hirose E. (2000). Plant rake and algal pouch of the larvae in the tropical ascidian *Diplosoma similis*: an adaptation for vertical transmission of photosynthetic symbionts *Prochloron* sp. *Zool Sci* **17**: 233–240.
- Hirose E, Hirose M. (2007). Morphological process of vertical transmission of photosymbionts in the colonial ascidian *Trididemnum miniatum* Kott, 1977. *Mar Biol* **150**: 359–367.
- Hirose E, Maruyama T. (2004). What are the benefits in the ascidian-*Prochloron* symbiosis? *Endocyt Cell Res* **15**: 51–62.
- Hirose E, Adachi R, Kuze K. (2006a). Sexual reproduction of the *Prochloron*-bearing ascidians, *Trididemnum cyclops* and *Lissoclinum bistratum*, in subtropical waters: seasonality and vertical transmission of photosymbionts. *J Mar Biol Ass UK* **86**: 175–179.
- Hirose E, Hirose M, Neilan BA. (2006b). Localization of symbiotic cyanobacteria in the colonial ascidian *Trididemnum miniatum* (Didemnidae, Ascidiacea). *Zool Sci* **23**: 435–442.
- Hirose E, Maruyama T, Cheng L, Lewin RA. (1996). Intracellular symbiosis of a photosynthetic prokaryote, *Prochloron* sp., in a colonial ascidian. *Invertebr Biol* **115**: 343–348.
- Hirose E, Ohtsuka K, Ishikura M, Maruyama T. (2004). Ultraviolet absorption in ascidian tunic and ascidian-*Prochloron* symbiosis. *J Mar Biol Assoc UK* **84**: 789–794.
- Hirose E, Turon X, López-Legentil S, Erwin PM, Hirose M. (2012). First records of didemnid ascidians harboring *Prochloron* from the Caribbean Panama: genetic relationships between Caribbean and Pacific photosymbionts and host ascidians. *Sys Biodivers* **10**: 435–445.
- Hoffmann F, Larsen O, Thiel V, Rapp HT, Pape T, Michaelis W *et al.* (2005). An anaerobic world in sponges. *Geomicrobiol J* **22**: 1–10.
- Hoffmann F, Radax R, Woebken D, Holtappels M, Lavik G, Rapp HT *et al.* (2009). Complex nitrogen cycling in the sponge *Geodia barretti*. *Environ Microb* **11**: 2228–2243.
- Hoffmann F, Røy H, Bayer K, Hentschel U, Pfannkuchen M, Brümmer F *et al.* (2008). Oxygen dynamics and

- transport in the Mediterranean sponge *Aplysina aerophoba*. *Mar Biol* **153**: 1257–1264.
- Huber JA, Mark Welch DB, Morrison HG, Huse SM, Neal PR, Butterfield DA *et al.* (2007). Microbial population structures in the deep marine biosphere. *Science* **318**: 97–100.
- Huse SM, Welch DM, Morrison HG, Sogin ML. (2010). Ironing out the wrinkles in the rare biosphere through improved OTU clustering. *Environ Microbiol* **12**: 1889–1898.
- Kojima A, Hirose E. (2012). Transmission of cyanobacterial symbionts during embryogenesis in the coral reef ascidians *Trididemnum nubilum* and *T. clinides* (Didemnidae, Ascidiacea, Chordata). *Biol Bull* **222**: 63–73.
- Kojima A, Hirose E. (2010). Transfer of prokaryotic algal symbionts from a tropical ascidian (*Lissoclinum punctatum*) colony to its larvae. *Zool Sci* **27**: 124–127.
- Kott P. (1980). Algal-bearing didemnid ascidians in the Indo-West-Pacific. *Mem Queensl Mus* **20**: 1–47.
- Kott P. (1982). Didemnid-algal symbioses: host species in the Western Pacific with notes on the symbiosis. *Micronesica* **18**: 95–127.
- Kott P. (2001). The Australian Ascidiacea. Part 4, Aplousobranchia (3), Didemnidae. *Mem Queensl Mus* **47**: 1–407.
- Kühl M, Behrendt L, Trampe E, Qvortrup K, Schreiber U, Borisov SM *et al.* (2012). Microenvironmental ecology of the chlorophyll b-containing symbiotic cyanobacterium *Prochloron* in the didemnid ascidian *Lissoclinum patella*. *Front Microbiol* **3**: 73–90.
- Kwan JC, Donia MS, Han AW, Hirose E, Haygood MG, Schmidt EW. (2012). Genome streamlining and chemical defense in a coral reef symbiosis. *Proc Natl Acad Sci USA* **109**: 20655–20660.
- Lambert G. (2005). Ecology and natural history of the protochordates. *Can J Zool* **83**: 34–50.
- Lema KA, Willis BL, Bourne DG. (2012). Corals form characteristic associations with symbiotic nitrogen-fixing bacteria. *Appl Environ Microbiol* **78**: 3136–3144.
- Lodwig EM, Hosie AHF, Bourdès A, Findlay K, Allaway D, Karunakaran R *et al.* (2003). Amino-acid cycling drives nitrogen fixation in the legume-Rhizobium symbiosis. *Nature* **422**: 722–726.
- López-Legentil S, Song B, Bosch M, Pawlik JR, Turon X. (2011). Cyanobacterial diversity and a new *Acarochloris*-like symbiont from Bahamian sea-squirrels. *PLoS One* **6**: e23938.
- López-Legentil S, Turon X. (2005). How do morphotypes and chemotypes relate to genotypes? The colonial ascidian *Cystodytes* (Polycitoridae). *Zool Scr* **34**: 3–14.
- Martínez-García M, Díaz-Valdés M, Wanner G, Ramos-Esplá A, Antón J. (2007). Microbial community associated with the colonial ascidian *Cystodytes dellechiaiei*. *Environ Microbiol* **9**: 521–534.
- Martínez-García M, Koblížek M, López-Legentil S, Antón J. (2011). Epibiosis of oxygenic phototrophs containing chlorophylls *a*, *b*, *c*, and *d* on the colonial ascidian *Cystodytes dellechiaiei*. *Microb Ecol* **61**: 13–19.
- Martínez-García M, Stief P, Díaz-Valdés M, Wanner G, Ramos-Esplá A, Dubilier N *et al.* (2008). Ammonia-oxidizing *Crenarchaeota* and nitrification inside the tissue of a colonial ascidian. *Environ Microbiol* **10**: 2991–3001.
- Merbt SN, Stahl DA, Casamayor EO, Martí E, Nicol GW, Prosser JL. (2012). Differential photoinhibition of bacterial and archaeal ammonia oxidation. *FEMS Microbiol Lett* **327**: 41–46.
- McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A *et al.* (2012). An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME J* **6**: 610–618.
- Moss C, Green DH, Pérez B, Velasco A, Henríquez R, McKenzie JD. (2003). Intracellular bacteria associated with the ascidian *Ecteinascidia turbinata*: phylogenetic and *in situ* hybridisation analysis. *Mar Biol* **143**: 99–110.
- Muscatine L, Porter JW. (1977). Reef corals: mutualistic symbioses adapted to nutrient-poor environments. *BioScience* **27**: 454–460.
- Münchhoff J, Hirose E, Maruyama T, Sunairi M, Burns BP, Neilan BA. (2007). Host specificity and phylogeography of the prochlorophyte *Prochloron* sp., an obligate symbiont in didemnid ascidians. *Environ Microbiol* **9**: 890–899.
- Pardy RL, Lewin RA. (1981). Colonial ascidians with Prochlorophyte symbionts: evidence for translocation of metabolites from alga to host. *Bull Mar Sci* **31**: 817–823.
- Paul VJ, Ritson-Williams R. (2008). Marine chemical ecology. *Nat Prod Rep* **25**: 662–695.
- Pawlik JR. (2011). The chemical ecology of sponges on Caribbean reefs: natural products shape natural systems. *Bio Science* **61**: 888–898.
- Preston CM, Wu KY, Molinski TF, DeLong EF. (1996). A psychrophilic crenarchaeon inhabits a marine sponge: *Cenarchaeum symbiosum* gen. nov., sp. nov. *Proc Natl Acad Sci USA* **93**: 6241–6246.
- Pérez-Matos AE, Rosado W, Govind NS. (2007). Bacterial diversity associated with the Caribbean tunicate *Ecteinascidia turbinata*. *A van Leeuw J Microb* **92**: 155–164.
- Pérez-Portela R, Bishop JDD, Davis AR, Turon X. (2009). Phylogeny of the families Pyuridae and Styelidae (Stolidobranchiata, Ascidiacea) inferred from mitochondrial and nuclear DNA sequences. *Mol Phylogenet Evol* **50**: 560–570.
- Raina JB, Dinsdale EA, Willis BL, Bourne DG. (2010). Do the organic sulfur compounds DMSP and DMS drive coral microbial associations? *Trends Microbiol* **18**: 101–108.
- Rath CM, Janto B, Earl J, Ahmed A, Hu FZ, Hiller L *et al.* (2011). Meta-omic characterization of the marine invertebrate microbial consortium that produces the chemotherapeutic natural product ET-743. *ACS Chem Biol* **6**: 1244–1256.
- Roesch LFW, Fulthorpe RR, Riva A, Casella G, Hadwin AKM, Kent AD *et al.* (2007). Pyrosequencing enumerates and contrasts soil microbial diversity. *ISME J* **1**: 283–290.
- Rowan R. (1998). Diversity and ecology of zooxanthellae on coral reefs. *J Phycol* **34**: 407–417.
- Schloss PD, Gevers D, Westcott SL. (2011). Reducing the effects of PCR amplification and sequencing artifacts on 16S rRNA-based studies. *PLoS One* **6**: e27310.
- Schloss PD, Westcott SL. (2011). Assessing and improving methods used in operational taxonomic unit-based approaches for 16S rRNA gene sequence analysis. *Appl Environ Microbiol* **77**: 3219–3226.
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB *et al.* (2009). Introducing mothur: open-source, platform-independent, community-supported

- software for describing and comparing microbial communities. *App Environ Microbiol* **75**: 7537–7541.
- Schmidt EW, Donia MS. (2010). Life in cellulose houses: symbiotic bacterial biosynthesis of ascidian drugs and drug leads. *Curr Opin Biotech* **21**: 827–833.
- Schmidt EW, Nelson JT, Rasko DA, Sudek S, Eisen JA, Haygood MG *et al.* (2005). Patellamide A and C biosynthesis by a microcin-like pathway in *Prochloron didemni*, the cyanobacterial symbiont of *Lissoclinum patella*. *Proc Natl Acad Sci USA* **102**: 7315–7320.
- Schmitt S, Tsai P, Bell J, Fromont J, Ilan M, Lindquist N *et al.* (2012). Assessing the complex sponge microbiota: core, variable and species-specific bacterial communities in marine sponges. *ISME J* **6**: 564–576.
- Schuett C, Doepke H, Groepler W, Wichels A. (2005). Diversity of intratunical bacteria in the tunic matrix of the colonial ascidian *Diplosoma migrans*. *Helgoland Mar Res* **59**: 136–140.
- Shenkar N, Swalla BJ. (2011). Global diversity of Ascidiacea. *PLoS One* **7**: e20657.
- Simister RL, Deines P, Botté ES, Webster NS, Taylor MW. (2012). Sponge-specific clusters revisited: a comprehensive phylogeny of sponge-associated microorganisms. *Environ Microbiol* **14**: 517–524.
- Smith EP, Van Belle G. (1984). Nonparametric estimation of species richness. *Biometrics* **40**: 119–129.
- Southwell MW, Weisz JB, Martens CS, Lindquist N. (2008). *In situ* fluxes of dissolved inorganic nitrogen from the sponge community on Conch Reef, Key Largo, Florida. *Limnol Oceanogr* **53**: 986–996.
- Spang A, Hatzenpichler R, Brochier-Armanet C, Rattei T, Tischler P, Spieck E *et al.* (2010). Distinct gene set in two different lineages of ammonia-oxidizing archaea supports the phylum Thaumarchaeota. *Trends Microbiol* **18**: 331–340.
- Stefaniak L, Lambert G, Gittenberger A, Zhang H, Lin S, Whitlatch RB. (2009). Genetic conspecificity of the worldwide populations of *Didemnum vexillum* Kott, 2002. *Aquat Invasions* **4**: 29–44.
- Sunagawa S, Woodley CM, Medina M. (2010). Threatened corals provide underexplored microbial habitats. *PLoS One* **5**: e9554.
- Tait E, Carman M, Sievert SM. (2007). Phylogenetic diversity of bacteria associated with ascidians in Eel Pond (Woods Hole, Massachusetts, USA). *J Exp Mar Biol Ecol* **342**: 138–146.
- Tarjuelo I, Posada D, Crandall KA, Pascual M, Turon X. (2004). Phylogeography and speciation of colour morphs in the colonial ascidian *Pseudodistoma crucigaster*. *Mol Ecol* **13**: 3125–3136.
- Taylor MW, Radax R, Steger D, Wagner M. (2007). Sponge-associated microorganisms: evolution, ecology, and biotechnological potential. *Microbiol Mol Biol Rev* **71**: 295–347.
- Taylor MW, Tsai P, Simister RL, Deines P, Botte E, Ericson G *et al.* (2013). “Sponge-specific” bacteria are widespread (but rare) in diverse marine environments. *ISME J* **7**: 438–443.
- Turon X, López-Legentil S, Banaigs B. (2005). Cell types, microsymbionts, and pyridocridine distribution in the tunic of three color morphs of the genus *Cystodytes* (Ascidiacea, Polycitoridae). *Invertebr Biol* **124**: 355–369.
- Usher KM, Toze S, Fromont J, Kuo J, Sutton DC. (2004). A new species of cyanobacterial symbiont from the marine sponge *Chondrilla nucula*. *Symbiosis* **36**: 183–192.
- Vermeij MJA, Bak RPM. (2002). How are coral populations structured by light? Marine light regimes and the distribution of Madracis. *Mar Ecol Prog Ser* **233**: 105–116.
- Ward BB, Capone DG, Zehr JP. (2007). What’s new in the nitrogen cycle? *Oceanography* **20**: 101–109.
- Webb KL, DuPaul WD, Wiebe W, Sottile W, Johannes RE. (1975). Enewetak (Eniwetok) Atoll: aspects of the nitrogen cycle on a coral reef. *Limnol Oceanogr* **20**: 198–219.
- Webster NS, Taylor MW. (2012). Marine sponges and their microbial symbionts: love and other relationships. *Environ Microbiol* **14**: 335–346.
- Webster NS, Taylor MW, Behnam F, Lückner S, Rattei T, Whalan S *et al.* (2010). Deep sequencing reveals exceptional diversity and modes of transmission for bacterial sponge symbionts. *Environ Microbiol* **12**: 2070–2082.
- Wilkinson CR. (1983). Net primary productivity in coral reef sponges. *Science* **219**: 410–412.
- Yokobori S, Kurabayashi A, Neilan BA, Maruyama T, Hirose E. (2006). Multiple origins of the ascidian-*Prochloron* symbiosis: molecular phylogeny of photosymbiotic and non-symbiotic colonial ascidians inferred from 18S rDNA sequences. *Mol Phylogenet Evol* **40**: 8–19.

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)