

## $\Delta R$ CORRECTION VALUES FOR THE NORTHERN INDIAN OCEAN

Koushik Dutta<sup>1</sup> • Ravi Bhushan • B L K Somayajulu

Physical Research Laboratory, Navrangpura, Ahmedabad – 380 009, India

**ABSTRACT.** Apparent marine radiocarbon ages are reported for the northern Indian Ocean region for the pre-nuclear period, based on measurements made in seven mollusk shells collected between 1930 and 1954. The conventional <sup>14</sup>C ages of these shells range from 693 ± 44 to 434 ± 51 BP in the Arabian Sea and 511 ± 34 to 408 ± 51 BP in the Bay of Bengal. These ages correspond to mean  $\Delta R$  correction values of 163 ± 30 yr for the northern Arabian Sea, 11 ± 35 yr for the eastern Bay of Bengal (Andaman Sea) and 32 ± 20 yr for the southern Bay of Bengal. Contrasting reservoir ages for these two basins are most likely due to differences in their thermocline ventilation rates.

### INTRODUCTION

The most common method for determining ages of marine sediments is radiocarbon dating of fossil calcareous tests of surface dwelling foraminifers and is frequently employed in high-resolution paleoclimate studies of the Late Quaternary period. The <sup>14</sup>C ages of marine fossils are on the average 400 years older than contemporary terrestrial wood, since the reservoir (seawater) from which these foraminifers derive carbon has lower <sup>14</sup>C/<sup>12</sup>C ratios compared to the atmosphere, due to mixing with deeper <sup>14</sup>C depleted water. Considerable spatial variability is seen in the apparent <sup>14</sup>C ages of marine calcareous shells due to variations in the regional ocean circulation patterns. For a given region (s) at any given time (t), the difference between the regional marine <sup>14</sup>C age  $R_s(t)$  and the global model marine <sup>14</sup>C age  $R_g(t)$ , is given by  $\Delta R(s)$  (Stuiver and Braziunas 1993). The  $\Delta R$  value accounts for regional deviations, necessary for calibrating <sup>14</sup>C ages of marine samples (Stuiver et al. 1986, 1993, 1998a) and is assumed to be approximately constant for a given region. This offset in the regional marine <sup>14</sup>C ages from the global modeled mean can be determined from <sup>14</sup>C dating of: 1) marine calcareous shells of known age from the pre-nuclear era, 2) growth bands in corals of pre-nuclear era, or 3) wood and marine shell pairs collected from same stratigraphic horizon.

The northern Indian Ocean region is an area of considerable interest to study the monsoon variations in southern Asia and Africa during the Quaternary period. The Arabian Sea and the Bay of Bengal sediments are archives of such climatic variations. The margin sediments, in particular, can record past monsoon variations with high temporal resolution (von Rad et al. 1999; Sarkar et al. 2000). Such studies obviously require a reliable <sup>14</sup>C chronology. Unfortunately, data on the <sup>14</sup>C reservoir ages for this part of the ocean are very meager. The present work is the first attempt to determine pre-nuclear  $\Delta^{14}C$  and  $\Delta R$  correction values for the northern Indian Ocean region, from <sup>14</sup>C measurements of archived marine shells. Here we report  $\Delta R$  values for the northern Arabian Sea, the northern and southern Bay of Bengal and for the Andaman Sea, based on <sup>14</sup>C measurements made on seven marine mollusk samples, collected between 1930 and 1954. The  $\Delta R$  values derived from <sup>14</sup>C measurements in annual bands of corals that grew between 1949 and 1954 in the Gulf of Kutch reported earlier (Chakraborty 1993; Bhushan et al. 1994) are also discussed.

### MATERIALS AND METHODS

The bivalve shells from Stewart Sound and Chilika Lake (Figure 1) were collected during 1935 and 1954, respectively, and procured from the archives of Zoological Survey of India (ZSI), Calcutta for <sup>14</sup>C measurements. Bivalve and gastropod samples from Port Okha, Dwarka and Rameswaram were

<sup>1</sup>Corresponding author. E-mail: koushik@prl.ernet.in.

obtained from different shell collectors. The coral *Favia speciosa* was collected from the Gulf of Kutch in June 1990. It was dated by identifying its annual growth bands using X-radiography, and sub-sampled for  $^{14}\text{C}$  measurements (Chakraborty 1993). The bivalve and gastropod samples were initially washed free of surface contamination with distilled water. They were then rinsed in 0.1N  $\text{HNO}_3$  and sonified in distilled water in an ultrasonic bath to dislodge surface organic coatings. The outer layers of the shells were etched off by dipping them in 2N HCl for about 2 minutes, to remove surface calcareous contaminants. They were then washed thoroughly with de-ionized water and oven dried. Before analysis, the shells were finely powdered in an agate mortar and homogenized. About 25–30 g of powdered samples were used for each analysis. For the two coral samples, two to three consecutive annual growth bands identified by X-radiograph were combined to get sufficient material for analysis (Chakraborty 1993). The powdered sample was hydrolyzed in a vacuum system using dilute (30% v/v)  $\text{H}_3\text{PO}_4$  and the  $\text{CO}_2$  liberated was converted to benzene using TASK Benzene Synthesizer. The benzene samples were assayed for  $^{14}\text{C}$  using Packard 2250C liquid scintillation counter in low level count mode (Bhushan et al. 1994). An aliquot of  $\text{CO}_2$  collected during the hydrolysis stage was analyzed for  $\delta^{13}\text{C}$  measurement using PDZ Europa Geo 20–20 mass spectrometer. The 1- $\sigma$  precisions for  $\Delta^{14}\text{C}$  and  $\delta^{13}\text{C}$  measurements are  $\pm 5\text{‰}$  and  $\pm 0.05\text{‰}$ , respectively.




Figure 1 Sampling sites from the northern Indian Ocean. Pooled mean  $\Delta R$  values in years ( $\pm 1\sigma$ ) are given for each region.

## RESULTS

The results of  $^{14}\text{C}$  measurements of the shell samples are given in Table 1. The  $^{14}\text{C}$  ages [ $R_s(t)$ ] are reported following the conventions of Stuiver and Polach (1977). The modeled global marine  $^{14}\text{C}$  ages [ $R_g(t)$ ] are from Stuiver et al. (1998b), reported for decadal samples. The  $\Delta R$  values are obtained as difference between  $R_s(t)$  and  $R_g(t)$ . Errors quoted on  $\Delta^{14}\text{C}$ ,  $R_s(t)$  and  $\Delta R$  are  $1\sigma$ . The

Table 1 Results of  $^{14}\text{C}$  analyses in archived pre-bomb marine shells and corals from the northern Indian Ocean<sup>a</sup>

Sample code (PRLCH-)	Species <sup>b</sup>	Sample location		Year of collection or growth (t)	$\delta^{13}\text{C}$ (‰)	$\Delta^{14}\text{C}$ (‰)	Conventional $^{14}\text{C}$ age (BP)		$\Delta R$ (years) $=R_s(t) - R_g(t)$
		Site	Lat., Long.				$R_s(t)$	$R_g(t)$	
<i>Arabian Sea coasts</i>									
319	<i>Turbo brunneus</i> sp.(g)	Port Okha, Gujarat	22°28'N, 69°05'E	1953	2.18	-83 ± 5	693 ± 44	473 ± 13	220 ± 46
543	<i>Turbo brunneus</i> sp.(g)	Dwarka, Gujarat	22°16'N, 68°57'E	1952	2.36	-72 ± 4	598 ± 35	473 ± 13	125 ± 37
542	<i>Conus monule</i> (g)	Dwarka, Gujarat	22°16'N, 68°57'E	1952	2.08	-173 ± 3	1524 ± 30 <sup>c</sup>	—	—
541	<i>Nerita oryzocem</i> (g)	Dwarka, Gujarat	22°16'N, 68°57'E	1952	3.39	-116 ± 4	988 ± 37 <sup>c</sup>	—	—
339	<i>Architectonica</i> sp. (g)	Dwarka, Gujarat	22°16'N, 68°57'E	1952	1.66	-141 ± 4	1219 ± 38 <sup>c</sup>	—	—
340	<i>Nassarius</i> sp. (g)	Dwarka, Gujarat	22°16'N, 68°57'E	1952	— <sup>d</sup>	-126 ± 4	1080 ± 37 <sup>c</sup>	—	—
87	<i>Favia speciosa</i> (c)	Pirotan Is., Gulf of Kutch	22°36'N, 70°E	1952–1954	-0.65	-53 ± 6 <sup>e</sup>	434 ± 51	473 ± 13	-39 ± 53
88	<i>Favia speciosa</i> (c)	Pirotan Is., Gulf of Kutch	22°36'N, 70°E	1949–1951	-0.79	-60 ± 6 <sup>e</sup>	497 ± 51	473 ± 13	24 ± 53
<i>Bay of Bengal coasts</i>									
320	<i>Marcia pinguis</i> (g)	Chilika Lake, Orissa	19°43'N, 85°37'E	1954	0.76	-50 ± 6	408 ± 51	473 ± 13	-65 ± 53
539	<i>Asaphis deflavata</i> (b)	Stewart Sd., N. Andaman	13°01'N, 92°58'E	1935	0.98	-55 ± 4	469 ± 34	458 ± 4	11 ± 35
545	<i>Cypraea</i> sp. (g)	Rameswaram, Tamilnadu	9°15'N, 79°29'E	1930	2.02	-56 ± 4	483 ± 34	458 ± 4	25 ± 35
544	<i>Cypraea</i> sp. (g)	Rameswaram, Tamilnadu	9°15'N, 79°29'E	1949	2.42	-60 ± 4	498 ± 34	465 ± 7	33 ± 35
526	<i>Cypraea</i> sp. (g)	Mandapam, Tamilnadu	9°12'N, 78°42'E	1953	2.36	-62 ± 4	511 ± 34	473 ± 13	38 ± 36

<sup>a</sup> $\Delta^{14}\text{C}$  and conventional  $^{14}\text{C}$  ages are defined by Stuiver and Polach (1977), normalized for isotopic fractionation to -25‰. Conventional  $^{14}\text{C}$  ages are calculated relative to AD 1950 using the Libby half-life.  $\Delta^{14}\text{C}$  values are corrected for  $^{14}\text{C}$  decay between the calendar year of the sample collection or sample growth (t) and AD 1950.

<sup>b</sup>b=bivalve, g=gastropod, c=coral.

<sup>c</sup>Not used for  $\Delta R$  calculation. See text for details.

<sup>d</sup> $\delta^{13}\text{C}$  not measured. 2‰ assumed (mean value of other similar samples).

<sup>e</sup>From Bhushan et al. (1994)

results are also summarized in Figure 1, where the  $\Delta R$  values quoted are error weighted pooled mean for a given region.

## DISCUSSION

### Arabian Sea

Two samples of *Turbobrunneus* collected from Port Okha and Dwarka from the northern margin of the Arabian Sea (see Figure 1), yielded mean  $^{14}\text{C}$  age of  $635 \pm 27$  BP ( $1 \sigma$ ), corresponding to mean  $\Delta R$  of  $163 \pm 30$  yr ( $1 \sigma$ ). In the northeastern Arabian Sea, vertical mixing is favored by 1) seasonal upwelling during southwest monsoon, and 2) convective processes associated with winter cooling (Madhupratap et al. 1996). The apparent  $^{14}\text{C}$  ages in the Arabian Sea are higher than that of the modeled world ocean, due to upwelling induced mixing with deeper  $^{14}\text{C}$ -depleted water. From accelerator mass spectrometry (AMS)  $^{14}\text{C}$  measurements of planktonic foraminifers in varved sediments off Pakistan, von Rad et al. (1999) estimated a local reservoir age of 640 yr for the Northern Arabian Sea. This is based on  $^{14}\text{C}$  ages of  $664 \pm 25$  BP and  $705 \pm 23$  BP for varves deposited in 1926 and 1898 AD, respectively. This core was collected from a location about 300 km northwest of our sampling site at Port Okha (Figure 1). The above ages correspond to  $\Delta R$  values of  $208 \pm 25$  and  $248 \pm 24$  yr, respectively, with a regional mean of  $229 \pm 27$  yr ( $1 \sigma$ ) (Reimer and Reimer 2001).

The coral *Favia speciosa* collected from the Pirotan Island within the Gulf of Kutch gives mean  $\Delta R$  of  $-8 \pm 37$  yr for the two coral bands, which is significantly younger than the open Arabian Sea samples. The average water depth of the gulf is less than 30 m (Chakraborty et al. 1994). The lower  $^{14}\text{C}$  age in the gulf compared to the open Arabian Sea appears to reflect faster equilibration time for the shallow gulf, with respect to atmospheric  $\text{CO}_2$  exchange. Even in the post-bomb era during late 1970s, the Gulf of Kutch recorded much higher level of  $^{14}\text{C}$  than the open Arabian Sea. The  $\Delta^{14}\text{C}$  of dissolved inorganic carbon (DIC) in the mixed layer at the GEOSECS station 416 in the northern Arabian Sea was  $59 \pm 4\text{‰}$  during 1978 (Stuiver and Ostlund 1983), while the mean  $\Delta^{14}\text{C}$  for the Pirotan Island coral sample for the year 1975 and 1978 is  $112 \pm 7\text{‰}$  (Chakraborty 1993). Four samples of gastropod collected from Dwarka yielded much older  $^{14}\text{C}$  ages, between 989 and 1524 BP (Table 1), and have been excluded from  $\Delta R$  calculations. The beaches of Port Okha and Dwarka in the western Gujarat from where these samples were collected are composed of calcareous sediments derived from Late Quaternary miliolite limestone and old coral reefs. The  $^{230}\text{Th}/^{234}\text{U}$  ages of these miliolite limestones range from  $\sim 30$  to  $\sim 235$  ka (Somayajulu 1993). The coral reefs from this region fall in two groups, one around 6 ka and the other between 118 to 176 ka (Somayajulu et al. 1985). Thus higher ages of the four gastropod samples from Dwarka are possibly due to incorporation of dead carbonate in their shells (Dye 1994).

### Bay of Bengal

The three samples from Tamilnadu were collected from the shallow Palk Bay area (water depth  $< 100$  m), yielded an average  $^{14}\text{C}$  age of  $497 \pm 20$  BP, corresponding to mean  $\Delta R$  of  $32 \pm 20$  yr. The  $^{14}\text{C}$  age of the sample collected from Stewart Sound in the Andaman Sea is  $469 \pm 34$  BP corresponding to  $\Delta R$  of  $11 \pm 35$  yr. This is statistically indistinguishable from that of the Palk Bay samples. The  $^{14}\text{C}$  age of the sample from Chilika Lake, a shallow lagoonal lake in the northern Bay of Bengal is  $408 \pm 51$  BP, the youngest compared to all the northern Indian Ocean samples analyzed in this study (Table 1).

The Bay of Bengal receives large amount of fresh water from the north, by seven major rivers (Milliman and Meade 1983), resulting in steep gradients of the isopycnal surfaces within the top 200 m.

This greatly reduces the vertical mixing rate, preventing advection of deeper  $^{14}\text{C}$  depleted water. As a result, relatively younger reservoir ages compared to the Arabian Sea are likely to be observed in this basin. However riverine DIC depleted in  $^{14}\text{C}$  will tend to counteract this effect (Little 1993). The  $^{14}\text{C}$  age of the Chilika Lake is slightly younger than that obtained for the southern Bay of Bengal region. Younger  $^{14}\text{C}$  reservoir ages in enclosed shallow lagoons are expected because of absence of upwelling of deeper  $^{14}\text{C}$  depleted waters and faster equilibration rate with the atmosphere. This is similar to the case for the Pirotan Island coral in the shallow Gulf of Kutch. More measurements from the northern Bay of Bengal are needed to ascertain its reservoir age.

## CONCLUSIONS

This is the first attempt to determine  $\Delta R$  correction values for the two northern Indian Ocean basins, the Arabian Sea and the Bay of Bengal, from apparent  $^{14}\text{C}$  ages of marine shells and corals formed prior to the nuclear era. The northern Arabian Sea has the mean  $\Delta R$  of  $163 \pm 30$  yr. For the enclosed Gulf of Kutch the  $\Delta R$  is  $-8 \pm 37$  yr obtained from  $^{14}\text{C}$  measurements in annual coral bands. The surface seawaters of the Bay of Bengal has younger apparent  $^{14}\text{C}$  age than that of the Arabian Sea, with  $\Delta R$  values of  $11 \pm 35$  yr for the Andaman Sea and  $32 \pm 20$  yr for the southern Bay of Bengal. The lowest  $\Delta R$  of  $-65 \pm 53$  yr is obtained for the Chilika Lake, a shallow lagoon in the northern Bay of Bengal.

## ACKNOWLEDGMENTS

We thank Dr J R B Alfred, Director, the Zoological Survey of India, Calcutta for providing the shell samples and Dr D K Sinha of BHU, Varanasi, for identification of some shell samples. We also thank Mr J C Drona, Mr G Rajagopal and Ms V Ganesh for providing samples from their collections. We are grateful to Prof S Krishnaswami for discussion and comments. We thank Dr John Southon for critically reviewing the manuscript and valuable suggestions. BLKS thanks CSIR and DOS for the Emeritus Scientistship and Hon. Professorship, respectively.

## REFERENCES

- Bhushan R, Chakraborty S, Krishnaswami S. 1994. Physical Research Laboratory (Chemistry) radiocarbon date list I. *Radiocarbon* 36(2):251–56.
- Chakraborty S. 1993. Environmental significance of isotopic and trace elemental variations in banded corals [unpublished PhD thesis]. MS University of Baroda, Vadodara, India.
- Chakraborty S, Ramesh R, Krishnaswami S. 1994. Air-sea exchange of  $\text{CO}_2$  in the Gulf of Kutch, northern Arabian Sea based on bomb-carbon in corals and tree rings. *Proceedings of the Indian Academy of Sciences* 103(2):231–42.
- Dye T. 1994. Apparent ages of marine shells: implications for archeological dating in Hawai'i. *Radiocarbon* 36(1):51–7.
- Little EA. 1993. Radiocarbon age calibration at archeological sites of coastal Massachusetts and vicinity. *Journal of Archeological Science* 20:457–71.
- Madhupratap M, Prasanna Kumar S, Bhattathiri PMA, Dileep Kumar M, Raghukumar S, Nair KKC, Ramiah, N. 1996. Mechanism of the biological response to the winter cooling in the northeastern Arabian Sea. *Nature* 384:549–52.
- Milliman JD, Mead RH. 1983. World delivery of riverine sediments to the oceans. *Journal of Geology* 21:1–21.
- Reimer PJ, Reimer RW. 2001. A marine reservoir correction database and on-line interface. *Radiocarbon*. This issue. [Supplemental material URL: <<http://calib.org/marine>>].
- Sarkar A, Ramesh R, Somayajulu BLK, Agnihotri R, Jull AJT, Burr GS. 2000. High resolution paleomonsoon record from the eastern Arabian Sea. *Earth and Planetary Science Letters* 177:209–18.
- Somayajulu BLK, Broecker WS, Goddard J. 1985. Dating Indian corals by U-decay-series methods. *Quaternary Research* 24:235–9.
- Somayajulu BLK. 1993. Age and mineralogy of the mollusks of Saurashtra and Kutch, Gujarat. *Current Science* 64(12):926–8.
- Stuiver M, Polach HA. 1977. Discussion: reporting of  $^{14}\text{C}$  data. *Radiocarbon* 19(3):355–63.
- Stuiver M, Ostlund HG. 1983. GEOSECS Indian Ocean and Mediterranean radiocarbon. *Radiocarbon* 25(1):1–29.

- Stuiver M, Pearson GW, Braziunas TF. 1986. Radiocarbon age calibration of marine samples back to 9000 cal BP. *Radiocarbon* 28(2B):980–1021.
- Stuiver M, Braziunas TF. 1993. Modeling atmospheric  $^{14}\text{C}$  influences and  $^{14}\text{C}$  ages of marine samples to 10,000 BC. *Radiocarbon* 35(1):137–89.
- Stuiver M, Reimer PJ, Braziunas TF. 1998a. High precision radiocarbon age calibration for terrestrial and marine samples. *Radiocarbon* 40(3):1127–51.
- Stuiver M, Reimer PJ, Bard E, Beck JW, Burr GS, Hughen KA, Kromer B, McCormac G, van der Plicht J, Spurk M. 1998b. INTCAL 98 radiocarbon age calibration, 24,000–0 cal BP. *Radiocarbon* 40(3):1041–83.
- von Rad U, Schaff M, Michels KH, Schulz H, Berger WH, Sirocko F. 1999. A 5000-yr record of climate change in varved sediments from the oxygen minimum zone off Pakistan, northeastern Arabian Sea. *Quaternary Research* 51:39–53.