

Washington University School of Medicine

Digital Commons@Becker

---

Open Access Publications

---

2017

## Draft genome sequence of a *Salmonella enterica* serovar Typhi strain resistant to fourth-generation cephalosporin and fluoroquinolone antibiotics

Danish Gul

*National University of Sciences and Technology*

Robert F. Potter

*Washington University School of Medicine in St. Louis*

Hurmat Riaz

*National University of Sciences and Technology*

Shifa Tariq Ashraf

*National University of Sciences and Technology*

Meghan A. Wallace

*Washington University School of Medicine in St. Louis*

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.wustl.edu/open\\_access\\_pubs](https://digitalcommons.wustl.edu/open_access_pubs)

---

### Recommended Citation

Gul, Danish; Potter, Robert F.; Riaz, Hurmat; Ashraf, Shifa Tariq; Wallace, Meghan A.; Munir, Tehmina; Ali, Amjad; Burnham, Carey-Ann; Dantas, Gautam; and Andleeb, Saadia, "Draft genome sequence of a *Salmonella enterica* serovar Typhi strain resistant to fourth-generation cephalosporin and fluoroquinolone antibiotics." *Genome Announcements*. 5,42. e00850-17. (2017).  
[https://digitalcommons.wustl.edu/open\\_access\\_pubs/6341](https://digitalcommons.wustl.edu/open_access_pubs/6341)

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact [vanam@wustl.edu](mailto:vanam@wustl.edu).

---

**Authors**

Danish Gul, Robert F. Potter, Hurmat Riaz, Shifa Tariq Ashraf, Meghan A. Wallace, Tehmina Munir, Amjad Ali, Carey-Ann Burnham, Gautam Dantas, and Saadia Andleeb



# Draft Genome Sequence of a *Salmonella enterica* Serovar Typhi Strain Resistant to Fourth-Generation Cephalosporin and Fluoroquinolone Antibiotics

Danish Gul,<sup>a</sup> Robert F. Potter,<sup>c</sup> Hurmat Riaz,<sup>a</sup> Shifa Tariq Ashraf,<sup>a</sup> Meghan A. Wallace,<sup>d</sup> Tehmina Munir,<sup>b</sup> Amjad Ali,<sup>a</sup> Carey-Ann Burnham,<sup>d,e,g</sup> Gautam Dantas,<sup>c,d,e,f</sup> Saadia Andleeb<sup>a</sup>

Atta-ur-Rahman School of Applied Biosciences, National University of Sciences and Technology (NUST), Islamabad, Pakistan<sup>a</sup>; Army Medical College, National University of Medical Sciences, Rawalpindi, Pakistan<sup>b</sup>; Center for Genome Sciences and Systems Biology, Washington University School of Medicine, St. Louis, Missouri, USA<sup>c</sup>; Department of Pathology and Immunology, Washington University School of Medicine, St. Louis, Missouri, USA<sup>d</sup>; Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, Missouri, USA<sup>e</sup>; Department of Biomedical Engineering, Washington University in St. Louis, St. Louis, Missouri, USA<sup>f</sup>; Department of Pediatrics, St. Louis Children's Hospital, St. Louis, Missouri, USA<sup>g</sup>

**ABSTRACT** Typhoid is endemic in developing countries. We report here the first draft genome sequence of a *Salmonella enterica* serovar Typhi clinical isolate from Pakistan exhibiting resistance to cefepime (a fourth-generation cephalosporin) and fluoroquinolone antibiotics, two of the last-generation therapies against this pathogen. The genome is ~4.8 Mb, with two putative plasmids.

Typhoid fever is one of the leading causes of mortality due to infectious diseases in developing Asian and African countries. Resistance to third-generation cephalosporin and fluoroquinolone antibiotics has been increasing in recent years in *Salmonella* spp. (1, 2), but there have been few reports of resistance to fourth-generation cephalosporins and fluoroquinolones in *Salmonella enterica* serovar Typhi.

In this study, we performed whole-genome sequencing of a multidrug-resistant *Salmonella enterica* subsp. *enterica* serovar Typhi clinical isolate obtained from a patient admitted to a tertiary care hospital in Rawalpindi, Pakistan (3). The isolate was phenotypically resistant to a fourth-generation cephalosporin (cefepime) and several fluoroquinolones (ciprofloxacin, levofloxacin, and moxifloxacin). Whole-genome sequencing (WGS) was carried out to identify the resistance determinants carried by this *S. Typhi* isolate.

Genomic DNA from 10 colonies of the isolate was extracted with the bacteremia DNA kit (Mo Bio) and used as input for sequencing libraries using a modification of the Nextera XT kit (Illumina) (4). A total of 10,307,350 paired-end 150-bp reads were generated from an Illumina NextSeq 2500 and processed using the High-Throughput Computing Facility at the Washington University School of Medicine (St. Louis, MO, USA). Illumina adapters and potentially contaminating human reads were removed using Trimmomatic and deconseq (5, 6). The resulting 10,288,264 paired-end reads were assembled into scaffolds using SPAdes 3.10.0 (7). Contigs were analyzed by QUAST (8). The assembly produced 95 contigs >500 bp, with an  $N_{50}$  value of 144,739 bp. The longest contig was 393,347 bp. Contigs were annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (<http://www.ncbi.nlm.nih.gov/genomes/static/Pipeline.html>), which identified 4,975 coding sequences (CDSs), 69 tRNA genes, and 4 rRNA genes.

Further analysis identified putative plasmid sequences and antimicrobial resistance genes. PlasmidFinder 1.3 (<https://cge.cbs.dtu.dk/services/PlasmidFinder/>) (9) identified

Received 31 July 2017 Accepted 18 September 2017 Published 19 October 2017

**Citation** Gul D, Potter RF, Riaz H, Ashraf ST, Wallace MA, Munir T, Ali A, Burnham C-A, Dantas G, Andleeb S. 2017. Draft genome sequence of a *Salmonella enterica* serovar Typhi strain resistant to fourth-generation cephalosporin and fluoroquinolone antibiotics. *Genome Announc* 5:e00850-17. <https://doi.org/10.1128/genomeA.00850-17>.

**Copyright** © 2017 Gul et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Saadia Andleeb, [saadia.andleeb@asab.nust.edu.pk](mailto:saadia.andleeb@asab.nust.edu.pk).

the presence of IncQ1 and IncY sequences. Antibiotic resistance genes (ARGs) were identified using ARDB (10), CARD (11), ResFinder (12), and ResFams (13). Annotation using these databases identified *bla*<sub>CTX-M-15</sub>, *bla*<sub>TEM-1</sub>, S83F *gyrA*, and *qnrS1*, which could potentially explain the observed resistance to cefepime and fluoroquinolones (ciprofloxacin, levofloxacin, and moxifloxacin).

The prevalence of *bla*<sub>CTX-M-15</sub>-positive *S. Typhi* isolates of Asian origin has been reported in Iraq, Kuwait, India, and Bangladesh (14–17). To the best of our knowledge, this study is the first draft genome sequence of a *bla*<sub>CTX-M-15</sub><sup>-</sup>, *bla*<sub>TEM-1</sub><sup>-</sup>, and *qnrS*-positive *S. Typhi* strain from Pakistan exhibiting resistance to cefepime and fluoroquinolones.

**Accession number(s).** This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [NIFP00000000](https://www.ncbi.nlm.nih.gov/nuccore/NIFP00000000). The version described in this paper is version NIFP01000000.

## ACKNOWLEDGMENTS

This work was supported in part by a United States Agency for International Development award (award number 3220-29047) to S.A., C.-A.B., and G.D. R.F.P. received support from an NIGMS training grant through award number T32 GM007067 (principal investigator [PI], James Skeath).

We acknowledge the pathology lab staff of AMC, NUMS Rawalpindi, for the clinical isolate. We also thank the Center for Genome Sciences & Systems Biology staff Eric Martin, Brian Koebbe, and Jessica Hoisington-López for their technical support and sequencing expertise.

## REFERENCES

- Kumar S, Rizvi M, Berry N. 2008. Rising prevalence of enteric fever due to multidrug-resistant *Salmonella*: an epidemiological study. *J Med Microbiol* 57:1247–1250. <https://doi.org/10.1099/jmm.0.2008/001719-0>.
- Eng S-K, Pusparajah P, Ab Mutalib N, Ser H, Chan K, Lee L. 2015. *Salmonella*: a review on pathogenesis, epidemiology and antibiotic resistance. *Front Life Sci* 8:284–293. <https://doi.org/10.1080/21553769.2015.1051243>.
- Munir T, Lodhi M, Ansari JK, Andleeb S, Ahmed M. 2016. Extended spectrum beta lactamase producing cephalosporin resistant *Salmonella* Typhi, reported from Rawalpindi, Pakistan. *J Pak Med Assoc* 66: 1035–1036.
- Baym M, Kryazhimskiy S, Lieberman TD, Chung H, Desai MM, Kishony R. 2015. Inexpensive multiplexed library preparation for megabase-sized genomes. *PLoS One* 10:e0128036. <https://doi.org/10.1371/journal.pone.0128036>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Schmieder R, Edwards R. 2011. Fast identification and removal of sequence contamination from genomic and metagenomic datasets. *PLoS One* 6:e17288. <https://doi.org/10.1371/journal.pone.0017288>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
- Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, Møller Aarestrup F, Hasman H. 2014. *In silico* detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 58:3895–3903. <https://doi.org/10.1128/AAC.02412-14>.
- Liu B, Pop M. 2009. ARDB—Antibiotic Resistance Genes Database. *Nucleic Acids Res* 37:D443–D447. <https://doi.org/10.1093/nar/gkn656>.
- McArthur AG, Waglechner N, Nizam F, Yan A, Azad MA, Baylay AJ, Bhullar K, Canova MJ, De Pascale G, Ejim L, Kalan L, King AM, Koteva K, Morar M, Mulvey MR, O'Brien JS, Pawlowski AC, Piddock LJ, Spanogiannopoulos P, Sutherland AD, Tang I, Taylor PL, Thaker M, Wang W, Yan M, Yu T, Wright GD. 2013. The comprehensive antibiotic resistance database. *Antimicrob Agents Chemother* 57:3348–3357. <https://doi.org/10.1128/AAC.00419-13>.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.
- Gibson MK, Forsberg KJ, Dantas G. 2015. Improved annotation of antibiotic resistance determinants reveals microbial resistomes cluster by ecology. *ISME J* 9:207–216. <https://doi.org/10.1038/ismej.2014.106>.
- Rotimi VO, Jamal W, Pal T, Sovenned A, Albert MJ. 2008. Emergence of CTX-M-15 type extended-spectrum beta-lactamase-producing *Salmonella* spp. in Kuwait and the United Arab Emirates. *J Med Microbiol* 57:881–886. <https://doi.org/10.1099/jmm.0.47509-0>.
- Pfeifer Y, Matten J, Rabsch W. 2009. *Salmonella enterica* serovar Typhi with CTX-M beta-lactamase, Germany. *Emerg Infect Dis* 15:1533–1535. <https://doi.org/10.3201/eid1509.090567>.
- Ahmed D, Hoque A, Mazumder R, Nahar K, Islam N, Gazi SA, Hossain MA. 2012. *Salmonella enterica* serovar Typhi strain producing extended-spectrum beta-lactamases in Dhaka, Bangladesh. *J Med Microbiol* 61: 1032–1033. <https://doi.org/10.1099/jmm.0.044065-0>.
- González-López JJ, Piedra-Carrasco N, Salvador F, Rodríguez V, Sánchez-Montalvá A, Planes AM, Molina I, Larrosa MN. 2014. ESBL-producing *Salmonella enterica* serovar Typhi in traveler returning from Guatemala to Spain. *Emerg Infect Dis* 20:1918–1920. <https://doi.org/10.3201/eid2011.140525>.