



# Drivers of multi-century trends in the atmospheric CO<sub>2</sub> mean annual cycle in a prognostic ESM

Jessica Liptak<sup>1</sup>, Gretchen Keppel-Aleks<sup>1</sup>, and Keith Lindsay<sup>2</sup>

<sup>1</sup>Department of Climate and Space Sciences and Engineering, University of Michigan, Ann Arbor, MI, USA

<sup>2</sup>Climate and Global Dynamics, National Center for Atmospheric Research, Boulder, CO, USA

Correspondence to: Jessica Liptak (liptak@umich.edu)

Received: 24 March 2016 – Discussion started: 5 April 2016

Revised: 10 February 2017 – Accepted: 21 February 2017 – Published: 20 March 2017

**Abstract.** The amplitude of the mean annual cycle of atmospheric CO<sub>2</sub> is a diagnostic of seasonal surface–atmosphere carbon exchange. Atmospheric observations show that this quantity has increased over most of the Northern Hemisphere (NH) extratropics during the last 3 decades, likely from a combination of enhanced atmospheric CO<sub>2</sub>, climate change, and anthropogenic land use change. Accurate climate prediction requires accounting for long-term interactions between the environment and carbon cycling; thus, analysis of the evolution of the mean annual cycle in a fully prognostic Earth system model may provide insight into the multi-decadal influence of environmental change on the carbon cycle.

We analyzed the evolution of the mean annual cycle in atmospheric CO<sub>2</sub> simulated by the Community Earth System Model (CESM) from 1950 to 2300 under three scenarios designed to separate the effects of climate change, atmospheric CO<sub>2</sub> fertilization, and land use change. The NH CO<sub>2</sub> seasonal amplitude increase in the CESM mainly reflected enhanced primary productivity during the growing season due to climate change and the combined effects of CO<sub>2</sub> fertilization and nitrogen deposition over the mid- and high latitudes. However, the simulations revealed shifts in key climate drivers of the atmospheric CO<sub>2</sub> seasonality that were not apparent before 2100. CO<sub>2</sub> fertilization and nitrogen deposition in boreal and temperate ecosystems were the largest contributors to mean annual cycle amplification over the mid-latitudes for the duration of the simulation (1950–2300). Climate change from boreal ecosystems was the main driver of Arctic CO<sub>2</sub> annual cycle amplification between 1950 and 2100, but CO<sub>2</sub> fertilization had a stronger effect on the Arctic CO<sub>2</sub> annual cycle amplitude during 2100–2300. Prior to 2100, the NH CO<sub>2</sub> annual cycle amplitude increased in con-

junction with an increase in the NH land carbon sink. However, these trends decoupled after 2100, underscoring that an increasing atmospheric CO<sub>2</sub> annual cycle amplitude does not necessarily imply a strengthened terrestrial carbon sink.

## 1 Introduction

The amplitude of the mean annual cycle of atmospheric CO<sub>2</sub>, an indicator of seasonal terrestrial and ocean carbon exchange, has increased over the Northern Hemisphere (NH) since observational records began in the late 1950s (Pearman and Hyson, 1981; Cleveland et al., 1983; Bacastow et al., 1985; Conway et al., 1994; Keeling et al., 1996; Randerson et al., 1997; Graven et al., 2013; Liu et al., 2015). The largest increases of 40–50 % were observed over the northern high latitudes via surface monitoring (Keeling et al., 1996) and from aircraft observations of the free troposphere (Graven et al., 2013). The amplification of the atmospheric CO<sub>2</sub> annual cycle primarily reflects enhanced net exchange of CO<sub>2</sub> with land surfaces rather than with the ocean (Manning, 1993). Land–atmosphere CO<sub>2</sub> exchange is highly seasonal, especially in the NH mid- and high latitudes where photosynthesis draws down CO<sub>2</sub> in the spring and summer, and net ecosystem respiration returns CO<sub>2</sub> to the atmosphere (e.g., Komhyr et al., 1985; Enting and Mansbridge, 1989; Nemry et al., 1996; Dettinger and Ghil, 1998).

Because atmospheric CO<sub>2</sub> observations are characterized by high precision and accuracy, the gradual, multi-decadal increase in the seasonal amplitude provides a unique observational target for Earth system models (ESMs) intended to predict the long-term coevolution of climate and the carbon

cycle. ESMs enable the study of long-term effects of natural and anthropogenic forcing on the terrestrial carbon cycle since they include mechanistic representations of the carbon cycle by coupling land surface models that explicitly resolve biogeochemical processes with models of the atmosphere, ocean, and other components of the climate system (Claussen et al., 2002). An advantage of using a coupled model is that feedbacks between the physical climate and biogeochemistry are represented in a self-consistent framework. This is crucial since carbon fluxes are inherently linked to the physical climate; for example, a change in gross primary productivity (GPP) will be associated with changes in evapotranspiration, which feeds back on metrics such as humidity, cloud cover, and precipitation. Moreover, in a fully prognostic model, both climate and carbon cycle diagnostics are free to evolve rather than being tied to input data sets that reflect the contemporary climate. The mechanisms embedded in ESMs to predict future carbon–climate interactions have been identified as the likely drivers of the observed mean annual cycle amplitude increase as described in the following paragraphs.

The magnitude of the amplitude increase suggests a dominant role for enhanced primary productivity during the growing season in addition to increased CO<sub>2</sub> release during the dormant season (Graven et al., 2013). Greater atmospheric CO<sub>2</sub> may facilitate plant carbon uptake through increased water use efficiency (Keenan et al., 2013), and results from Kohlmaier et al. (1989) and McGuire et al. (2001) suggest that CO<sub>2</sub> fertilization contributes at least 10 % to the CO<sub>2</sub> mean annual cycle amplitude trend. Human activity has not only increased the atmospheric concentration of atmospheric CO<sub>2</sub>, but also modified reactive nitrogen deposition (N deposition) in ecosystems. Deposition of nitrogen oxides (NO<sub>x</sub>) and ammonia from combustion, livestock, agriculture, and industrial sources may augment the supply of soil nitrogen available for fixation by plants (Prentice et al., 2001), alleviating a limitation on terrestrial GPP (Lloyd, 1999; Norby et al., 2010).

Climate-change-induced warming and lengthening of the growing season may also stimulate GPP and increase the seasonality of net exchange. Keeling et al. (1996) proposed that increased terrestrial CO<sub>2</sub> uptake from a longer high-latitude growing season has driven the amplification of the CO<sub>2</sub> annual cycle since the trends in the CO<sub>2</sub> annual cycle amplitude strengthen moving northward, and the greatest warming has occurred during the winter and spring over the northern high latitudes. Findings by Randerson et al. (1999), McDonald et al. (2004), and Barichivich et al. (2013) support the hypothesis that longer growing seasons enhance spring CO<sub>2</sub> uptake and annual cycle amplitudes over the Arctic. This effect may be counteracted by the fact that early growing season onset may lead to growing season moisture deficits that reduce terrestrial productivity later in the growing season (Angert et al., 2005; Buermann et al., 2013; Parida and Buermann, 2014). Model evidence suggests that climate-driven shifts in vegetation cover can also enhance GPP. Forkel et al.

(2016) showed that the interaction of vegetation dynamics and climate change leads to increased GPP over NH boreal and Arctic regions that in turn drive the observed increases in NH high-latitude seasonal CO<sub>2</sub> amplitudes.

In short, there are several potential drivers of the CO<sub>2</sub> annual cycle amplification that feed back into the climate system. Despite the representation of these mechanisms by ESMs, Graven et al. (2013) showed that none of the CMIP5 carbon cycle models were able to simulate the magnitude of the observed increase in atmospheric CO<sub>2</sub> seasonality. Since understanding the drivers of the CO<sub>2</sub> seasonality is crucial for model development, we used the Community Earth System Model (CESM) to study the contribution of natural drivers of variability in CO<sub>2</sub> fluxes to the increasing amplitude by separating the effects of CO<sub>2</sub> radiative forcing (climate change), CO<sub>2</sub> fertilization and N deposition, and land use change on the atmospheric CO<sub>2</sub> annual cycle amplitude over the NH subtropics, NH midlatitudes, and the NH high latitudes before and after 2100 in an extension of the high-emission RCP8.5 scenario (Meinshausen et al., 2011).

In addition to revealing potential effects of continued increases in CO<sub>2</sub> emissions, anthropogenic nitrogen, and land use change up to 2100, extending the RCP8.5 scenario to 2300 allowed us to assess the behavior of the mean annual CO<sub>2</sub> cycle in a warmer climate following stabilization of atmospheric CO<sub>2</sub> mole fraction and a shift in the terrestrial biosphere from a CO<sub>2</sub> sink to a source as shown by Randerson et al. (2015).

The questions guiding our analysis of CESM extended concentration pathway simulations are as follows:

1. Does the relative importance of drivers of the CO<sub>2</sub> amplitude trend change after 2100? For example, do we see evidence of saturation of the CO<sub>2</sub> fertilization effect or evidence of a climatic tipping point after which the CO<sub>2</sub> amplitude declines?
2. Do the regional contributions to CO<sub>2</sub> mean annual cycle trends change in response to large changes in climate?
3. Does the CO<sub>2</sub> annual cycle amplitude scale with the hemispheric carbon sink from net ecosystem productivity (NEP) as climate and atmospheric conditions evolve in the future?

The CESM provides a unique platform for exploring these questions in that it is one of the few prognostic ESMs to include coupled carbon–nitrogen biogeochemistry and diagnostic atmospheric CO<sub>2</sub> variability. This paper is organized as follows: first, we discuss the ability of the CESM to capture observed changes in the mean CO<sub>2</sub> annual cycle amplitude throughout the NH. Second, we describe how climate change, CO<sub>2</sub> fertilization and N deposition, and land use change impact the NH CO<sub>2</sub> annual cycle amplitude in the CESM before and after 2100. Third, we examine how forcing from different regions contributes to the amplitude changes

attributed to each driver. Finally, we discuss our results and provide recommendations for future analysis.

## 2 Methods

### 2.1 Model

We analyzed simulations from the Community Earth System model with coupled biogeochemistry (CESM1(BGC); Hurrell et al., 2013) to explore the role of environmental change on land–atmosphere carbon exchange. The Community Atmosphere Model (CAM, version 4; Neale et al., 2013) and the Community Land Model (CLM, version 4; Lawrence et al., 2012) were the most important components for our research, but all components of the model, including physical and biogeochemical ocean processes and sea ice processes, were interactive in the model configuration. The CAM4 was run on a  $0.94^\circ \times 1.25^\circ$  finite volume grid with 26 vertical levels. The model simulated climate conditions, including temperature, precipitation, and humidity, that provide important boundary conditions for land biogeochemistry. Moreover, the CAM4 directly simulates three-dimensional transport of atmospheric CO<sub>2</sub>, as well as separate CO<sub>2</sub> tracers derived from fossil fuel emissions, land exchange, and ocean exchange.

The CLM4 exchanged fluxes of sensible and latent heat, momentum, moisture, radiation, and terrestrial carbon with the CAM4 and was run at the same horizontal resolution. Biogeochemistry is represented in CLM4 by a prognostic carbon–nitrogen model (CLM4CN, Thornton et al., 2007) and fire model adapted from the Thonicke et al., 2001 model. We note that important high-latitude processes, such as permafrost carbon dynamics, were not simulated in the CLM4, meaning that the model may have underestimated both the seasonal dynamics of soil carbon fluxes and the long-term dynamics of permafrost melt and the subsequent radiative feedback into the climate system (Koven et al., 2015). In our analysis, we used the CLM4 NEP, defined as the difference between GPP and total respiration (autotrophic + heterotrophic), to calculate the atmospheric CO<sub>2</sub> annual cycle amplitudes described in Sect. 2.3.

### 2.2 Experiments


Three CESM simulations were run from 1850 to 2300 to separate the effects of climate change, CO<sub>2</sub> fertilization and N deposition, and land use change. The mole fraction of CO<sub>2</sub> in the atmosphere is prescribed according to the RCP8.5 and ECP8.5 scenarios described by Meinshausen et al. (2011), and it is this value that controls both radiative forcing and CO<sub>2</sub> fertilization. However, the CESM retains a separate, spatially varying CO<sub>2</sub> tracer that is a diagnostic passive tracer of land, ocean, and fossil fuel carbon fluxes; the additional carbon exported from the land and ocean to the atmosphere does not exert any radiative forcing on the climate.

The degree of coupling between CO<sub>2</sub> biogeochemistry and radiative forcing differed across the three runs. In the first simulation, denoted FullyCoupled, the imposed CO<sub>2</sub> was radiatively active, and additional anthropogenic radiative forcing resulted from prescribed CH<sub>4</sub>, chlorofluorocarbons, ozone, and aerosols. In this simulation, the increasing CO<sub>2</sub> was also biogeochemically active, meaning it contributed to CO<sub>2</sub> fertilization. Transient land use change (LUC) from agriculture and wood harvest, and land and ocean N deposition (Lamarque et al., 2010) were applied through 2100, then held at 2100 values through 2300. Keppel-Aleks et al. (2013) and Lindsay et al. (2014) provide additional descriptions of the model configuration and analyses of the FullyCoupled simulation during the 20th century. In the second simulation (NoRad), radiative forcing from CO<sub>2</sub> and other species was fixed at 1850 values, but the changing CO<sub>2</sub> mole fraction interacted with biogeochemistry via CO<sub>2</sub> fertilization. LUC and N deposition were likewise prescribed as in FullyCoupled. Randerson et al. (2015) also details the design of the FullyCoupled and NoRad (referred to as “NoCO<sub>2</sub>Forcing”) simulations through 2300. We isolated the impact of climate change on the mean annual CO<sub>2</sub> cycle by taking the difference between the FullyCoupled and NoRad simulations. The third simulation (NoLUC), was configured identically to FullyCoupled with the exception that LUC was held constant at 1850 values; therefore, LUC effects on terrestrial carbon exchange were determined from the difference between FullyCoupled and NoLUC.

Variations in fractional coverage, albedo, nutrient limitations, and surface energy fluxes among trees, grasses, and crops may enhance or oppose the effects of climate change and CO<sub>2</sub> fertilization on the atmospheric CO<sub>2</sub> mean annual cycle amplitude. These changes based on plant functional type (PFT) were approximated by prescribing transient land cover change through 2100 in FullyCoupled and NoRad based on annual fractional transition among primary vegetation, secondary vegetation, pasture (grazing land), and crops described by Hurtt et al. (2006), with CESM PFTs detailed in Lawrence et al. (2011). The crop model was inactive in the CESM simulations, and the crop PFT in Hurtt et al. (2006) data was specified as unmanaged grass (Lindsay et al., 2014). Therefore, our CESM results do not include anthropogenic influences on CO<sub>2</sub> seasonality or agricultural intensification. Moreover, these simulations were run without dynamic vegetation, which potentially damps feedbacks that could contribute to changes in the CO<sub>2</sub> annual cycle through 2300.

### 2.3 Mapping atmospheric CO<sub>2</sub> from surface fluxes

Although the CESM simulated the three-dimensional structure of atmospheric CO<sub>2</sub>, we used a pulse-response transport operator to separate imprints of CO<sub>2</sub> fluxes from different regions on the atmospheric CO<sub>2</sub> variations. The transport operator was developed using the GEOS-Chem transport model (version 9.1.2, Nassar et al., 2010). GEOS-Chem was config-



**Figure 1.** Map of stations where CO<sub>2</sub> from the pulse-response code is computed. Table 1 lists the station identifiers and locations. The GEOS-Chem regions defined in the pulse-response code (see Table 2) are shaded and grouped according to the Arctic (blue shades), boreal (orange shades), NH temperate (green shades), NH subtropical (purple shades), tropical (red shades), and SH land (yellow shades) ecoclimate regions. Greenland and Antarctica (gray shades) were excluded from the analysis. Gray circles indicate a subset of stations from which the atmospheric annual CO<sub>2</sub> amplitudes were computed from 1985 to 2013 monthly mean observations (bold stations in Table 1).

ured as in Keppel-Aleks et al. (2014) on a  $4^\circ \times 5^\circ$  horizontal grid with 47 vertical layers and was forced with meteorological fields from the 3–6 hourly Modern Era Retrospective-Analysis for Research and Applications (MERRA) reanalysis data set (Rienecker et al., 2011). A tagged  $1 \text{ Pg C month}^{-1}$  pulse was released for each of the 20 terrestrial source regions in Fig. 1 for each calendar month and was allowed to decay for 60 subsequent months. Each  $1 \text{ Pg C month}^{-1}$  pulse was distributed spatially according to monthly fluxes from the Carnegie–Ames–Stanford approach (CASA; see list of acronyms in Table 2) from Olsen and Randerson (2004).

At a given location, the magnitude and phasing of the atmospheric CO<sub>2</sub> response of the pulse depends on the characteristics of atmospheric transport (Fig. 2). For example, at Barrow (BRW) in northern Alaska, a  $1 \text{ Pg}$  pulse released in boreal North America (NBNA) in the winter months (December–January) has a large impact on atmospheric CO<sub>2</sub> during the first 1–2 months after a pulse is released (2 ppm, Fig. 2a), but more vigorous vertical mixing in the summer months reduces the imprint to 0.5 ppm. In contrast, when the pulse is released from temperate North America (ETNA, WTNA), there is a phase lag of 2–3 months (Fig. 2b, c), and when the pulse is released from the Amazon (AMZN), there is a delay in the peak response at BRW of at least 4 months (Fig. 2d). Following the 12-month period in which pulses were released, the signals were allowed to decay for 60 subsequent months, at which point CO<sub>2</sub> was well-mixed in the atmosphere (Fig. 2a–d). We then sampled GEOS-Chem at the locations of 41 NOAA cooperative CO<sub>2</sub> flask sample sites (Dlugokencky et al., 2013; Table 1, Fig. 1) for each of the 72 total months simulated. This resulted in a CO<sub>2</sub> transport operator matrix with the dimensions  $N_{\text{reg.}} \times N_{\text{obs.}} \times N_{\text{mon.}}$ .

We used monthly mean NEP from the CESM to derive atmospheric CO<sub>2</sub> from the pulse-response function. We aggregated NEP fluxes from CLM4 to the spatial scale of the 20 source regions (Fig. 1) and used matrix multiplication to propagate these fluxes to atmospheric CO<sub>2</sub>. We calculated the monthly mean CO<sub>2</sub> mole fraction at the observation sites (e.g., blue lines in Fig. 3) by summing over the instantaneous contributions from all regions and the background contributions from fluxes released during the 60 previous months to get a CO<sub>2</sub> response matrix with the dimensions  $N_{\text{obs.}} \times N_{\text{mon.}}$ . We analyzed both the CO<sub>2</sub> fields from global fluxes and the CO<sub>2</sub> patterns influenced only by larger regions representing Arctic, boreal, temperate, subtropical, tropical, and Southern Hemisphere (SH) ecosystems. We calculated the CO<sub>2</sub> annual cycle amplitude values as the peak-to-trough differences in CO<sub>2</sub> summed over each component region (e.g., the CO<sub>2</sub> annual cycle amplitude at a given station from pulses emitted from the Arctic was calculated as the peak-to-trough difference in the sum of CO<sub>2</sub> from pulses emitted by the blue regions in Fig. 1). We note that our analysis focuses on surface observations of atmospheric CO<sub>2</sub> and does not include aircraft measurements.


The advantage of the pulse-response method is that we can efficiently compute the regional contribution to changes in atmospheric CO<sub>2</sub>; it would be prohibitively expensive to run a full atmospheric transport model for each of the regions separately for 350 years. However, using this simplified transport operator introduces errors. To evaluate the pulse-response method, we show a comparison in which we have generated CO<sub>2</sub> using net ecosystem exchange (NEE), which includes fire, harvest, and land use fluxes (Fig. 4), since the land CO<sub>2</sub> tracer in the CAM4 is derived from NEE (despite using NEP

**Table 1.** Latitude band, station location, station ID, latitude, and longitude of CO<sub>2</sub> sample locations. Observations from bold stations were analyzed over the 1985–2013 period.

Latitude band	Station location	Station ID	Latitude	Longitude
NH high latitudes	<b>Alert, Nunavut</b>	<b>ALT</b>	<b>82.45</b>	<b>297.49</b>
	Ny-Ålesund, Svalbard	ZEP	78.90	11.90
	<b>Barrow, Alaska</b>	<b>BRW</b>	<b>71.30</b>	<b>203.40</b>
NH midlatitudes	Baltic Sea	BAL	55.35	17.22
	<b>Shemya Island, Alaska</b>	<b>SHM</b>	<b>52.70</b>	<b>174.10</b>
	Hohenpeissenberg, Germany	HPB	47.80	11.02
	Hegyhátsál, Hungary	HUN	46.95	16.65
	Ulaan-Uul, Mongolia	UUM	44.45	111.10
	Trinidad Head, California	THD	41.10	235.80
	Shangdianzi, China	SDZ	40.65	117.12
NH subtropics	Tae-ahn Peninsula, South Korea	TAP	36.70	126.10
	Mt. Waliguan, China	WLG	36.29	100.90
	Lampedusa, Italy	LMP	35.52	12.62
	Tudor Hill, Bermuda	BMW	32.30	295.10
	Weizmann Institute of Science Station, Negev Desert, Israel	WIS	29.97	35.06
	Izana, Tenerife, Canary Islands	IZO	28.31	343.50
	<b>Sand Island, Midway, USA</b>	<b>MID</b>	<b>28.21</b>	<b>182.62</b>
	<b>Key Biscayne, Florida</b>	<b>KEY</b>	<b>25.67</b>	<b>279.84</b>
	Lulin, Taiwan	LLN	23.47	120.87
NH tropics	<b>Mauna Loa, Hawaii</b>	<b>MLO</b>	<b>19.50</b>	<b>204.42</b>
	Pacific Ocean (15° N)	POCN15	15.00	215.00
	<b>Mariana Islands, Guam</b>	<b>GMI</b>	<b>13.39</b>	<b>144.66</b>
	Ragged Point, Barbados	RPB	13.17	300.57
	Pacific Ocean (10° N)	POCN10	10.00	211.00
	<b>Christmas Island, Rep. Kiribati</b>	<b>CHR</b>	<b>1.700</b>	<b>202.85</b>
SH tropics	Bukit Kototabang, Indonesia	BKT	−0.20	100.32
	<b>Mahé Island, Seychelles</b>	<b>SEY</b>	<b>−4.68</b>	<b>55.53</b>
	Maxaranguape, Brazil	NAT	−5.52	324.74
	<b>Ascension Island, UK</b>	<b>ASC</b>	<b>−7.97</b>	<b>345.60</b>
	Pacific Ocean (10° S)	POCS10	−10.00	199.00
	Tutuila, American Samoa	SMO	−14.25	189.44
	Pacific Ocean (20° S)	POCS20	−20.00	186.00
SH	Gobabeb, Namibia	NMB	−23.58	15.03
	<b>Easter Island, Chile</b>	<b>EIC</b>	<b>−27.16</b>	<b>250.57</b>
	Pacific Ocean (35° S)	POCS35	−35.00	180.00
	<b>Cape Grim, Tasmania, Australia</b>	<b>CGO</b>	<b>−40.68</b>	<b>144.69</b>
	Baring Head Station, NZ	BHD	−41.41	174.87
	<b>Palmer Station, Antarctica</b>	<b>PSA</b>	<b>−64.92</b>	<b>296.00</b>
	Syowa Station, Antarctica	SYO	−69.01	39.59
	<b>Halley Station, Antarctica</b>	<b>HBA</b>	<b>−75.61</b>	<b>333.79</b>
South Pole, Antarctica	SPO	−89.90	335.20	

for subsequent analyses). The errors are generally less than 2 ppm between the full transport and pulse-response calculations due to different model boundary layer schemes and atmospheric transport (Fig. 4c). We note that the largest differences were during the last century of the simulation, which was likely due to shifts in atmospheric transport in response to the dramatic climate change in the CAM4. The fact that long-term trends in transport are not simulated by the pulse-

response approach is one of the major sources of bias. In a site-by-site comparison (Fig. 3), the increasing bias through 2300 appears to be due to amplification of existing biases in the pulse-CO<sub>2</sub> compared to the full transport CO<sub>2</sub>. A second source of uncertainty is that the spatial distribution of fluxes within each region is different in CLM compared to CASA. We expect that this has a minimal impact based on results from Nevison et al. (2012), who showed that a simi-



**Figure 2.** The imprints of 1 PgC pulses emitted in individual months (colored curves) on atmospheric CO<sub>2</sub> at Barrow (BRW). Imprints on the atmosphere sampled at BRW are shown from selected regions: (a) northern boreal North America (NBNA), (b) eastern temperate North America (ETNA), (c) western temperate North America (WTNA), and (d) the Amazon (AMZN). For clarity, we plot only the first calendar year (months 13–24) after the pulses were released.

lar pulse-response code using different transport models did a reasonable job ( $r^2 = 0.8$ ) of simulating the fossil fuel influence on CO<sub>2</sub> despite that fossil fuel emissions show a vastly different spatial configuration than do ecosystem fluxes. In our analysis, we aggregate the sites into high-, mid-, subtropical, and tropical latitude belts to minimize local effects at individual sites and instead to focus on large-scale trends owing to broad patterns of changing fluxes.

We assessed the validity of ignoring ocean contributions to trends in the mean annual cycle of CO<sub>2</sub> by calculating the CO<sub>2</sub> amplitudes in the CAM land and ocean tracers. We found that the contemporary peak-to-trough amplitude in the ocean tracer averaged across our high-latitude stations was 2 ppm (in contrast to 10 ppm in the land tracer). Although both the land and ocean amplitudes grow with time, by 2300 the high-latitude ocean tracer had an amplitude of 3 ppm, only 18 % of the land amplitude for this time period.

## 2.4 Atmospheric CO<sub>2</sub> time series analysis


To place the CO<sub>2</sub> annual cycle amplitudes simulated by the CESM in the context of observations, we quantified observed and simulated CO<sub>2</sub> annual cycle amplitude at NOAA observatories before aggregating amplitudes across four latitude bands spanning 60–90° N (NH high latitudes), 40–60° N (NH midlatitudes), 20–40° N (NH subtropics), and 0–20° N (NH tropics). We identified a subset of stations in the NOAA Global Monitoring Division (Conway et al., 1994) and Scripps Institute of Oceanography (Keeling et al., 2005) networks with better than 95 % temporal coverage of monthly mean values from 1985 to 2013 (gray circles in Fig. 1). The trends at these stations were calculated iteratively as a second-order polynomial, as described by Keppel-Aleks et al. (2013). After subtracting the trend from the raw observations, we calculated the peak-to-trough amplitude ( $A^{\text{Obs}}$ ) for each calendar year in which observations existed. We then aggregated  $A^{\text{Obs}}$  from all stations within the specified latitude bands to determine a regionally averaged amplitude.

We calculated the regional CO<sub>2</sub> amplitudes for the Fully-Coupled simulation ( $A^{\text{FC}}$ ) using a nearly identical methodology. However, due to the length of the simulated time series, we detrended the data in 10-year increments. For CESM output, we used only the sampling locations with greater than 95 % temporal coverage for comparison with the observations (Fig. 1, gray circles) but aggregated amplitudes at a larger set of marine boundary layer observatories when assessing future trends (Fig. 1, black circles). Due to the flexible transport operator, we separately calculated amplitudes from NoRad ( $A^{\text{NoRad}}$ ) and NoLUC ( $A^{\text{NoLUC}}$ ) simulations and were further able to simulate only the contribution from specified ecosystem types. The contribution of climate change to the CO<sub>2</sub> mean annual cycle amplitude ( $A^{\text{Clim}}$ ) was calculated from the difference between  $A^{\text{FC}}$  and  $A^{\text{NoRad}}$ . Likewise, the LUC contribution to the annual cycle amplitude was calculated from the difference between  $A^{\text{FC}}$  and  $A^{\text{NoLUC}}$ .

## 3 Results

### 3.1 Trends in observed and modeled CO<sub>2</sub> annual cycle amplitudes

Throughout the NH, the CESM simulated both smaller mean annual cycle amplitudes and smaller trends in amplitude relative to observations (Fig. 5). The CESM underestimated the magnitudes of  $A^{\text{Obs}}$  by roughly 50 % (Fig. 5b, c), and the 16 % relative increase in the hemispheric-average amplitude between 1985 and 2013 estimated by the CESM was lower than the observed increase of 24 %. The 1985–2013 mean  $A^{\text{Obs}}$  averaged over the whole NH was 10.5 ppm, while  $A^{\text{FC}}$  was 5.8 ppm. At high latitudes, the observed 1985–



**Figure 3.** Mean annual cycles of atmospheric CO<sub>2</sub> derived from (blue curves) NEE run through the pulse-response function and (black curves) the CESM land CO<sub>2</sub> tracer for (a–d) Barrow (BRW), (e–h) Shemya Island (SHM), (i–l) Key Biscayne (KEY), and (m–p) Mauna Loa (MLO) in 1990–1999, 2090–2099, 2190–2199, and 2290–2299.

2013 amplitude value was 15.9 ppm, but only 10.8 ppm in the CESM, which is broadly consistent with Keppel-Aleks et al. (2013), who showed that the CESM1(BGC) underestimated seasonality by 25–40 % at NH surface monitoring sites. Although the CESM simulates low mean annual cycle amplitude throughout the NH, we note that many land models have a low bias in their simulated flux seasonality. For example, TRENDY land models show a 40 % deficit in the magnitude of the seasonal cycle (Zhao et al., 2016).

Consistent with the observations, the CESM simulated an increasing amplitude trend with latitude over the NH. However, the meridional gradient in the trend was too weak, leading to small absolute increases over the high latitudes. Both the modeled and observed trends in the CO<sub>2</sub> annual cycle amplitude were calculated from individual sites whose records date to 1985 (gray circles in Fig. 1). The modeled trend in the CO<sub>2</sub> annual cycle amplitude over the high latitudes was 0.05 ppm yr<sup>-1</sup> (0.43 % yr<sup>-1</sup>) for the 1985–2013 period, while the observed trend was 0.09 ppm yr<sup>-1</sup> (0.57 % yr<sup>-1</sup>). Midlatitude and subtropical trends simulated by the CESM were around 0.03 ppm yr<sup>-1</sup> (0.43 and 0.46 % yr<sup>-1</sup>, respec-


tively), and the trends in the magnitudes were closer to the observed midlatitude trend of 0.04 ppm yr<sup>-1</sup> (0.22 % yr<sup>-1</sup>) and subtropical trend of 0.05 ppm yr<sup>-1</sup> (0.61 % yr<sup>-1</sup>).

We note that the potential drivers of the amplitude increase during 1985–2013 were simulated to different levels of fidelity by the CESM: the 1985–2013 NH atmospheric temperature increase over land (1.02 K) was near the NCEP–NCAR Reanalysis (Kalnay et al., 1996) value (0.95 K), but the 2010 annual mean NH atmospheric CO<sub>2</sub> mole fraction in CESM was too high (425 ppm vs. 391 ppm). Previous analysis of CESM shows that this high bias in simulated CO<sub>2</sub> is attributable to persistent weak uptake in both land and ocean (Keppel-Aleks et al., 2013; Long et al., 2013).

## 3.2 Future CO<sub>2</sub> annual cycle amplitude changes

### 3.2.1 Total amplitude changes


Given the weak atmospheric CO<sub>2</sub> seasonal amplitude response in the CESM for the contemporary period, we examined the response of flux seasonality to stronger forc-



**Figure 4.** CO<sub>2</sub> annual cycle amplitudes in the FullyCoupled simulation derived from (a) the CESM land CO<sub>2</sub> tracer and (b) running NEE from the CLM4 through the pulse-response code. (c) The difference between the land tracer and NEE-derived CO<sub>2</sub> annual cycle amplitudes.

ings in the FullyCoupled simulation run to 2300. Both near-surface atmospheric temperature and the mean atmospheric CO<sub>2</sub> mole fraction increased markedly in the CESM (Fig. 6a, b). The accumulated mean NH atmospheric CO<sub>2</sub> mole fraction (Fig. 6a, solid black line) increased from approximately 320 to 2350 ppm between 1950 and 2300. In the NoRad simulation, where radiative forcing was held fixed at 1850 levels, the atmospheric mole fraction followed a similar pattern of increase but plateaued at a lower mean value by 2200. We note that the atmospheric CO<sub>2</sub> mole fraction values were diagnostic only, and the biogeochemical and radiative processes in the CESM responded to the lower mole fraction values prescribed according to the ECP8.5 forcing scenario indicated by the dashed black line in Fig. 6a.

In the FullyCoupled simulation, the increases in CO<sub>2</sub> and other radiative forcing agents resulted in a 6 K temperature increase by 2100 and an 11 K temperature increase by 2300 relative to the 1950–1959 baseline mean (Fig. 6b). Temperature in the NoRad simulation only increased by ~1.5 K through 2300, which can be traced to changes in albedo and surface energy balance. For small temperature changes and high levels of CO<sub>2</sub> fertilization, the NoRad experiment was able to maintain a steady carbon sink of 4–5 Pg C yr<sup>-1</sup> between 2100 and 2300 (Fig. 6c). In contrast,




**Figure 5.** (a) Observed atmospheric CO<sub>2</sub> annual cycle amplitudes ( $A^{\text{Obs}}$ ), (b) FullyCoupled atmospheric CO<sub>2</sub> annual cycle amplitudes ( $A^{\text{FC}}$ ), and (c) the difference between  $A^{\text{Obs}}$  and  $A^{\text{FC}}$  during 1985–2013.

the sink in the FullyCoupled simulation reached a maximum of 4.8 Pg C yr<sup>-1</sup> by 2120 and then declined to 3–4 Pg C yr<sup>-1</sup> in the last 100 years of the simulation, suggesting that the extreme climate change in this simulation reduced the efficiency of the global terrestrial sink with time. LUC partly offset the weakening of the land carbon sink due to climate change (likely as a result of increased crop cover prescribed in the FullyCoupled simulation), enhancing net terrestrial carbon uptake by up to 2.8 Pg C yr<sup>-1</sup> after 2100.

The NH mean CO<sub>2</sub> annual cycle amplitude increased by 3.4 ppm (65 %) by 2100 and 5.0 ppm (96 %) by 2300 from the 1950–1959 baseline in the FullyCoupled simulation (Table 3). Consistent with the observed annual cycle amplification, the magnitudes of the  $A^{\text{FC}}$  increases strengthened with increasing latitude (Fig. 7).  $A^{\text{FC}}$  increases between 1950 and 2300 ranged from 1.4 ppm (56 %) over the NH tropics to 10.6 ppm (122 %) over the high latitudes (Table 3). Peak  $A^{\text{FC}}$  magnitudes occurred between 2230 and 2250 and ranged from 4.2 ppm over the NH tropics to 19.5 ppm over the high latitudes (Table 4).

Given the weak baseline seasonal exchange in the CESM, simulated CO<sub>2</sub> annual cycle amplitudes did not approach the contemporary mean observed mean  $A^{\text{Obs}}$  (2009–2013) until about 2240 in the NH high latitudes (Fig. 7a, black filled square, Table 4). Over the NH midlatitudes, subtropics, and tropics, peak  $A^{\text{FC}}$  occurred by 2240, but values were still





**Figure 6.** (a) Annual mean atmospheric CO<sub>2</sub>, (b) annual mean bottom-level atmospheric *T* over land, and (c) annual mean NEP averaged over the NH (0–90° N) in the CESM FullyCoupled (solid black curves), NoRad (blue curves), and NoLUC (magenta curves) simulations. Negative NEP indicates net annual CO<sub>2</sub> uptake by the land surface. Values are filtered using a 10-year running median. The CO<sub>2</sub> mole fraction values in (a) result from the net contributions of land, ocean, and fossil fuel tracers calculated from NEE as described in Sect. 2.3, and they differ from the atmospheric CO<sub>2</sub> mixing ratio that is prescribed according to the ECP8.5 scenario (dashed black curve).

0.4–2.5 ppm below current  $A^{Obs}$ . In the tropics, the discrepancy between CO<sub>2</sub> seasonality inferred only from simulated NEP and from observations reflects the nontrivial contributions of ocean and fossil fuel fluxes to the CO<sub>2</sub> annual cycle (Randerson et al., 1997, Table 4 and Lindsay et al., 2014, Fig. 15). When high-latitude CO<sub>2</sub> amplitudes reached peak values, climate change and nonradiative forcing from CO<sub>2</sub> fertilization and N deposition each contributed about half of the 9.6 ppm increase relative to the 1950–1959 baseline (Fig. 8). At this point, Arctic temperatures were 16 K higher than the baseline, midlatitude temperatures were 11 K higher (Fig. 7b, c), and the mean NH atmospheric CO<sub>2</sub> mole fraction was approximately 2330 ppm (Fig. 6a).

The relationship between the atmospheric CO<sub>2</sub> annual cycle amplitude and the NH annual net land carbon sink changed over the course of the FullyCoupled simulation. Several recent papers have hypothesized that the mean atmospheric CO<sub>2</sub> annual cycle may be a diagnostic of net terrestrial carbon uptake since these variables tend to correlate positively in model simulations (e.g., Ito et al., 2016). While we found that the NH net land carbon sink and atmo-

**Table 2.** GEOS-Chem regions that comprise the ecoclimate regions shown in Fig. 1.

Ecoclimate region	Pulse region	Acronym
Arctic	Northern boreal North America	NBNA
	Eastern boreal Asia	EABA
	Western boreal Asia	WEBA
Boreal	Southern boreal North America	SBNA
	Southern boreal Asia	SOBA
NH temperate	Eastern temperate North America	ETNA
	Western temperate North America	WTNA
	Europe	EURO
	Central Asia	CNAS
NH subtropics	Central America	CNAM
	Sahara and Middle East	SAME
	Southeast Asia	SEAS
NH tropics	Amazon	AMZN
	Middle Africa	MDAF
	African rainforest	AFRF
	Equatorial Asia	EQAS
SH land	Eastern South America	EASA
	Western South America	WESA
	South Africa	SOAF
	Australia	AUST
Polar	Greenland	GNLD
	Antarctica	ATCA

spheric CO<sub>2</sub> annual cycle amplitude were positively correlated through 2100 (Fig. 9), the net land carbon sink began to decrease before the CO<sub>2</sub> annual cycle amplitude decreased. In the FullyCoupled simulation, the NH decadal mean CO<sub>2</sub> annual cycle amplitude peaked near 10 ppm by 2240 (Table 4). However, the magnitude of the NH land sink began to decrease (Fig. 6c), likely reflecting enhanced subtropical respiration. This underscores that while amplification of the CO<sub>2</sub> annual cycle may reflect enhanced seasonality of land carbon uptake, it does not necessitate enhanced annually integrated land carbon uptake. Moreover, in the 23rd century, both the CO<sub>2</sub> annual cycle amplitude and the NH net land carbon sink declined, but there was evidence of hysteresis between the two quantities. In the latter part of the simulation when both the amplitude and net land carbon sink declined, the slope of their linear relationship was shallower than when both quantities were increasing before 2100. We hypothesize that the change in the relationship between the CO<sub>2</sub> annual cycle amplitude and the NH net land carbon sink resulted from respiration catching up to increased land carbon uptake after atmospheric CO<sub>2</sub> concentration, and thus the fertilization effect, leveled off in the last 100 years of the FullyCoupled simulation.

**Table 3.** Cumulative changes in mean CO<sub>2</sub> annual cycle amplitudes (ppm) from the 1950–1959 baseline in the FullyCoupled simulation, and the contributions of nonradiative forcing (CO<sub>2</sub> fertilization and N deposition), climate change, and LUC to the FullyCoupled amplitude changes averaged over the NH high latitudes, midlatitudes, subtropics, tropics, and the NH.

Year	FullyCoupled			NoRad			Climate change			LUC		
	2100	2200	2300	2100	2200	2300	2100	2200	2300	2100	2200	2300
High latitudes	6.4	9.6	10.6	2.4	4.5	5.3	4.0	5.1	5.2	−1.1	−1.3	−1.7
Midlatitudes	5.2	8.3	8.9	3.3	5.7	6.8	1.9	2.6	2.1	−0.9	−0.7	−1.3
Subtropics	2.8	3.6	3.4	1.8	3.0	3.5	1.0	0.6	−0.1	−0.7	−0.9	−0.9
NH tropics	1.0	1.5	1.4	0.9	1.5	1.8	−0.1	−0.1	−0.3	−0.5	−0.4	−0.4
NH	3.4	4.7	5.0	2.0	3.4	4.1	1.4	1.3	1.0	−0.6	−0.7	−0.9

**Table 4.** Observed and FullyCoupled 2009–2013 mean CO<sub>2</sub> annual cycle amplitudes and corresponding decades when peak FullyCoupled amplitudes occurred in each latitude band shown in Fig. 5a.

Latitude	A <sup>Obs</sup> (ppm) 2009–2013	A <sup>FC</sup> (ppm) 2009–2013	Mean A <sup>FC</sup> (ppm) during peak decade	Center year of decade of peak A <sup>FC</sup>
High latitude	17.3	11.3	19.5	2241
Midlatitude	19.5	8.7	17.8	2236
Subtropics	9.3	5.9	9.0	2236
NH tropics	6.7	3.7	4.2	2241
NH mean	11.3	6.2	10.4	2236

### 3.3 Contributions of nonradiative forcing, climate change, and LUC to amplitude trends


#### 3.3.1 Effects of nonradiative forcing from CO<sub>2</sub> fertilization and N deposition

In the CESM, CO<sub>2</sub> fertilization was the largest driver of CO<sub>2</sub> annual cycle amplification through 2300 over much of the NH (Fig. 8a), contributing 4.1 to the 5.0 ppm increase in NH A<sup>FC</sup> ( $\Delta A^{\text{FC}}$ ) between 1950 and the end of the 23rd century (Table 3). Results from Devaraju et al. (2016) suggest that global net primary productivity (NPP) is influenced equally by CO<sub>2</sub> fertilization and N deposition over the historical period in the CESM. Therefore, trends in the CO<sub>2</sub> mean annual cycle amplitude likely responded to both drivers prior to 2100. While we cannot fully separate the influence of CO<sub>2</sub> fertilization and N deposition given the experimental design, N deposition was held fixed at 2100 values for the last 200 years of the simulations; thus, we expect that amplitude trends after 2100 mainly reflect enhanced CO<sub>2</sub> fertilization.

The amplitude increase owing to CO<sub>2</sub> fertilization and N deposition (NoRad simulation) originated mainly from NH temperate regions (Fig. 10a), which accounted for 2.2 ppm (53 %) of  $\Delta A^{\text{NoRad}}$  at the end of the 23rd century (Table 3). Boreal regions (Fig. 11a) made the second greatest contribution (32 %, 1.3 ppm) to NH  $\Delta A^{\text{NoRad}}$ . The remaining 15 % (0.6 ppm) of the NH  $\Delta A^{\text{NoRad}}$  increase came from the Arctic (Fig. 12a) or subtropical and tropical ecosystems. Temperate ecosystems had the largest response to CO<sub>2</sub> fertilization

and N deposition. Consistent with this large temperate response, the increase in A<sup>FC</sup> from nonradiative forcing was largest over the NH midlatitudes (Table 3). The effects of CO<sub>2</sub> fertilization and N deposition in boreal ecosystems contributed another 25 % to the high-latitude CO<sub>2</sub> A<sup>FC</sup> increase, 22 % to the midlatitude CO<sub>2</sub> A<sup>FC</sup> increase, and 60 % to the subtropical A<sup>FC</sup> increase.

In the CESM, the impact of CO<sub>2</sub> fertilization on the amplitude trend roughly scales to the magnitude of overall GPP, consistent with hypotheses from Tans et al. (1990) and Schimel et al. (2015) that the fertilization effect on the land carbon sink is proportional to productivity. Thus, CO<sub>2</sub> fertilization and N deposition effects on A<sup>FC</sup> were smallest in the Arctic, the region with the smallest GPP for the contemporary period. Figure 13 shows that nonradiative forcing from boreal and temperate regions together constituted at least 35 % of the increase in high-latitude, midlatitude, and subtropical A<sup>FC</sup> from the beginning of the 21st century through the end of the 23rd century. Furthermore, temperate CO<sub>2</sub> fertilization was the primary driver of midlatitude CO<sub>2</sub> annual cycle amplification in all periods and subtropical amplification from 2050 onward. In temperate and boreal regions, where CO<sub>2</sub> fertilization was the dominant driver of increases in the mean CO<sub>2</sub> annual cycle amplitude, increases in GPP seasonality outpaced increase in respiration. For example, in eastern temperate North America (ETNA), FullyCoupled GPP seasonal amplitudes increased from 6.8 Pg C in 1950 to 11 Pg C in 2250, while heterotrophic respiration (HR) amplitudes increased from 0.85 to 1 Pg C, and autotrophic res-




**Figure 7.** (a) From 1950–1959 to 2291–2300, 10-year moving averages of FullyCoupled atmospheric CO<sub>2</sub> annual cycle amplitude ( $A^{FC}$ ). The black square indicates the decade in which the CESM reached observed CO<sub>2</sub> annual cycle amplitudes averaged over 2009–2013 in the NH high latitudes (Table 4). Open squares indicate the decades when peak amplitudes occurred. In the NH midlatitudes, subtropics, and tropics, peak amplitudes did not reach present observed values. (b) Atmospheric temperatures over land in the FullyCoupled simulation ( $T^{FC}$ ). (c) The change in atmospheric temperature with respect to the 1950–1959 mean caused by climate change ( $\Delta T^{Clim}$ ).


piration (AR) amplitudes increased from 4 to 7.6 Pg C. The strong fertilization effect on the amplification of the NH CO<sub>2</sub> annual cycle is surprising given that nitrogen limitation in the CLM4 produced weaker fertilization in the CESM compared to other CMIP5 models (Thornton and Zimmermann, 2007; Piao et al., 2013; Peng and Dan, 2015).

### 3.3.2 Climate change effects

During the early part of the simulation, boreal (Fig. 11b) and Arctic (Fig. 12b) climate change drove high-latitude atmospheric CO<sub>2</sub> annual cycle amplification, increasing  $A^{Clim}$  by 5 ppm by 2200 (Fig. 13a). After 2200, CO<sub>2</sub> fertilization and climate change contributed nearly equally to high-latitude CO<sub>2</sub> annual cycle amplification. The increase in high-latitude  $A^{Clim}$  largely reflected warmer growing season temperatures that led to larger peak GPP values and a longer growing season. In the pulse regions that comprised the Arctic ecoclimate regions, annual mean GPP continued to increase with temperature ( $r = 0.7 \text{ PgC K}^{-1}$ ,  $r^2 = 0.99$ ) until annual mean near-surface air temperatures surpassed 284 K.




**Figure 8.** Changes in the decadal mean atmospheric CO<sub>2</sub> annual cycle amplitudes from 1950–1959 values from (a) CO<sub>2</sub> fertilization and N deposition ( $\Delta A^{NoRad}$ ), (b) climate change ( $\Delta A^{Clim}$ ), and (c) land use change ( $\Delta A^{LUC}$ ) from all land regions in each latitude bin. Note that the color scale in (c) is different from the scale in (a, b).




**Figure 9.** NH mean CO<sub>2</sub> annual cycle amplitude (ppm) as a function of annual mean NH NEP (PgC). NEP < 0 indicates net carbon uptake by land, and circles are shaded by simulation year. The relationship between amplitude and net land carbon exchange shows evidence of hysteresis, with different slopes before and after 2100.


By 2100, Arctic and boreal climate change increased high-latitude  $A^{FC}$  by about 1.3 and 2.5 ppm (39%) from the 1950–1959 mean, respectively, outweighing the combined contributions of boreal and temperate nonradiative forcing (2.2 ppm, 23%) (Fig. 13a). However, the effects of temperate and boreal CO<sub>2</sub> fertilization outweighed climate change ef-



**Figure 10.** As in Fig. 8 but for (a) CO<sub>2</sub> fertilization + N deposition, (b) climate change, and (c) LUC from NH temperate regions. Note that the color scales are different in each panel.



**Figure 12.** As in Fig. 8 but for (a) CO<sub>2</sub> fertilization + N deposition, (b) climate change, and (c) LUC from the Arctic. Note that the color scales are different in each panel.




**Figure 11.** As in Fig. 8 but for (a) CO<sub>2</sub> fertilization + N deposition, (b) climate change, and (c) LUC from boreal regions.

fects on CO<sub>2</sub> annual cycle amplification at the end of the simulation. CO<sub>2</sub> fertilization and N deposition added 5.3 ppm (62 %) to high-latitude, 6.8 ppm (81 %) to midlatitude, and 3.5 ppm (69 %) to the subtropical base period A<sup>FC</sup>. It is worth noting that climate change effects from lower latitudes made up most of the high-latitude and subtropical residuals after 2000 and most of the midlatitude residual after 2200 (gray bars in Fig. 13).

### 3.3.3 LUC effects

Land use change in the CESM decreased the seasonality of terrestrial CO<sub>2</sub> exchange (Figs. 8c, 10c–12c). Between 1850 (the year used for PFT fraction boundary conditions in the NoLUC simulation) and 2100, crop cover increased at the expense of grass and tree cover in the NH, reducing the seasonal amplitude of NH mean GPP for the duration of the simulation. As a result, LUC decreased the hemisphere-average atmospheric CO<sub>2</sub> annual cycle amplitude by 0.9 ppm (17 %) from the 1950–1959 baseline in 2300 (Table 3).

This finding contrasts recent results suggesting that agriculture intensification contributes significantly to positive trends in the mean annual cycle amplitude (Gray et al., 2014; Zeng et al., 2014) and may reflect that croplands were treated as unmanaged grasslands in the CESM. Zeng et al. (2014) suggests that up to 45 % of the observed trend may be due to land use practices, with the remainder partitioned roughly equally between climate and fertilization effects. We note




**Figure 13.** Contributions of temperate, boreal, and Arctic combined CO<sub>2</sub> fertilization and N deposition (NoRad), climate change (Climate), land use change (LUC) to the change in the FullyCoupled mean atmospheric CO<sub>2</sub> annual cycle amplitude from 1950–1959 values ( $\Delta A^{FC}$ , black circles) in the decades ending in 2000, 2050, 2100, 2150, 2200, 2250, and 2300 over (a) the NH high latitudes, (b) the NH midlatitudes, and (c) the NH subtropics. Residual values are the sums of the contributions of LUC, climate change, CO<sub>2</sub> fertilization, and N deposition in the subtropical, tropical, and SH GEOS-Chem land regions. Negative values indicate that the forcing decreased the atmospheric CO<sub>2</sub> annual cycle amplitude. Units are ppm.

that future LUC and land management will affect on the CO<sub>2</sub> annual cycle, but that the overall impact the CO<sub>2</sub> annual cycle will depend on economic and social drivers in addition to climate feedbacks.

### 3.3.4 Changes in growing season length


The growing season, defined as months with negative NEP (net terrestrial carbon uptake), increased for all NH terrestrial regions by about 1 month. The overall lengthened growing seasons accounted for 1–1.3 % yr<sup>-1</sup> of the high-latitude net terrestrial carbon uptake after 2050 and up to 5 % yr<sup>-1</sup> of the midlatitude terrestrial carbon uptake after 2100. Thus, while this is an important contributor, it is secondary to increased midsummer GPP.

The driver of the increased growing season length was different for different regions. For regions north of 30° N, climate change was the driver of increased growing season length, with boreal and temperate growing seasons increasing by 1 month after 2100 (Fig. 14a). Climate change extended the growing season for an additional month in the fall in the Arctic and facilitated an earlier start to the midlatitude growing season in the spring (Fig. 14c).



**Figure 14.** Growing season length (10-year running median of months with NEP < 0) in each latitude band in the (a) FullyCoupled and (b) NoRad simulations. (c) The contribution of climate change to growing season length.

CO<sub>2</sub> fertilization increased the growing season length in the subtropics (Fig. 14b) by increasing the water use efficiency (WUE, calculated as the ratio of canopy transpiration to GPP). For example, in subtropical Central America (Fig. 15), the period of net carbon uptake increased by 1 month on average in the NoRad simulation (Fig. 15a, magenta curve). In contrast, climate change altered the phasing of the mean annual cycle in the FullyCoupled simulation, with the growing season onset occurring earlier (Fig. 15a, black curve). GPP declined later in the FullyCoupled simulation despite WUE increases (Fig. 15b), suggesting that CO<sub>2</sub> fertilization effects were counteracted by climate-change-driven reductions in total soil water (Fig. 15c, d) and increases in temperature in the FullyCoupled simulation. Conversely, during the latter part of the NoRad simulation, the growing season extended into June, for which WUE increased from around 2 g C kg<sup>-1</sup> H<sub>2</sub>O in the 20th century to around 6 g C kg<sup>-1</sup> H<sub>2</sub>O by 2300, and GPP likewise increased from 0.2 to 0.45 Pg C month<sup>-1</sup>, despite essentially no change in soil water content (Fig. 15e, g). Increased GPP outweighed changes in respiration and led to a shift toward net land carbon uptake during June (Fig. 15f, g).



**Figure 15.** (a) Mean annual cycles of NEP in Central America averaged over 1950–1959 and 2290–2299 in the FullyCoupled (black dashed and solid lines) and NoRad (magenta dashed and solid lines) simulations. (b, e) GPP as a function of WUE, (c, f) NEP as a function of WUE, and (d, g) NEP as a function of volumetric soil water in the top layer (0.71 cm depth) in the month of June over Central America in the (b–d) FullyCoupled simulation and (e–g) the NoRad simulation. NEP < 0 indicates net carbon uptake by land.

## 4 Discussion and conclusions

### 4.1 CO<sub>2</sub> annual cycle amplitude trends and applications for model evaluation

By analyzing CESM simulations run to 2300 with the ECP boundary conditions, we identified notable carbon cycle interactions that were not apparent before 2100, the nominal end date for CMIP5 runs. We found that the mean NH atmospheric CO<sub>2</sub> annual cycle amplitude increased by 65 % from 1950 to 2100 and by an additional 30 % by 2300 in the CESM1(BGC). Despite significant changes in climate and atmospheric CO<sub>2</sub> mole fractions in the extended simulation, the sensitivity of the CO<sub>2</sub> annual cycle amplitude to climate and fertilization drivers was remarkably similar both before and after 2100. We likewise found that regional contributions to the NH CO<sub>2</sub> seasonal amplitude trend were generally consistent throughout the simulation, in response to the second question posed in the introduction. CO<sub>2</sub> fertilization was the dominant driver of CO<sub>2</sub> annual cycle amplification for most of the NH, with the notable exception of Arctic ecosystems where temperature increases drove amplification prior to 2100. GPP leveled off in boreal high latitudes when annual mean temperatures exceeded 284 K, which contributed to small amplitude declines in the 23rd century. CO<sub>2</sub> fertilization increased the CO<sub>2</sub> annual cycle amplitude globally for the duration of the simulation even after the growth rate of CO<sub>2</sub> slowed during the 23rd century, suggesting that the

CO<sub>2</sub> fertilization effect had not saturated in the CESM when CO<sub>2</sub> mole fractions were around 2000 ppm.

We saw evidence of hysteresis in the relationship between carbon cycle diagnostics that was only apparent after 2100, with respect to our third research question. For example, we found that the relationship between the NH mean CO<sub>2</sub> annual cycle amplitude and the NH net land carbon sink changed after 2100. A strong positive relationship between the CO<sub>2</sub> annual cycle amplitude and terrestrial carbon sink, which was previously noted by Ito et al. (2016), became decoupled in the mid-22nd century when sink strength began to decrease after 2100, while the CO<sub>2</sub> annual cycle amplitude continued to increase. When the NH CO<sub>2</sub> annual cycle amplitude began to decrease in the 23rd century, the correlation between the amplitude and land carbon sink weakened considerably compared to the relationship between the two quantities during the 20th and 21st century correlations.

Our analysis of extended concentration pathway simulations suggests several approaches for how to leverage multi-decadal atmospheric CO<sub>2</sub> observations for model evaluation. For example, consideration of only the trend in the hemispheric mean annual cycle, rather than trends in the latitudinally resolved mean annual cycle of CO<sub>2</sub>, may obscure model biases. The Extended Concentration Pathway simulations run in the CESM with coupled biogeochemistry show that the NH mean annual cycle amplitude of atmospheric CO<sub>2</sub> increased by 16 % from 1985 to 2013. The relative increase was in line with the observed 24 % increase over the

NH during the same time period. However, the spatial pattern of the relative amplitude change was more uniform throughout the NH than observed. Furthermore, the trend in the absolute magnitude of the amplitude at high latitudes was about half of the observed trend (0.05 vs. 0.09 ppm yr<sup>-1</sup>). This result highlights the importance of considering meridionally resolved atmospheric CO<sub>2</sub> data that explicitly account for the role of transport since analysis of only hemispheric spatial patterns obscures incorrect spatial patterns simulated by the CESM. Analysis of the long-term CESM simulations are suggestive of patterns we might see in ongoing monitoring of the CO<sub>2</sub> annual cycle amplitude. CESM simulations show that the major drivers of the mean annual cycle amplification impact differential imprints on atmospheric CO<sub>2</sub> in different latitude bands. For example, CO<sub>2</sub> fertilization leaves the largest imprint in both absolute and relative terms on midlatitude CO<sub>2</sub>, whereas climate change may amplify high-latitude CO<sub>2</sub> while having a near-neutral impact on CO<sub>2</sub> annual cycle amplitudes south of 60° N (Fig. 13). These fingerprints may be useful for developing hypotheses regarding observed trends and determining future observational strategies to monitor carbon–climate feedbacks.

Several recent papers have considered how the amplitude of NH net carbon exchange has changed over the historical period in different categories of prognostic models. Ito et al. (2016) analyzed MsTMIP terrestrial ecosystem models to determine how atmospheric CO<sub>2</sub>, climate change, and land use affect the NH flux amplitude for the historical period, and Zhao et al. (2016) analyzed the net terrestrial flux to the atmosphere in TRENDY models. Both of these studies find that CO<sub>2</sub> fertilization is the strongest driver of increasing ecosystem productivity and thus the amplitude of the net carbon exchange in the NH, consistent with our results. A significant difference between the approach used by these papers and our study is that they consider the net flux amplitude, whereas we propagate fluxes using an atmospheric transport operator to determine the influence on latitudinally resolved atmospheric CO<sub>2</sub> fields. Given the importance of atmospheric transport to the mean annual cycle of atmospheric CO<sub>2</sub> (e.g., Barnes et al., 2016), and the small biases induced by the simplified pulse-response transport operator, we recommend that future studies explicitly simulate the full atmospheric CO<sub>2</sub> field.

#### 4.2 Uncertainties and future model needs

The mean annual cycle of atmospheric CO<sub>2</sub> is a first-order diagnostic of terrestrial carbon exchange and its trend with time integrates a range of environmental and human factors (Randerson et al., 1997). An active area of carbon cycle research is determining the extent to which coupled ESMs provide predictive skill for future carbon–climate feedbacks. We note that many of the methods used to evaluate the carbon cycle in ESMs rely on benchmarking short-term responses to either seasonal or interannual climate variability (Keppel-

Aleks et al., 2013), or they rely on extrapolating future behavior based on some mechanistic link between short-term and long-term variability (Cox et al., 2013; Hoffman et al., 2014). The changing CO<sub>2</sub> annual cycle provides a unique opportunity to gauge a model's sensitivity to slow-varying climate and environmental changes since we observed large trends in this quantity over the instrumental record (Graven et al., 2013).

However, biases in seasonality in the CESM1(BGC) that lead to smaller increases in NH atmospheric CO<sub>2</sub> seasonal amplitudes in the CESM compared to observations during 1950–2010 prompt further model development. Moreover, the relatively smooth response to extreme changes in temperature and CO<sub>2</sub> suggests that the CESM may not parameterize processes that could cause nonlinear carbon cycle feedbacks. CESM1(BGC) did not include parameterizations for permafrost carbon dynamics, which have since been improved in CESM (Koven et al., 2015). The lack of permafrost dynamics likely has a large impact on CO<sub>2</sub> annual cycle trends, especially later in the simulation when global mean temperature has increased by over 10 K in the Fully-Coupled simulation. Thus, short soil carbon turnover time in CLM4 may have contributed to the amplitude underestimation by damping ecosystem respiration outside of the growing season (Keppel-Aleks et al., 2013; Koven et al., 2013) and would affect both baseline values and trends. Ongoing model development in the CESM includes improved representation of permafrost carbon (Koven et al., 2015), and thus future model configurations will provide an improved tool for investigating a process that may provide one of the tipping points we identified in our key science questions.

In addition, Forkel et al. (2016) found that interaction between climate change and changes in vegetation cover over northern high latitudes was the primary driver of the north–south gradient in observed NH atmospheric CO<sub>2</sub> seasonal amplitude trends, indicating that the lack of dynamic vegetation in the CLM4 likely contributes to underestimation of the seasonal amplitudes by the CESM. Tree cover is expected to expand further northward with climate change (e.g., Lloyd, 2005), which may contribute to the long-term increase in NEP flux amplitude within high-latitude ecosystems. In contrast, drying at lower latitudes may lead to replacement of trees with grasses and subsequent decreases in NEP amplitude. An ecosystem demography version (CLM-ED) that will permit successional patterns in response to environmental change is presently under development. We consider the documentation of trends in the static-vegetation configuration presented in this paper to be a crucial first step toward eventually determining the sensitivity of land–atmosphere biogeochemical couplings in more sophisticated future configurations of the CESM model.

Development is also underway to represent irrigation and fertilization in croplands in future versions of the CLM. Gray et al. (2014) and Zeng et al. (2014) suggest that agricultural amplification, facilitated by irrigation and fertilization, may

be an important driver of the observed mean annual cycle trend. In the CESM, however, crop cover is currently treated as unmanaged grass. Thus, these agricultural practices are not explicitly modeled and do not mitigate the reduction in tree cover in the FullyCoupled simulation. Our results indicate that explicit consideration of human modifications may be necessary for prognostic models both to match observations and to provide realistic predictions of future changes. After accounting for land management contributions to the amplitude increase, the sensitivity of the CO<sub>2</sub> amplitude to natural factors may be reasonable. Our results suggest that model development focused on human modification of carbon fluxes (e.g., by agriculture, Levis et al., 2014, or by disturbance, Kloster et al., 2012) may facilitate improved comparison both of mean behavior and trends.

*Data availability.* The post-processed monthly CESM data may be obtained by submitting a request to NCAR CISL at <https://www2.cisl.ucar.edu/resources/storage-and-file-systems/using-data-access-nodes>.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* This research was supported in part by the Biogeochemistry – Climate Feedbacks Scientific Focus Area (SFA), which is sponsored by the Regional and Global Climate Modeling (RGCM) Program in the Climate and Environmental Sciences Division (CESD) of the Biological and Environmental Research Program in the US Department of Energy Office of Science.

The National Center for Atmospheric Research (NCAR) is sponsored by the National Science Foundation.

Computing resources were provided by the Climate Simulation Laboratory at NCAR's Computational and Information Systems Laboratory (CISL), sponsored by the National Science Foundation and other agencies.

We gratefully acknowledge Ernesto Munoz for running and processing the CESM simulations.

We thank P. Peylin and one anonymous referee for their thoughtful comments that greatly improved the paper.

Edited by: L. Bopp

Reviewed by: P. Peylin and one anonymous referee

## References

- Angert, A., Biraud, S., Bonfils, C., Henning, C. C., Buermann, W., Pinzon, J., Tucker, C. J., and Fung, I.: Drier summers cancel out the CO<sub>2</sub> uptake enhancement induced by warmer springs, *P. Natl. Acad. Sci. USA*, 102, 10823–10827, 2005.
- Bacastow, R. B., Keeling, C. D., and Whorf, T. P.: Seasonal amplitude increase in atmospheric CO<sub>2</sub> concentration at Mauna Loa,

Hawaii, 1959–1982, *J. Geophys. Res.-Atmos.*, 90, 10529–10540, doi:10.1029/JD090iD06p10529, 1985.

- Barichivich, J., Briffa, K. R., Myneni, R. B., Osborn, T. J., Melvin, T. M., Ciais, P., Piao, S., and Tucker, C.: Large-scale variations in the vegetation growing season and annual cycle of atmospheric CO<sub>2</sub> at high northern latitudes from 1950 to 2011, *Glob. Change Biol.*, 19, 3167–3183, doi:10.1111/gcb.12283, 2013.
- Barnes, E. A., Parazoo, N., Orbe, C., and Denning, A. S.: Isentropic transport and the seasonal cycle amplitude of CO<sub>2</sub>, *J. Geophys. Res.-Atmos.*, 121, 8106–8124, doi:10.1002/2016JD025109, 2016.
- Buermann, W., Bikash, P. R., Jung, M., Burn, D. H., and Reichstein, M.: Earlier springs decrease peak summer productivity in North American boreal forests, *Environ. Res. Lett.*, 8, 024027, doi:10.1088/1748-9326/8/2/024027, 2013.
- Claussen, M., Mysak, L., Weaver, A., Crucifix, M., Fichet, T., Loutre, M.-F., Weber, S., Alcamo, J., Alexeev, V., Berger, A., Calov, R., Ganopolski, A., Goosse, H., Lohmann, G., Lunkeit, F., Mokhov, I., Petoukhov, V., Stone, P., and Wang, Z.: Earth system models of intermediate complexity: closing the gap in the spectrum of climate system models, *Clim. Dynam.*, 18, 579–586, doi:10.1007/s00382-001-0200-1, 2002.
- Cleveland, W. S., Freeny, A. E., and Graedel, T. E.: The seasonal component of atmospheric CO<sub>2</sub>: Information from new approaches to the decomposition of seasonal time series, *J. Geophys. Res.-Oceans*, 88, 10934–10946, doi:10.1029/JC088iC15p10934, 1983.
- Conway, T. J., Tans, P. P., Waterman, L. S., Thoning, K. W., Kitzis, D. R., Masarie, K. A., and Zhang, N.: Evidence for interannual variability of the carbon cycle from the National Oceanic and Atmospheric Administration/Climate Monitoring and Diagnostics Laboratory global air sampling network, *J. Geophys. Res.-Atmos.*, 99, 22831–22855, 1994.
- Cox, P. M., Pearson, D., Booth, B. B., Friedlingstein, P., Huntingford, C., Jones, C. D., and Luke, C. M.: Sensitivity of tropical carbon to climate change constrained by carbon dioxide variability, *Nature*, 494, 341–344, doi:10.1038/nature11882, 2013.
- Dettinger, M. D. and Ghil, M.: Seasonal and interannual variations of atmospheric CO<sub>2</sub> and climate, *Tellus B*, 50, 1–24, 1998.
- Devaraju, N., Bala, G., Caldeira, K., and Nemani, R.: A model based investigation of the relative importance of CO<sub>2</sub>-fertilization, climate warming, nitrogen deposition and land use change on the global terrestrial carbon uptake in the historical period, *Clim. Dynam.*, 47, 173–190, doi:10.1007/s00382-015-2830-8, 2016.
- Dlugokencky, E. J., Lang, P. M., Masarie, K., Crotwell, A. M., and Crotwell, M. J.: Atmospheric carbon dioxide dry air mole fractions from the NOAA ESRL carbon cycle cooperative global air sampling network, 1968–2012, Version: 2013-08-27, available at: [ftp://afsp.cmdl.noaa.gov/data/trace\\_gases/co2/flask/surface/](ftp://afsp.cmdl.noaa.gov/data/trace_gases/co2/flask/surface/) (last access: 9 March 2017), 2013.
- Enting, I. G. and Mansbridge, J. V.: Seasonal sources and sinks of atmospheric CO<sub>2</sub>: Direct inversion of filtered data, *Tellus B*, 41, 111–126, doi:10.1111/j.1600-0889.1989.tb00129.x, 1989.
- Forkel, M., Carvalhais, N., Rödenbeck, C., Keeling, R., Heimann, M., Thonicke, K., Zaehle, S., and Reichstein, M.: Enhanced seasonal CO<sub>2</sub> exchange caused by amplified plant productivity in northern ecosystems, *Science*, 696–699, doi:10.1126/science.aac4971, 2016.



- Graven, H. D., Keeling, R. F., Piper, S. C., Patra, P. K., Stephens, B. B., Wofsy, S. C., Welp, L. R., Sweeney, C., Tans, P. P., Kelley, J. J., Daube, B. C., Kort, E. A., Santoni, G. W., and Bent, J. D.: Enhanced Seasonal Exchange of CO<sub>2</sub> by Northern Ecosystems Since 1960, *Science*, 341, 1085–1089, doi:10.1126/science.1239207, 2013.
- Gray, J. M., Frolking, S., Kort, E. A., Ray, D. K., Kucharik, C. J., Ramankutty, N., and Friedl, M. A.: Direct human influence on atmospheric CO<sub>2</sub> seasonality from increased cropland productivity, *Nature*, 515, 398–401, doi:10.1038/nature13957, 2014.
- Hoffman, F. M., Randerson, J. T., Arora, V. K., Bao, Q., Cadule, P., Ji, D., Jones, C. D., Kawamiya, M., Khatriwala, S., Lindsay, K., Obata, A., Shevliakova, E., Six, K. D., Tjiputra, J. F., Volodin, E. M., and Wu, T.: Causes and implications of persistent atmospheric carbon dioxide biases in Earth System Models, *J. Geophys. Res.-Biogeophys.*, 119, 141–162, doi:10.1002/2013JG002381, 2014.
- Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., Lamarque, J.-F., Large, W. G., Lawrence, D., Lindsay, K., Lipscomb, W. H., Long, M. C., Mahowald, N., Marsh, D. R., Neale, R. B., Rasch, P., Vavrus, S., Vertenstein, M., Bader, D., Collins, W. D., Hack, J. J., Kiehl, J., and Marshall, S.: The Community Earth System Model: A Framework for Collaborative Research, *B. Am. Meteorol. Soc.*, 94, 1339–1360, doi:10.1175/BAMS-D-12-00121.1, 2013.
- Hurtt, G. C., Frolking, S., Fearon, M. G., Moore, B. B., Shevliakova, E. E., Malyshev, S., Pacala, S. W., and Houghton, R. A.: The underpinnings of land-use history: three centuries of global gridded land-use transitions, wood-harvest activity, and resulting secondary lands, *Glob. Change Biol.*, 12, 1208–1229, doi:10.1111/j.1365-2486.2006.01150.x, 2006.
- Ito, A., Inatomi, M., Huntzinger, D. N., Schwalm, C., Michalak, A. M., Cook, R., King, A. W., Mao, J., Wei, Y., Post, W. M., Wang, W., Arain, M. A., Huang, S., Hayes, D. J., Ricciuto, D. M., Shi, X., Huang, M., Lei, H., Tian, H., Lu, C., Yang, J., Tao, B., Jain, A., Poulter, B., Peng, S., Ciais, P., Fisher, J. B., Parazoo, N., Schaefer, K., Peng, C., Zeng, N., and Zhao, F.: Decadal trends in the seasonal-cycle amplitude of terrestrial CO<sub>2</sub> exchange resulting from the ensemble of terrestrial biosphere models, *Tellus B*, 28, 28968, doi:10.3402/tellusb.v68.28968, 2016.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, G., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., J. Janowiak, Mo, K. C., Ropelewski, C., Wang, J., Leetma, A. A., Reynolds, R., Jenne, R., and Joseph, D.: The {NCEP/NCAR} 40-year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–470, 1996.
- Keeling, C. D., Chin, J. F. S., and Whorf, T. P.: Increased activity of northern vegetation inferred from atmospheric CO<sub>2</sub> measurements, *Nature*, 382, 146–149, doi:10.1038/382146a0, 1996.
- Keeling, C. D., Piper, S. C., Bacastow, R. B., Wahlen, M., Whorf, T. P., Heimann, M., and Meijer, H. A.: Atmospheric CO<sub>2</sub> and 13CO<sub>2</sub> Exchange with the Terrestrial Biosphere and Oceans from 1978 to 2000: Observations and Carbon Cycle Implications, in: *Ecological Studies*, edited by: Baldwin, I. T., Caldwell, M. M., Heldmaier, G., Jackson, R. B., Lange, O. L., Mooney, H., Schulze, E. D., Sommer, U., Ehleringer, Dearing, M. D., and Cerling, T. E., Vol. 177, 83–113, doi:10.1007/0-387-27048-5\_5, Springer, New York, 2005.
- Keenan, T. F., Hollinger, D. Y., Bohrer, G., Dragoni, D., Munger, J. W., Schmid, H. P., and Richardson, A. D.: Increase in forest water-use efficiency as atmospheric carbon dioxide concentrations rise, *Nature*, 499, 324–327, doi:10.1038/nature12291, 2013.
- Keppel-Aleks, G., Randerson, J. T., Lindsay, K., Stephens, B. B., Moore, J. K., Doney, S. C., Thornton, P. E., Mahowald, N. M., Hoffman, F. M., Sweeney, C., Tans, P. P., Wennberg, P. O., and Wofsy, S. C.: Atmospheric carbon dioxide variability in the Community Earth System Model: Evaluation and transient dynamics during the 20th and 21st centuries, *J. Climate*, 26, 4447–4475, doi:10.1175/JCLI-D-12-00589.1, 2013.
- Keppel-Aleks, G., Wolf, A. S., Mu, M., Doney, S. C., Morton, D. C., Kasibhatla, P. S., Miller, J. B., Dlugokencky, E. J., and Randerson, J. T.: Separating the influence of temperature, drought, and fire on interannual variability in atmospheric CO<sub>2</sub>, *Global Biogeochem. Cy.*, 28, 1295–1310, doi:10.1002/2014GB004890, 2014.
- Kloster, S., Mahowald, N. M., Randerson, J. T., and Lawrence, P. J.: The impacts of climate, land use, and demography on fires during the 21st century simulated by CLM-CN, *Biogeosciences*, 9, 509–525, doi:10.5194/bg-9-509-2012, 2012.
- Kohlmaier, G. H., Siré, E.-O., Janecek, A., Keeling, C. D., Piper, S. C., and Revelle, R.: Modelling the seasonal contribution of a CO<sub>2</sub> fertilization effect of the terrestrial vegetation to the amplitude increase in atmospheric CO<sub>2</sub> at Mauna Loa Observatory, *Tellus B*, 41, 487–510, doi:10.1111/j.1600-0889.1989.tb00137.x, 1989.
- Komhyr, W. D., Gammon, R. H., Harris, T. B., Waterman, L. S., Conway, T. J., Taylor, W. R., and Thoning, K. W.: Global atmospheric CO<sub>2</sub> distribution and variations from 1968–1982 NOAA/GMCC CO<sub>2</sub> flask sample data, *J. Geophys. Res.-Atmos.*, 90, 5567–5596, doi:10.1029/JD090iD03p05567, 1985.
- Koven, C. D., Riley, W. J., Subin, Z. M., Tang, J. Y., Torn, M. S., Collins, W. D., Bonan, G. B., Lawrence, D. M., and Swenson, S. C.: The effect of vertically resolved soil biogeochemistry and alternate soil C and N models on C dynamics of CLM4, *Biogeosciences*, 10, 7109–7131, doi:10.5194/bg-10-7109-2013, 2013.
- Koven, C. D., Lawrence, D. M., and Riley, W. J.: Permafrost carbon-climate feedback is sensitive to deep soil carbon decomposability but not deep soil nitrogen dynamics, *P. Natl. Acad. Sci. USA*, 112, 3752–3757, doi:10.1073/pnas.1415123112, 2015.
- Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039, doi:10.5194/acp-10-7017-2010, 2010.
- Lawrence, D. M., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., Zeng, X., Yang, Z.-L., Levis, S., Sakaguchi, K., Bonan, G. B., and Slater, A. G.: Parameterization improvements and functional and structural advances in Version 4 of the Community Land Model, *J. Adv. Model. Earth Syst.*, 3, M03001, doi:10.1029/2011MS000045, 2011.
- Lawrence, D. M., Oleson, K. W., Flanner, M. G., Fletcher, C. G., Lawrence, P. J., Levis, S., Swenson, S. C., and Bonan, G. B.:

- The CCSM4 land simulation, 1850–2005: Assessment of surface climate and new capabilities, *J. Climate*, 25, 2240–2260, doi:10.1175/JCLI-D-11-00103.1, 2012.
- Levis, S., Hartman, M. D., and Bonan, G. B.: The Community Land Model underestimates land-use CO<sub>2</sub> emissions by neglecting soil disturbance from cultivation, *Geosci. Model Dev.*, 7, 613–620, doi:10.5194/gmd-7-613-2014, 2014.
- Lindsay, K., Bonan, G. B., Doney, S. C., Hoffman, F. M., Lawrence, D. M., Long, M. C., Mahowald, N. M., Keith Moore, J., Randerson, J. T., and Thornton, P. E.: Preindustrial-Control and Twentieth-Century Carbon Cycle Experiments with the Earth System Model CESM1(BGC), *J. Climate*, 27, 8981–9005, doi:10.1175/JCLI-D-12-00565.1, 2014.
- Liu, M., Wu, J., Zhu, X., He, H., Jia, W., and Xiang, W.: Evolution and variation of atmospheric carbon dioxide concentration over terrestrial ecosystems as derived from eddy covariance measurements, *Atmos. Environ.*, 114, 75–82, 2015.
- Lloyd, A. H.: Ecological histories from Alaskan tree lines provide insight into future change, *Ecology*, 86, 1687–1695, doi:10.1890/03-0786, 2005.
- Lloyd, J.: The CO<sub>2</sub> dependence of photosynthesis, plant growth responses to elevated CO<sub>2</sub> concentrations and their interaction with soil nutrient status, II. Temperate and boreal forest productivity and the combined effects of increasing CO<sub>2</sub> concentrations and increased nitrogen deposition at a global scale, *Funct. Ecol.*, 13, 439–459, doi:10.1046/j.1365-2435.1999.00350.x, 1999.
- Long, M. C., Lindsay, K., Peacock, S., Moore, J. K., and Doney, S. C.: Twentieth-Century Oceanic Carbon Uptake and Storage in CESM1(BGC), *J. Climate*, 26, 6775–6800, doi:10.1175/JCLI-D-12-00184.1, 2013.
- Manning, M. R.: Seasonal cycles in atmospheric CO<sub>2</sub> concentrations, in: NATO ASI Series, edited by: Heimann, M., Vol. 15, 65–94, doi:10.1007/978-3-642-84608-3\_3, Springer, Berlin, Heidelberg, 1993.
- McDonald, K. C., Kimball, J. S., Njoku, E., Zimmermann, R., and Zhao, M.: Variability in Springtime Thaw in the Terrestrial High Latitudes: Monitoring a Major Control on the Biospheric Assimilation of Atmospheric CO<sub>2</sub> with Spaceborne Microwave Remote Sensing, *Earth Interact.*, 8, 1–23, doi:10.1175/1087-3562(2004)8<1:VISTIT>2.0.CO;2, 2004.
- McGuire, A. D., Sitch, S., Clein, J. S., Dargaville, R., Esser, G., Foley, J., Heimann, M., Joos, F., Kaplan, J., Kicklighter, D. W., Meier, R. A., Melillo, J. M., Moore, B., Prentice, I. C., Ramankutty, N., Reichenau, T., Schloss, A., Tian, H., Williams, L. J., and Wittenberg, U.: Carbon balance of the terrestrial biosphere in the Twentieth Century: Analyses of CO<sub>2</sub>, climate and land use effects with four process-based ecosystem models, *Global Biogeochem. Cy.*, 15, 183–206, doi:10.1029/2000GB001298, 2001.
- Meinshausen, M., Smith, S., Calvin, K., Daniel, J., Kainuma, M., Lamarque, J.-F., Matsumoto, K., Montzka, S., Raper, S., Riahi, K., Thomson, A., Velders, G., and van Vuuren, D.: The RCP greenhouse gas concentrations and their extensions from 1765 to 2300, *Climatic Change*, 109, 213–241, doi:10.1007/s10584-011-0156-z, 2011.
- Nassar, R., Jones, D. B. A., Suntharalingam, P., Chen, J. M., Andres, R. J., Wecht, K. J., Yantosca, R. M., Kulawik, S. S., Bowman, K. W., Worden, J. R., Machida, T., and Matsueda, H.: Modeling global atmospheric CO<sub>2</sub> with improved emission inventories and CO<sub>2</sub> production from the oxidation of other carbon species, *Geosci. Model Dev.*, 3, 689–716, doi:10.5194/gmd-3-689-2010, 2010.
- Neale, R. B., Richter, J., Park, S., Lauritzen, P. H., Vavrus, S. J., Rasch, P. J., and Zhang, M.: The Mean Climate of the Community Atmosphere Model (CAM4) in forced SST and Fully Coupled Experiments, *J. Climate*, 26, 5150–5168, doi:10.1175/JCLI-D-12-00236.1, 2013.
- Nemry, B., Francois, L., Warnant, P., Robinet, F., and Gerard, J. C.: The seasonality of the CO<sub>2</sub> exchange between the atmosphere and the land biosphere: A study with a global mechanistic vegetation model, *J. Geophys. Res.-Atmos.*, 101, 7111–7125, doi:10.1029/95JD03656, 1996.
- Nevison, C. D., Baker, D. F., and Gurney, K. R.: A methodology for estimating seasonal cycles of atmospheric CO<sub>2</sub> resulting from terrestrial net ecosystem exchange (NEE) fluxes using the Transcom T3L2 pulse-response functions, *Geosci. Model Dev. Discuss.*, 5, 2789–2809, doi:10.5194/gmdd-5-2789-2012, 2012.
- Norby, R. J., Warren, J. M., Iversen, C. M., Medlyn, B. E., and McMurtrie, R. E.: CO<sub>2</sub> enhancement of forest productivity constrained by limited nitrogen availability, *P. Natl. Acad. Sci. USA*, 107, 19368–19373, doi:10.1073/pnas.1006463107, 2010.
- Olsen, S. C. and Randerson, J. T.: Differences between surface and column atmospheric CO<sub>2</sub> and implications for carbon cycle research, *J. Geophys. Res.*, 109, D02301, doi:10.1029/2003JD003968, 2004.
- Parida, B. R. and Buermann, W.: Increasing summer drying in North American ecosystems in response to longer nonfrozen periods, *Geophys. Res. Lett.*, 41, 5476–5483, doi:10.1002/2014GL060495, 2014.
- Pearman, G. I. and Hyson, P.: The annual variation of atmospheric CO<sub>2</sub> concentration observed in the Northern Hemisphere, *J. Geophys. Res.*, 86, 9839–9843, doi:10.1029/JC086iC10p09839, 1981.
- Peng, J. and Dan, L.: Impacts of CO<sub>2</sub> concentration and climate change on the terrestrial carbon flux using six global climate-carbon coupled models, *Ecol. Model.*, 304, 69–83, 2015.
- Piao, S., Sitch, S., Ciais, P., Friedlingstein, P., Peylin, P., Wang, X., Ahlstrom, A., Anav, A., Canadell, J. G., Cong, N., Huntingford, C., Jung, M., Levis, S., Levy, P. E., Li, J., Lin, X., Lomas, M. R., Lu, M., Luo, Y., Ma, Y., Myneni, R. B., Poulter, B., Sun, Z., Wang, T., Viovy, N., Zaehle, S., and Zeng, N.: Evaluation of terrestrial carbon cycle models for their response to climate variability and to CO<sub>2</sub> trends, *Glob. Change Biol.*, 19, 2117–2132, doi:10.1111/gcb.12187, 2013.
- Prentice, I. C., Farquhar, G. D., Fasham, M. J. R., Goulden, M. L., Heimann, M., Jaramillo, V. J., Khesghi, H. S., Le Quere, C., Scholes, R. J., and Wallace, D. W. R.: *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, 881 pp., Cambridge University Press, Cambridge, UK and New York, NY, USA, 2001.
- Randerson, J. T., Thompson, M. V., Conway, T. J., Fung, I. Y., and Field, C. B.: The contribution of terrestrial sources and sinks to trends in the seasonal cycle of atmospheric carbon dioxide, *Global Biogeochem. Cy.*, 11, 535–560, doi:10.1029/97GB02268, 1997.
- Randerson, J. T., Field, C. B., Fung, I. Y., and Tans, P. P.: Increases in early season ecosystem uptake explain re-

- cent changes in the seasonal cycle of atmospheric CO<sub>2</sub> at high northern latitudes, *Geophys. Res. Lett.*, 26, 2765–2768, doi:10.1029/1999GL900500, 1999.
- Randerson, J. T., Lindsay, K., Munoz, E., Fu, W., Moore, J. K., Hoffman, F. M., Mahowald, N. M., and Doney, S. C.: Multicentury changes in ocean and land contributions to the climate-carbon feedback, *Global Biogeochem. Cy.*, 29, 744–759, doi:10.1002/2014GB005079, 2015.
- Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA: NASA's Modern-Era Retrospective Analysis for Research and Applications, *J. Climate*, 24, 3624–3648, doi:10.1175/JCLI-D-11-00015.1, 2011.
- Schimel, D., Stephens, B. B., and Fisher, J. B.: Effect of increasing CO<sub>2</sub> on the terrestrial carbon cycle, *P. Natl. Acad. Sci. USA*, 112, 436–441, doi:10.1073/pnas.1407302112, 2015.
- Tans, P. P., Fung, I. Y., and Takahashi, T.: Observational Constraints on the Global Atmospheric CO<sub>2</sub> Budget, *Science*, 247, 1431–1438, doi:10.1126/science.247.4949.1431, 1990.
- Thonicke, K., Venevsky, S., Sitch, S., and Cramer, W.: The Role of Fire Disturbance for Global vegetation Dynamics: Coupling Fire into a Dynamic Global Vegetation Model, *Global Ecol. Biogeogr.*, 10, 661–677, 2001.
- Thornton, P. E. and Zimmermann, N. E.: An improved Canopy Integration Scheme for a Land Surface Model with Prognostic Canopy Structure, *J. Climate*, 20, 3902–3923, doi:10.1175/JCLI4222.1, 2007.
- Thornton, P. E., Lamarque, J.-F., Rosenbloom, N. A., and Mahowald, N. M.: Influence of carbon-nitrogen cycle coupling on land model response to CO<sub>2</sub> fertilization and climate variability, *Glob. Biogeochem. Cy.*, 21, GB4018, doi:10.1029/2006GB002868, 2007.
- Zeng, N., Zhao, F., Collatz, G. J., Kalnay, E., Salawitch, R. J., West, T. O., and Guanter, L.: Agricultural Green Revolution as a driver of increasing atmospheric CO<sub>2</sub> seasonal amplitude, *Nature*, 515, 394–397, doi:10.1038/nature13893, 2014.
- Zhao, F., Zeng, N., Asrar, G., Friedlingstein, P., Ito, A., Jain, A., Kalnay, E., Kato, E., Koven, C. D., Poulter, B., Rafique, R., Sitch, S., Shu, S., Stocker, B., Viovy, N., Wiltshire, A., and Zaehle, S.: Role of CO<sub>2</sub>, climate and land use in regulating the seasonal amplitude increase of carbon fluxes in terrestrial ecosystems: a multimodel analysis, *Biogeosciences*, 13, 5121–5137, doi:10.5194/bg-13-5121-2016, 2016.