

# *Drosophila* PTB promotes formation of high-order RNP particles and represses *oskar* translation

Florence Besse,<sup>1,2</sup> Sonia López de Quinto,<sup>1,3</sup> Virginie Marchand, Alvar Trucco, and Anne Ephrussi<sup>4</sup>

Developmental Biology Unit, European Molecular Biology Laboratory, 69117 Heidelberg, Germany

Local translation of asymmetrically enriched mRNAs is a powerful mechanism for functional polarization of the cell. In *Drosophila*, exclusive accumulation of Oskar protein at the posterior pole of the oocyte is essential for development of the future embryo. This is achieved by the formation of a dynamic *oskar* ribonucleoprotein (RNP) complex regulating the transport of *oskar* mRNA, its translational repression while unlocalized, and its translational activation upon arrival at the posterior pole. We identified the nucleo–cytoplasmic shuttling protein PTB (polypyrimidine tract-binding protein)/hnRNP I as a new factor associating with the *oskar* RNP in vivo. While PTB function is largely dispensable for *oskar* mRNA transport, it is necessary for translational repression of the localizing mRNA. Unexpectedly, a cytoplasmic form of PTB can associate with *oskar* mRNA and repress its translation, suggesting that nuclear recruitment of PTB to *oskar* complexes is not required for its regulatory function. Furthermore, PTB binds directly to multiple sites along the *oskar* 3' untranslated region and mediates assembly of high-order complexes containing multiple *oskar* RNA molecules in vivo. Thus, PTB is a key structural component of *oskar* RNP complexes that dually controls formation of high-order RNP particles and translational silencing.

[Keywords: *Drosophila*; PTB; *oskar*; translation; RNP assembly; hnRNP]

Supplemental material is available at <http://www.genesdev.org>.

Received September 5, 2008; revised version accepted November 20, 2008.

In eukaryotic cells, transcription represents the first step of gene expression. However, a variety of nuclear and cytoplasmic post-transcriptional events determine the final fate of RNAs and thus regulate gene product diversity as well as the spatio-temporal pattern of gene expression.

In recent years, subcellular targeting of mRNAs, coupled to localized translation, has emerged as a powerful mechanism to spatially and temporally restrict protein synthesis. Furthermore, a genome-wide in situ hybridization analysis in *Drosophila* embryos suggests that RNA localization could represent a general mechanism for the establishment of cell polarity (Lecuyer et al. 2007). Consistent with this, functional studies have shown that local translation of asymmetrically enriched mRNAs is used by differentiated cells to generate functionally distinct compartments, or by developing organisms to partition cell fate determinants (St Johnston 2005; Du

et al. 2007). In several species, the asymmetric distribution in unfertilized eggs of maternal RNAs encoding cytoplasmic determinants controls embryonic body axis specification. In *Drosophila*, *oskar* mRNA encodes the posterior determinant and is transported to the posterior pole of the oocyte, where it is specifically translated. This precise spatio-temporal control is critical for embryonic patterning, as mutant oocytes in which *oskar* is not expressed at the posterior pole develop into embryos lacking abdominal structures and germ cells (Ephrussi et al. 1991; Kim-Ha et al. 1991). Conversely, ectopic translation of unlocalized *oskar* causes patterning defects characterized by a loss of anterior structures, and in extreme cases, duplication of posterior structures (Ephrussi and Lehmann 1992; Kim-Ha et al. 1995).

RNA localization relies on the formation of functional ribonucleoprotein (RNP) complexes that precisely control and coordinate multiple steps including motor-based transport of the mRNA along the cytoskeleton and translational repression of the localizing mRNA, as well as its translation activation and anchoring upon arrival at the final destination (St Johnston 2005). These complexes contain different RNA-associated proteins, including heterogeneous nuclear RNPs (hnRNPs) that regulate various RNA processing events (Dreyfuss et al. 2002; Glisovic et al. 2008). Furthermore, RNPs seem to assemble in vivo

<sup>1</sup>These authors contributed equally to this work.

Present addresses: <sup>2</sup>Institute of Developmental Biology and Cancer/UMR6543, University Nice-Sophia Antipolis, Parc Valrose, 06108 Nice Cedex 2, France.

<sup>3</sup>Cardiff University, School of Biosciences, Museum Avenue, Cardiff CF10 3AX, Wales, UK.

<sup>4</sup>Corresponding author.

E-MAIL [ephrussi@embl.de](mailto:ephrussi@embl.de); FAX 49-6221-387-8166.

Article published online ahead of print. Article and publication date are online at <http://www.genesdev.org/cgi/doi/10.1101/gad.505709>.

Besse et al.

into large particles containing several RNA molecules and associated proteins, as evidenced by light microscopy visualization or by biochemical sedimentation techniques (Kiebler and Bassell 2006; Lange et al. 2008). Although the *in vivo* functional relevance of such high-order structures remains unknown, their formation has been proposed to represent a crucial step in the establishment of transport-competent RNP complexes.

The composition, architecture, and dynamics of RNP complexes dictate the specific behavior of target RNAs within the cell. Importantly, RNP complexes undergo an extensive remodeling upon export into the cytoplasm, yet their cytoplasmic fate is highly connected to their nuclear history (Kress et al. 2004; St Johnston 2005; Giorgi and Moore 2007; Lewis and Mowry 2007). For example, in the case of *oskar* mRNA, nuclear recruitment of the exon-junction complex upon splicing is necessary for *oskar* mRNA transport to the posterior pole of the oocyte (Hachet and Ephrussi 2001, 2004). Furthermore, *oskar* translational repression is controlled by nucleo-cytoplasmic shuttling RNA-binding proteins such as Bruno and Hrp48 (Kim-Ha et al. 1995; Yano et al. 2004; Snee et al. 2008). While these proteins likely associate with the RNA in the nucleus, how important the nuclear recruitment of these proteins to their target RNA is remains to be functionally tested.

In a visual protein-trap screen, we identified *Drosophila* polypyrimidine tract-binding protein (PTB) as a protein colocalizing with *oskar* mRNA. PTB belongs to the hnRNP family of RNA-binding proteins and regulates various nuclear and cytoplasmic RNA processing events in vertebrates (Auweter and Allain 2008). Specifically, PTB plays a predominant role in the regulation of alternative splicing (Valcarcel and Gebauer 1997; Spellman et al. 2005), and has also been shown to regulate internal ribosome entry site (IRES)-dependent translation initiation (Stoneley and Willis 2004; Jang 2006; Semler and Waterman 2008) and mRNA localization (Cote et al. 1999; Ma et al. 2007). In our *in vivo* study, we show that *Drosophila* PTB is a new component of the *oskar* RNP complex and that PTB regulates translational repression of *oskar* mRNA. Surprisingly, while PTB strongly accumulates in germ cell nuclei and thus, potentially, associates with *oskar* in this compartment, its recruitment to *oskar* RNP in the germ cell cytoplasm is sufficient for efficient repression. Consistent with a direct role in *oskar* translation regulation, PTB binds to multiple binding sites in the *oskar* 3' untranslated region (UTR). Finally, we show that PTB is a *trans*-acting factor required *in vivo* for *oskar* mRNA oligomerization. Our study thus reveals that *Drosophila* PTB is a key structural component of *oskar* RNP complexes and suggests a functional link between the assembly of high-order RNP particles and the mechanism of *oskar* translational repression.

## Results

### *PTB colocalizes with oskar mRNA during Drosophila oogenesis*

To identify new proteins involved in *oskar* mRNA regulation, we performed a protein-trap screen that relies


on GFP-tagging of proteins by random mobilization of a GFP-containing transposable element in the genome (Morin et al. 2001; Bonin and Mann 2004). Among the recovered GFP fusion proteins colocalizing with *oskar* mRNA, one showed an enrichment in the oocyte cytoplasm during early oogenesis, followed by an exclusive accumulation at the posterior pole from mid-oogenesis onward (Fig. 1A,B; see Spradling 1993 for a detailed description of ovarian cell types and developmental stages). Additionally, the GFP-tagged protein localizes to the nuclei of both somatic and germ cells, as well as in the cytoplasm of the follicular epithelial cells surrounding the oocyte, at late stages of oogenesis.

Using inverse PCR, we mapped the insertion site of the protein-trap cassette to the *hephaestus* locus (CG31000), encoding the unique *Drosophila* homolog of mammalian PTB, also known as hnRNP-I (Dansereau et al. 2002; Davis et al. 2002). Similar to its mammalian ortholog, *Drosophila* PTB contains a putative N-terminal bipartite nuclear localization signal (NLS) and four conserved RNA recognition motifs (RRM1–4) (Fig. 1C). Western-blot analysis using anti-PTB antibody raised against the *Drosophila* protein revealed a 65-kDa doublet in wild-type ovarian extracts (Fig. 1D), which is shifted to 95 kDa in the protein-trap line as a result of the GFP insertion upstream of RRM1 (Fig. 1C).

Staining of wild-type ovaries with anti-PTB antibodies revealed a localization pattern of endogenous PTB identical to that of GFP-PTB fusions (Fig. 1E–G). Similarly to the *oskar*-associated protein Staufen, PTB accumulates in the cytoplasm of young oocytes as early as in the germarium (Fig. 1E–G'; data not shown), transiently in the center of the oocyte at stage 7/8 (Fig. 1F–F'), and at the posterior pole of the oocyte from stage 9 onward (Fig. 1G–G'; see Supplemental Fig. S1C for a specificity control). The perfect colocalization of PTB with Staufen in the oocyte cytoplasm throughout oogenesis suggested that PTB is transported together with *oskar* mRNA to the posterior pole of the oocyte.

### *PTB is a component of the oskar RNP complex*

To confirm the *in vivo* association of PTB with *oskar* mRNA, we tested if the distribution of PTB in the oocyte depends on *oskar* mRNA. Strikingly, the cytoplasmic enrichment of PTB in young oocytes is no longer detected in egg chambers lacking *oskar* mRNA (*osk<sup>A87</sup>/Df(3R)p<sup>XT03</sup>*) (Fig. 2B), indicating that the initial transport of PTB into the oocyte cytoplasm requires the presence of *oskar* mRNA. As *oskar* mRNA-null egg chambers do not develop past stage 7 (Jenny and Hachet et al. 2006), we analyzed the *oskar* dependence of PTB localization during mid-oogenesis, following the distribution of PTB in *gurken* (*grk*) and *staufer* mutant oocytes. In *grk<sup>2E12</sup>/grk<sup>2B6</sup>* females, microtubule (MT) polarity is altered such that MT plus ends are enriched in the center of the oocyte instead of at the posterior pole (González-Reyes et al. 1995). In these mutants, *oskar* mRNA and Staufen are detected in the center of the oocyte, together with PTB (Fig. 2D–D'). Furthermore, PTB fails to localize at the posterior pole in *stau<sup>D3</sup>* oocytes (Fig. 2E), which appear to



**Figure 1.** Distribution of PTB during *Drosophila* oogenesis. (A,B) GFP expression pattern in an ovariole (A) and a stage 10 egg chamber (B) from homozygous GFP-PTB protein-trap females. An ovariole contains egg chambers of different developmental stages, from stage 1 in the germarium, until stage 14 (not shown). The oocyte and its 15 sibling nurse cells develop as a syncytium. The 16-cell germline cyst is surrounded by a layer of somatic epithelial cells. Egg chambers are oriented anterior to the *left*, posterior to the *right*. The oocyte, whose size dramatically increases as oogenesis proceeds, is the most posterior cell of the germline cyst. Bars, 60  $\mu$ m. (C) Schematic representation of *Drosophila* PTB domain organization indicating the percentage of amino acid identity of the RRM domains compared with PTB orthologs, as well as the insertion site of the GFP sequence (red triangle) in the protein-trap line. (D) Western blot of ovarian extracts from *w* females (first lane), females heterozygous (protein-trap/+; second lane) or homozygous (protein-trap/-; third lane) for the protein-trap insertion, probed with rat anti-PTB antibodies. (E–G) *w* stage 5 (E), stage 8, (F) and stage 9 (G) egg chambers triple-stained with anti-PTB antibodies (*top* panel, red in the overlay), anti-Staufen antibodies (*middle* panel, blue in the overlay), and phalloidin (green in the overlay). Note that PTB is expressed both in the germline and in the somatic epithelium. In contrast, Staufen exclusively accumulates in the oocyte. Bar, 60  $\mu$ m.

have normal MT polarity but fail to enrich *oskar* mRNA at the posterior pole (St Johnston et al. 1991; Palacios and St Johnston 2002; S. López de Quinto, unpubl.). Thus, PTB accumulation at the posterior of the oocyte relies on that of *oskar* mRNA, suggesting that PTB is part of *oskar* RNP in vivo.

Consistent with this, semiquantitative RT-PCR showed that *oskar* mRNA is enriched in the fraction precipitated from GFP-PTB protein-trap ovaries using anti-GFP antibodies, but not in immunoprecipitates from *w* control females (Fig. 2F). No specific enrichment of mRNAs highly expressed in ovaries (*rp49*, *tubulin67C*) or of other mRNAs asymmetrically localized in the oocyte (*bicoid*) was observed in the GFP-PTB precipitated fractions. Altogether, these results indicate that PTB is a bona fide component of *oskar* RNP in vivo.


*oskar* mRNA localization to the posterior pole is delayed in *ptb* mutant oocytes

To investigate the role of PTB in *oskar* mRNA regulation, we analyzed the distribution of *oskar* mRNA in *ptb*

mutant oocytes. Four different lethal alleles belonging to the same complementation group were used (Supplemental Fig. S1A). *heph*<sup>e1</sup> and *heph*<sup>e2</sup> are EMS-induced mutations, the first harboring an amino acid substitution (G  $\rightarrow$  Q) in RRM1, and the second corresponding to a deletion covering RRM1, RRM2, and part of RRM3 (Dansereau et al. 2002). *heph*<sup>03429</sup> and *heph*<sup>1545</sup> are insertions of a *P*-element and a *piggyBac* transposon into the *heph* locus, respectively. *heph*<sup>e2</sup> behaves as a null allele (Dansereau et al. 2002), and *heph*<sup>e1</sup>, *heph*<sup>03429</sup>, and *heph*<sup>1545</sup> as hypomorphic mutants strongly affecting PTB expression levels (Supplemental Fig. S1A,B). Germ cells homozygous for *heph*<sup>e2</sup> stop developing after formation of 16-cell cysts and never produce differentiated egg chambers, suggesting that PTB function is required for early germ cell maintenance (data not shown).

Three other alleles—*heph*<sup>03429</sup>, *heph*<sup>1545</sup>, and, to a lesser extent, *heph*<sup>e1</sup>—support development to later stages. While *oskar* mRNA accumulates normally in these mutant oocytes during early stages of oogenesis (data not shown), its transport to the posterior pole at mid-oogenesis

is affected (Fig. 3B,C). In wild-type oocytes, *oskar* mRNA and Staufen transiently localize in the center of the oocyte at stages 7/8 and accumulate as a tight posterior




**Figure 2.** PTB colocalizes and associates with *oskar* mRNA. (A,B) Stage 5 egg chambers from *osk<sup>A87/+</sup>* (A) and *osk<sup>A87/Df</sup>* (*3R*)*p<sup>XT03</sup>* (B) females expressing GFP-PTB constructs under the control of the *mat- $\alpha$ 4-tub* promoter, and triple-stained for GFP (green in the overlay), DNA (blue, DAPI), and the oocyte cytoplasm marker Orb (red). Bar, 15  $\mu$ m. (C–E) Distribution of the GFP-PTB protein-trap fusion in wild-type (C), *grk<sup>2E12</sup>/grk<sup>2B6</sup>* (D), and *stau<sup>D3</sup>* (E) oocytes. (Left) GFP signal. (Middle) Staufen protein. (Right) Overlay. Bar, 60  $\mu$ m. (F, left panel) RT-PCR amplification of mRNAs recovered in fractions immunoprecipitated from GFP-PTB protein-trap or *w* control ovarian extracts using anti-GFP antibodies. (Right panel) Amplifications from unbound fractions are used as controls.

crescent from early stage 9 onward (Figs. 1E–G, 3A; data not shown). In contrast, *oskar* mRNA persists in the center of the oocyte or as a cloud near the posterior cortex, in  $\sim$ 20% of early stage 9 *ptb* oocytes (Fig. 3B,C,F). This phenotype seems to reflect a transient delay in *oskar* mRNA transport to the posterior pole, as it is rarely observed in late stage 9 or stage 10 oocytes (Fig. 3E,F; data not shown). Since *oskar* mRNA localization relies on the polarity of the MT cytoskeleton, we assessed the distribution of the MT plus-end marker kin- $\beta$ -gal (Clark et al. 1994) in *ptb* mutant oocytes at stage 9. In these oocytes, *oskar* mRNA mislocalization correlates with a strong reduction in the posterior accumulation of kin- $\beta$ -gal (Fig. 3H–H'), suggesting that *oskar* mRNA localization defects may result from a delay in focusing the MT plus-ends at the posterior pole.

As shown in Figure 3I, the *oskar* localization phenotype was rescued upon expression of a wild-type GFP-PTB fusion in *ptb* mutant germ cells. Interestingly, a mutant version of PTB unable to enter the nucleus, but still able to associate with *oskar* mRNA in the cytoplasm (GFP-PTB- $\Delta$ NLS) (Supplemental Fig. S2), failed to rescue this phenotype (Fig. 3I). Our results thus indicate that PTB function is required in germ cell nuclei for efficient establishment of MT polarity at stages 8–9 of oogenesis, ultimately controlling localization of *oskar* mRNA. Given that mammalian PTB has been extensively implicated in the regulation of alternative splicing (Valcarcel and Gebauer 1997; Spellman et al. 2005), it is possible that nuclear *Drosophila* PTB regulates the splicing of genes involved in establishment of MT polarity in the oocyte.

#### *oskar* is prematurely translated in *ptb* mutant oocytes

In wild-type females, *oskar* translation is repressed in early egg chambers and is activated only once the mRNA reaches the posterior pole of the oocyte (Markussen et al. 1995; Rongo et al. 1995). Thus, Oskar protein is not detected before late stage 8 (Fig. 4A) and accumulates exclusively at the oocyte posterior pole in later stages (Fig. 4F). Interestingly, *oskar* is prematurely translated in *ptb* mutant females, and ectopic Oskar is detected as early as in stages 5–6 *ptb* oocytes (data not shown). At stage 7,  $\sim$ 50% of *heph<sup>03429</sup>* and 30% of *heph<sup>1545</sup>* mutant oocytes display ectopic Oskar expression (Fig. 4B–E), the protein accumulating as a poorly defined cloud in the center of the oocyte (Fig. 4B), more rarely as cytoplasmic aggregates (Fig. 4C), or close to the oocyte nucleus at the anterior-dorsal corner (Fig. 4D). Notably, premature *oskar* translation is suppressed by germline expression of wild-type GFP-PTB, as well as by expression of the GFP-PTB- $\Delta$ NLS mutant protein (Fig. 4E). These results indicate that PTB represses *oskar* mRNA translation during early oogenesis and that assembly of translationally silenced *oskar* RNP complexes does not require the recruitment of PTB in the nuclei of germ cells. Furthermore, the differential phenotypic rescue obtained when expressing GFP-PTB- $\Delta$ NLS rules out that the premature translation of *oskar* observed in *ptb* mutant oocytes is a consequence of MT polarity defects.




**Figure 3.** PTB modulates *oskar* mRNA localization and oocyte MT polarity. (A–E) Double staining of early stage 9 (A–C) or stage 10 (D,E) oocytes with anti-Staufen antibodies (red) and DAPI (cyan). (A,D) w. (B,E) *Ovo*<sup>D</sup>-selected *heph*<sup>e1</sup> germline clone. (C) *Ovo*<sup>D</sup>-selected *heph*<sup>1545</sup> germline clone. Bars, 60 μm. (F) Percentage of early stage 9 or stage 10 oocytes in which Staufen was mislocalized (at least 56 oocytes were scored per genotype). (G,H) Stage 9 oocytes heterozygous (G) and homozygous (H) for *heph*<sup>1545</sup>, and triple-stained with anti-Staufen (G,H, red in the overlay), anti-βgal (G',H', green in the overlay), and DAPI (white in the overlay). Bar, 60 μm. (I) Graph showing the percentages of early stage 9 oocytes with mislocalized *oskar* mRNA in different mutant and rescue contexts. GFP-PTB and GFP-PTBΔNLS constructs were expressed under the control of the germline-specific promoter *mat-α4-tub*. Expression levels of the insertions chosen for the rescue experiments were similar, and comparable with endogenous levels (data not shown). The number of egg chambers counted per genotype is control (*n* = 62); *heph*<sup>1545</sup> (*n* = 86); GFP-PTB; *heph*<sup>1545</sup> (*n* = 56); GFP-PTBΔNLS; *heph*<sup>1545</sup> (*n* = 56).

Ectopic Oskar accumulation is also detected in later stage 10 *ptb* oocytes, either at the lateral or at the anterior cortex (Fig. 4G,H). Since such a phenotype has been associated with embryonic anterior patterning defects, ranging from deletion of the head to duplication of posterior structures (Kim-Ha et al. 1995; Yoshida et al. 2004), we examined the cuticle pattern of embryos derived from *heph*<sup>03429</sup> germline clones. The vast majority of these embryos stopped developing before cuticle formation (57.4%; 251/437) or exhibited severe, seemingly *oskar*-independent morphological defects that prevented the analysis of their head (27.7%; 121/437) (Supplemental

Fig. S3). However, 12% (8/63) of the analyzable embryos lacked head structures, consistent with ectopic accumulation of Oskar protein.

To test if PTB might act by regulating the recruitment of other translational repressors, we analyzed the binding of the two best-characterized *oskar* repressors, namely, Bruno and Hrp48 (Kim-Ha et al. 1995; Gunkel et al. 1998; Yano et al. 2004), using both UV-cross-linking and affinity pull-down assays. As illustrated in Supplemental Figure S4, binding of these proteins to *oskar* 3'UTR is preserved in *ptb* mutant extracts. Taken together, these results suggest that PTB is a new *oskar* translational repressor

Besse et al.



**Figure 4.** *ptb* mutant oocytes display ectopic Oskar protein accumulation. (A–D) *w* (A), *heph*<sup>1545</sup> (B), and *heph*<sup>03429</sup> (C,D) stage 7 egg chambers double-stained with anti-Oskar antibodies (green) and DAPI (white). Bar, 30  $\mu$ m. The different patterns of ectopic Oskar accumulation reflect the dynamic distribution of *oskar* mRNA at this stage, which is not altered in *ptb* mutant oocytes compared with wild-type oocytes (data not shown). (E) Graph showing the percentage of stage 7 oocytes with premature Oskar translation in different mutant and rescue backgrounds (see the legend for Fig. 3I for a description of the rescue constructs). (\*\*\*)  $P < 0.005$ ; (\*\*\*\*)  $P < 0.001$  in a  $\chi^2$  test. (F–H) *w* (F), *heph*<sup>1545</sup> (G), and *heph*<sup>03429</sup> (H) stage 10 oocytes double-stained with anti-Oskar antibodies (green) and DAPI (white). Bar, 60  $\mu$ m.

and that it does not affect the interaction of the two best described *oskar* repressors with the RNA.

#### *PTB binds directly and with high affinity to the oskar 3'UTR*

PTB is an RNA-binding protein that regulates several aspects of RNA metabolism through direct interaction with its target RNAs. Depending on the target, PTB binds either to coding sequences or to untranslated regions


(Auweter and Allain 2008). We thus analyzed the capacity of endogenous PTB to interact with different regions of *oskar* mRNA. As shown in Figure 5A, ovarian PTB preferentially associates with the *oskar* 3'UTR, and to a lesser extent with the *oskar* 5' regulatory region separating the two functional AUGs (M1M2) (Gunkel et al. 1998). In this assay, PTB poorly associated with *oskar* coding sequence, and no significant interaction with a nonrelated RNA (*y14* coding sequence) was observed.

To test if PTB can interact with *oskar* mRNA directly, we analyzed the binding of recombinant PTB to the *oskar* M1M2 and 3'UTR regions, in electrophoretic mobility shift assays (EMSA). Two complexes (RNP1 and RNP2) form upon incubation of M1M2 RNA with increasing amounts of maltose-binding protein (MBP)-PTB (Fig. 5B). This interaction is specific, as it is not observed when using another MBP-tagged RNA-binding protein (MS2-binding protein) (data not shown), or an unrelated RNA (*y14* coding sequence) (Fig. 5D). However, the interaction of PTB with *oskar* M1M2 RNA differs from that observed with the 3'UTR. Under native conditions, the *oskar* 3'UTR runs as two bands, likely corresponding to RNA monomers and dimers (Fig. 5C). With increasing amounts of PTB, the discrete 3'UTR complexes detected at low protein concentrations (asterisks) shift into more heterogeneous complexes, and finally into higher-order RNP complexes (Fig. 5C), which are efficiently out-competed by addition of increasing amounts of unlabeled *oskar* 3'UTR competitor to the binding reaction, but not upon addition of unlabeled *y14* RNA (Supplemental Fig. S5A). A similar binding profile was obtained using His-tagged recombinant protein (data not shown).

Filter-binding assays in the presence of increasing amounts of recombinant protein revealed that PTB/*oskar* 3'UTR complexes are formed very rapidly, within 1 min of incubation (data not shown), and indicate that PTB binds with an “apparent” dissociation constant (Kd) of  $\sim 280$  nM (Supplemental Fig. S5B). Furthermore, the sigmoidal shape of the binding curve, together with the Hill coefficient (2.4) obtained after Hill transformation, shows that the binding of PTB to the *oskar* 3'UTR is cooperative. In conclusion, this analysis indicates that PTB can interact directly with *oskar* mRNA and binds with high affinity, cooperatively, and specifically to the 3'UTR.

#### *The oskar 3'UTR contains multiple PTB-binding sites*

The results of the EMSA suggested the presence of multiple PTB-binding sites in the *oskar* 3'UTR. Although no strict consensus sequence for PTB binding has so far been defined, previous reports showed that mammalian PTB binds with high affinity to pyrimidine-rich sequences (Singh et al. 1995; Perez et al. 1997). Consistent with this, short RNA sequences containing two pyrimidine-rich motifs efficiently compete the formation of PTB/*oskar* 3'UTR complexes in EMSA assays (Supplemental Fig. S5A). Strikingly, we found a high number of phylogenetically conserved pyrimidine-rich sequences distributed



**Figure 5.** Characterization of PTB binding to *oskar* RNA. (A) RNA affinity pull-down assay using biotinylated RNAs covering the entire *oskar* mRNA, and *w* ovarian extracts. The coding sequence of *y14* was used as an unrelated RNA control. (Top) Diagram showing the three different probes used in the assay. Proteins in the bound or unbound fractions were visualized after Western blot analysis using the indicated antibodies. Bruno exclusively binds to the *oskar* 3'UTR, whereas Khc, a component of the *oskar* transport machinery without RNA-binding capacity, is not detected in the precipitated fraction. (B–D) EMSA analysis using 50 nM fluorescently labeled *oskar* M1M2 region (B), *oskar* 3'UTR (C), or *y14* coding sequence (D) in the presence of increasing amounts of MBP-PTB (from 0 to 1500 nM) and of 5  $\mu$ M of tRNA. The position of the unbound RNA (Free-RNA) and of the discrete RNP complexes is marked on the *left* and *right*, respectively.

throughout the *oskar* 3'UTR (Supplemental Fig. S6A). To see if PTB could interact *in vivo* with several polypyrimidine tract-containing subregions of the *oskar* 3'UTR, we divided the 3'UTR into five domains and used the corresponding RNAs in affinity pull-down assays. As shown in Figure 6A, PTB associates with each individual domain, although less strongly than with the entire 3'UTR. This binding pattern is distinct from that of other *oskar* regulatory proteins that bind to specific regions of the 3'UTR (see Bruno-binding pattern in Fig. 6A; Kim-Ha et al. 1995; Gunkel et al. 1998; Chang et al. 1999).

Consistent with these data, PTB associates with *oskar* mRNA *in vivo*, most obviously in the context of the entire 3'UTR. Indeed, the accumulation of endogenous PTB in the cytoplasm of early oocytes is restored to wild-type levels upon sole expression of the *oskar* 3'UTR in *oskar* RNA-null ovaries (Fig. 6B). However, expression of subfragments of the 3'UTR at similar levels resulted in only mild accumulation of PTB in the oocyte (data not shown). Together, these results indicate that, both *in vitro* and *in vivo*, PTB binds to multiple sites distributed throughout the *oskar* 3'UTR.


#### *PTB mediates in vivo oligomerization of oskar mRNA molecules*

Previous studies showed that in wild-type oocytes, hybrid *LacZ-oskar* 3'UTR RNAs localize to the posterior pole by hitchhiking on endogenous *oskar* molecules (Hachet and Ephrussi 2004). This property seems to be mediated by the *oskar* 3'UTR, as a *LacZ* transcript fused to the *oskar* coding sequence, but lacking most of the *oskar* 3'UTR region, does not localize at the posterior pole (Kim-Ha et al. 1993). We therefore hypothesized that, *in vivo*, *oskar* mRNA might assemble via its 3'UTR into RNP particles containing multiple mRNA molecules.

Mammalian PTB has been proposed to act as a chaperone promoting intra- and intermolecular RNA interactions (Mitchell et al. 2005; Oberstrass et al. 2005; Song et al. 2005; Auweter and Allain 2008). The binding of PTB to multiple regions of the *oskar* 3'UTR therefore prompted us to test if PTB might mediate the multimerization of *oskar* molecules *in vivo*. To do so, we analyzed the efficiency with which intronless *oskar* 3'UTR-containing RNAs hitchhike on endogenous *oskar* in oocytes expressing low levels of PTB. As shown in Figure 7, the amount of endogenous *oskar* RNA accumulating at the posterior pole of the oocyte (Fig. 7B,D), as well as the total amount of chimeric reporter *EGFP-oskar* 3'UTR RNA expressed in the ovary (Fig. 7F), are comparable in *heph*<sup>1545</sup> and the control. However, while the *EGFP-oskar* 3'UTR reporter RNA accumulates at the posterior pole of control oocytes at late stage 9 (Fig. 7A), only low levels of *EGFP-oskar* 3'UTR RNA are detected at the posterior pole of *heph*<sup>1545</sup> oocytes at this stage (Fig. 7A–E). A similar reduction in the amount of M1M2-*LacZ-oskar* 3'UTR mRNA (Gunkel et al. 1998) hitchhiking to the posterior pole was observed in *heph*<sup>1545</sup> oocytes (Supplemental Fig. S7). These results indicate that the *oskar* 3'UTR is not sufficient for efficient multimerization of *oskar* mRNA *in vivo* and that the *trans*-acting factor PTB plays an essential role in this process.

To assess the role of PTB in assembly of endogenous *oskar* RNP particles, we analyzed their size in early stage 9 *ptb* oocytes using *in situ* hybridization coupled to immuno-electron microscopy (IEM). As illustrated in Figure 7, G and H, this ultrastructural analysis revealed that *oskar* particles are significantly smaller in *heph*<sup>03429</sup> oocytes compared with wild type ( $52 \pm 1$  nm and  $149 \pm 6$  nm, respectively), indicative of a defective copackaging of *oskar* mRNA molecules. Thus, both the presence of

Besse et al.



**Figure 6.** PTB binds to multiple sites along the entire *oskar* 3'UTR. (A) Affinity pull-down assays comparing the binding of PTB to the whole *oskar* 3'UTR and to individual 3'UTR subdomains (see Supplemental Fig. S5). The conditions used were similar to those described in Figure 5A. (B) PTB (green) and actin (red) staining showing the accumulation of PTB in the cytoplasm of stage 6 *w* oocytes, or oocytes either lacking endogenous *oskar* mRNA (*osk<sup>A87</sup>/Df*) or expressing full-length *oskar* 3'UTR (see the Materials and Methods for complete genotypes). Bar, 15  $\mu$ m.

*oskar* 3'UTR *cis*-regulatory sequences and the activity of the *trans*-acting factor PTB are required *in vivo* for the formation of high-order RNP complexes.

## Discussion

### *PTB promotes formation of high-order oskar RNP complexes*

The finding that exogenous RNAs fused to the *oskar* 3'UTR hitchhike on endogenous *oskar* molecules for their localization at the posterior pole of the oocyte revealed the capacity of *oskar* to oligomerize *in vivo* and assemble into high-order RNP particles containing multiple mRNA molecules (Hachet and Ephrussi 2004). Importantly, the *oskar* 3'UTR is not only sufficient, but also required for *in vivo* oligomerization, as exogenous RNAs harboring deletions in this region fail to hitchhike on endogenous *oskar* (Kim-Ha et al. 1993).

We identified PTB as a *trans*-acting factor required for formation of high-molecular-weight complexes *in vitro* (Fig. 5C), and for efficient copackaging of both 3'UTR-containing reporters and endogenous *oskar* mRNAs *in vivo* (Fig. 7). This property correlates with the strong binding of PTB to multiple sites dispersed throughout the


3'UTR. Interestingly, a chaperone activity has been proposed for the vertebrate PTB, based on its capacity to bridge two separate regions of the FMDV IRES (Song et al. 2005), on the conformational changes in RNA induced upon its binding (Mitchell et al. 2003; Pickering et al. 2004), as well as on its role in remodeling of the Vg1 RNP complex (Lewis et al. 2008). This function is further supported by a structural analysis revealing that RRM3 and RRM4 of human PTB adopt a fixed and atypical orientation in which the RNA-binding surfaces of these domains are positioned away from each other (Oberstrass et al. 2005). As a consequence, RRM3 and RRM4 have the capacity to bring distantly located tracts into close proximity and thus, induce looping of the bound RNA. Given that the amino acid composition of these two RNA-binding domains and their linker region is highly conserved (F. Besse and S. López de Quinto, unpubl.), *Drosophila* PTB likely folds and functions similarly to its mammalian counterpart. In the context of *oskar* mRNA, PTB binding may therefore induce specific RNA folding required to establish the RNA–RNA or RNA–protein interactions essential for multimerization of *oskar* mRNA. Alternatively, PTB may itself bridge different *oskar* RNA molecules and nucleate the assembly of multimolecular complexes.

### *PTB represses oskar translation*

Repression of *oskar* mRNA translation is a complex process involving both cap-dependent and cap-independent mechanisms (Nakamura et al. 2004; Chekulaeva et al. 2006), as well as the presence of 5' and 3' regulatory regions (Gunkel et al. 1998). The phenotype of oocytes with reduced PTB levels indicates that *Drosophila* PTB, while dispensable for the transport of *oskar* mRNA, is required for translational repression of the localizing mRNA. Mammalian PTB is already known to regulate translation, mainly by promoting cap-independent translation initiation. PTB binds the IRES located on the 5'UTR of cellular and viral RNAs, thus enhancing the recruitment of *trans*-acting factors and ribosomes (Stoneley and Willis 2004; Jang 2006; Semler and Waterman 2008). Recent reports indicate that PTB can also promote translation of specific mRNAs when bound to their 3'UTRs (Reyes and Izquierdo 2007; Galban et al. 2008). However, PTB does not exclusively act as a translational activator, as its interaction with the IRES present in *unr* and *bip* mRNAs has been shown to down-regulate their activity (Kim et al. 2000; Cornelis et al. 2005). Our results suggest that *Drosophila* PTB also acts as a translational repressor when bound to the 3'UTR of a target mRNA.

The mechanism by which PTB enhances or represses translation of target mRNAs remains elusive. The most accepted hypothesis, which may also explain the multiple roles of PTB in RNA regulation, is its capacity to act as a chaperone molecule, promoting the folding of RNAs into specific conformations, thereby modulating the binding of other regulatory proteins (Mitchell et al. 2003; Pickering et al. 2004; Song et al. 2005; Lewis et al. 2008). Thus, depending on the specific structure adopted





**Figure 7.** PTB promotes copackaging of *oskar* mRNA molecules in vivo. In situ hybridization of heterozygous (A,B) and homozygous (C,D) *heph*<sup>1545</sup> late stage 9 egg chambers expressing a UAS-EGFP-*osk* 3'UTR construct under the control of the *mat-α4-tub*-Gal4 driver. (A,C) Egg chambers double-stained for EGFP transcripts (red) and DAPI (white). Images A and C were taken using identical settings. (B,D) Egg chambers double-stained for endogenous *oskar* transcripts (green) and DAPI (white). Endogenous *oskar* was detected using a probe exclusively covering the coding region. Pictures B and D were taken using identical settings. Bar, 60 μm. (E) Quantification of the in situ hybridization signal detected at the posterior pole of late stage 9 oocytes. Intensity values obtained using ImageJ are expressed as a percentage of control oocytes. (\*\*\*)  $P < 0.001$  in a Student's *t*-test. (F) Expression levels of *rp49*, endogenous *oskar*, and EGFP-*oskar* 3'UTR transcripts in both heterozygous and homozygous *heph*<sup>1545</sup> backgrounds, as measured by semiquantitative RT-PCR. (G,H) Electron micrographs of *w* (G) and *heph*<sup>03429</sup> (H) early stage 9 oocytes hybridized with an *oskar* antisense probe. *oskar* mRNA was detected using 10-nm gold particles, and the central region of the oocyte is shown. Note that in *heph*<sup>03429</sup> oocytes, the slightly electron-dense *oskar* RNP complexes are smaller and more numerous. Bar, 150 nm.

by the RNA, the cellular context, and the binding of other regulatory *trans*-acting factors, PTB may either promote or inhibit recruitment of the translation machinery. In the case of *oskar* mRNA, however, our results show that PTB is not required for the binding of the two best-characterized *oskar* translational repressors, namely, Bruno and Hrp48 (Kim-Ha et al. 1995; Gunkel et al. 1998; Yano et al. 2004). Although we cannot exclude that PTB influences the activity of these proteins, PTB function seems beyond the simple recruitment of *oskar* main translation repressors.

Significantly, our results establish an in vivo link between *oskar* translation repression and multimerization of *oskar* 3'UTR-containing RNAs. An attractive possibility is that PTB promotes the formation of densely packed *oskar* RNP particles, thereby rendering the mRNA inaccessible to the translation machinery. A similar cap-independent function has been proposed for the translational repressor Bruno, based on its ability to promote in vitro oligomerization of a 3'UTR fragment containing duplicated Bruno response elements (BREs), together with its capacity to nucleate the assembly

of heavy silencing particles (Chekulaeva et al. 2006). However, we observed that BREs are neither strictly required nor sufficient for the hitchhiking of *oskar* 3'UTR-containing RNAs on endogenous *oskar* in vivo (M. Chekulaeva and F. Besse, unpubl.). Therefore, other *trans*-acting factors and *cis*-acting sequences likely contribute to the formation of high-order *oskar* RNPs in the oocyte. In this context, the chaperone activity of PTB may be essential to promote multimerization of *oskar* molecules, which would ultimately contribute to their complete translational repression. In contrast, the *oskar* phenotype of *ptb* mutant oocytes suggests that the assembly of high-order RNP complexes does not play a significant role in *oskar* mRNA posterior transport.

#### *Association of PTB with oskar in the cytoplasm is sufficient to repress translation*

The assembly of RNP complexes competent for mRNA localization and precise translational control has been suggested to occur in a stepwise manner, some factors associating with the mRNA in the nucleus, and being essential for the subsequent recruitment of other components

in the cell cytoplasm (Hachet and Ephrussi 2001, 2004; Huynh et al. 2004; Kress et al. 2004; St Johnston 2005; Giorgi and Moore 2007; Lewis and Mowry 2007; Snee et al. 2008). For example, nucleolar association of the RNA-binding protein She2p with its mRNA target *ash1* was recently proposed to be an essential step in the assembly of translationally silenced localizing *ash1* RNP complexes in yeast (Du et al. 2008). Some *oskar* translation repressors have been shown to localize both in the nucleus and in the cytoplasm of germ cells (Huynh et al. 2004; Yano et al. 2004; Snee et al. 2008). However, whether these regulators are recruited to the *oskar* complex in the nucleus and whether nuclear association of these factors is required for subsequent translation silencing have not been tested so far.

PTB belongs to the hnRNP family of nucleo-cytoplasmic shuttling RNA-binding proteins, which regulate different aspects of RNA metabolism both in the nucleus and in the cytoplasm of eukaryotic cells (Dreyfuss et al. 2002; Glisovic et al. 2008). Consistent with this, we observed that *Drosophila* PTB not only colocalizes with *oskar* in the oocyte cytoplasm, but also strongly accumulates in the nuclei of germ cells. Given that the nuclear association of PTB with *Vg1* mRNA has been proposed to control the subsequent localization of these transcripts in the cytoplasm of *Xenopus* oocytes (Kress et al. 2004), we tested if the association of *Drosophila* PTB with *oskar* mRNA in the nuclei of germ cells is required for its translation repression activity. Notably, we found that a cytoplasmic version of PTB localizes to the posterior pole of wild-type and *ptb* mutant oocytes and that this localization is *oskar*-dependent (Supplemental Fig. S2; data not shown), strongly suggesting that it is still able to associate with *oskar* mRNA. More importantly, the mutant GFP-PTB-ΔNLS is competent in *oskar* translation repression (Fig. 4E). Although it is possible that endogenous PTB is loaded onto *oskar* RNP complexes in the nucleus of germ cells, our data suggest that nuclear recruitment of PTB is not a prerequisite for the formation of translationally silenced complexes. Our analysis supports a model in which the complex behavior of RNP particles is controlled by the independent association of specific protein modules in different cell compartments. It also provides further evidence for the reorganization of RNP complexes upon translocation into the cytoplasm.

## Materials and methods

### Identification of GFP-PTB protein-trap line

The GFP protein-trap screen and inverse PCR mapping were described previously (Besse et al. 2007). In the PTB protein-trap line, the *piggyBac* transposon is inserted at position 27,699,986 in the genomic scaffold AE014297. Females homozygous for the protein-trap insertion are homozygous viable and do not show *oskar*-related phenotypes.

### Fly stocks

*w<sup>1118</sup>* flies (*w*) served as wild-type controls. The P[PZ]heph [03429] (Bloomington Stock Center #11589) and PBac[WH]-

heph[f01545] (Exelixis Collection, Harvard Medical School) were recombined onto the FRT<sup>82B</sup> chromosome. The FRT<sup>82B</sup>-recombined EMS mutant alleles *heph<sup>e1</sup>* and *heph<sup>e2</sup>* were characterized by Dansereau et al. (2002). Germline clone analysis was carried out using the dominant female sterile technique (Chou et al. 1993). Other mutant stocks used were *osk<sup>A87</sup>* and *Df(3R)p<sup>XTO3</sup>* (Jenny and Hachet et al. 2006), *grk<sup>2B6</sup>* (Schüpbach 1987), *grk<sup>2E12</sup>* (Neuman-Silberberg and Schüpbach 1993), and *stau<sup>D3</sup>* (Schüpbach and Wieschaus 1986). The MT polarity marker *kin-β-gal* was obtained from the stock *y w KZ32; Ly/TM3Sb* (Clark et al. 1994). Rescue of the *oskar* RNA-null phenotype using an UAS-*oskar* 3'UTR transgene was performed as described (Jenny and Hachet et al. 2006). For the hitchhiking assay described in Figure 7, a *mat-α4-tub*-Gal4VP16 insertion (Bloomington stock center #7062) was recombined with a UASp-EGFP-*oskar* 3'UTR transgene (H. Jambor and A. Ephrussi, unpubl.).

### Transgenes

For rescue analyses, a full-length clone (RE56755) was N-terminal-tagged with EGFP and expressed using the germline-specific *tubulin67c* promoter. Briefly, the PTB coding sequence was PCR-amplified using the PTB-sense and PTB-antisense primers (Supplemental Table 1) and ligated into pCR II-TOPO (Invitrogen). To generate the PTBΔNLS protein, lysines at positions 56, 57, and 59 were mutated into glutamic acid using a nested-PCR site-directed mutagenesis approach with the mutagenic primer Mut-NLS-II together with PTB-sense and Internal-as as external primers (Supplemental Table 1). The SspI-NruI fragment covering the mutated positions was exchanged in the wild-type pCR II-TOPO-PTB construct, and both wild-type and ΔNLS-mutant pCR II-TOPO-PTB clones were fully sequenced. To generate EGFP fusions of PTB and PTBΔNLS, the 1.45-kb BamHI-XhoI fragment was introduced into a GFP-containing BamHI-XhoI backbone of GFP-par-1(N1S) plasmid (Shulman et al. 2000).

### Generation of Drosophila anti-PTB antisera and Western blot analysis

The BDGP clone LD03185 was PCR-amplified using PTB-GST-sense and PTB-GST-as primers (Supplemental Table 1). The PCR fragment was subsequently ligated into pCR II-TOPO (Invitrogen) and fully sequenced, and the ~1.4 BamHI-XhoI fragment was inserted into the pGEX5-5X-2 plasmid (GenBank accession no. U13857) resulting in the N-terminal fusion of GST to PTB coding sequence. The recombinant protein was expressed in *Escherichia coli*, purified using standard conditions, and injected into rabbits and rats.

### Immunofluorescence

Ovaries from females mated for 1–2 d at 25°C on fresh food were dissected in PBS and processed as described (Vanzo and Ephrussi 2002). Samples were incubated overnight at 4°C with mouse anti-βGal (Promega; 1:2000), rabbit anti-Staufen (St Johnston et al. 1991), rabbit anti-Oskar (Vanzo and Ephrussi 2002), rat anti-PTB (1:1000), and rabbit anti-Bruno (1:5000) [R. Matthiesen and A. Ephrussi, unpubl.]. Alexa-conjugated secondary antibodies (1:750; Invitrogen) were used. For detection of ectopic Oskar, we used the protocol for in situ hybridization coupled to immunofluorescence described already (Vanzo and Ephrussi 2002) in combination with preadsorbed anti-Oskar and Alexa488-conjugated antibodies. Ovaries were stained with phalloidin to label F-actin (Invitrogen) and/or with 3 μg/mL DAPI to label nuclei. Images were captured by a confocal microscope (Leica TCS SP2 AOBs) using a 40× PL APO oil-immersion lens (N.A. 1.25) and processed with Adobe Photoshop.

### RNA in situ hybridization

RNA in situ hybridization was performed as described previously (Vanzo and Ephrussi 2002). Probes corresponding to either *oskar* or EGFP coding sequences were detected using sheep HRP-conjugated anti-DIG antibody (1:200; Roche) followed by Cy3-tyramide signal amplification reaction (Perkin Elmer).

### Immunoprecipitation and RT-PCR analysis

The protocol was adapted from Munro et al. (2006) with the following modifications: protein extracts were precleared with protein A-agarose beads (Roche) and incubated without prior cross-linking with 1.5  $\mu$ g of mouse anti-GFP antibodies (Molecular Probes) coupled to protein A-agarose beads (Roche) for 3 h at 4°C. RNAs were released in 100  $\mu$ L of extraction buffer by incubation for 20 min at 65°C. Total RNA was extracted using Trizol (Invitrogen), DNase-treated, and used as template for RT. All RNA recovered from the bound fraction and 2.5% of that in the unbound fraction was used for cDNA synthesis in combination with Superscript III (Invitrogen) and Oligo(dT). One percent of the RT product was then PCR-amplified. Twenty-two cycles were carried out at an annealing temperature of 55°C using specific primers (Supplemental Table 1).

### Affinity pull-down assays

Ovaries of *w<sup>1118</sup>* females were dissected in cold PBS and extracted using a pestle in 20  $\mu$ L of hypotonic buffer [10 mM HEPES at pH 7.4, 10 mM KOAc, 1.5 mM Mg(OAc)<sub>2</sub>, 2.5 mM DTT]. Typically, 20 pairs of ovaries were used per condition. The protein extract was cleared by centrifugation at 12,000 rpm for 5 min, and the supernatant was incubated with magnetic streptavidin beads (Roche) previously coupled to equimolecular amounts of in vitro transcribed UTP-biotinylated-RNA probes (ranging from 0.5 to 2  $\mu$ g) in binding buffer (10 mM HEPES-KOH at pH 7.9, 3 mM MgCl<sub>2</sub>, 40 mM KCl, 5 mM EDTA, 5% glycerol, 2 mM DTT, 0.5% IGEPAL, 3  $\mu$ g/ $\mu$ L Heparin, and 0.5  $\mu$ g/ $\mu$ L tRNA) during 1 h at 4°C. After two washes in binding buffer followed by two more washes in binding buffer containing 150 mM KCl, the bound proteins were eluted in SDS-PAGE loading buffer and subjected to electrophoresis and Western blotting using rabbit anti-PTB (1:2000), rat anti-Bruno (1:4000) (R. Matthiesen and A. Ephrussi) and rabbit anti-Khc (1:10,000; Cytoskeleton).

### Subcloning of different *oskar* mRNA regions

All *oskar* regions were cloned into pBS-II-KS+ (Stratagene) downstream from the T7 promoter. Nucleotides flanking each fragment described below have been numbered based on their position in the *oskar* cDNA sequence (*osk*-RA; Flybase), and all primer sequences are specified in Supplemental Table 1. Using a genomic *oskar* clone as template (Ephrussi et al. 1991), the M1M2 (nucleotides 1–434) and 3'UTR (nucleotides 1834–2890) regions were PCR-amplified with primers M1M2-sense, and M1M2-as, or 3UTR-s and 3UTR-as, respectively, and subsequently inserted into the SacI–HindIII pBS fragment. A similar strategy was used for subcloning the individual 3'UTR domains with the following pair of primers: Domain1 (nucleotides 1850–2068), Dom1-sense and Dom1-as; Domain2 (nucleotides 2065–2211), Dom2-Long-s and Dom2-as; Domain3 (nucleotides 2210–2334), Dom3-Long-s and Dom3-as; Domain4 (nucleotides 2322–2619), Dom4-sense and Dom4-as and Domain5 (nucleotides 2609–2890), Dom5-sense and 3UTR-as. The region corresponding to Short Oskar coding sequence (nucleotides 427–1838) was PCR-amplified using ShortOsk-Not and ShortOsk-as as primers and an

*oskar* cDNA clone as template (Ephrussi et al. 1991). The PCR fragment was ligated into pCR II-TOPO (Invitrogen), and the EcoRI fragment was inserted into EcoRI-linearized pBS-II-SK+ (Stratagene).

### Recombinant PTB protein

Full-length PTB was N-terminal-tagged by MBP via PCR amplification of the DGC clone RE56755 with GW-PTB-sense and GW-PTB-as primers (Supplemental Table 1). The PCR product was ligated into pENTRY/D-TOPO vector (Invitrogen), fully sequenced, and subsequently recombined into the Gateway destination vector pETG-40A (A. Geerloff). The protein was expressed in *E. coli* (BL21 DE3) and purified under standard conditions. The MS2-binding protein—MBP-expressing plasmid (Zhou et al. 2002) was a gift from Reinhard Lührmann.

### EMSA

RNA transcripts were fluorescently labeled with ATTO-680 Aminoallyl-UTP (Jena Bioscience) via in vitro transcription (T7 megascript kit; Ambion). Full-length RNA molecules were gel-purified and incubated with different amounts of recombinant MBP-PTB in the presence of 20 mM Tris-HCl (pH 8.0), 60 mM NaCl, 3 mM MgCl<sub>2</sub>, 1 mM DTT, 8% glycerol, 5  $\mu$ M tRNA during 15 min at 4°C. Complexes were resolved in native 0.8% agarose gels containing 0.5 $\times$  TBE buffer and 5% (v/v) glycerol, run at 100 V constant at 4°C. Gels were visualized and quantitated using the Odyssey Infrared Imaging System (LI-COR).

### IEM

Ovaries were fixed in 4% paraformaldehyde and processed as described (Delanoue and Herpers et al. 2007), with minor modifications. Digoxigenin-labeled *oskar* antisense probes were detected using sheep anti-Dig antibody (1:650; Roche), followed by rabbit anti-sheep secondary antibody (1:500; Dako) and protein A-coupled gold particles (10 nm). For quantification, the largest diameter of electron-dense *oskar*-containing particles present in the center of the oocyte was measured using Image J. At least 200 particles from three different oocytes were analyzed for each genotype.

### Acknowledgments

We thank Helena Jambor, William Brook, Daniel St Johnston, Reinhard Lührmann, and Paul McDonald for flies and reagents. We also thank the EMBL Laboratory Animal Resources for antibody production and the EMBL Electron Microscopy Core Facility for technical support. We thank Eva Löser and Anna Cyrklaff for excellent technical assistance. We are grateful to members of the Ephrussi group for discussions and to Vladimir Rybin for advice. F.B. was supported by fellowships from the Federation of European Biochemical Societies (FEBS) and the Human Frontier Science Program Organization. S.L.Q. was supported by a Long Term EMBO Fellowship, a Marie Curie Intra-European Fellowship, and a RCUK Fellowship in Translational Medicine. V.M. was supported by a fellowship from the Fondation pour la Recherche Médicale and A.T. was supported by a FEBS fellowship.

### References

Auweter, S.D. and Allain, F.H. 2008. Structure–function relationships of the polypyrimidine tract binding protein. *Cell. Mol. Life Sci.* **65**: 516–527.

Besse et al.

- Besse, F., Mertel, S., Kittel, R.J., Wichmann, C., Rasse, T.M., Sigrist, S.J., and Ephrussi, A. 2007. The Ig cell adhesion molecule Basigin controls compartmentalization and vesicle release at *Drosophila melanogaster* synapses. *J. Cell Biol.* **177**: 843–855.
- Bonin, C.P. and Mann, R.S. 2004. A piggyBac transposon gene trap for the analysis of gene expression and function in *Drosophila*. *Genetics* **167**: 1801–1811.
- Chang, J.S., Tan, L., and Schedl, P. 1999. The *Drosophila* CPEB homolog, Orb, is required for Oskar protein expression in oocytes. *Dev. Biol.* **215**: 91–106.
- Chekulaeva, M., Hentze, M.W., and Ephrussi, A. 2006. Bruno acts as a dual repressor of oskar translation, promoting mRNA oligomerization and formation of silencing particles. *Cell* **124**: 521–533.
- Chou, T.-B., Noll, E., and Perrimon, N. 1993. Autosomal *P[ovo<sup>D1</sup>]* dominant female-sterile insertions in *Drosophila* and their use in generating germ-line chimeras. *Development* **119**: 1359–1369.
- Clark, I., Giniger, E., Ruohola-Baker, H., Jan, L.Y., and Jan, Y.N. 1994. Transient posterior localization of a kinesin fusion protein reflects anteroposterior polarity of the *Drosophila* oocyte. *Curr. Biol.* **4**: 289–300.
- Cornelis, S., Tinton, S.A., Schepens, B., Bruynooghe, Y., and Beyaert, R. 2005. UNR translation can be driven by an IRES element that is negatively regulated by polypyrimidine tract binding protein. *Nucleic Acids Res.* **33**: 3095–3108.
- Cote, C.A., Gautreau, D., Denegre, J.M., Kress, T.L., Terry, N.A., and Mowry, K.L. 1999. A *Xenopus* protein related to hnRNPI has a role in cytoplasmic RNA localization. *Mol. Cell* **4**: 431–437.
- Dansereau, D.A., Lunke, M.D., Finkielstein, A., Russell, M.A., and Brook, W.J. 2002. Hephaestus encodes a polypyrimidine tract binding protein that regulates Notch signalling during wing development in *Drosophila melanogaster*. *Development* **129**: 5553–5566.
- Davis, M.B., Sun, W., and Standiford, D.M. 2002. Lineage-specific expression of polypyrimidine tract binding protein (PTB) in *Drosophila* embryos. *Mech. Dev.* **111**: 143–147.
- Delanoue, R., Hoppers, B., Soetaert, J., Davis, I., and Rabouille, C. 2007. *Drosophila* Squid/hnRNP helps Dynein switch from a gurken mRNA transport motor to an ultrastructural static anchor in sponge bodies. *Dev. Cell* **13**: 523–538.
- Dreyfuss, G., Kim, V.N., and Kataoka, N. 2002. Messenger-RNA-binding proteins and the messages they carry. *Nat. Rev. Mol. Cell Biol.* **3**: 195–205.
- Du, T.G., Schmid, M., and Jansen, R.P. 2007. Why cells move messages: The biological functions of mRNA localization. *Semin. Cell Dev. Biol.* **18**: 171–177.
- Du, T.G., Jellbauer, S., Muller, M., Schmid, M., Niessing, D., and Jansen, R.P. 2008. Nuclear transit of the RNA-binding protein She2 is required for translational control of localized ASH1 mRNA. *EMBO Rep.* **9**: 781–787.
- Ephrussi, A. and Lehmann, R. 1992. Induction of germ cell formation by oskar. *Nature* **358**: 387–392.
- Ephrussi, A., Dickinson, L.K., and Lehmann, R. 1991. oskar organizes the germ plasm and directs localization of the posterior determinant nanos. *Cell* **66**: 37–50.
- Galban, S., Kuwano, Y., Pullmann Jr., R., Martindale, J.L., Kim, H.H., Lal, A., Abdelmohsen, K., Yang, X., Dang, Y., Liu, J.O., et al. 2008. RNA-binding proteins HuR and PTB promote the translation of hypoxia-inducible factor 1 $\alpha$ . *Mol. Cell. Biol.* **28**: 93–107.
- Giorgi, C. and Moore, M.J. 2007. The nuclear nurture and cytoplasmic nature of localized mRNPs. *Semin. Cell Dev. Biol.* **18**: 186–193.
- Glisovic, T., Bachorik, J.L., Yong, J., and Dreyfuss, G. 2008. RNA-binding proteins and post-transcriptional gene regulation. *FEBS Lett.* **582**: 1977–1986.
- González-Reyes, A., Elliott, H., and St Johnston, D. 1995. Polarization of both major body axes in *Drosophila* by *gurken-torpedo* signalling. *Nature* **375**: 654–658.
- Gunkel, N., Yano, T., Markussen, F.-H., Olsen, L.C., and Ephrussi, A. 1998. Localization-dependent translation requires a functional interaction between the 5' and 3' ends of oskar mRNA. *Genes & Dev.* **12**: 1652–1664.
- Hachet, O. and Ephrussi, A. 2001. *Drosophila* Y14 shuttles to the posterior of the oocyte and is required for oskar mRNA transport. *Curr. Biol.* **11**: 1666–1674.
- Hachet, O. and Ephrussi, A. 2004. Splicing of oskar RNA in the nucleus is coupled to its cytoplasmic localization. *Nature* **428**: 959–963.
- Huynh, J.R., Munro, T.P., Smith-Litiere, K., Lepesant, J.A., and St Johnston, D. 2004. The *Drosophila* hnRNPA/B homolog, Hrp48, is specifically required for a distinct step in osk mRNA localization. *Dev. Cell* **6**: 625–635.
- Jang, S.K. 2006. Internal initiation: IRES elements of picornaviruses and hepatitis c virus. *Virus Res.* **119**: 2–15.
- Jenny, A., Hachet, O., Zavorszky, P., Cyrklaff, A., Weston, M.D., Johnston, D.S., Erdelyi, M., and Ephrussi, A. 2006. A translation-independent role of oskar RNA in early *Drosophila* oogenesis. *Development* **133**: 2827–2833.
- Kiebler, M.A., and Bassell, G.J. 2006. Neuronal RNA granules: Movers and makers. *Neuron* **51**: 685–690.
- Kim, Y.K., Hahm, B., and Jang, S.K. 2000. Polypyrimidine tract-binding protein inhibits translation of bip mRNA. *J. Mol. Biol.* **304**: 119–133.
- Kim-Ha, J., Smith, J.L., and Macdonald, P.M. 1991. oskar mRNA is localized to the posterior pole of the *Drosophila* oocyte. *Cell* **66**: 23–35.
- Kim-Ha, J., Webster, P.J., Smith, J.L., and Macdonald, P.M. 1993. Multiple RNA regulatory elements mediate distinct steps in localization of oskar mRNA. *Development* **119**: 169–178.
- Kim-Ha, J., Kerr, K., and Macdonald, P.M. 1995. Translational regulation of oskar mRNA by bruno, an ovarian RNA-binding protein, is essential. *Cell* **81**: 403–412.
- Kress, T.L., Yoon, Y.J., and Mowry, K.L. 2004. Nuclear RNP complex assembly initiates cytoplasmic RNA localization. *J. Cell Biol.* **165**: 203–211.
- Lange, S., Katayama, Y., Schmid, M., Burkacky, O., Brauchle, C., Lamb, D.C., and Jansen, R.P. 2008. Simultaneous transport of different localized mRNA species revealed by live-cell imaging. *Traffic* **9**: 1256–1267.
- Lecuyer, E., Yoshida, H., Parthasarathy, N., Alm, C., Babak, T., Cerovina, T., Hughes, T.R., Tomancak, P., and Krause, H.M. 2007. Global analysis of mRNA localization reveals a prominent role in organizing cellular architecture and function. *Cell* **131**: 174–187.
- Lewis, R.A., and Mowry, K.L. 2007. Ribonucleoprotein remodeling during RNA localization. *Differentiation* **75**: 507–518.
- Lewis, R.A., Gagnon, J.A., and Mowry, K.L. 2008. PTB/hnRNP I is required for RNP remodeling during RNA localization in *Xenopus* oocytes. *Mol. Cell. Biol.* **28**: 678–686.
- Ma, S., Liu, G., Sun, Y., and Xie, J. 2007. Relocalization of the polypyrimidine tract-binding protein during PKA-induced neurite growth. *Biochim. Biophys. Acta* **1773**: 912–923.
- Markussen, F.-H., Michon, A.-M., Breitwieser, W., and Ephrussi, A. 1995. Translational control of oskar generates Short OSK, the isoform that induces pole plasm assembly. *Development* **121**: 3723–3732.

- Mitchell, S.A., Spriggs, K.A., Coldwell, M.J., Jackson, R.J., and Willis, A.E. 2003. The Apaf-1 internal ribosome entry segment attains the correct structural conformation for function via interactions with PTB and unr. *Mol. Cell* **11**: 757–771.
- Mitchell, S.A., Spriggs, K.A., Bushell, M., Evans, J.R., Stoneley, M., Le Quesne, J.P., Spriggs, R.V., and Willis, A.E. 2005. Identification of a motif that mediates polypyrimidine tract-binding protein-dependent internal ribosome entry. *Genes & Dev.* **19**: 1556–1571.
- Morin, X., Daneman, R., Zavortink, M., and Chia, W. 2001. A protein trap strategy to detect GFP-tagged proteins expressed from their endogenous loci in *Drosophila*. *Proc. Natl. Acad. Sci.* **98**: 15050–15055.
- Munro, T.P., Kwon, S., Schnapp, B.J., and St Johnston, D. 2006. A repeated IMP-binding motif controls *oskar* mRNA translation and anchoring independently of *Drosophila melanogaster* IMP. *J. Cell Biol.* **172**: 577–588.
- Nakamura, A., Sato, K., and Hanyu-Nakamura, K. 2004. *Drosophila* cup is an eIF4E binding protein that associates with Bruno and regulates *oskar* mRNA translation in oogenesis. *Dev. Cell* **6**: 69–78.
- Neuman-Silberberg, F.S. and Schüpbach, T. 1993. The *Drosophila* dorsoventral patterning gene *gurken* produces a dorsally localized RNA and encodes a TGF $\alpha$ -like protein. *Cell* **75**: 165–174.
- Oberstrass, F.C., Auweter, S.D., Erat, M., Hargous, Y., Henning, A., Wenter, P., Reymond, L., Amir-Ahmady, B., Pitsch, S., Black, D.L., et al. 2005. Structure of PTB bound to RNA: Specific binding and implications for splicing regulation. *Science* **309**: 2054–2057.
- Palacios, I.M., and St Johnston, D. 2002. Kinesin light chain-independent function of the Kinesin heavy chain in cytoplasmic streaming and posterior localisation in the *Drosophila* oocyte. *Development* **129**: 5473–5485.
- Perez, I., Lin, C.H., McAfee, J.G., and Patton, J.G. 1997. Mutation of PTB binding sites causes misregulation of alternative 3' splice site selection in vivo. *RNA* **3**: 764–778.
- Pickering, B.M., Mitchell, S.A., Spriggs, K.A., Stoneley, M., and Willis, A.E. 2004. Bag-1 internal ribosome entry segment activity is promoted by structural changes mediated by poly(rC) binding protein 1 and recruitment of polypyrimidine tract binding protein 1. *Mol. Cell Biol.* **24**: 5595–5605.
- Reyes, R., and Izquierdo, J.M. 2007. The RNA-binding protein PTB exerts translational control on 3'-untranslated region of the mRNA for the ATP synthase  $\beta$ -subunit. *Biochem. Biophys. Res. Commun.* **357**: 1107–1112.
- Rongo, C., Gavis, E.R., and Lehmann, R. 1995. Localization of *oskar* RNA regulates *oskar* translation and requires Oskar protein. *Development* **121**: 2737–2746.
- Schüpbach, T. 1987. Germ line and soma cooperate during oogenesis to establish the dorsoventral pattern of egg shell and embryo in *Drosophila melanogaster*. *Cell* **49**: 699–707.
- Schüpbach, T., and Wieschaus, E. 1986. Germline autonomy of maternal-effect mutations altering the embryonic body pattern of *Drosophila*. *Dev. Biol.* **113**: 443–448.
- Semler, B.L., and Waterman, M.L. 2008. IRES-mediated pathways to polysomes: Nuclear versus cytoplasmic routes. *Trends Microbiol.* **16**: 1–5.
- Shulman, J., Benton, R., and St Johnston, D. 2000. The *Drosophila* homolog of *C.elegans* PAR-1 organizes the oocyte cytoskeleton and directs *oskar* mRNA localization to the posterior pole. *Cell* **101**: 377–388.
- Singh, R., Valcarcel, J., and Green, M.R. 1995. Distinct binding specificities and functions of higher eukaryotic polypyrimidine tract-binding proteins. *Science* **268**: 1173–1176.
- Snee, M., Benz, D., Jen, J., and Macdonald, P.M. 2008. Two distinct domains of Bruno bind specifically to the *oskar* mRNA. *RNA Biol.* **5**: 1–9.
- Song, Y., Tzima, E., Ochs, K., Bassili, G., Trusheim, H., Linder, M., Preissner, K.T., and Niepmann, M. 2005. Evidence for an RNA chaperone function of polypyrimidine tract-binding protein in picornavirus translation. *RNA* **11**: 1809–1824.
- Spellman, R., Rideau, A., Matlin, A., Gooding, C., Robinson, F., McGlincy, N., Grellscheid, S.N., Southby, J., Wollerton, M., and Smith, C.W. 2005. Regulation of alternative splicing by PTB and associated factors. *Biochem. Soc. Trans.* **33**: 457–460.
- Spradling, A.C. 1993. Developmental genetics of oogenesis. In *The development of Drosophila melanogaster* (eds. M. Bate and A. Martinez-Arias), pp. 1–70. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- St Johnston, D. 2005. Moving messages: The intracellular localization of mRNAs. *Nat. Rev. Mol. Cell Biol.* **6**: 363–375.
- St Johnston, D., Beuchle, D., and Nüsslein-Vorhard, C. 1991. *staufer*, a gene required to localize maternal RNAs in *Drosophila* eggs. *Cell* **66**: 51–63.
- Stoneley, M., and Willis, A.E. 2004. Cellular internal ribosome entry segments: Structures, trans-acting factors and regulation of gene expression. *Oncogene* **23**: 3200–3207.
- Valcarcel, J. and Gebauer, F. 1997. Post-transcriptional regulation: The dawn of PTB. *Curr. Biol.* **7**: R705–R708.1016/S0960-9822(06)00361-7.
- Vanzo, N.F., and Ephrussi, A. 2002. Oskar anchoring restricts pole plasm formation to the posterior of the *Drosophila* oocyte. *Development* **129**: 3705–3714.
- Yano, T., López de Quinto, S., Matsui, Y., Shevchenko, A., and Ephrussi, A. 2004. Hrp48, a *Drosophila* hnRNPA/B homolog, binds and regulates translation of *oskar* mRNA. *Dev. Cell* **6**: 637–648.
- Yoshida, S., Muller, H.A., Wodarz, A., and Ephrussi, A. 2004. PKA-R1 spatially restricts Oskar expression for *Drosophila* embryonic patterning. *Development* **131**: 1401–1410.
- Zhou, Z., Sim, J., Griffith, J., and Reed, R. 2002. Purification and electron microscopic visualization of functional human spliceosomes. *Proc. Natl. Acad. Sci.* **99**: 12203–12207.



## ***Drosophila* PTB promotes formation of high-order RNP particles and represses *oskar* translation**

Florence Besse, Sonia López de Quinto, Virginie Marchand, et al.

*Genes Dev.* 2009, **23**: originally published online January 8, 2009  
Access the most recent version at doi:[10.1101/gad.505709](https://doi.org/10.1101/gad.505709)

---

**Supplemental Material** <http://genesdev.cshlp.org/content/suppl/2009/01/08/gad.505709.DC1>

**Related Content** **A splicer that represses (translation)**  
Robin P. Wharton  
[Genes Dev. January , 2009 23: 133-137](#)

**References** This article cites 67 articles, 23 of which can be accessed free at:  
<http://genesdev.cshlp.org/content/23/2/195.full.html#ref-list-1>  
Articles cited in:  
<http://genesdev.cshlp.org/content/23/2/195.full.html#related-urls>

### **License**

**Email Alerting Service** Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

---

horizon  
a PerkinElmer company

Streamline your research with  
**Horizon Discovery's ASO tool**

The advertisement features a dark blue background with a glowing DNA double helix structure in shades of red, orange, and yellow. The text is white and blue, with 'horizon' in a bold, lowercase font and 'a PerkinElmer company' in a smaller font below it. To the right, the text 'Streamline your research with Horizon Discovery's ASO tool' is displayed in a white, sans-serif font.