



# *Drosophila* Studies on Autism Spectrum Disorders

Yao Tian<sup>1</sup> · Zi Chao Zhang<sup>1</sup> · Junhai Han<sup>1</sup>

Received: 28 February 2017 / Accepted: 23 April 2017 / Published online: 9 August 2017  
© Shanghai Institutes for Biological Sciences, CAS and Springer Nature Singapore Pte Ltd. 2017

**Abstract** In the past decade, numerous genes associated with autism spectrum disorders (ASDs) have been identified. These genes encode key regulators of synaptogenesis, synaptic function, and synaptic plasticity. *Drosophila* is a prominent model system for ASD studies to define novel genes linked to ASDs and decipher their molecular roles in synaptogenesis, synaptic function, synaptic plasticity, and neural circuit assembly and consolidation. Here, we review *Drosophila* studies on ASD genes that regulate synaptogenesis, synaptic function, and synaptic plasticity through modulating chromatin remodeling, transcription, protein synthesis and degradation, cytoskeleton dynamics, and synaptic scaffolding.

**Keywords** Autism spectrum disorders · *Drosophila* · Chromatin remodeling · Synaptic scaffolding · Synaptic transmission

## Introduction

Autism spectrum disorders (ASDs) are complex developmental disabilities, whose prevalence is estimated to be 1 in 68 children under 8 years of age in the USA, and they differ substantially between boys (1 in 42) and girls (1 in 189) [1]. The core diagnostic features are impaired social interaction, and repetitive and restrictive behaviors [2]. In addition, children with ASDs frequently present with a host

of associated behavioral issues, such as motor deficits (hypotonia, apraxia, or motor delay), sleep abnormalities, gastrointestinal disturbances, and epilepsy [3–7]. In the past decade, numerous mutations in ASD-associated genes have been identified and various mouse models of monogenic forms of ASDs have been generated and characterized [8]. So far, a variety of standard assessments for ASD-related behavioral phenotypes have been established in mouse models [9]. Meanwhile, ASD models in other animals such as nonhuman primates and *Drosophila* are necessary complements in ASD studies, and are valuable in translating genetic findings and deciphering the shared molecular pathways and phenotypes in ASDs [8, 10].

The fruit fly *Drosophila melanogaster* is a prominent model system in neuroscience. As a model system, it has a wide range of practical and genetic advantages, such as a short generation time (~10 days at room temperature) and a large number of offspring for rapid large-scale analysis (females can lay up to 100 eggs per day). In addition, *Drosophila* has some unique aspects for genetic studies, including the lack of meiotic recombination in males and the use of balancer chromosomes that carry visible genetic markers to facilitate the maintenance of mutant lines [10]. *Drosophila* is also useful for defining gene interaction networks and identifying novel regulatory connections. It offers efficient and high-throughput genetic manipulation, and greatly facilitates the discovery of single gene functions, neurogenetic events, and advanced behaviors [10, 11]. Despite the low anatomical conservation, the biological processes are highly conserved between *Drosophila* and humans at the molecular, cellular, and synaptic levels. About 75% of human disease genes have identifiable homologs in *Drosophila*, 44% of which are sufficiently conserved for functional study [12, 13]. Here, we review the studies in *Drosophila* that characterize the

✉ Junhai Han  
junhaihan@seu.edu.cn

<sup>1</sup> Institute of Life Sciences, The Key Laboratory of Developmental Genes and Human Disease, Southeast University, Nanjing 210096, China

genetic and molecular pathology of ASDs. These studies involve many ASD-associated genes that influence the structure and the turnover of synapses at different levels, including chromatin remodeling, transcription, protein synthesis and degradation, actin cytoskeleton dynamics, and synaptic transmission (Fig. 1).

## Chromatin Remodeling and Transcription


Some important regulators of chromatin remodeling and transcription are promising genetic factors for ASDs. However, how changes in these genes affect neuronal morphology and activity is unclear. Several studies in *Drosophila* have revealed the underlying molecular mechanisms of chromatin remodeling and transcription regulators in neural development and ASD-related behaviors (Figs. 1, 2).

Mutations in *POGZ* have been reported in individuals with ASDs, intellectual disability, and schizophrenia [14–16]. *POGZ* encodes a heterochromatin protein 1  $\alpha$ -binding protein and is hypothesized to function as a transcriptional regulator in molecular networks crucial for neuronal function [17]. Downregulation of *row* (*Drosophila* ortholog of *POGZ*) in neurons leads to deficits in habituation, a form of non-associative learning that is relevant to both intellectual disability and ASDs [18]. Euchromatin histone methyltransferase (*EHMT*) is another

ASD risk gene, which encodes a member of the evolutionarily-conserved protein family methylates histone 3 at lysine 9 [19–21]. In *Drosophila*, loss of *Ehmt* results in a significant decrease of dendrite end number, higher-order branching, and dendritic field complexity, as well as learning and memory deficits [22].

Disrupted-in-schizophrenia-1 (*DISC1*) is associated with a wide range of mental illnesses, including ASDs [23]. *DISC1* interacts with and activates transcription factor 4 (ATF4)/CREB2 in the nucleus [24]. A fly model expressing human *DISC1* has shown that accumulation of exogenous human *DISC1* in the nucleus disturbs sleep homeostasis, implying a deficit in neuronal activity. This function is modulated by interaction with ATF4/CREB2 and recruitment of a co-repressor, N-CoR, to the CRE-mediated transcriptional machinery [25].

MicroRNA (miRNA) is another way to post-transcriptionally regulate gene expression. The autism susceptibility gene *A2bp1* has been identified in *Drosophila* as an mRNA target of miR-980 [26]. MiR-980 inhibition enhances olfactory learning and memory stability, while its overexpression in the mushroom bodies impairs 3-h memory. Overexpression of its target *A2bp1* in the mushroom bodies enhances memory. These defects may be attributed to the role of miR-980 in inhibiting excitability, as projection neurons overexpressing miR-980 exhibit a strong trend for a lower mean firing frequency with an injected current at 40–50 pA [26].




**Fig. 1** ASD-associated genes regulate synaptic function and neural circuits through various cellular events.

## Protein Synthesis and Degradation

Neuronal activity and function are partially determined by synaptic protein levels, which are strictly regulated by protein synthesis and degradation. On the other hand, the levels of synaptic proteins are also influenced by neuronal activity [27]. Mutations of the genes involved in such homeostatic regulation have been found in ASD patients [28]. Numerous studies in *Drosophila* have illustrated that dysfunction of ASD-related genes affects protein synthesis and degradation, and subsequently results in deficits in synaptogenesis and synaptic function, as well as synaptic plasticity (Fig. 2).

The fragile X mental retardation 1 gene (*FMR1*) encodes a pan-neuronal RNA-binding protein, FMR1, that associates with specific mRNAs to repress their translation [29–33]. So far, >800 distinct mRNA targets of FMR1 have been found, and it has been implicated in many aspects of brain development and function [34]. Fragile X Syndrome (FXS), the leading monogenetic cause of autism, is caused by transcriptional silencing of the *FMR1* gene due to a trinucleotide repeat expansion in its 5'-UTR [35, 36]. Since the generation of the first *Fmr1*-knockout

**Fig. 2** Functions of ASD-associated genes in different cellular processes.



mouse model to study FXS, other animal models including the fly FXS model have been further developed and studied, providing more cellular and molecular clues to explain this complex syndrome. There is only one *FMR1* homolog, named *dfmr1* in *Drosophila*, which encodes the dFmr1 protein that shares high identity with mammalian FMR1 in the functional domains [37]. The *Drosophila* neuromuscular junction (NMJ) is a glutamatergic synapse characterized by stereotypic innervation patterns of motor neurons into well-defined target body-wall muscles, making it easier to study synaptogenesis, synaptic transmission, and plasticity [38]. *Drosophila dfmr1* loss-of-function mutants show synapse overelaboration (overgrowth, overbranching, and excess synaptic boutons) in peripheral NMJs [39] as well as in the mushroom body (MB) of the central nervous system [40], accompanied by altered neurotransmission. The hypermorph mutants of *dfmr1* show opposite defects. A further rescue study indicated a pre-synaptic requirement of dFMR1 for synapse structuring, along with both a pre- and post-synaptic requirement for functional neurotransmission [41]. Furthermore, *dfmr1* loss-of-function mutants exhibit more dendritic branching in dendritic arborization neurons and its role in dendrite development is partially mediated by Rac1 as well as microRNA-124a [42, 43]. In addition, deficits in axonal targeting have been extensively reported in *dfmr1* mutants [40, 44–47].

Loss of Fmr1 up-regulates the translation of its target mRNAs [39, 42, 48, 49]. In the fly NMJ, *Adar* acts downstream of *Fmr1* for proper NMJ architecture [50]. The other two dFmr1 targets, the synaptic heparan sulfate proteoglycans glycosylphosphatidyl inositol-anchored Dally-like protein (*Dlp*) and transmembrane Syndecan

(*Sdc*), play important roles in modulating synaptic structure and function [51]. The expression of these two proteins is markedly elevated in *dfmr1*-null NMJs [52]. Bone morphogenetic protein type II receptor (*BMPR2*) is also one of the targets of FMR1. The structural defects at the NMJ in loss-of-function *dfmr1* mutants can be rescued by reducing *Wit*, the *Drosophila* ortholog of *BMPR2* [53]. The role of FMR1 in mRNA translation is regulated by polyglutamine-binding protein 1 (PQBP1), whose mutations cause Renpenning syndrome [54]. *Drosophila* Pqbp1 interacts with Fmr1 and facilitates target mRNA assembly into ribosomes [54]. Two of the common mutations found in Renpenning syndrome, PQBP1 c.459\_462delAGAG and c.463\_464dupAG, encode a distinct C-terminal epitope that preferentially binds non-phosphorylated FMRP and promotes its ubiquitin-mediated degradation [55]. Therefore, PQBP1 c.463\_464dupAG transgenic flies show remarkable defects of synaptic over-growth similar to *dfmr1* mutants, which can be rescued by exogenously-expressed dFmr1 [55].

The mTOR pathway controls global mRNA translation and has been shown to play roles in many fundamental cellular processes including autophagy, transcription, and cytoskeletal dynamics [56]. It is a key regulator of neuronal differentiation [57, 58], and hyper-activation of this pathway increases the risk of ASDs [59]. In *Drosophila*, the gene *unkempt* (*unk*) has been identified as a novel negative regulator of photoreceptor differentiation acting downstream of mTORC1. Unk together with its binding partner headcase (*Hdc*) negatively regulates the InR/mTOR pathway and controls the timing of neurogenesis [60, 61]. Tuberous sclerosis complex (TSC) is an autosomal dominant disorder that shows clinical features of epilepsy and

autism. It is caused by mutations in the *TSC1* or *TSC2* genes. TSC protein regulates synaptic growth *via* the TORC2-Akt pathway, and the mTOR pathway is upregulated by TSC2 protein [59]. In the *Drosophila* NMJ, *Tsc2* mutants show increased synaptic growth [62].

Duplications of the genomic region encompassing *UBE3A* (15q11–q13) are the second most common genetic lesions found in autism (3%–5% of cases) [63]. *UBE3A* is a maternally-expressed gene and its product, the ubiquitin protein ligase E3A, is present in many regions of the brain, with highest expression in the hippocampus and cerebellum [64]. In *Drosophila*, loss of *Ube3a* significantly increases the number of both total and satellite boutons in conjunction with compromised endocytosis in the NMJs [65]. *Ube3a* specifically ubiquitinates the type I BMP receptor *Tkv*, and promotes its proteasomal degradation. Therefore *Ube3a* has a critical role in regulating NMJ development by repressing BMP signaling [65]. Over-expression of *Ube3a* in *Drosophila* results in a decreased number of active zones per bouton and vesicle area, and 40%–50% of larvae intermittently fail to evoke junction potentials at rapid stimulation rates (15 Hz) [66]. Both loss-of-function and over-expression of *Ube3a* decrease dendritic branching, suggesting that the proper level of *Ube3a* is critically important for normal dendritic patterning [67]. Moreover, improper expression of *Ube3a* appears to be detrimental to learning, thereby recapitulating the learning deficits in autism [68]. Over-expression of human *Ube3a* in *Drosophila* that mimics the gene duplication in autism patients has been used to screen *UBE3A* substrates [69]. A key regulator of monoamine synthesis, the gene *Punch* or *GCH1* encodes the enzyme GTP cyclohydrolase I [70]. It is interesting that the mRNA and protein levels of the *Drosophila* vesicular monoamine transporter *dVMAT* are also elevated in the absence of *dFmr1* [71]. The altered monoamine (dopamine/serotonin) synthesis pathway found both in *dUBE3A* and *dFMR1* mutants may provide a potential explanation for the repetitive behaviors and hyperactivity associated with autism and also explain why some individuals with ASDs respond better to selective serotonin reuptake inhibitors than others.

*Highwire* is another highly-conserved E3 ligase associated with ASDs, and it functions presynaptically to negatively regulate synaptic growth at the *Drosophila* NMJ [72]. Mutations of *wallenda*, which encodes an MAP kinase kinase homologous to the vertebrate dual leucine zipper-bearing kinases *DLK* and *LZK*, completely suppress the synaptic overgrowth phenotype in *highwire* mutants [73]. In addition, *Rae1* has been identified as a *Highwire* cofactor to prevent the autophagy-mediated degradation of *Highwire* protein in post-mitotic neurons [74].

## Cell Adhesion Molecules

ASDs also involve many proteins mediating neuronal connectivity and synaptic transmission, such as the synaptic adhesion molecules *neurexin* (*NRX*) and *neuroligin* (*NLG*) in synaptogenesis, various neurotransmitters and proteins associated with synaptic vesicle recycling in synaptic transmission, *contactin-associated proteinlike 2* (*CNTNAP2*) in neuronal conduction, and  $\text{Ca}^{2+}$  channels in ion permeability (Fig. 1). Given that the *Drosophila* genome is relatively less redundant than the human genome, a single mutation in the homolog of an ASD-related gene is more likely to avoid compensatory effects and yield a measurable phenotype.

The well-known autism candidate genes *NRX* and *NLG* are classical trans-synaptic partners that play critical roles in synaptogenesis and synaptic transmission (Fig. 2). Loss of *Nlgs* and *Nrx* results in reduced bouton numbers, aberrant presynaptic and postsynaptic development at NMJs, and impaired synaptic transmission [75–79]. It has been found that *Drosophila* *neuroligin 1* (*dNlg1*) and *dNlg3* act predominantly in pre-synaptic terminals, while *dNlg2* functions both pre- and post-synaptically [75–78]. *Nrx* and *Nlgs* also play critical roles in synaptic transmission. *dNrx* has been shown to functionally couple with  $\text{Ca}^{2+}$  channels to regulate synaptic transmission. And *dNlg4* modulates GABA transmission in large ventral lateral neurons through recruiting  $\text{GABA}_A$  receptors resistance to dieldrin [80]. Interestingly, both *dnrx* mutants and *dnlg4* mutants exhibit reduced night-time sleep, even though they function in different brain regions [80, 81]. In addition, impaired *dNrx* and *dNlgs* result in neuronal plasticity defects [80, 82]. *dNrx* has been shown to interact with N-ethylmaleimide-sensitive factor (NSF), an enzyme that mediates disassembly of the soluble NSF attachment protein receptor (SNARE) complex, and plays an important role in synaptic vesicle release [83]. Besides, *dNrx* is expressed beginning from an early neurodevelopmental stage prior to synaptogenesis. *dNrx* plays an essential role in columnar restriction during L4 axon branching in the *Drosophila* visual system through clustering of one of the classical axon guidance molecules, *Ephrin*, which implies a novel role of *dNrx* in early neural development [84].

*Neurexin IV* (*NrxIV*) is an ortholog of the autism gene *CNTNAP2* [85]. *NrxIV* homozygous-null mutants display reduced bouton numbers, while heterozygous-nulls do not observably differ from the wild-type [86]. The polygenic causes of ASDs have also been investigated in *Drosophila* models using orthologs to human ASD genes with different copy number variants (CNVs) [87]. Two ASD candidate genes encoding adherens junction proteins, *NOTCH1* and *p120ctn* (orthologs of human *NOTCH1* and *catenin delta 2* (*CTNND2*)) that show gain or loss of CNVs, respectively,



were tested pairwise and have shown synergistic effects on NMJ bouton number [87]. These studies provide evidence for synergistic interactions between CNV candidate gene sets, supporting shared and distinct genetic etiologies of ASDs.

## Synaptic Receptors and Ion Channels

The *Drosophila* NMJ is an asymmetric glutamatergic synapse formed between motor neurons and muscle cells. Since it displays some advantageous features, including structural accessibility, stereotypic features, and amenability to genetic manipulations as well as electrophysiological and microscopic analyses, it is considered to be a convenient and useful model for elucidating the mechanisms underlying synapse formation, synaptic transmission and plasticity, and synaptic degeneration. Many mutants of genes encoding the components in synaptic transmission are also associated with ASDs, including synaptic receptors, components in synaptic vesicle cycling, and ion channels (Figs. 1, 2).

Mutations in the *Drosophila* group II metabolic glutamate receptor gene (*DmGluRA*) increase neuronal excitability by preventing PI3 kinase activation and consequently hyper-activating the transcription factor *Foxo* [88]. Loss of *dFmr1* also results in excessive activity of metabotropic glutamate receptors (mGluRs) and learning and memory deficits, which are related to the inhibition of cAMP signaling reported in patients and animal models [36]. These deficits can be rescued by pharmacological inhibition of mGluRs, which provides further support for the agonistic effects of *dFmr1* and mGluR signaling [89].

Several genes involved in the dopamine (DA) network are also associated with ASDs, including the plasma membrane protein syntaxin 1 (*STX1*) [90] and the DA transporter (*DAT*) [91]. A novel *de novo* missense mutation in the human *DAT* (*hDAT*) gene results in a Thr-to-Met substitution at site 356 (*hDAT-T356M*). Expression of *hDAT-T356M* in DA neurons with the *Drosophila DAT*-null allele leads to hyperlocomotion, indicating that alterations in DA homeostasis may confer risks for ASDs and related neuropsychiatric conditions [91]. Another two missense variants, *SLC6A3 R/W* and *STX1A R/Q* [17, 92, 93], disrupt the reverse transport of DA, resulting in DA dysfunction and associated locomotor behavioral abnormalities [94].

Acetylcholine is the major excitatory neurotransmitter in the central nervous system of insects [95]. The  $\alpha 7$  subunit of the nicotinic acetylcholine receptor is one of the most prevalent receptors that are involved in neurological pathologies including autism [96]. In *Drosophila*, *D $\alpha$ 7* protein is enriched at the dendrites of the giant fiber that integrates sensory input, activates flight motor neurons, and mediates synaptic transmission [97].

To date, three GABA receptor subunit classes have been cloned in *Drosophila*. They are *Rdl* (resistant to dieldrin), *Grd* (GABA and glycine-like receptor of *Drosophila*) and *Lcch3* (ligand-gated Cl<sup>-</sup> channel homolog 3) [98]. A significant reduction has been found in all three subunits and glutamic acid decarboxylase in *dfmr1* mutants [99, 100].

Normal synaptic vesicle recycling contributes to functional neurotransmission. Synaptotagmin (*Synj*) is a phosphoinositide phosphatase known to play an important role in synaptic vesicle recycling [101]. Dyrk1A, also known as Minibrain (*Mnb*), is a serine/threonine kinase implicated in ASDs [15, 102]. The protein encoded by the *Drosophila Mnb* gene has been shown to interact with the *IN1* ortholog *Snr1*, which is a chromatin-remodeling factor involved in the morphogenesis of dendritic arbors in *Drosophila* sensory neurons [103, 104]. Phosphorylation of *Synj* by *Mnb* kinase enhances *Synj* activity and is required for reliable synaptic vesicle recycling [105]. Hence, it is not surprising that *Synj* and *Mnb* mutations have been linked to autism [15, 102].

The  $\alpha 2\delta$  gene family plays roles in Ca<sup>2+</sup> channel trafficking and membrane stabilization-dependent synaptic morphogenesis [106], and is associated with a wide range of neurological diseases, including ASDs [17]. Presynaptic homeostatic potentiation is disturbed when  $\alpha 2\delta$ -3 is lost, due to a failure to potentiate presynaptic Ca<sup>2+</sup> influx and the Rab-3 interacting molecule-dependent readily-releasable vesicle pool [107].

## Scaffolding Proteins and the Actin Cytoskeleton

The correct positioning of cell-adhesion molecules, receptors, and channels at the synapse requires the complex assembly of scaffolding proteins and the actin cytoskeleton. Many mutations in these genes are also found in ASD patients (Figs. 1, 2).

The *SHANK* family gene *SHANK3* is considered to be one of the most prevalent causes of ASDs [108, 109]. Prosap/Shank family proteins have multi-domains including ankyrin repeats, SH3, PDZ, proline-rich, and SAM domains, and are the key organizers of the postsynaptic density (PSD) [110]. One possible molecular pathogenesis is an imbalance between excitatory and inhibitory receptors linked with the Nlgs-PSD-95-SHANK complex *via* PDZ binding. In mammals, the Shank family binds Nlgs and functions to coordinate pre/postsynaptic signaling through Neurexin-Neuroigin signaling complexes [111, 112]. There are three *Shank* family genes in mammals but only a single homolog of *Shank* in *Drosophila*, which makes it easier to implement *in vivo* null-mutant studies in *Drosophila* [113]. Both loss and overexpression of *Shank* decrease synaptic bouton number and maturity, and result in defects in the organization of the sub-

synaptic reticulum, a complex system of folding the postsynaptic membrane at the NMJ. Furthermore, Shank regulates a non-canonical Wnt signaling pathway in postsynaptic cells by modulating the internalization of the Wnt receptor Fz2 [114]. These findings imply that *Shank* dosage is critical for synaptic development and establish a novel connection between Shank and synaptic Wnt signaling. The Neurobeachin (*NBEA*) gene has been shown to be disrupted in patients with idiopathic autism [115]. *NBEA* encodes a neuron-specific multi-domain signal scaffold protein that is predominantly expressed in the brain during development [116]. Loss of *Rugose* (*rg*), the *Drosophila* homolog of human *NBEA*, results in abnormal synaptic architecture and physiology [117].

Cytoskeleton dysregulation is one of the major problems in *Fmr1* mutant neurons. The microtubule (MT) network is apparently altered in both loss- and gain-of-function *dfmr1* mutants [48, 118]. The number and transport of mitochondria in axons are also affected by *dFmr1* [118]. Besides, live imaging shows that *dFmr1*-associated mRNA granules are less motile and show decreased directional movement in cultured *dfmr1* mutant neurons, demonstrating that FMR1 indeed regulates the association between mRNA cargos and microtubules [119]. *Drosophila* *Fmr1* targets like *Futsch* and *Profilin* regulate MT- or actin-dependent synaptic growth and function [39, 49]. *Drosophila* *Ank2* is the closest homolog of human *ANK2* and *ANK3*. It is expressed specifically in the nervous system and associates with the presynaptic membrane cytoskeleton, similar to the human autism gene *ANK3* [120]. *Ank2* functions downstream of Spectrin in the anchorage of synaptic microtubules. As a consequence, synaptic stability is severely disrupted in *Ank2* mutants, resulting in a reduction in overall terminal size, withdrawal of synaptic boutons, and disassembly of presynaptic active zones [120, 121]. Knock-down of *Ank2* in mushroom bodies shows normal learning but a significant reduction in short-term memory, suggesting a specific role of *Ank2* in cognition [122]. CNVs at 16p11.2 have recently been implicated in the pathogenesis of ASDs [123], but the genes responsible for the increased risk of ASDs are currently unknown. A Kinesin-2-encoding gene *klp68D* closely related to human *KIF22* at the 16p11.2 locus has been identified by genetic screening using the *Drosophila* NMJ system. Disruption of *klp68D* induces ectopic targeting of motor axons [124], suggesting that Kinesin proteins are important for synaptic connectivity.

## Others

Children born to older parents are at a higher risk for disorders such as schizophrenia and autism. Studies in *Drosophila* models have provided evidence for parental

age-related memory impairment [125]. Bisphenol A (BPA), a widely-used chemical in plastic containers, has been suggested to play a role in developmental disorders including autism [126]. Flies exposed to BPA show many autistic-like behaviors, which emphasizes the importance of environment etiology for neurodevelopmental disorders such as autism.

## Conclusions and Perspective

In the past decade, ASDs have undergone considerable diagnostic evolution. To discover the novel biomarkers and molecular pathology underlying these disorders, several animal models have made contributions. While fruit-flies and humans have very different body plans, *Drosophila* has been a prominent model system in neuroscience since the 1960s for the remarkable similarity at the biological process level, such as a similar origin of the central nervous system [127] and several similar neurobiological processes including membrane excitability, neuronal signaling, and classes of neurotransmitters [128]. Several landmark studies have made discoveries ranging from single genes and neurogenetic events to advanced behaviors. Neuroscientists have taken advantage of *Drosophila* for its relatively simple nervous system, a powerful genetic toolkit, high throughput, and low cost. One of the best successes in *Drosophila* is the Fragile X story, reviewed above. In this case, the fly model can phenocopy FXS patients and FMR1-knockout mice, while pharmacological tests have been used to develop a potential treatment. Moreover, taking advantage of high-throughput screening, several *Fmr1* targets have been screened in *Drosophila*, which help to explain the pathogenesis of FXS.

However, with the increasing thorough research on ASDs and the use of primates in ASD studies, what can be further contributed by research in *Drosophila*? Growing evidence supports a causal role for combinatorial contributions of multiple loci in ASD pathogenesis. Thus, it is necessary to investigate the roles of multiple gene interactions in ASD cases. As a good *in vivo* model with a powerful genetic toolkit, accompanied by the development of informatics-targeted screening, *Drosophila* will definitely show its power in identifying multiple candidate interactions, revealing distinct molecular etiologies underlying ASDs.

**Acknowledgements** We apologize to those whose works have contributed greatly to our knowledge but were not sufficiently reviewed or were not cited owing to space limitations. We thank members of the Han laboratory for proof-reading. This review was supported by the National Natural Science Foundation of China (31471031, 31400927, and 31671045), and the Natural Science Foundation of Jiangsu Province, China (BK20140623).

## References

- Developmental Disabilities Monitoring Network Surveillance Year 2010 Principal Investigators; Centers for Disease Control and Prevention. Prevalence of autism spectrum disorder among children aged 8 years-autism and developmental disabilities monitoring network, 11 sites, United States, 2010. *MMWR Surveill Summ* 2014, 63: 1–21.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. DSM-5 5th ed. Arlington, Virginia: American Psychiatric Association, 2013.
- Schreck KA, Mulick JA. Parental report of sleep problems in children with autism. *J Autism Dev Disord* 2000, 30: 127–135.
- Honomichl RD, Goodlin-Jones BL, Burnham M, Gaylor E, Anders TF. Sleep patterns of children with pervasive developmental disorders. *J Autism Dev Disord* 2002, 32: 553–561.
- Malow BA, Marzec ML, McGrew SG, Wang L, Henderson LM, Stone WL. Characterizing sleep in children with autism spectrum disorders: a multidimensional approach. *Sleep* 2006, 29: 1563–1571.
- Souders MC, Mason TB, Valladares O, Bucan M, Levy SE, Mandell DS, *et al.* Sleep behaviors and sleep quality in children with autism spectrum disorders. *Sleep* 2009, 32: 1566–1578.
- Geschwind DH. Advances in autism. *Ann Rev Med* 2009, 60: 367–380.
- de la Torre-Ubieta L, Won H, Stein JL, Geschwind DH. Advancing the understanding of autism disease mechanisms through genetics. *Nat Med* 2016, 22: 345–361.
- Silverman JL, Yang M, Lord C, Crawley JN. Behavioural phenotyping assays for mouse models of autism. *Nature Rev Neurosci* 2010, 11: 490–502.
- McCammon JM, Sive H. Addressing the genetics of human mental health disorders in model organisms. *Ann Rev Genomics Hum Genet* 2015, 16: 173–197.
- Konopka RJ, Benzer S. Clock mutants of *Drosophila melanogaster*. *Proc Natl Acad Sci U S A* 1971, 68: 2112–2116.
- Reiter LT, Potocki L, Chien S, Gribskov M, Bier E. A systematic analysis of human disease-associated gene sequences in *Drosophila melanogaster*. *Genome Res* 2001, 11: 1114–1125.
- Adams MD, Celniker SE, Holt RA, Evans CA, Gocayne JD, Amanatides PG, *et al.* The genome sequence of *Drosophila melanogaster*. *Science* 2000, 287: 2185–2195.
- Gilissen C, Hehir-Kwa JY, Thung DT, van de Vorst M, van Bon BW, Willemsen MH, *et al.* Genome sequencing identifies major causes of severe intellectual disability. *Nature* 2014, 511: 344–347.
- Iossifov I, Ronemus M, Levy D, Wang Z, Hakker I, Rosenbaum J, *et al.* *De novo* gene disruptions in children on the autistic spectrum. *Neuron* 2012, 74: 285–299.
- Fromer M, Pocklington AJ, Kavanagh DH, Williams HJ, Dwyer S, Gormley P, *et al.* *De novo* mutations in schizophrenia implicate synaptic networks. *Nature* 2011, 506: 179–184.
- Silvia DR, Xin H, Goldberg AP, Poultney CS, Kaitlin S, A Erucment Cicek, *et al.* Synaptic, transcriptional and chromatin genes disrupted in autism. *Nature* 2014, 515: 209–215.
- Stessman HAF, Willemsen MH, Fenckova M, Penn O, Hoischen A, Xiong B, *et al.* Disruption of POGZ is associated with intellectual disability and autism spectrum disorders. *Am J Hum Genet* 2016, 98: 541–552.
- Ogawa H, Ishiguro K, Gaubatz S, Livingston DM, Nakatani Y. A complex with chromatin modifiers that occupies E2F- and Myc-responsive genes in G0 cells. *Science* 2002, 296: 1132–1136.
- Mis J, Ner SS, Grigliatti TA. Identification of three histone methyltransferases in *Drosophila*: dG9a is a suppressor of PEV and is required for gene silencing. *Mol Genet Genomics* 2006, 275: 513–526.
- Tachibana M, Sugimoto K, Nozaki M, Ueda J, Ohta T, Ohki M, *et al.* G9a histone methyltransferase plays a dominant role in euchromatic histone H3 lysine 9 methylation and is essential for early embryogenesis. *Genes Dev* 2002, 16: 1779–1791.
- Kramer JM, Kochinke K, Oortveld MA, Marks H, Kramer D, de Jong EK, *et al.* Epigenetic regulation of learning and memory by *Drosophila* EHMT/G9a. *PLoS Biol* 2011, 9: e1000569.
- Kilpinen H, Ylisaukko-Oja T, Hennah W, Palo OM, Varilo T, Vanhala R, *et al.* Association of DISC1 with autism and Asperger syndrome. *Mol Psychiatr* 2008, 13: 187–196.
- Morris JA, Kandpal G, Ma L, Austin CP. DISC1 (Disrupted-In-Schizophrenia 1) is a centrosome-associated protein that interacts with MAP1A, MIPT3, ATF4/5 and NUDEL: regulation and loss of interaction with mutation. *Hum Mol Genet* 2003, 12: 1591–1608.
- Sawamura N, Ando T, Maruyama Y, Fujimuro M, Mochizuki H, Honjo K, *et al.* Nuclear DISC1 regulates CRE-mediated gene transcription and sleep homeostasis in the fruit fly. *Mol Psychiatr* 2008, 13: 1138–1148.
- Guen-Ozkan T, Busto GU, Schutte SS, Cervantes-Sandoval I, O'Dowd DK, Davis RL. MiR-980 is a memory suppressor microRNA that regulates the autism-susceptibility gene A2bp1. *Cell Rep* 2016, 14: 1698–1709.
- Ma XM, Blenis J. Molecular mechanisms of mTOR-mediated translational control. *Nat Rev Mol Cell Biol* 2009, 10: 307.
- Iii RJK, Bear MF. The autistic neuron: troubled translation? *Cell* 2008, 135: 401–406.
- Ashley CT, Jr., Wilkinson KD, Reines D, Warren ST. FMR1 protein: conserved RNP family domains and selective RNA binding. *Science* 1993, 262: 563–566.
- Brown V, Jin P, Ceman S, Darnell JC, O'Donnell WT, Tenenbaum SA, *et al.* Microarray identification of FMRP-associated brain mRNAs and altered mRNA translational profiles in fragile X syndrome. *Cell* 2001, 107: 477–487.
- Ascano M, Jr., Mukherjee N, Bandaru P, Miller JB, Nusbaum JD, Corcoran DL, *et al.* FMRP targets distinct mRNA sequence elements to regulate protein expression. *Nature* 2012, 492: 382–386.
- O'Donnell WT, Warren ST. A decade of molecular studies of fragile X syndrome. *Ann Rev Neurosci* 2002, 25: 315–338.
- Siomi H, Siomi MC, Nussbaum RL, Dreyfuss G. The protein product of the fragile-X gene, FMR1, has characteristics of an rna-binding protein. *Cell* 1993, 74: 291–298.
- Darnell JC, Van Driesche SJ, Zhang C, Hung KY, Mele A, Fraser CE, *et al.* FMRP stalls ribosomal translocation on mRNAs linked to synaptic function and autism. *Cell* 2011, 146: 247–261.
- Pieretti M, Zhang FP, Fu YH, Warren ST, Oostra BA, Caskey CT, *et al.* Absence of expression of the FMR-1 gene in fragile X syndrome. *Cell* 1991, 66: 817–822.
- Kanellopoulos AK, Semelidou O, Kotini AG, Anezaki M, Skoulakis EM. Learning and memory deficits consequent to reduction of the fragile X mental retardation protein result from metabotropic glutamate receptor-mediated inhibition of cAMP signaling in *Drosophila*. *J Neurosci* 2012, 32: 13111–13124.
- Wan L, Dockendorff TC, Jongens TA, Dreyfuss G. Characterization of dFMR1, a *Drosophila melanogaster* homolog of the fragile X mental retardation protein. *Mol Cell Biol* 2000, 20: 8536–8547.
- Schwarz TL. Transmitter release at the neuromuscular junction. *Int Rev Neurobiol* 2006, 75: 105–144.
- Zhang YQ, Bailey AM, Matthies HJ, Renden RB, Smith MA, Speese SD, *et al.* *Drosophila* fragile X-related gene regulates the

- MAP1B homolog Futsch to control synaptic structure and function. *Cell* 2001, 107: 591–603.
40. Pan L, Zhang YQ, Woodruff E, Broadie K. The *Drosophila* fragile X gene negatively regulates neuronal elaboration and synaptic differentiation. *Curr Biol* 2004, 14: 1863–1870.
  41. Gatto CL, Broadie K. Temporal requirements of the fragile X mental retardation protein in the regulation of synaptic structure. *Development* 2008, 135: 2637–2648.
  42. Lee A, Li W, Xu K, Bogert BA, Su K, Gao FB. Control of dendritic development by the *Drosophila* fragile X-related gene involves the small GTPase Rac1. *Development* 2003, 130: 5543–5552.
  43. Xu XL, Li Y, Wang F, Gao FB. The steady-state level of the nervous-system-specific microRNA-124a is regulated by dFMR1 in *Drosophila*. *J Neurosci* 2008, 28: 11883–11889.
  44. Dockendorff TC, Su HS, McBride SM, Yang Z, Choi CH, Siwicki KK, *et al.* *Drosophila* lacking *dfmr1* activity show defects in circadian output and fail to maintain courtship interest. *Neuron* 2002, 34: 973–984.
  45. Morales J, Hiesinger PR, Schroeder AJ, Kume K, Verstreken P, Jackson FR, *et al.* *Drosophila* fragile X protein, DFXR, regulates neuronal morphology and function in the brain. *Neuron* 2002, 34: 961–972.
  46. Tessier CR, Broadie K. *Drosophila* fragile X mental retardation protein developmentally regulates activity-dependent axon pruning. *Development* 2008, 135: 1547–1557.
  47. Michel CI, Kraft R, Restifo LL. Defective neuronal development in the mushroom bodies of *Drosophila* fragile X mental retardation 1 mutants. *J Neurosci* 2004, 24: 5798–5809.
  48. Lu R, Wang H, Liang Z, Ku L, O'Donnell W T, Li W, *et al.* The fragile X protein controls microtubule-associated protein 1B translation and microtubule stability in brain neuron development. *Proc Natl Acad Sci U S A* 2004, 101: 15201–15206.
  49. Reeve SP, Bassetto L, Genova GK, Kleyner Y, Leyssen M, Jackson FR, *et al.* The *Drosophila* fragile X mental retardation protein controls actin dynamics by directly regulating profilin in the brain. *Curr Biol* 2005, 15: 1156–1163.
  50. Bhogal B, Jepson JE, Savva YA, Pepper AS, Reenan RA, Jongsens TA. Modulation of dADAR-dependent RNA editing by the *Drosophila* fragile X mental retardation protein. *Nat Neurosci* 2011, 14: 1517–1524.
  51. Johnson KG, Tenney AP, Ghose A, Duckworth AM, Higashi ME, Parfitt K, *et al.* The HSPGs Syndecan and Dallylike bind the receptor phosphatase LAR and exert distinct effects on synaptic development. *Neuron* 2006, 49: 517–531.
  52. Friedman SH, Dani N, Rushton E, Broadie K. Fragile X mental retardation protein regulates trans-synaptic signaling in *Drosophila*. *Dis Model Mech* 2013, 6: 1400–1413.
  53. Kashima R, Roy S, Ascano M, Martinez-Cerdeno V, Ariza-Torres J, Kim S, *et al.* Augmented noncanonical BMP type II receptor signaling mediates the synaptic abnormality of fragile X syndrome. *Sci Signal* 2016, 9: ra58.
  54. Wan D, Zhang ZC, Zhang X, Li Q, Han J. X chromosome-linked intellectual disability protein PQBP1 associates with and regulates the translation of specific mRNAs. *Hum Mol Genet* 2015, 24: 4599–4614.
  55. Zhang XY, Qi J, Shen YQ, Liu X, Liu A, Zhou Z, *et al.* Mutations of PQBP1 in Renpenning syndrome promote ubiquitin-mediated degradation of FMRP and cause synaptic dysfunction. *Hum Mol Genet* 2017 (in press).
  56. Laplante M, Sabatini DM. mTOR signaling in growth control and disease. *Cell* 2012, 149: 274–293.
  57. Bateman JM, McNeill H. Insulin/IGF signalling in neurogenesis. *Cell Mol Life Sci* 2006, 63: 1701–1705.
  58. Fishwick KJ, Li RA, Halley P, Deng P, Storey KG. Initiation of neuronal differentiation requires PI3-kinase/TOR signalling in the vertebrate neural tube. *Dev Biol* 2010, 338: 215–225.
  59. Orlova KA, Crino PB. The tuberous sclerosis complex. Springer, Vienna, 2010.
  60. Avet-Rochex A, Carvajal N, Christoforou CP, Yeung K, Maierbrugger KT, Hobbs C, *et al.* *Unkempt* is negatively regulated by mTOR and uncouples neuronal differentiation from growth control. *PLoS Genet* 2014, 10: e1004624.
  61. Bateman JM. Mechanistic insights into the role of mTOR signaling in neuronal differentiation. *Neurogenesis (Austin, TX)* 2015, 2: e1058684.
  62. Natarajan R, Trivedi-Vyas D, Wairkar YP. Tuberous sclerosis complex regulates *Drosophila* neuromuscular junction growth via the TORC2/Akt pathway. *Hum Mol Genet* 2013, 22: 2010–2023.
  63. Moreno-De-Luca D, Sanders SJ, Willsey AJ, Mulle JG, Lowe JK, Geschwind DH, *et al.* Using large clinical data sets to infer pathogenicity for rare copy number variants in autism cohorts. *Mol Psychiatr* 2013, 18: 1090–1095.
  64. Dindot SV, Antalffy BA, Bhattacharjee MB, Beaudet AL. The Angelman syndrome ubiquitin ligase localizes to the synapse and nucleus, and maternal deficiency results in abnormal dendritic spine morphology. *Hum Mol Genet* 2008, 17: 111–118.
  65. Li W, Yao A, Zhi H, Kaur K, Zhu YC, Jia M, *et al.* Angelman syndrome protein Ube3a regulates synaptic growth and endocytosis by inhibiting BMP signaling in *drosophila*. *PLoS Genet* 2016, 12: e1006062.
  66. Valdez C, Scroggs R, Chassen R, Reiter LT. Variation in *Dube3a* expression affects neurotransmission at the *Drosophila* neuromuscular junction. *Biol Open* 2015, 4: 776–782.
  67. Lu Y, Wang F, Li Y, Ferris J, Lee JA, Gao FB. The *Drosophila* homologue of the Angelman syndrome ubiquitin ligase regulates the formation of terminal dendritic branches. *Hum Mol Genet* 2009, 18: 454–462.
  68. Chakraborty M, Paul BK, Nayak T, Das A, Jana NR, Bhutani S. The E3 ligase *ube3a* is required for learning in *Drosophila melanogaster*. *Biochem Biophys Res Commun* 2015, 462: 71–77.
  69. Reiter LT, Seagroves TN, Bowers M, Bier E. Expression of the Rho-GEF Pbl/ECT2 is regulated by the UBE3A E3 ubiquitin ligase. *Hum Mol Genet* 2006, 15: 2825–2835.
  70. Ferdousy F, Bodeen W, Summers K, Doherty O, Wright O, Elsisni N, *et al.* *Drosophila* *ube3a* regulates monoamine synthesis by increasing GTP cyclohydrolase I activity via a non-ubiquitin ligase mechanism. *Neurobiol Dis* 2011, 41: 669–677.
  71. Tauber JM, Vanlandingham PA, Zhang B. Elevated levels of the vesicular monoamine transporter and a novel repetitive behavior in the *Drosophila* model of fragile X syndrome. *PLoS One* 2011, 6: e27100.
  72. Wan HI, DiAntonio A, Fetter RD, Bergstrom K, Strauss R, Goodman CS. Highwire regulates synaptic growth in *Drosophila*. *Neuron* 2000, 26: 313–329.
  73. Collins CA, Wairkar YP, Johnson SL, DiAntonio A. Highwire restrains synaptic growth by attenuating a MAP kinase signal. *Neuron* 2006, 51: 57–69.
  74. Tian X, Li J, Valakh V, DiAntonio A, Wu C. *Drosophila* *Rael* controls the abundance of the ubiquitin ligase Highwire in post-mitotic neurons. *Nat Neurosci* 2011, 14: 1267–1275.
  75. Chen YC, Lin YQ, Banerjee S, Venken K, Li J, Ismat A, *et al.* *Drosophila* *neuroligin 2* is required presynaptically and postsynaptically for proper synaptic differentiation and synaptic transmission. *J Neurosci* 2015, 32: 16018–16030.



76. Zeng X, Sun M, Liu L, Chen F, Wei L, Xie W. Neurexin-1 is required for synapse formation and larvae associative learning in *Drosophila*. *FEBS Lett* 2007, 581: 2509–2516.
77. Li J, Ashley J, Budnik V, Bhat MA. Crucial role of *Drosophila* neurexin in proper active zone apposition to postsynaptic densities, synaptic growth, and synaptic transmission. *Neuron* 2007, 55: 741–755.
78. Sun MK, Xing GL, Yuan LD, Gan GM, Knight D, With SI, *et al.* Neuroligin 2 is required for synapse development and function at the *Drosophila* neuromuscular junction. *J Neurosci* 2011, 31: 687–699.
79. Xing G, Gan G, Chen D, Sun M, Yi J, Lv H, *et al.* *Drosophila* neuroligin3 regulates neuromuscular junction development and synaptic differentiation. *J Biol Chem* 2014, 289: 31867–31877.
80. Li Y, Zhou ZK, Zhang XW, Tong HW, Li PP, Zhang ZC, *et al.* *Drosophila* neuroligin 4 regulates sleep through modulating GABA transmission. *J Neurosci* 2013, 33: 15545–15554.
81. Tong H, Li Q, Zhang ZC, Li Y, Han J. Neurexin regulates nighttime sleep by modulating synaptic transmission. *Sci Rep* 2016, 6: 38246.
82. Larkin A, Chen MY, Kirszenblat L, Reinhard J, Swinderen BV, Claudianos C. Neurexin-1 regulates sleep and synaptic plasticity in *Drosophila melanogaster*. *Eur J Neurosci* 2015, 42: 2455–2466.
83. Li T, Tian Y, Li Q, Chen H, Lv H, Xie W, *et al.* The neurexin/N-ethylmaleimide-sensitive factor (NSF) interaction regulates short term synaptic depression. *J Biol Chem* 2015, 290: 17656–17667.
84. Liu L, Tian Y, Zhang XY, Zhang X, Li T, Xie W, *et al.* Neurexin restricts axonal branching in columns by promoting ephrin clustering. *Dev Cell* 2017, 41: 94–106 e104.
85. Poliak S. Caspr2, a new member of the neurexin superfamily, is localized at the juxtaparanodes of myelinated axons and associates with K<sup>+</sup> channels. *Neuron* 2000, 24: 1037–1047.
86. Sun MK, Liu LJ, Zeng XK, Xu M, Liu L, Fang M, *et al.* Genetic interaction between Neurexin and CAKI/CMG is important for synaptic function in *Drosophila* neuromuscular junction. *Neurosci Res* 2009, 64: 362–371.
87. Grice SJ, Liu JL, Webber C. Synergistic interactions between *Drosophila* orthologues of genes spanned by *de novo* human CNVs support multiple-hit models of autism. *PLoS Genet* 2015, 11: e1004998.
88. Howlett E, Lin CJ, Lavery W, Stern M. A PI3-Kinase-mediated negative feedback regulates neuronal excitability. *PLoS Genet* 2008, 4: 212–220.
89. McBride SM, Choi CH, Wang Y, Liebelt D, Braunstein E, Ferreiro D, *et al.* Pharmacological rescue of synaptic plasticity, courtship behavior, and mushroom body defects in a *Drosophila* model of fragile X syndrome. *Neuron* 2005, 45: 753–764.
90. Nakamura K, Anitha A, Yamada K, Tsujii M, Iwayama Y, Hattori E, *et al.* Genetic and expression analyses reveal elevated expression of syntaxin 1A (STX1A) in high functioning autism. *Int J Neuropsychopharmacol* 2008, 11: 1073–1084.
91. Hamilton PJ, Campbell NG, Sharma S, Erreger K, Herborg Hansen F, Saunders C, *et al.* De novo mutation in the dopamine transporter gene associates dopamine dysfunction with autism spectrum disorder. *Mol Psychiatr* 2013, 18: 1315–1323.
92. Iossifov I, O’Roak BJ, Sanders SJ, Ronemus M, Krumm N, Levy D, *et al.* The contribution of de novo coding mutations to autism spectrum disorder. *Nature* 2014, 515: 216–221.
93. Neale BM, Kou Y, Liu L, Ma’ayan A, Samocha KE, Sabo A, *et al.* Patterns and rates of exonic de novo mutations in autism spectrum disorders. *Nature* 2012, 485: 242–245.
94. Cartier E, Hamilton PJ, Belovich AN, Shekar A, Campbell NG, Saunders C, *et al.* Rare autism-associated variants implicate syntaxin 1 (STX1 R26Q) phosphorylation and the dopamine transporter (hDAT R51W) in dopamine neurotransmission and behaviors. *EBioMedicine* 2015, 2: 135–146.
95. Gauthier M. State of the art on insect nicotinic acetylcholine receptor function in learning and memory. *Adv Exp Med Biol* 2010, 683: 97–115.
96. Valles AS, Barrantes FJ. Chaperoning alpha 7 neuronal nicotinic acetylcholine receptors. *Biochim Biophys Acta* 2012, 1818: 718–729.
97. Fayyazuddin A, Zaheer MA, Hiesinger PR, Bellen HJ. The nicotinic acetylcholine receptor D $\alpha$ 7 is required for an escape behavior in *Drosophila*. *PLoS Biol* 2006, 4: e63.
98. Hosie AM, Aronstein K, Sattelle DB. Molecular biology of insect neuronal GABA receptors. *Trends Neurosci* 1997, 20: 578–583.
99. D’Hulst C, De Geest N, Reeve SP, Van Dam D, De Deyn PP, Hassan BA, *et al.* Decreased expression of the GABAA receptor in fragile X syndrome. *Brain Res* 2006, 1121: 238–245.
100. Gatto CL, Pereira D, Broadie K. GABAergic circuit dysfunction in the *Drosophila* Fragile X syndrome model. *Neurobiol Dis* 2014, 65: 142–159.
101. Harris TW, Hartweg E, Horvitz HR, Jorgensen EM. Mutations in synaptojanin disrupt synaptic vesicle recycling. *J Cell Biol* 2000, 150: 589–600.
102. O’Roak BJ, Vives L, Fu W, Egertson JD, Stanaway IB, Phelps IG, *et al.* Multiplex targeted sequencing identifies recurrently mutated genes in autism spectrum disorders. *Science* 2012, 338: 1619–1622.
103. Kinstrie R, Lochhead PA, Sibbet G, Morrice N, Cleghon V. dDYRK2 and Minibrain interact with the chromatin remodelling factors SNR1 and TRX. *Biochem J* 2006, 398: 45–54.
104. Parrish JZ, Kim MD, Jan LY, Jan YN. Genome-wide analyses identify transcription factors required for proper morphogenesis of *Drosophila* sensory neuron dendrites. *Genes Dev* 2006, 20: 820–835.
105. Chen CK, Bregere C, Paluch J, Lu JF, Dickman DK, Chang KT. Activity-dependent facilitation of Synaptojanin and synaptic vesicle recycling by the Minibrain kinase. *Nat Commun* 2014, 5: 4246.
106. Kurshan PT, Oztan A, Schwarz TL. Presynaptic  $\alpha$ 2 $\delta$ -3 is required for synaptic morphogenesis independent of its Ca<sup>2+</sup>-channel functions. *Nat Neurosci* 2009, 12: 1415–1423.
107. Wang TT, Jones RT, Whippen JM, Davis GW.  $\alpha$ 2 $\delta$ -3 is required for rapid transsynaptic homeostatic signaling. *Cell Rep* 2016, 16: 2875–2888.
108. Betancur C, Buxbaum JD. SHANK3 haploinsufficiency: a “common” but underdiagnosed highly penetrant monogenic cause of autism spectrum disorders. *Mol Autism* 2013, 4: 17–19.
109. Kolevzon A, Cai G, Soorya L, Takahashi N, Grodberg D, Kajiwara Y, *et al.* Analysis of a purported SHANK3 mutation in a boy with autism: clinical impact of rare variant research in neurodevelopmental disabilities. *Brain Res* 2011, 1380: 98–105.
110. Naisbitt S, Kim E, Tu JC, Xiao B, Sala C, Valtchanoff J, *et al.* Shank, a novel family of postsynaptic density proteins that binds to the NMDA receptor/PSD-95/GKAP complex and cortactin. *Neuron* 1999, 23: 569–582.
111. Meyer G, Varoqueaux F, Neeb A, Oschlies M, Brose N. The complexity of PDZ domain-mediated interactions at glutamatergic synapses: a case study on neuroligin. *Neuropharmacology* 2004, 47: 724–733.
112. Arons MH, Thynne CJ, Grabrucker AM, Li D, Schoen M, Cheyne JE, *et al.* Autism-associated mutations in ProSAP2/Shank3 impair synaptic transmission and neurexin-neuroligin-mediated transsynaptic signaling. *J Neurosci* 2012, 32: 14966–14978.
113. Liebl FL, Featherstone DE. Identification and investigation of *Drosophila* postsynaptic density homologs. *Bioinform Biol Insights* 2007, 2: 375–387.

114. Harris KP, Akbergenova Y, Cho RW, Baas-Thomas MS, Littleton JT. Shank modulates postsynaptic Wnt signaling to regulate synaptic development. *J Neurosci* 2016, 36: 5820–5832.
115. Castermans D, Wilquet V, Parthoens E, Huysmans C, Steyaert J, Swinnen L, *et al.* The neurobeachin gene is disrupted by a translocation in a patient with idiopathic autism. *J Med Genet* 2003, 40: 352–356.
116. Savelyeva L, Sagulenko E, Schmitt JG, Schwab M. The neurobeachin gene spans the common fragile site FRA13A. *Hum Genet* 2006, 118: 551–558.
117. Wise A, Tenezaca L, Fernandez RW, Schatoff E, Flores J, Ueda A, *et al.* *Drosophila* mutants of the autism candidate gene neurobeachin (*rugose*) exhibit neuro-developmental disorders, aberrant synaptic properties, altered locomotion, and impaired adult social behavior and activity patterns. *J Neurogenet* 2015, 29: 135–143.
118. Yao A, Jin S, Li X, Liu Z, Ma X, Tang J, *et al.* *Drosophila* FMRP regulates microtubule network formation and axonal transport of mitochondria. *Hum Mol Genet* 2011, 20: 51–63.
119. Estes PS, O’Shea M, Clasen S, Zarnescu DC. Fragile X protein controls the efficacy of mRNA transport in *Drosophila* neurons. *Mol Cell Neurosci* 2008, 39: 170–179.
120. Koch I, Schwarz H, Beuchle D, Goellner B, Langegger M, Aberle H. *Drosophila* ankyrin 2 is required for synaptic stability. *Neuron* 2008, 58: 210–222.
121. Pielage J, Cheng L, Fetter RD, Carlton PM, Sedat JW, Davis GW. A novel presynaptic giant ankyrin stabilizes the NMJ through regulation of presynaptic microtubules and trans-synaptic cell adhesion. *Neuron* 2008, 58: 195–209.
122. Iqbal Z, Vandeweyer G, van der Voet M, Waryah AM, Zahoor MY, Besseling JA, *et al.* Homozygous and heterozygous disruptions of ANK3: at the crossroads of neurodevelopmental and psychiatric disorders. *Hum Mol Genet* 2013, 22: 1960–1970.
123. Kumar RA, KaraMohamed S, Sudi J, Conrad DF, Brune C, Badner JA, *et al.* Recurrent 16p11.2 microdeletions in autism. *Hum Mol Genet* 2008, 17: 628–638.
124. Park SM, Littleton JT, Park HR, Lee JH. *Drosophila* homolog of human KIF22 at the autism-linked 16p11.2 loci influences synaptic connectivity at larval neuromuscular junctions. *Exp Neurol* 2016, 25: 33–39.
125. Burns JG, Mery F. Transgenerational memory effect of ageing in *Drosophila*. *J Evol Biol* 2010, 23: 678–686.
126. Cock MD, Maas YGH, Bor MVD. Does perinatal exposure to endocrine disruptors induce autism spectrum and attention deficit hyperactivity disorders? Review. *Acta Paediatr* 2012, 101: 811–818.
127. Hirth F, Reichert H. Conserved genetic programs in insect and mammalian brain development. *Bioessays* 1999, 21: 677–684.
128. O’Kane CJ. *Drosophila* as a model organism for the study of neuropsychiatric disorders. *Curr Top Behav Neurosci* 2011, 7: 37–60.