



OPEN

Drought stress triggers proteomic changes involving lignin, flavonoids and fatty acids in tea plants

Honglian Gu¹, Yu Wang¹, Hui Xie¹, Chen Qiu¹, Shuning Zhang¹, Jun Xiao², Hongyan Li³, Liang Chen⁴, Xinghui Li⁵ & Zhaotang Ding¹✉

Drought stress triggers a series of physiological and biochemical changes in tea plants. It is well known that flavonoids, lignin and long-chain fatty acids play important roles in drought resistance. However, changes in proteins related to these three metabolic pathways in tea plants under drought stress have not been reported. We analysed the proteomic profiles of tea plants by tandem mass tag and liquid chromatography-tandem mass spectrometry. A total of 4789 proteins were identified, of which 11 and 100 showed up- and downregulation, respectively. The proteins related to the biosynthesis of lignin, flavonoids and long-chain fatty acids, including phenylalanine ammonia lyase, cinnamoyl-CoA reductase, peroxidase, chalcone synthase, flavanone 3-hydroxylase, flavonol synthase, acetyl-CoA carboxylase 1, 3-ketoacyl-CoA synthase 6 and 3-ketoacyl-CoA reductase 1, were downregulated. However, the contents of soluble proteins, malondialdehyde, total phenols, lignin and flavonoids in the tea plants increased. These results showed that tea plants might improve drought resistance by inhibiting the accumulation of synthases related to lignin, flavonoids and long-chain fatty acids. The proteomic spectrum of tea plants provides a scientific basis for studying the pathways related to lignin, flavonoid and long-chain fatty acid metabolism in response to drought stress.

Tea plants, which require relatively humid environment, are often confronted with drought stress throughout their lifecycle. Drought stress is one of the most important environmental stresses that adversely affects the growth and quality of tea plants. It has been reported that drought stress can reduce tea production by 14–33% and can increase tea plant mortality by 6–19%¹. Tea plants undergo a series of complex morphological, physiological and molecular changes to resist drought stress. Previous studies have revealed several key features of tea plants under water deficit stress, including small leaves, well-developed root systems, relatively thick cuticles and palisade tissue. Moreover, their stomata become closed, there is a loss of cell wall semipermeability, respiration and photosynthesis decrease, proteolysis is accelerated, and carbohydrate synthesis is reduced. Free radical content, antioxidative systems, and osmo-protectant contents were enhanced. In addition, the contents of some characteristic tea plant components, such as the contents of caffeine, polyphenols, and theanine, which are components composing tea aroma, and the corresponding synthetic genes, together with the expression of many related genes, are also affected by drought stress^{2–8}. Moreover, other plant species also show a series of complex changes related to oxidative stress and antioxidant defence to resist drought stress^{9–11}. Many studies have shown that flavonoids, lignin and long-chain fatty acids (LCFAs) play important roles in drought resistance^{12–14}. Nevertheless, the biosynthesis pathways of these components in response to drought stress are poorly understood.

With the improvement in proteomics technology, the study of stress proteins involved in key pathways of tea plants under drought stress has become possible. Previous proteomic analyses showed that many photosynthetic proteins were significantly downregulated in tea plants under drought stress¹⁵. Previous proteomic analysis also showed that 23 proteins involved in redox status, metabolism and the defence response were upregulated in tea plant seeds under desiccation¹⁶. However, to date, little research has focused on proteins related to the biosynthesis of lignin, flavonoids and fatty acids in tea plants in response to drought stress. Therefore, we used tandem mass tag (TMT) and liquid chromatography–mass spectrometry (LC–MS) to study the global profile of

¹Tea Research Institute, Qingdao Agricultural University, Qingdao 266109, Shandong, China. ²School of Biological Science and Winery Engineering, Taishan University, Taian 271000, Shandong, China. ³Haiyang Fruit Technology Promotion Station, Yantai 265100, Shandong, China. ⁴Tea Research Institute, Chinese Academy of Agricultural Sciences, Hangzhou 310008, Zhejiang, China. ⁵Tea Research Institute, Nanjing Agricultural University, Nanjing 210095, Jiangsu, China. ✉email: dzttea@163.com

proteomes of tea plants under drought stress. We then used a bioinformatics method to analyse the Gene Ontology (GO) terms, Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways and protein–protein interactions (PPIs). These results provide insights into the complex molecular mechanisms associated with lignin, flavonoid and fatty acid biosynthesis in tea plants under drought stress and provide an important basis for improving the quality of tea plants under drought conditions.

Materials and methods

Plant materials, stress treatments and physiological determinations. Two-year-old *Camellia sinensis* (L.) O. Kuntze ‘Zhongcha 108’ plants were cultivated under conditions of 12 h light (25 °C)/12 h dark (20 °C) in a growth chamber with 1800 $\mu\text{E m}^{-2} \text{s}^{-1}$ light intensity and 75% humidity for 2 weeks. Drought stress was imposed by withholding watering for 96 h (DT). Moreover, well-watered plants (CK) were also included. Finally, the third and/or fourth mature leaf was taken from the terminal bud for further experiments. The samples were frozen in liquid nitrogen and then stored at -80 °C for physiological and proteomic analyses. Three biological replicates were included.

According to the manufacturer’s instructions (Suzhou Comin Biotechnology Co., Ltd., China), the contents of Cpr, malondialdehyde (MDA), total phenols (TP), flavonoids, and lignin and the activity of phenylalanine ammonia lyase (PAL) were determined on a microplate reader using a BCA protein content kit (serial number: BCAP-1-W), an MDA kit (serial number: MDA-1-Y), a TP kit (serial number: TP-1-G), a plant flavonoid kit (serial number: LHT-1-G), a lignin content kit (serial number: MZS-1-G), and a PAL kit (serial number: PAL-1-Y), respectively. These physiological measurements were determined according to previously reported methods, with slight modifications.

Protein extraction and trypsin digestion. The protein extraction and digestion of the samples were performed according to previous methods^{17,18}. In brief, four volumes of lysis buffer were added to the samples, followed by sonication and centrifugation. The lysis buffer consisted of 8 M urea, 1% Triton-100, 10 mM dithiothreitol, 1% protease inhibitor cocktail, 3 μM TSA, 50 mM NAM inhibitor and 2 mM EDTA. Finally, the precipitate was reconstituted with 8 M urea, and the protein concentration was measured by a BCA kit. For digestion, the protein solution was reduced with 5 mM dithiothreitol at 56 °C for 30 min and then alkylated with 11 mM iodoacetamide at room temperature for 15 min in darkness.

TMT labelling and HPLC fractionation. The peptides were desalted through a Strata X C18 SPE column (Phenomenex) and freeze dried under vacuum. The samples were then reconstituted in 0.5 M TEAB and processed according to the TMT kit instructions. The specific procedure was performed in accordance with the methods of previous research¹⁸.

For HPLC fractionation, the peptides were fractionated by high-pH reversed-phase HPLC using an Agilent 300 Extend C18 column. The specifications and dimensions of the C18 column were as follows: 5 μm and 4.6 \times 250 mm. The operation was as follows: the peptide gradient involved 8–32% acetonitrile (pH 9) for 60 min, divided into 60 fractions, after which the peptides were combined into 18 fractions.

LC–MS/MS analysis. The tryptic peptides were dissolved in 0.1% formic acid (solvent A) and separated using an EASY-nLC 1000 system. The gradient elution procedure was as follows: a linear gradient involving an increase from 7 to 25% of solvent B (0.1% formic acid in 98% acetonitrile) for 26 min, an increase from 25 to 36% of solvent B for 8 min, an increase to 80% of solvent B for 3 min, after which it was for 3 min. The flow rate was constant at 700 nL/min.

The peptides were subjected to an NSI source followed by tandem mass spectrometry (MS/MS) on an OrbitrapFusion™ instrument (Thermo). The electrospray voltage applied was 2.0 kV. Both the peptide precursor ion and its secondary fragments were detected and analysed using a high-resolution Orbitrap device. The primary mass spectrometer scan range was 350–1550 m/z , and intact peptides were detected in the Orbitrap at a resolution of 60,000. The fragments were detected in the Orbitrap at a resolution of 15,000. A data-dependent procedure alternated between one MS scan followed by 20 MS/MS scans, with a 15.0 s dynamic exclusion. Automatic gain control was set to 5E4. The mass spectrometry proteomic data are available via ProteomeXchange under identifier PXD011688.

PRM analyses. PRM mass spectrometric analysis was performed using a Q Exactive™ Plus tandem MS/MS instrument (Thermo). The LC parameters, electrospray voltage, scan range, and Orbitrap resolution were the same as those of the TMT methods. The AGC was set at 3E6 for full MS and 1E5 for MS/MS. The maximum IT was set to 50 ms for full MS and ‘auto’ for MS/MS. The isolation window for MS/MS was set at 1.6 m/z . The enzyme was set as trypsin [KR/P], and the max missed cleavage was set as 0. The peptide length was set to 7–25. The product ions were set from ion 3 to the last ion, and the ion match tolerance was set to 0.02 Da.

Database search and bioinformatic analysis. The MS/MS data were processed using the MaxQuant search engine (v.1.5.2.8). The tandem mass spectra were searched against the *Camellia sinensis* genome database (https://www.plantkingdomgdb.com/tea_tree/) concatenated with the reverse decoy database. Trypsin/P was specified as a cleavage enzyme allowing up to 1 missing cleavage. The minimum length of the peptide was set to 7 amino acid residues, and the maximum number of decoration settings was 5. The mass tolerance for precursor ions was set to 20 ppm in the first search range, 5 ppm in the main search and 0.02 Da for the fragment ions. The variable modification was set to ‘oxidation of methionine, acetylation of the N-terminus of the protein, Iodo




Figure 1. Morphological and physiological analyses of tea plants under drought stress. (A) The phenotypes of CK and DT, (B) the content of Cpr, (C) the content of MDA, (D) the activity of PAL, (E) the content of TP, (F) the content of flavonoids, and (G) the content of lignin were determined, with standard error bars from three replicates shown. The different letters within each column indicate significant differences between treatments at the $p < 0.05$ level.

Sample	Cpr	MDA	PAL	TP	Flavonoids	Lignin
CK	126.25 ± 2.16b	46.85 ± 1.85b	137.06 ± 4.28a	21.01 ± 1.29b	11.20 ± 0.42b	91.24 ± 3.17b
DT	196.83 ± 3.92a	53.78 ± 1.31a	138.17 ± 7.51a	32.23 ± 1.66a	14.03 ± 0.47a	120.83 ± 8.55a

Table 1. Statistical analysis of the physiological parameters of tea plants. The mean values ± standard deviations (n = 6). The values with the same letter are not significantly different ($p < 0.05$). CK control experiment, DT drought stress.

TMT-6plex var'. The false discovery rate of protein identification and peptide spectrum match identification was set to 1%.

GO annotation and enrichment analyses were performed by the UniProt-GOA database ([www.http://www.ebi.ac.uk/GOA/](http://www.ebi.ac.uk/GOA/)), and the KEGG database was adopted for the enrichment of pathways by the DAVID functional annotation tool against the background of tea plant. A PPI network analysis was performed via the STRING database (<https://string-db.org/>).

Results

Identification and quantification of proteins in tea plant. To analyse the physiological and biochemical responses to drought, we analysed the contents of soluble proteins (Cpr), MDA, TP, flavonoids, and lignin and the activity of PAL in tea plants under drought stress from 0 to 96 h (Fig. 1) (Table 1). In turn, the tea plants became wrinkled and shrivelled during drought stress, which was especially severe at 96 h (Fig. 1A). The contents of Cpr and MDA increased under DT (Fig. 1B, C). The activity of PAL in the tea plants increased in response to drought stress (Fig. 1D). Accordingly, the contents of polyphenols, flavonoids and lignin increased under DT (Fig. 1E–G). These results showed that drought stress caused damage to tea plants and induced the synthesis of flavonoids and lignin.

To analyse the changes in protein accumulation in tea plants under drought stress, we performed an analysis of the global proteome using TMT isobaric labelling technology. In total, 51,213 unique spectra were generated, which matched to 23,854 unique peptides (Supplementary Table S1). Correspondingly, 4789 proteins were identified, and 4242 proteins were quantified (Supplementary Table S2), of which 11 proteins were upregulated and 100 proteins downregulated, with high repeatability ($FC > 1.5$ and $p < 0.05$).

Functional classification and enrichment of differentially expressed proteins (DEPs). To classify the functions of proteins in tea plant under drought stress, we performed GO functional classifications of




Figure 2. Functional classification and enrichment of DEPs. (A) Biological processes, (B) molecular functions, (C) cellular components, (D) subcellular localization, (E) GO enrichment and (F) KEGG enrichment.

DEPs based on their associated biological processes, molecular functions and cellular components (Fig. 2A–C). The results from bioinformatic predictions showed that the functions of the DEPs were mainly involved in metabolic processes involving catalytic activity and the cell. However, except for the metabolic processes and catalytic activity, the GO functional classifications of DEPs were mainly enriched in cellular processes involving binding and organelles.

WoLF PSORT software was used to predict and classify the subcellular structure of the differentially expressed proteins. Of the 111 DEPs, 39 were distributed in chloroplasts, 33 were distributed in the nucleus, 26 were in the cytoplasm, and the other proteins were distributed in the plasma membrane and vacuolar membrane. The subcellular localization for the global proteome was also calculated for comparison (Fig. 2D). According to the data, the subcellular localization of proteins in tea plants under drought stress and the global proteome showed no significant difference.

To obtain information on the functional enrichment of proteins in tea plants under drought stress, we conducted GO enrichment of DEPs based on clustering analysis (Fig. 2E; Supplementary Table S3). In the category of molecular function, the DEPs were mainly associated with 3-dehydroquininate dehydratase activity, 3-deoxy-7-phosphoheptulonate synthase activity and shikimate 3-dehydrogenase (NADP+) activity. In terms of biological processes, a large portion of the proteins mainly participated in the L-phenylalanine metabolic process, erythrose 4-phosphate/phosphoenolpyruvate family amino acid metabolic process and aromatic amino acid family metabolic process. The analysis of cellular components showed that the DEPs were mainly involved in intracellular nonmembrane-bound organelles and nonmembrane-bound organelles.

KEGG pathway analysis showed that all the identified proteins were downregulated. These proteins were strongly associated with secondary metabolite biosynthesis, such as flavonoid biosynthesis, phenylpropanoid biosynthesis and phenylalanine metabolism (Fig. 2F; Supplementary Table S4). The pathways were mainly related to the biosynthesis of polyphenols and lignin.

Protein–protein interaction networks. PPIs reflect the process by which two or more protein molecules form protein complexes through noncovalent bonds. To predict interactions between proteins, we generated PPI networks for all DEPs against the STRING database (Fig. 3; Supplementary Table S5). We extracted several interactive clusters from the entire interaction network by means of the MCODE plug-in tool kit. Among these proteins, 42 mapped to the PPI networks, and they were mainly clustered into 6 subnetworks. These subnetworks clearly show that there are different degrees of interaction between proteins. It can be predicted that DEPs, which are related to catechin biosynthesis, interact more strongly in many major metabolic processes of tea plants in response to drought stress. We will further study the specific interaction relationship in future work.




Figure 3. PPI networks of DEPs. The red circle represents clusters 1–4.

Protein accession	DT/CK ratio	DT/CK <i>p</i> value	DT/CK ratio
C4H	0.52	0.00031045	0.58
CCR	0.48	0.00112761	0.56
F3H	0.35	0.00033684	0.41
FLS	0.33	0.00017023	0.44
DFR	0.32	0.00001117	0.38
ANR	0.33	0.00095506	0.42
LDOX	0.28	0.00006974	0.42
GSTU	0.41	0.00206034	0.48
EMB	0.44	0.00060575	0.56
XTH	0.53	0.00004320	0.57
GWD	0.53	0.00017301	0.65

Table 2. PRM analysis of eleven candidate proteins in tea plants under drought stress.

Validation of DEPs by PRM. To verify the results of the TMT analysis at the protein level, PRM was performed among key enzymes that presented significant differences in tea plant (Table 2). In this study, 11 candidate proteins with potential drought tolerance-related functions were chosen; for each protein, two or more unique peptides with anticipated chemical stability were selected. C4H, cinnamoyl-CoA reductase (CCR), flavanone 3-hydroxylase (F3H), flavonol synthase (FLS), DFR, ANR, LDOX, GSTU, EMB, XTH, and GWD were downregulated to certain degrees, similar to the TMT results. Thus, our PRM assay revealed that the TMT results were credible, allowing further analysis.




Figure 4. General biosynthesis pathway of lignin in plants. PAL, phenylalanine ammonia lyase; C4H, trans-cinnamate 4-monooxygenase; 4CL, 4-coumarate-CoA ligase; CCR, cinnamoyl-CoA reductase; POD, peroxidase. The orange colour indicates a downregulated protein.

Discussion

Drought stimuli induce the expression of a series of genes and the synthesis of proteins related to the drought response; this occurs via the cell perception, signalling and transport ability of plants, after which their physiological and biochemical metabolism is optimized to cope with the drought stress¹⁹. In this study, the accumulation of enzymes related to the biosynthesis of lignin, flavonoids and fatty acids was inhibited by drought stress, which provides a basis for further elucidating the drought resistance mechanism of tea plants.

The accumulation of proteins related to lignin biosynthesis was inhibited by drought stress. As one of the important components of the plant cell wall, lignin is of great significance for plant growth and environmental adaptability. To determine the accumulation levels of proteins related to the lignin pathway, we evaluated the response of the phenylpropanoid pathway to drought stress (Fig. 4). The enzymatic steps leading to the synthesis of lignin monomers have been extensively reported^{20–22}.

As the first step of the phenylpropanoid pathway, PAL mainly breaks down phenylalanine into cinnamic acid²³. A previous study showed that the activity of PAL was significantly correlated with the content of lignin²⁴. Additionally, inhibition of PAL activity can lead to a decrease in lignin biosynthesis^{25–27}. In addition, a strong positive correlation between the expression of both *CsPAL* and *CsC4H* and catechin content in tea plants has been shown²⁸. However, there is little research about the correlation of PAL accumulation with lignin content in tea plants under drought stress. In our study, the accumulation of PAL was downregulated, while the activity of PAL increased. The content of lignin also increased. Therefore, we speculated that the increase in lignin content under drought was not only positively correlated with the activity of PAL but also negatively correlated with the accumulation of PAL. In our previous study, we found that PAL in tea plants was ubiquitinated in response to drought stress, which was related to the L-phenylalanine metabolic process¹⁸. Whether ubiquitinated PAL or other factors are involved in the synthesis of lignin need further study.

As the first specific step in the synthesis of lignin monomers, CCR catalyses the conversion of cinnamoyl-CoA esters to their corresponding cinnamaldehydes²⁹. A previous study showed that when CCR activity is severely inhibited, the lignin content decreases accordingly²⁷. The present study showed that the lignin content increased during the development of leaves in both tea plant cultivars Fudingdabai and Suchazao, while the expression level of *CsCCR* was not uniform³⁰. Recent research has shown that the accumulation of CCR increases in the roots and decreases in the leaves of tea plants under high concentrations of Al stress, while the lignin content decreases in both organs³¹. In our study, the accumulation of CCR was downregulated in tea plants under drought stress, while the content of lignin increased. These data indicate that drought stress may promote the accumulation of lignin by inhibiting the accumulation of CCR in tea plants.

As a specific step in the late stages of lignin synthesis, the polymerization of monolignols catalysed by peroxidase (POD) and/or laccase enzymes yields lignin³². Previous studies have shown that abiotic stresses generally




Figure 5. General biosynthesis pathway of flavonoids in tea plants. PAL, phenylalanine ammonia lyase; C4H, trans-cinnamate 4-monooxygenase; 4CL, 4-coumarate-CoA ligase; CHS, chalcone synthase; CHI, chalcone-flavanone isomerase; F3H, flavanone 3-hydroxylase; FLS, flavonol synthase; ANR, anthocyanidin reductase; LAR, leucoanthocyanidin reductase; LDOX, leucoanthocyanidin dioxygenase; F3'H, flavonoid 3' hydroxylase; F3'5'H, flavonoid 3',5'-hydroxylase; DFR, dihydroflavonol-4-reductase. The orange colour indicates a downregulated protein.

induce POD activity until it reaches a specific threshold, leading to lignin accumulation^{33,34}. A previous study also showed that overexpression of *POD* increases the contents of phenols and lignin in plants³⁵. An above-mentioned study showed that the accumulation of POD increased in the roots but decreased in the leaves of tea plants under high concentrations of Al stress, while the lignin content decreased in both organs³¹. In our study, the accumulation of POD was downregulated in tea plants under drought stress, while the content of lignin increased. We speculate that drought stress may promote the accumulation of lignin by inhibiting the accumulation of POD in tea plants.

The accumulation of proteins related to flavonoid biosynthesis was inhibited by drought stress. Flavonoids are important components of tea products and are closely related to the taste, flavour, and health benefits of tea. Recent studies have revealed that plants can improve their drought, salt and other abiotic resistance by increasing the accumulation of flavonoids^{36–39}. In our study, to determine the influence of proteins related to catechin biosynthesis under drought stress, we mapped the pathway of proteins involved in the phenylpropanoid and flavonoid pathways (Fig. 5). The enzymatic steps leading to the synthesis of these flavonoid monomers have been extensively reviewed⁴⁰.

As the first committed step of the flavonoid pathway, chalcone synthase (CHS) utilizes a tetraketide intermediate by condensation with one 4-coumaroyl-CoA and three malonyl-CoA compounds to form naringenin chalcone^{41,42}. A previous study has showed that the activities of C4H, a key enzyme that functions upstream of baicalin biosynthesis, are consistent with the accumulation of baicalin⁴³. In addition, the expression of multi-copy flavonoid pathway genes (*Chs1*, *Chs2*, and *Chs3*) coincides with anthocyanin, flavonol and flavan-3-ol accumulation in grapevine⁴⁴. In the present study, the expression level of *CHS* was not related to the concentration of catechin in tea leaves but was related to the concentration of O-glycosylated flavonols⁴⁵. The expression levels of unigenes such as *CHS* tended to first decrease but then increase in tea plants in response to drought stress, while the total flavonoid content was upregulated⁷. However, relatively little is known about the regulation of CHS at the protein level of tea plants under drought stress. In this study, it is noteworthy that the accumulation of CHS under drought stress was lower than that in the controls, while the content of flavonoids increased. Taken together, these results demonstrate that drought stress may promote the accumulation of flavonoids by inhibiting the accumulation of CHS in tea plants.

F3H is an abundant enzyme in tea plants that catalyses the stereo-specific hydroxylation of (2S)-naringenin to form (2R,3R)-dihydrokaempferol⁴⁶. A previous study showed that drought stress induced an increase in the expression of the *RsF3H* gene, enzyme activity and the contents of total flavonoid⁴⁷. A recent study showed that

the expression level of *F3H* was not related to the concentration of catechin in tea leaves but was related to the concentration of *Pas*⁴⁵. In tea plants, the expression of *CsF3H* is downregulated in response to drought, abscisic acid and gibberellic acid treatment but upregulated in response to wounding, and the concentration of catechins paralleled the expression data⁴⁶. Additionally, negative correlations between *F3H* and *anthocyanidin synthase* expression levels and catechin content were identified in spring tea plants, whereas the correlations were positive in autumn tea plants⁴⁸. However, previous studies have primarily focused on the cloning, expression and regulation of the gene at the transcriptional level, and little is known *F3H* accumulation at the protein level. In this study, the accumulation of *F3H* was downregulated, and the contents of flavonoids increased in response to drought stress, indicating that drought stress may promote the accumulation of flavonoids by inhibiting the accumulation of *F3H* in tea plants.

FLS catalyses dihydroflavonols into flavonols, and *FLS*, as the key enzyme involved in flavonol synthesis, determines the final contents of flavonols in plants⁴⁹. A previous study suggested that the activity of *FLS* was highly positively correlated with the total contents of flavonols during grape berry development, and *FLS* accumulation was essentially consistent with *FLS* activity at most stages⁴⁹. Another study showed that the unigenes *FLS* and *flavone synthase* were continuously upregulated in tea plants under drought stress, which may be the reason for the increase in total flavonoid contents in response to drought stress⁷. However, in tea plants, there is little information about the regulation of *FLS* at the protein level in response to drought stress. In our study, the accumulation of *FLS* was downregulated in the tea plants, while the content of lignin increased, indicating that drought stress could affect the accumulation of flavonoids by inducing *FLS* accumulation.

Response of LCFAs to drought stress. Fatty acids are the main components of cell membrane lipids and the precursor of some signalling molecules in plants. They are widely distributed on the cell surface to prevent moisture and heat loss and are closely related to cell recognition specificity and tissue immunity. However, relatively little is known about long-chain fatty acid amides: most of the well-defined physiological functions, the metabolism underlying their biosynthesis, their degradation, and their cellular transport remain elusive^{50–52}. In this study, we focused on the accumulation of essential enzymes related to the biosynthesis of lignin, flavonoids and fatty acids in tea plants in response to drought stress at the protein level.

The first step of fatty acid biosynthesis catalysed by plastid ACCase activity is the main determinant of the overall rate of fatty acid synthesis⁵³. Previous studies have shown that acetyl-CoA carboxylase 1 (*ACC1*) is essential for very long-chain fatty acid elongation and embryo development in *Arabidopsis*⁵⁴. *P. simonii* *Populus simonii* *ACC1*, which is highly similar to the *Arabidopsis acc1* gene, increased in response to drought stress, which suggests that the biosynthesis of LCFAs may play an important role in the drought resistance of *Populus simonii*⁵⁵. However, in tea plants, there is no information about the regulation of *ACC1* in response to drought stress. In our study, the accumulation of *ACC1* was downregulated, which indicated that tea plants could improve their drought resistance by inhibiting the accumulation of *ACC1*.

3-Ketoacyl-CoA synthase (*KCS*) catalyses a condensation reaction to form 3-ketoacyl-CoA during very long-chain fatty acid synthesis⁵². A previous study showed that the phenotypic changes of *kcs1-1* mutants included relatively thin stems and reduced resistance to low-humidity stress at a young age⁵⁶. Moreover, a recent study showed that the transcription of most lipid biosynthesis genes, such as *KCS6*, increases during the rapid elongation stage and is maintained at a high level; moreover, the fatty acid biosynthetic pathway is also significantly induced in AKR2A-57 fibres⁵⁷. In the present study, *KCS-6* accumulation was downregulated, which indicated that tea plants could improve drought resistance by inhibiting the accumulation of *KCS-6*.

A subsequent reaction involves the reduction to 3-hydroxyacyl-CoA catalysed by 3-ketoacyl-CoA reductase (*KCR*)⁵⁸. Previous studies have shown that *GhKCRs* are involved in endoplasmic reticulum-associated very long-chain fatty acid elongation during cotton fibre development⁵⁹. At present, there is no evidence on the response of proteins related to the LCFA synthesis pathway in response to drought stress in tea plants. Our data indicated that the accumulation of *ACC1*, *KCS-6*, and *KCR* in the LCFA synthesis pathway was downregulated, which indicated that tea plants could improve drought resistance by inhibiting the accumulation of essential enzymes related to the biosynthesis of lignin, flavonoids and fatty acids. With global climate change and increasing water scarcity, in-depth studies on the relationships between long-chain fatty acids and key enzymes involved in the biosynthesis pathway are of great significance to improve the drought resistance of tea plants, which needs further research.

Data availability

The datasets generated during the current study are available via ProteomeXchange under identifier PXD011688, (<https://www.ebi.ac.uk/pride>).

Received: 28 January 2020; Accepted: 31 August 2020

Published online: 23 September 2020

References

1. Cheruiyot, E. K., Mumera, L. M., Ng'etich, W. K., Hassanali, A. & Wachira, F. N. High fertilizer rates increase susceptibility of tea to water stress. *J. Plant Nutr.* **33**, 115–129. <https://doi.org/10.1080/01904160903392659> (2009).
2. Upadhyaya, H., Panda, S. K. & Dutta, B. K. Variation of physiological and antioxidative responses in tea cultivars subjected to elevated water stress followed by rehydration recovery. *Acta Physiol. Plant.* **30**, 457–468. <https://doi.org/10.1007/s11738-008-0143-9> (2008).
3. Scott, E. R. *et al.* Interactive effects of drought severity and simulated herbivory on tea (*Camellia sinensis*) volatile and non-volatile metabolites. *Environ. Exp. Bot.* **157**, 283–292. <https://doi.org/10.1016/j.envexpbot.2018.10.025> (2019).
4. Nyarukowa, C., Koech, R., Loots, T. & Apostolides, Z. SWAPDT: a method for short-time withering assessment of probability for drought tolerance in *Camellia sinensis* validated by targeted metabolomics. *J. Plant Physiol.* **198**, 39–48. <https://doi.org/10.1016/j.jplph.2016.04.004> (2016).

5. Liu, S. C. *et al.* Physiological changes and differential gene expression of tea plant under dehydration and rehydration conditions. *Sci. Hortic.* **184**, 129–141. <https://doi.org/10.1016/j.scienta.2014.12.036> (2015).
6. Lin, S. K. *et al.* Time-course of photosynthesis and non-structural carbon compounds in the leaves of tea plants (*Camellia sinensis* L.) in response to deficit irrigation. *Agric. Water Manag.* **144**, 98–106. <https://doi.org/10.1016/j.agwat.2014.06.005> (2014).
7. Wang, W. *et al.* Transcriptomic analysis reveals the molecular mechanisms of drought-stress-induced decreases in *Camellia sinensis* leaf quality. *Front. Plant Sci.* **7**, 385–385. <https://doi.org/10.3389/fpls.2016.00385> (2016).
8. Chao, Z., Yu, W., Ding, Z. & Lei, Z. Global transcriptional analysis reveals the complex relationship between tea quality, leaf senescence and the responses to cold-drought combined stress in *Camellia sinensis*. *Front. Plant Sci.* **7**, 1858. <https://doi.org/10.3389/fpls.2016.01858> (2016).
9. Ahammed, G. J., Li, X., Wan, H., Zhou, G. & Cheng, Y. SlWRKY81 reduces drought tolerance by attenuating proline biosynthesis in tomato. *Sci. Hortic.* **270**, 109444. <https://doi.org/10.1016/j.scienta.2020.109444> (2020).
10. Ahammed, G. J. *et al.* Tomato WRKY81 acts as a negative regulator for drought tolerance by modulating guard cell H₂O₂-mediated stomatal closure. *Environ. Exp. Bot.* **171**, 103960. <https://doi.org/10.1016/j.envexpbot.2019.103960> (2020).
11. Li, H. *et al.* Alkanes (C29 and C31)-mediated intracellular wax accumulation contributes to melatonin- and ABA-induced drought tolerance in watermelon. *J. Plant Growth Regul.* <https://doi.org/10.1007/s00344-020-10099-z> (2020).
12. Hernandez, L., Alegre, L. & Munne-Bosch, S. Enhanced oxidation of flavan-3-ols and proanthocyanidin accumulation in water-stressed tea plants. *Phytochemistry* **67**, 1120–1126. <https://doi.org/10.1016/j.phytochem.2006.04.002> (2006).
13. Hu, Y., Li, W. C., Xu, Y. Q., Li, G. J. & Fu, F. L. Differential expression of candidate genes for lignin biosynthesis under drought stress in maize leaves. *J. Appl. Genet.* **50**, 213–223. <https://doi.org/10.1007/BF03195675> (2009).
14. Kuiper, P. J. C. & Stuijver, B. Cyclopropane fatty acids in relation to earliness in spring and drought tolerance in plants. *Plant Physiol.* **49**, 307–309. <https://doi.org/10.1104/pp.49.3.307> (1972).
15. Wang, Y. *et al.* Proteomic analysis of *Camellia sinensis* (L.) reveals a synergistic network in the response to drought stress and recovery. *J. Plant Physiol.* **219**, 91–99. <https://doi.org/10.1016/j.jplph.2017.10.001> (2017).
16. Chen, Q., Yang, L., Ahmad, P., Wan, X. & Hu, X. Proteomic profiling and redox status alteration of recalcitrant tea (*Camellia sinensis*) seed in response to desiccation. *Planta* **233**, 583–592. <https://doi.org/10.1007/s00425-010-1322-7> (2011).
17. Sun, J. *et al.* Ammonium triggered the response mechanism of lysine crotonylome in tea plants. *BMC Genom.* **20**, 340. <https://doi.org/10.1186/s12864-019-5716-z> (2019).
18. Xie, H. *et al.* Global ubiquitome profiling revealed the roles of ubiquitinated proteins in metabolic pathways of tea leaves in responding to drought stress. *Sci. Rep.* **9**, 4286. <https://doi.org/10.1038/s41598-019-41041-3> (2019).
19. Umezawa, T. *et al.* Molecular basis of the core regulatory network in ABA responses: sensing, signaling and transport. *Plant Cell Physiol.* **51**, 1821–1839. <https://doi.org/10.1093/pcp/pcq156> (2010).
20. Bonawitz, N. D. & Chapple, C. The genetics of lignin biosynthesis: connecting genotype to phenotype. *Annu. Rev. Genet.* **44**, 337–363. <https://doi.org/10.1146/annurev-genet-102209-163508> (2010).
21. Boerjan, W., Ralph, J. & Baucher, M. Lignin biosynthesis. *Annu. Rev. Plant Biol.* **54**, 519–546. <https://doi.org/10.1146/annurev.arplant.54.031902.134938> (2003).
22. Vanholme, R., Demedts, B., Morreel, K., Ralph, J. & Boerjan, W. Lignin biosynthesis and structure. *Plant Physiol.* **153**, 895–905. <https://doi.org/10.1104/pp.110.155119> (2010).
23. Toscano, S., Ferrante, A., Leonardi, C. & Romano, D. PAL activities in asparagus spears during storage after ammonium sulfate treatments. *Postharvest Biol. Technol.* **140**, 34–41. <https://doi.org/10.1016/j.postharvbio.2018.02.010> (2018).
24. Terzi, R., Saruhan Güler, N., Kutlu Çalişkan, N. & Kadioğlu, A. Lignification response for rolled leaves of *Ctenanthe setosa* under long-term drought stress. *Turk. J. Biol.* **37**, 614–619. <https://doi.org/10.3906/biy-1210-27> (2013).
25. Chen, M. & McClure, J. W. Altered lignin composition in phenylalanine ammonia-lyase-inhibited radish seedlings: implications for seed-derived sinapoyl esters as lignin precursors. *Phytochemistry* **53**, 365–370. [https://doi.org/10.1016/S0031-9422\(99\)00531-2](https://doi.org/10.1016/S0031-9422(99)00531-2) (2000).
26. Kawaoka, A., Kaohien, P., Yoshida, K., Endo, S. & Ebinuma, H. Functional analysis of tobacco LIM protein Ntlm1 involved in lignin biosynthesis. *Plant J.* **22**, 289–301. <https://doi.org/10.1046/j.1365-313x.2000.00737.x> (2000).
27. Anterola, A. M. & Lewis, N. G. Trends in lignin modification: a comprehensive analysis of the effects of genetic manipulations/mutations on lignification and vascular integrity. *Phytochemistry* **61**, 221–294. <https://doi.org/10.1002/chin.200303272> (2002).
28. Singh, K., Kumar, S., Rani, A., Gulati, A. & Ahuja, P. S. Phenylalanine ammonia-lyase (PAL) and cinnamate 4-hydroxylase (C4H) and catechins (flavan-3-ols) accumulation in tea. *Funct. Integr. Genom.* **9**, 125–134. <https://doi.org/10.1007/s10142-008-0092-9> (2009).
29. Lacombe, E., Hawkins, S., Doorselaere, J. V., Piquemal, J. & Grima-Pettenati, J. Cinnamoyl CoA reductase, the first committed enzyme of the lignin branch biosynthetic pathway: cloning, expression and phylogenetic relationships. *Plant J.* **11**, 429–441. <https://doi.org/10.1046/j.1365-313x.1997.11030429.x> (1997).
30. Wang, Y. *et al.* Identification of genes revealed differential expression profiles and lignin accumulation during leaf and stem development in tea plant (*Camellia sinensis* (L.) O. Kuntze). *Protoplasma* **256**, 359–370. <https://doi.org/10.1007/s00709-018-1299-9> (2019).
31. Xu, Q. *et al.* Aluminum induced physiological and proteomic responses in tea (*Camellia sinensis*) roots and leaves. *Plant Physiol. Biochem.* **115**, 141–151. <https://doi.org/10.1016/j.plaphy.2017.03.017> (2017).
32. Govender, N. T., Mahmood, M., Seman, I. A. & Wong, M. Y. The phenylpropanoid pathway and lignin in defense against ganoderma boninense colonized root tissues in oil palm (*Elaeis guineensis* Jacq.). *Front. Plant Sci.* **8**, 1395–1395. <https://doi.org/10.3389/fpls.2017.01395> (2017).
33. Liu, W. *et al.* Analysis of CmCADs and three lignifying enzymes in oriental melon ('CaiHong7') seedlings in response to three abiotic stresses. *Sci. Hortic.* **237**, 257–268. <https://doi.org/10.1016/j.scienta.2018.04.024> (2018).
34. Lee, B. R. *et al.* Peroxidases and lignification in relation to the intensity of water-deficit stress in white clover (*Trifolium repens* L.). *J. Exp. Bot.* **58**, 1271–1279. <https://doi.org/10.1093/jxb/erl280> (2007).
35. Kim, Y. H. *et al.* Overexpression of sweetpotato swpa4 peroxidase results in increased hydrogen peroxide production and enhances stress tolerance in tobacco. *Planta* **227**, 867–881. <https://doi.org/10.1007/s00425-007-0663-3> (2008).
36. Shojaie, B., Mostajeran, A. & Ghanadian, M. Flavonoid dynamic responses to different drought conditions: Amount, type, and localization of flavonols in roots and shoots of *Arabidopsis thaliana* L. *Turk. J. Biol.* **40**, 612–622. <https://doi.org/10.3906/biy-1505-2> (2016).
37. Li, X. *et al.* Salicylic acid acts upstream of nitric oxide in elevated carbon dioxide-induced flavonoid biosynthesis in tea plant (*Camellia sinensis* L.). *Environ. Exp. Bot.* **161**, 367–374. <https://doi.org/10.1016/j.envexpbot.2018.11.012> (2019).
38. Li, X. *et al.* RBOH1-dependent apoplastic H₂O₂ mediates epigallocatechin-3-gallate-induced abiotic stress tolerance in *Solanum lycopersicum* L. *Environ. Exp. Bot.* **161**, 357–366. <https://doi.org/10.1016/j.envexpbot.2018.11.013> (2019).
39. Ahammed, G. J., Li, Y., Li, X., Han, W.-Y. & Chen, S. Epigallocatechin-3-gallate alleviates salinity-retarded seed germination and oxidative stress in tomato. *J. Plant Growth Regul.* **37**, 1349–1356. <https://doi.org/10.1007/s00344-018-9849-0> (2018).
40. Ferreyra, M. L. F., Rius, S. P. & Casati, P. Flavonoids: biosynthesis, biological functions, and biotechnological applications. *Front. Plant Sci.* **3**, 222–222. <https://doi.org/10.3389/fpls.2012.00222> (2012).
41. Pang, Y. *et al.* Characterization and expression of chalcone synthase gene from Ginkgo biloba. *Plant Sci.* **168**, 1525–1531. <https://doi.org/10.1016/j.plantsci.2005.02.003> (2005).

42. Yu, C. K. Y. *et al.* A stilbene synthase gene (SbSTS1) is involved in host and nonhost defense responses in sorghum. *Plant Physiol.* **138**, 393–401. <https://doi.org/10.2307/4629834> (2005).
43. Cheng, L. *et al.* Changes in the physiological characteristics and baicalin biosynthesis metabolism of *Scutellaria baicalensis* Georgi under drought stress. *Ind. Crops Prod.* **122**, 473–482. <https://doi.org/10.1016/j.indcrop.2018.06.030> (2018).
44. Jeong, S., Goto-Yamamoto, N., Hashizume, K. & Esaka, M. Expression of multi-copy flavonoid pathway genes coincides with anthocyanin, flavonol and flavan-3-ol accumulation of grapevine. *Vitis* **47**, 135–140 (2008).
45. Wang, Y. *et al.* Influence of shade on flavonoid biosynthesis in tea (*Camellia sinensis* (L.) O Kuntze). *Sci. Hortic.* **141**, 7–16. <https://doi.org/10.1016/j.scienta.2012.04.013> (2012).
46. Singh, K. *et al.* An early gene of the flavonoid pathway, flavanone 3-hydroxylase, exhibits a positive relationship with the concentration of catechins in tea (*Camellia sinensis*). *Tree Physiol.* **28**, 1349–1356. <https://doi.org/10.1093/treephys/28.9.1349> (2008).
47. Liu, M., Li, X., Liu, Y. & Cao, B. Regulation of flavanone 3-hydroxylase gene involved in the flavonoid biosynthesis pathway in response to UV-B radiation and drought stress in the desert plant, *Reaumuria soongorica*. *Plant Physiol. Biochem.* **73**, 161–167. <https://doi.org/10.1016/j.plaphy.2013.09.016> (2013).
48. Liu, M. *et al.* Relationship between gene expression and the accumulation of catechin during spring and autumn in tea plants (*Camellia sinensis* L.). *Hortic. Res.* **2**, 15011–15011. <https://doi.org/10.1038/hortres.2015.11> (2015).
49. Fang, F., Tang, K. & Huang, W.-D. Changes of flavonol synthase and flavonol contents during grape berry development. *Eur. Food Res. Technol.* **237**, 529–540. <https://doi.org/10.1007/s00217-013-2020-z> (2013).
50. Witkamp, R. F. Fatty acids, endocannabinoids and inflammation. *Eur. J. Pharmacol.* **785**, 96–107. <https://doi.org/10.1016/j.ejphar.2015.08.051> (2016).
51. Liu, J. J., Zhang, C. & Lu, W. Biosynthesis of long-chain ω -hydroxy fatty acids by engineered *Saccharomyces cerevisiae*. *J. Agric. Food Chem.* **67**, 4545–4552. <https://doi.org/10.1021/acs.jafc.9b00109> (2019).
52. Wang, Q. *et al.* Functional identification of ELO-like genes involved in very long chain fatty acid synthesis in *Arabidopsis thaliana*. *Russ. J. Plant Physiol.* **61**, 853–861. <https://doi.org/10.1134/S1021443714060193> (2014).
53. Schulte, W., Topfer, R., Stracke, R., Schell, J. & Martini, N. Multi-functional acetyl-CoA carboxylase from *Brassica napus* is encoded by a multi-gene family: Indication for plastidic localization of at least one isoform. *Proc. Natl. Acad. Sci. U.S.A.* **94**, 3465–3470. <https://doi.org/10.1073/pnas.94.7.3465> (1997).
54. Baud, S., Guyon, V., Kronenberger, J., Wuillème, S. & Rochat, C. Multifunctional acetyl-CoA carboxylase 1 is essential for very long chain fatty acid elongation and embryo development in *Arabidopsis*. *Plant J.* **33**, 75–86. <https://doi.org/10.1046/j.1365-313X.2003.016010.x> (2003).
55. Chen, J., Song, Y., Zhang, H. & Zhang, D. Genome-wide analysis of gene expression in response to drought stress in *populus simonii*. *Plant Mol. Biol. Rep.* **31**, 946–962. <https://doi.org/10.1007/s11105-013-0563-6> (2013).
56. Todd, J., Post-Beittenmiller, D. & Jaworski, J. G. KCS1 encodes a fatty acid elongase 3-ketoacyl-CoA synthase affecting wax biosynthesis in *Arabidopsis thaliana*. *Plant J.* **17**, 119–130. <https://doi.org/10.1046/j.1365-313X.1999.00352.x> (1999).
57. Hu, W. *et al.* AKR2A participates in the regulation of cotton fibre development by modulating biosynthesis of very-long-chain fatty acids. *Plant Biotechnol. J.* **18**, 526–539. <https://doi.org/10.1111/pbi.13221> (2020).
58. Cinti, D. L., Cook, L., Nagi, M. N. & Suneja, S. K. The fatty acid chain elongation system of mammalian endoplasmic reticulum. *Prog. Lipid Res.* **31**, 1–51. [https://doi.org/10.1016/0163-7827\(92\)90014-A](https://doi.org/10.1016/0163-7827(92)90014-A) (1992).
59. Qin, Y. M. *et al.* Cloning and functional characterization of two cDNAs encoding NADPH-dependent 3-ketoacyl-CoA reductase from developing cotton fibers. *Cell Res.* **15**, 465–473. <https://doi.org/10.1038/sj.cr.7290315> (2005).

Acknowledgements

This research was subsidized by the following: 1. the Special Foundation for Distinguished Taishan Scholar of Shandong Province (No. ts201712057), 2. The Significant Application Projects of Agriculture Technology Innovation in Shandong Province (SD2019ZZ010), 3. The Technology System of Modern Agricultural Industry in Shandong Province (SDAIT-19-01) and the China Earmarked Fund for Modern Agro-industry Technology Research System (CARS-19), 4. The Livelihood Project of Qingdao City (19-6-1-64-nsh), 5. The Project of Agricultural Science and Technology Fund in Shandong Province (2019LY002, 2019YQ010), and 6. The Key Research and Development Project of Shandong Province (2019LYXZ009).

Author contributions

Z.D., H.G., Y.W., and H.X. designed the study, and H.G. wrote the initial draft of the manuscript. Z.D., Y.W., H.X., C.Q., and S.Z. contributed to the analysis and interpretation of the data and assisted in the preparation of the manuscript. All the other authors contributed to the data collection and interpretation and critically reviewed the manuscript. All the authors approved the final version of the manuscript and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41598-020-72596-1>.

Correspondence and requests for materials should be addressed to Z.D.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020