

<https://helda.helsinki.fi>

Characterization of *Leuconostoc gasicomitatum* sp. nov.,
Associated with Spoiled Raw Tomato-Marinaded Broiler Meat
Strips Packaged under Modified-Atmosphere Conditions

Björkroth, Johanna

American Society for Microbiology (ASM)

2000

A p p l i e d a n d E n v i r o n m e n t a l M i c r o b i o l o g y . 2 0

<http://hdl.handle.net/1975/525>

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Characterization of *Leuconostoc gasicomitatum* sp. nov., Associated with Spoiled Raw Tomato-Marinaded Broiler Meat Strips Packaged under Modified-Atmosphere Conditions

K. JOHANNA BJORKROTH,^{1*} ROLF GEISEN,² ULRICH SCHILLINGER,² NORBERT WEISS,³ PAUL DE VOS,⁴ WILHELM H. HOLZAPFEL,² HANNU J. KORKEALA,¹ AND PETER VANDAMME⁴

Department of Food and Environmental Hygiene, University of Helsinki,¹ Helsinki, Finland, and Toxicology, Federal Research Centre for Nutrition and Consumer Protection,² Karlsruhe, Germany; Collection of Microorganisms and Cell Cultures, Braunschweig,³ Germany; and Laboratory of Microbiology, University of Ghent,⁴ Ghent, Belgium

Received 15 February 2000/Accepted 7 June 2000

Lactic acid bacteria (LAB) associated with gaseous spoilage of modified-atmosphere-packaged, raw, marinated broiler meat strips were identified on the basis of a restriction fragment length polymorphism (RFLP) (ribotyping) database containing DNAs coding for 16S and 23S rRNAs (rDNAs). A mixed LAB population dominated by *Leuconostoc* species resembling *Leuconostoc gelidum* caused the spoilage of the product. *Lactobacillus*, *Lactobacillus curvatus* gram-positive rod phenotypically similar to heterofermentative *Lactobacillus* species were the other main organisms detected. An increase in pH together with extreme bulging of packages suggested a rare LAB spoilage type called "protein swell." This spoilage characterized by excessive production of gas due to amino acid decarboxylation, and the rise in pH is associated with the subsequent deamination of amino acids. Protein swell has not previously been associated with meat product. A polyphasic approach, including classical phenotyping, whole-cell protein electrophoresis, and 23S rDNA RFLP, 16S rDNA sequence analysis, and DNA-DNA reassociation analysis, was used for the identification of the dominant *Leuconostoc* species. In addition to the RFLP analysis, phenotyping, whole-cell protein analysis, and 16S rDNA sequence homology indicated that the most similar to the spoilage-associated species. The two spoilage strains studied possessed 98.8 and 99.0% 16S rDNA sequence homology with the *gelidum* type strain. DNA-DNA reassociation, however, clearly distinguished the two species. The same strains showed only 22 and 34% hybridization with the *gelidum* strain. These results warrant a separate species status, and we propose the name *Leuconostoc gasicomitatum* sp. nov. for this spoilage-associated *Leuconostoc* species.

Lactic acid bacteria (LAB) are the dominant spoilage organisms, which have also been set as the retail shelf life. In vacuum or modified-atmosphere (MA)-packaged meat products (1, 2, 8, 10, 23, 31). Spoilage is mainly caused by bulging due to gas formation 5 days after packaging (3, 4, 26, 28; W. H. Holzappel and E. S. Gerber, Abstr. 32nd Eur. Meet. Meat Res. Workers, p. 26, 1986) that time, only this tomato-marinaded product was spoiled by *Leuconostoc* (7, 14, 28, 43, 49) species. The activities of these organisms at stationary phase produce the compounds associated with sensory spoilage (22). Depending on the product, this quality deterioration usually starts after packaging, and it is manifested mainly as fermentation and sour or cheesy off odors and/or off tastes. Provided the shelf life of the product has been estimated correctly, changes do not occur before the sell-by day. However, in poultry products, our study set out to characterize of potent spoilage LAB and/or poor production and to identify these spoilage LAB to the species level. Severe quality faults (5, 7, 26, 27) have occurred in product recalls.

The consumption of marinated, ready-to-cook, raw meat products has been increasing in Europe. As easy-to-eat and low-fat food, they are favored by many consumers. In this study, we describe and characterize an unusual spoilage type of MA-packaged, tomato-marinaded, raw broiler meat strips. The product was manufactured at a modern large-scale processing plant, and normally, good quality was maintained at the time of spoilage. Based on these results, the dominant species of the spoilage population was considered novel, and we propose the name *Leuconostoc gasicomitatum* sp. nov. for it.

* Corresponding author. Mailing address: Department of Food and Environmental Hygiene, Faculty of Veterinary Medicine, University of Helsinki, P.O. Box 57, FIN-00014 Helsinki University, Finland. Phone: 358-9-19149705. Fax: 358-9-19149718. E-mail: kjorkroth@helsinki.fi.
Description of the product and pH, sensory, and microbiological analysis. The product was manufactured from raw, skinned broiler meat, which was cut into strips, marinated with the marinade, and packaged under MA as ca. 500-g consumer packages. The marinade contained plant oil, tomatoes, paprika, cayenne,

TABLE 1. *Leuconostoc sensu stricto* reference strains

Species	Strain
<i>L. carnosum</i>	LMG 11498 ^{Ta}
<i>L. citreum</i>	LMG 11417
<i>L. fallax</i>	CCUG 30061 ^{Tb}
<i>L. gelidum</i>	LMG 9850 ^T
<i>L. lactis</i>	CCUG 30064 ^T
	LMG 7940
<i>L. mesenteroides</i> ssp. <i>premoris</i>	LMG 13562
<i>L. mesenteroides</i> ssp. <i>dextranicum</i>	LMG 7954
	LMG 11318
<i>L. mesenteroides</i> ssp. <i>mesenteroides</i>	LMG 7939
<i>L. pseudomesenteroides</i>	LMG 11482 ^T
	LMG 11483
	LMG 11499

^a BCCM/LMG Belgian Coordinated Collections of Microorganisms.
^b Culture Collection of the University of Gothenburg.

salt, protein hydrosylates, starch and modified starch, natural aromatics, preservatives, and a buffering additive. The pH of the normal product varied from 4.0 to 4.5 with pH 4.5 set as the optimal target value. The expected shelf life was 30 days, with the day of manufacture regarded as day 0.

Six unopened packages showing clear bulging were analyzed on the shelf life. LAB were enumerated from serial 10-fold dilutions of the spoiled product on Rogosa selenite F agar (Oxoid, Basingstoke, United Kingdom) and Rogosa lactobacillus agar (Orion Diagnostica, Espoo, Finland) as described by Korkeala et al. (3). The plates were incubated at 25 C in an anaerobic jar and then generating kit (Oxoid) for 5 days. The pH was measured directly on homogenized samples. Evaluation of odor, color, appearance, and texture of spoiled product was performed by three trained judges, as described by Lindroth (21).

Bacterial strains and the use of strains in different phases of the study. A total of 120 spoiled packages recovered from the six spoiled packages (20 isolates from each package) were purified. During the course of the study, strains were assessed in the different phases of the study as described below. The 120 spoilage isolates were all subjected to basic phenotypic characterization and ribotyping. The ribopatterns were compared with the corresponding patterns in the LAB database of the Department of Food and Environmental Hygiene, University of Helsinki, Helsinki, Finland. These comprise patterns of the spoilage LAB in the genus *Lactobacillus*, *Leuconostoc*, *Enterococcus*, and *Weissella* (4, 6, 7, 25). Before Southern blotting, the restriction endonuclease analysis (REA) patterns of the main spoilage species were inspected visually. From these strains, those with different REA patterns, four strains representing both pattern types (two from each) were chosen for further taxonomic studies. These isolates were given the following strain numbers: LMG 18811, LMG 18812, LMG 18813, and LMG 18889. The strains presented in Table 1 were used during the more detailed taxonomic study dealing with the main spoilage species, and the LAB ribotyping database contained the ribopatterns of these strains.

All of the strains were maintained in MRS broth (Difco, Detroit, Mich.) at 27 C and cultured using MRS broth or MRS agar (Oxoid).

Phenotypic characterization. All 120 isolates were Gram stained, and tested, streaked on Rogosa selenite F agar, and studied for the production of gas from glucose (44). Further phenotypic characterization of the main spoilage species was done with strains LMG 18811, LMG 18812, LMG 18813, and LMG 18889. Production of ammonia from arginine was determined by the method of Briggs (11). Dextran formation was studied on agar containing 5% sucrose (20). Fermentation of carbohydrates was determined by the API 50 CHL Lactobacillus identification system (Biomerieux, Marcy l'Etoile, France). The ability to produce different lactic acid isomers was tested enzymatic method (48) utilizing Boehringer Mannheim GmbH (Mannheim, Federal Republic of Germany)-lactate dehydrogenases. The four strains were also tested for growth in MRS broth at 4, 10, 15, and 37 C. No growth was observed or at least for 21 days.

Enzymatic activity. The proteolytic activity of *L. gasicomitatum* nov. isolates was tested on MRS agar supplemented with sterile skim milk (2% concentration). API ZYM (Biomerieux) was also used for the characterization of the enzymatic activities of the LMG 18811 and LMG 18812 strains.

Peptidoglycan analysis. Preparation of cell walls and determination of peptidoglycan structure of LMG 18811 was carried out by the methods

by Schleifer and Kandler (41) with the modification of using thin-layer chromatography on cellulose sheets instead of paper chromatography. Briefly, freeze-dried cell walls was hydrolyzed in 100 C for 16 h (total hydrolysate) and 45 min (partial hydrolysate). Diamino acids were determined from total hydrolysate by one-dimensional chromatography in the system methanol-pyridine-water-10 N HCl (320:40:70:10 [vol/vol/vol/vol]). Diamino acids and peptides from total and partial hydrolysates were identified by two-dimensional chromatography in the systems published by Schleifer and Kandler (41), by their mobilities and staining characteristics with ninhydrin. The resulting "fingerprints" were compared with known peptidoglycan structures. Whole-cell protein analysis. Similarity of the main spoilage species (*L. mesenteroides* ssp. *premoris*, LMG 18812, LMG 18813, and LMG 18889) *Leuconostoc sensu stricto* species (Table 1) was studied by means of whole-cell protein analysis. Strains were grown for 5 days on MRS agar at 25 C in a microaerobic atmosphere (approximately 5% O₂, 10% CO₂, and 85% N₂). Preparation of cellular protein extracts and polyacrylamide gel electrophoresis were performed as described previously (36). Briefly, discontinuous gels were run overnight at constant temperature in a vertical slab apparatus. The separation gel was long and contained 12% total acrylamide (the monomer solution contained 2.67% cross-linking in 0.375 M Tris-HCl [pH 8.8] and 0.1% sodium dodecyl sulfate). The stacking gel was 12 mm long and contained 30% total acrylamide (the monomer solution contained 2.67% cross-linking in 0.125 M Tris-HCl [pH 6.8] and 0.1% sodium dodecyl sulfate). Protein bands were stained with Coomassie blue R-250 in 50% (v/v) methanol and 10% (vol/vol) acetic acid. These conditions allowed the separation of proteins and peptides in the molecular weight range of 14,000 to 116,000.

Isolation of DNA, REA, and 16 and 23S rDNA RFLP (ribotyping). EcoRI, and HindIII restriction enzymes (New England Biolabs, Beverly, Mass.) were used for ribotyping. DNA was isolated by the guanidium thiocyanate method of Pitcher et al. (35) as modified by Bax and Korkeala (3) by the use of lysozyme and mutanolysin (Sigma, St. Louis, Mo.) treatment. Restriction endonuclease treatment of DNA was done as specified by the manufacturer (New England Biolabs), and REA was performed as described previously (3). Before Southern blotting, the REA patterns were inspected in order to obtain preliminary information about clonal variation. Genomic blots were made using a vacuum device (Vacugene; Pharmacia, Uppsala, Sweden), and the rDNA probe for ribotyping was labeled by reverse transcription (AMV-RT [Promega, Madison, Wis.] and Dig DNA labeling kit [Roche Molecular Biochemicals, Mannheim, Germany]) as previously described (Korkeala et al. (9)). Membranes were hybridized at 68 C as described by Bax and Korkeala (3).

Pattern analysis. The ClaI, EcoRI, and HindIII ribopatterns were compared with the corresponding patterns in the previously established LAB database. For cluster analysis, ribopatterns were scanned using a Hewlett-Packard Jet 4c/T scanner and analyzed using the BioNumerics 1.0 software package (Applied Maths, Kortrijk, Belgium). The similarities between pairs of patterns were assessed by Dice coefficient correlation, and unweighted pair-group method with arithmetic averages clustering was used for the construction of dendrograms. The protein profiles were scanned using a 2202 UltraScan laser densitometer (LKB, Bromma, Sweden). The densitometric analysis, normalization, and correlation of the protein profiles were performed with the GelComp software package (Applied Maths). Numerical analysis was performed using the BioNumerics 1.0 software package. The similarities between all pairs of patterns were expressed by the Pearson product moment correlation coefficient converted to a percent value.

Different types of banding patterns were integrated in a single dendrogram by means of the three different enzymes were performed by using the BioNumerics 1.0 software package. In these combined analyses, equal weight was given to each of the three banding patterns.

The 16S rRNA gene sequence analysis. The 16S rRNA gene was amplified with a universal primer pair (45): primer A (5'-GAGTATGATCTGGCTCAG-3') and primer B (5'-GAAAGGAGGTGATCCAGCC-3'). Two strains, LMG 18811 and LMG 18812, were studied. They represented the two groups detected in ribotyping. Chromosomal DNA was isolated as for ribotyping. Amplification was performed using 200 ng of chromosomal DNA as a template. The PCR mixture contained 500 U of Taq polymerase (Promega), 100 U of Taq polymerase buffer (BioLabs), 5% sucrose (20), and of nucleotide mixture (dATP, dCTP, dTTP, and dGTP; 2.5 mM each), 4 mM of MgCl₂ (25 mM), 1.25 M of primers A and B (120 μM), and 10 μM of DMSO. The cycles used for amplification were as described previously (36).

Sequencing of the purified PCR product (Quantum Prep PCR Kleen spin columns, Bio-Rad Laboratories, Hercules, Calif.) was performed by Sanger dideoxynucleotide chain termination method using an ABI PRISM sequencer (Perkin-Elmer Corp., Norwalk, Conn.) according to the manufacturer's recommendations. Sequencing was performed as two long reactions, and complementary sequences were joined by the DNASIS program (Hitachi Software, Yokohama, Japan).

Phylogenetic analysis was performed by using the GeneCompar 2.0 software package (Applied Maths). The consensus sequence and the sequences of

TABLE 2. Recovery of LAB on MRS and Rogosa selective Lactobacillus agar and pH values analyzed from six spoiled packages showing clear bulging due to gas formation

Package no.	No. of LAB (CFU/g)		pH
	MRS	Rogosa selective Lactobacillus agar	
1	23 10 ¹⁰	13 10 ¹⁰	4.9
2	43 10 ¹⁰	43 10 ¹⁰	4.8
3	43 10 ¹⁰	13 10 ¹⁰	4.7
4	13 10 ¹⁰	23 10 ⁸	4.8
5	13 10 ¹⁰	33 10 ¹⁰	4.8
6	93 10 ¹⁰	33 10 ¹⁰	5.0

belonging to the same phylogenetic group (retrieved from the National Center for Biotechnology Information GenBank data library) were aligned. The accession numbers of the 16S rDNA sequences used are as follows: *Leuconostoc argentinum* LMG 18543^T, AF175403; *Leuconostoc carnosum* LMG 11498^T, X95997; *Leuconostoc citreum* LMG 9824^T, X53963 and S78399; *Leuconostoc fallax* LMG 13177^T, S63851; *Leuconostoc gelidum* LMG 9850^T, S63851; *Leuconostoc lactis* LMG 8894^T, M23031 and M23032; *Leuconostoc mesenteroides* subsp. cremoris LMG 6909^T, M23034; *Leuconostoc mesenteroides* subsp. mesenteroides LMG 6893^T, M23035; *Leuconostoc pseudomesenteroides* LMG 6852^T, X95979; and *Weissella paramesenteroides* LMG 6852^T, X95982.

DNA base composition and DNA-DNA hybridization was isolated from two spoilage isolates (LMG 18811 and LMG 18813) and the type strain NCFB 2775, and the *Leuconostoc mesenteroides* subsp. dextranictype strain DSM 20484. *L. gelidum* was selected because it had the highest similarity to the main spoilage species according to some phenotyping schemes, RFLP analysis, whole-cell protein analysis, and 16S rDNA sequence analysis. *L. gelidum* was chosen on the basis of API characterization results, and it was also used as a control species in the present study.

For large-scale DNA isolation, the modified (3) guanidium thiocyanate method of Pitcher et al. (35) was scaled up 10-fold. Cells from 200 ml of well-grown MRS broth culture were used for each isolation batch. In one batch was dissolved overnight in 1 ml of TE 10:1 (10 mM Tris, 1 mM EDTA, pH 8.0). RNase A (Sigma) was added to provide a concentration of 125 mg/ml, and the solution was incubated at 37 °C with gentle shaking for 1 h. Following the 1-h incubation, proteinase K (Sigma) was added to provide a concentration of 1 mg/ml, and incubation at 37 °C was continued for at least 6 h. DNA was isolated as described by Pitcher (35) and dissolved in SSC 130. SSC 1 is 0.15 M NaCl plus 0.015 M sodium citrate). When dissolved, the SSC concentration of a sample was adjusted to 20 × SSC.

Purified DNA was dialyzed twice overnight at 4 °C using a 12,000–14,000 Da-pore-size membrane (Mediatech International Ltd., London, United Kingdom). The first dialysis was carried out against DEPC (10 mM), and the second was carried out against SSC 1. DNA was fragmented two times in a French pressure cell (SML Aminco; Colora Messtechnik GMBH, Lorch, Germany) at about 13,500 Pa. Before reassociation, it was dialyzed overnight at 4 °C against SSC 2.

The DNA base composition (moles percent) was estimated by the thermal-denaturation method (12), and the DNA homology values were determined from renaturation rates using a Gilford Response spectrophotometer (Giba Corning Diagnostics Corp., Gilford Systems, Oberlin, Ohio).

Nucleotide sequence accession numbers for approximately 1,500-bp sequences of the 16S ribosomal genes of strains LMG 18811 and LMG 18813 have been deposited in the GenBank data library with accession numbers AF231131 and AF231132, respectively.

RESULTS

Microbiological and sensorial qualities of the products according to Table 2 shows the results of microbial enumeration on MRS and Rogosa selective Lactobacillus agars and corresponding pH values obtained from the six packages. An increase in the pH of the product, very atypical for LAB spoilage, was detected. Instead of the normal pH values, ranging from 4.2 to 4.4, values from 4.7 to 5.0 were detected. All of the packages were deemed unfit for human consumption by all three judges. They were all described as clearly bulged, and the smell of the product was described as pungent and very unpleasant. The consistency and texture of the product was, however, normal, and no color changes were visible.

LAB population associated with the spoiled product shows the division of the 120 isolates into different and groups of species based on the LAB ribotyping database. An organism possessing typical lower molecular bands leuconostoc HindIII ribopatterns (Fig. 1C) was found to dominate (67 of 120) in the product. These isolates were gram-positive, catalase-negative oval cocci, produced glucose, and did not grow on Rogosa agar. They possessed identical ribopatterns, showing, however, two different the REA patterns. The distribution of the isolates between these two REA types was almost even.

The two other major species associated with the product were *Lactobacillus curvatus* (32 of 120) and *Lactobacillus sakei* (16 of 120). Isolates possessing ribotypes identical to *L. sakei* and *L. curvatus* patterns in the database (no new patterns were detected) were gram-positive rods or coccoid. The ones grown on Rogosa agar, were catalase negative, and did not produce gas from glucose. Three of the 120 isolates identified as *Leuconostoc sensu stricto* species, *Leuconostoc summatum* and *L. gelidum*. They all shared identical ribopatterns with the corresponding *Leuconostoc* type strain, were oval cocci, produced gas from glucose, and did not grow on Rogosa selective Lactobacillus agar. Twelve isolates could not be identified with the existing ribotyping database. They were gram-positive rods growing on Rogosa selective Lactobacillus agar, produced gas from glucose, and shared identical ribopatterns. They did not have any similarity to the ribopatterns of *Leuconostoc brevis*, *Leuconostoc buchneri*, *Leuconostoc colliformis*, *Leuconostoc fermentum*, *Leuconostoc fructivorans*, *Leuconostoc hilgardii*, or *L. sibiricus*. This was also the case with respect to the patterns of *Carnobacterium divergens*, *Carnobacterium piscium*, *Carnobacterium mobile*, and *Carnobacterium valinartum* type strains.

Phenotypic reactions of the main spoilage species, LMG 18811, LMG 18812, LMG 18813, and LMG 18889 strains showed typical reactions for the genus *Leuconostoc*. They did not produce ammonia from arginine, did not grow in the presence of 6.5 to 12% NaCl, and synthesized only lactic acid from glucose. They all grew at 4 and 15 °C but not at 25 °C. Growth was already slower at 30 °C, and during the study it was observed as an optimum temperature for growth on MRS. All four strains produced excessive slime from sucrose and fermented arabinose, ribose, glucose, fructose, mannose, a-methyl-glucoside, acetyl-glucosamine, esculetin, cellobiose, maltose, melibiose, sucrose, trehalose, gentiobiose, turanose, and 5-keto-gluconate. LMG 18811, LMG 18813, and LMG 18889 also fermented galactose and gluconate. Glycerol, erythritol, xylitol, xylose, adonitol, a-methyl-xyloside, sorbose, rhamnose, dulcitol, inositol, mannitol, sorbitol, a-methyl-mannoside, amygdalin, arbutin,

TABLE 3. Species division of 120 LAB isolates originating from packages (20/package) of MA-packaged raw marinated broiler meat products according to the EcoRI, and HindIII ribopattern database

Package no.	No. of isolates					
	<i>Leuconostoc</i> spp.	<i>L. carnosum</i>	<i>L. gelidum</i>	<i>L. sakei</i>	<i>L. curvatus</i>	Unidentified
1	4	1	2	4	5	4
2	9			4	5	2
3	10			3	5	2
4	15			4		1
5	5			1	6	1
6	12				11	2
Total	57	1	2	16	32	12

FIG. 1. Ribopatterns and numerical analysis of the patterns presented as dendrograms. Patterns and dendrograms (A) and (B), by using HindIII (C) restriction enzymes are shown. The left sides of the banding patterns show large molecular sizes (23 Kbp), and the right (50 Kbp) show small sizes (10 Kbp). The dendrograms show the similarity values for the patterns. The dendrogram (A) shows that the Argentine strain showed ribopatterns identical to those of the Chilean strain. Scales from 30 to 100 show percentile similarity values for the patterns.

TABLE 4. Main differences in sugar fermentation between *L. carnosum*, *L. gelidum*, and *L. gasicomitatum* nov. sp. nov.

Sugar	Fermentation		
	<i>L. carnosum</i>	<i>L. gelidum</i>	<i>L. gasicomitatum</i>
Amygdalin	2	1	2
L-Arabinose	2	1	1
Arbutin	2	1	2
Raffinose	2	1	1
D-Xylose	2	1	1
Salicin	2/1 ^c	1	2

^a Reactions adapted from Shaw and Harding (43).
^b 2, no fermentation; 1, fermentation takes place.
^c Most strains do not ferment salicin.

salicin, lactose, inulin, melezitose, starch, glycogen, xyli-
 lyxose, tagatose, fucose, arabinol, and 2-keto-gluconate were
 not fermented. The API Chem System identified these isolates with an extremely good identification level (99.9%) as *L. mesenteroides* p. 2. Table 4 shows the key carbohydrate fermentation reactions among the phylogenetically associated *L. gelidum* and the strains representing the main spoilage group.

Peptidoglycan type of the main spoilage species. Cell walls of LMG 18811 contain, besides muramic acid and glucosamine, the amino acids lysine, glutamic acid, and alanine in a molar ratio of 1:1:4, respectively. The amino acid composition of the partial hydrolysate were compatible only with type A, 3-Lys-Ala-Ala.

Enzymatic activities. None of the *L. gasicomitatum* nov. isolates changed the appearance of the skim milk on MRS agar. According to API ZYM analysis, both LMG 18811 and LMG 18812 showed the presence of alkaline phosphatase, lipase, α-galactosidase, α-glucosidase, α-amylase, and naphthol-AS-BI-phosphohydrolase activities.

Numerical analyses of the main spoilage species based on the HindIII ribotypes and whole-cell protein patterns and whole-cell protein patterns. Figure 2 shows the dendrograms and banding patterns of the main spoilage species and the reference strains *C. casei* ATCC 35061 and *L. lactis* ATCC 7940. Figure 3 shows a dendrogram obtained by combining the pattern information of all ribotypes into one numerical analysis. The result of the numerical analysis of the whole-cell protein patterns is shown in Figure 4, and the combined information from all of the ribotypes and whole-cell protein analysis is presented as a dendrogram in Figure 5.

The three spoilage isolates and the reference strains *Leuconostoc* species formed distinct clusters in the dendrograms based on the HindIII ribotypes (Fig. 1C) and the protein patterns (Fig. 3), indicating that these techniques generated species-specific patterns. In the dendrograms generated from numerical comparison of HindIII ribotypes, only *Citream. pseudomesenteroides* and the spoilage isolates formed distinct species-specific clusters (Fig. 1C). When equal weight is given to all three types of ribotypes, the analysis combining this information resulted in distinct species-specific clusters (Fig. 2). The *L. gelidum* type strain had the highest similarity to spoilage isolates in the numerical analyses of combined patterns (Fig. 2) as well as in the whole-cell protein patterns (Fig. 3). In respect to the subdivision of the genus, the numerical analyses performed correlated with the interspecies division of the genus. The type strain of *L. lactis* ATCC 7940 was found to possess the same ribotypes as the LMG 7940 strain, and in the dendrogram based on the numerical analysis of whole-cell protein patterns the LMG 7940 strain formed a tight cluster with other strains (Fig. 3). Such a close association was only seen among strains of a single species. Phylogenetic analyses based on 16S rDNA sequences shows the sequence homologies of strains LMG 18811 and LMG 18812 compared with the *Leuconostoc sensu stricto* spe-

FIG. 2. Dendrogram obtained by combining the pattern information of HindIII ribotypes and whole-cell protein patterns into one numerical analysis. The scale from 40 to 100 shows percentile similarity values.

FIG. 3. Numerical analysis of whole-cell protein patterns presented as a dendrogram. The scale from 60 to 100 shows percentile si

cies. The two strains, LMG 18811 and 18812, showed DNA-DNA reassociation: LMG 18811/LMG 18812, 100%; sequence homology of 99.3%, and the highest homology with LMG 18811 3 L. gelidum LMG 18297T, 22%; LMG 188123 and 98.8%, respectively, was exhibited with the type L. gelidum LMG 18297^T, 34%; and LMG 188123 L. mesen- teroides LMG 6908^T, 33%. The DNA G1 C DNA base composition and DNA-DNA hybridization contents of strains LMG 18811 and LMG 18812 are 37 and 38 sults. The following DNA homology values were obtained: LMG 18811, 37%; LMG 18812, 38%; respectively.

FIG. 4. Combined information from EcoRI, and HindIII ribopatterns and whole-cell protein patterns presented as a dendrogram. The strain showed ribopatterns identical to LMG 7940. The scale from 40 to 100 shows percentile similarity values.

TABLE 5. Homology values for a 1,491-nucleotide region of 16S rDNA

Species	Homology (%) to:											
	1	2	3	4	5	6	7	8	9	10	11	12
1. <i>Leuconostoc</i> sp. strain LMG 18811	100											
2. <i>Leuconostoc</i> sp. strain LMG 18812	99.3	100										
3. <i>L. argentinum</i> LMG 18534 ^T	97.1	97.1	100									
4. <i>L. carnosum</i> LMG 11498 ^T	98.3	98.5	97.5	100								
5. <i>L. citreum</i> LMG 9824 ^T	97.5	97.6	98.7	97.4	100							
6. <i>L. fallax</i> LMG 13177 ^T	93.6	93.7	92.4	94.3	94.5	100						
7. <i>L. gelidum</i> LMG 9850 ^T	98.8	99.0	97.1	98.3	97.4	93.5	100					
8. <i>L. lactis</i> LMG 8894 ^T	97.3	97.6	99.3	97.3	98.5	93.6	97.6	100				
9. <i>L. mesenteroides</i> sp. nov. LMG 6909 ^T	97.7	98.0	97.7	97.8	97.7	94.6	97.9	98.2	100			
10. <i>L. mesenteroides</i> sp. nov. LMG 6893 ^T	97.7	98.0	97.5	97.8	97.7	94.6	97.9	98.2	100	100		
11. <i>L. pseudomesenteroides</i> LMG 1482 ^T	97.9	98.0	97.8	97.8	97.5	94.6	98.0	98.0	99.5	99.5	100	
12. <i>W. paramesenteroides</i> LMG 9852 ^T	91.2	91.4	90.3	91.0	91.2	92.4	90.9	91.3	91.8	91.8	91.7	

DISCUSSION

Gaseous deterioration caused by LAB has mainly been associated with highly acidic foods, such as fermented vegetables (16, 29) or acetic acid preserves (6, 24), but it may also affect meat products (10). Even though LAB have also been found as the dominant spoilage organisms in vacuum- or MA-packaged poultry products (34, 39, 40), the strains have never been identified to the species level. Gaseous deterioration as a LAB spoilage type in vacuum- or MA-packaged poultry products has not been reported previously. It was not surprising to find *L. sakei* and *L. curvatum* strains in the poultry product studied here. These species are very typical for all meat products (3, 4, 26, 28; Holzapfel and Gerber, Abstr. 32nd Eur. Meet. Meat Res. Workers) and might have been a major component of the spoilage population detected in the previous studies dealing with vacuum- or MA-packaged poultry products (17, 33, 39, 40). *L. carnosum*, *L. gelidum* and also quite common species occurring in vacuum- or MA-packaged cold-stored meat products (43, 49). Two species occurring in the spoiled poultry product were unusual LAB species for meat products. *U. comitatus* sp. nov. was the main spoilage species characterized in this study, and the identification of the gram-positive rod shaped organism will be carried out as a separate study. In addition to the novel species, this LAB spoilage product also showed unique properties. Normally, in a case of clear LAB spoilage, the pH of the product decreases due to lactic acid formation, but in this case an increase of pH was detected. This type of LAB spoilage was first reported by Meyer (50) for some LAB (50). The plant was simultaneously processed in canned fish marinades. He called it "protein swell" and distinguished it from "carbohydrate swell," where increased CO₂ and CO₂ formation result from heterofermentative utilization of glucose. In protein swell, proteins are decomposed by proteolytic enzyme action, and the subsequent decarboxylation of amino acids leads to enhanced CO₂ production. Therefore, the LAB having an effect on gas production in protein swell may also possess homofermentative glucose metabolism. An increase in acidity related to protein swell has been attributed to production of ammonia by bacterial deamination of amino acids. Protein swell has also been reported to affect stuffed olives (19), but to our knowledge there are no reports of this type of spoilage affecting any type of meat product. The previous studies of protein swell have assigned the main component of the spoilage LAB with the decarboxylase activity to *U. comitatus* sp. nov. (19, 30) and considered protein hydrolysis to be due to endogenous fish enzymes. The initial hydrolysis of muscle proteins has also been attributed to endogenous enzymes, mainly cathepsins, and the bacterial activity was associated with the degradation of oligopeptides and

amino acids (32, 46). The proteolytic systems of various related LAB are poorly known, and the ability of *L. curvatum* to degrade myofibrillar proteins has only recently been studied (15, 37). These species have been shown to possess peptidase activity and also to express strong amino acid metabolism (15, 33, 37, 38), and even though they were not major components of the spoilage population, they may have played a major role in this case. *Leuconostocs* have not yet been detected as dominant species in protein swell, which makes their predominance in *U. comitatus* species produce gas during normal glucose fermentation, and in this case, the extreme bloom may have resulted from complicated interaction between various LAB species and the endogenous muscle-associated enzymes. Whether the *U. comitatus* component alone, or in association with the endogenous muscle proteinases, could cause the gaseous spoilage remains unknown. There are no data on the proteolytic systems of meat-associated leuconostocs. The main component did not show proteolytic activity on the skim milk-supplemented MRS agar due to the substrate specificity of proteolytic systems. Complicated techniques should be used for the evaluation of the proteolytic effect on myofibrillar proteins. In the product, the LAB counts were exceptionally high (10⁹ g). Two factors, the marinade and the small rise in pH, have played major roles in facilitating the growth of LAB. The marinade had a tomato base, which contains growth stimulating factors. The plant was simultaneously processed in poultry strips in other marinades, such as honey based, but the carbohydrate-marinated product showed gaseous deterioration. The carbohydrates and protein hydrolysates in the marinade may have provided nutrients facilitating pronounced gaseous spoilage. The spoilage problem was overcome by stabilizing the pH with another type of additive. Apparently this could have an effect on the growth of leuconostocs, which are generally as acid tolerant as lactic acid bacteria species. *Leuconostoc sensu stricto* comprises *L. argentinum*, *L. carnosum*, *L. citreum*, *L. gelidum*, *L. lactis*, *L. mesenteroides* (three subspecies), *L. pseudomesenteroides* and *U. comitatus* showing 97 to 99% 16S rDNA sequence homology. In addition, an atypical leuconostoc strain assigned to 95% 16S rDNA homology with the other sensu stricto species, has been described. Our results show high sequence homology between *U. comitatus* and *L. fallax* (93.6 and 93.7%). According to these data

