

## Invited review

## Dual roles of NF- $\kappa$ B in cell survival and implications of NF- $\kappa$ B inhibitors in neuroprotective therapy<sup>1</sup>

Zheng-hong QIN<sup>2,4,5</sup>, Lu-yang TAO<sup>3,4</sup>, Xin CHEN<sup>2,4</sup>

Departments of <sup>2</sup>Pharmacology and <sup>3</sup>Forensic Medicine and <sup>4</sup>Laboratory of Aging and Nervous Diseases, Soochow University School of Medicine, Suzhou 215123, China

### Key words

NF- $\kappa$ B; transcription factor; apoptosis; neuroprotection; neurodegeneration

<sup>1</sup> Project supported by a grant from the National Natural Science Foundation of China (No 30772560).

<sup>5</sup> Correspondence to Prof Zheng-hong QIN. Phn/Fax 86-512-6588-0406. E-mail zhqin5@hotmail.com

Received 2007-09-02

Accepted 2007-09-24

doi: 10.1111/j.1745-7254.2007.00741.x

### Abstract

NF- $\kappa$ B is a well-characterized transcription factor with multiple physiological and pathological functions. NF- $\kappa$ B plays important roles in the development and maturation of lymphoids, regulation of immune and inflammatory response, and cell death and survival. The influence of NF- $\kappa$ B on cell survival could be protective or destructive, depending on types, developmental stages of cells, and pathological conditions. The complexity of NF- $\kappa$ B in cell death and survival derives from its multiple roles in regulating the expression of a broad array of genes involved in promoting cell death and survival. The activation of NF- $\kappa$ B has been found in many neurological disorders, but its actual roles in pathogenesis are still being debated. Many compounds with neuroprotective actions are strongly associated with the inhibition of NF- $\kappa$ B, leading to speculation that blocking the pathological activation of NF- $\kappa$ B could offer neuroprotective effects in certain neurodegenerative conditions. This paper reviews the recent developments in understanding the dual roles of NF- $\kappa$ B in cell death and survival and explores its possible usefulness in treating neurological diseases. This paper will summarize the genes regulated by NF- $\kappa$ B that are involved in cell death and survival to elucidate why NF- $\kappa$ B promotes cell survival in some conditions while facilitating cell death in other conditions. This paper will also focus on the effects of various NF- $\kappa$ B inhibitors on neuroprotection in certain pathological conditions to speculate if NF- $\kappa$ B is a potential target for neuroprotective therapy.

### Introduction

NF- $\kappa$ B is ubiquitously expressed in peripheral and brain cells and regulates the expression of a wide variety of genes involved in cell survival, growth, stress responses, and immune and inflammatory processes<sup>[1–3]</sup>. This factor was first described by Sen and Baltimore in 1986 as a NF, that when activated by agents, such as bacterial lipopolysaccharide, bound to a 10 bp sequence in the enhancer region of the gene encoding the  $\kappa$  light chain ( $\kappa$ ) of antibody molecules in B cells (B)<sup>[4]</sup>. NF- $\kappa$ B family members have been implicated in the development of the nervous system and plasticity of synapses<sup>[5–7]</sup>. NF- $\kappa$ B is persistently activated in cancer, chronic inflammation, neurodegenerative diseases, stress, stroke, trauma, heart disease, and other disease conditions<sup>[8,9]</sup>.

As NF- $\kappa$ B is an important regulator in programmed cell death<sup>[10]</sup>, it has been speculated that NF- $\kappa$ B may play important roles in normal brain function and neurodegenerative disorders<sup>[11,12]</sup>.

### NF- $\kappa$ B biology

**c-Rel/NF- $\kappa$ B family** NF- $\kappa$ B is composed of 5 members of the c-Rel (Rel) family, including NF- $\kappa$ B1 (p50), NF- $\kappa$ B2 (p52), RelA (p65), RelB, and Rel. All the Rel proteins contain a conserved N-terminal region, called the Rel homology domain (RHD). The N-terminal part of the RHD contains the DNA-binding domain, whereas the dimerization domain is located from C-terminal region of the RHD<sup>[13]</sup>. Close to the C-terminal end of the RHD lies the nuclear localization signal (NLS), which is essential for the transport of active NF- $\kappa$ B

complexes into the nucleus<sup>[14]</sup>. NF- $\kappa$ B family proteins are divided into 2 groups based on C-terminal sequences of the RHD. The members of group 1 include NF- $\kappa$ B proteins p105 and p100, which are precursors of p50 and p52. Limited proteolysis is required to produce p50 and p52. The second group (the Rel proteins) mainly includes c-Rel (and its retroviral homologue v-Rel), RelB, and RelA (p65)<sup>[15]</sup>. All vertebrate NF- $\kappa$ B proteins can form homodimers or heterodimers, except for RelB, which can only form heterodimers. These homodimers and heterodimers that exhibit differential binding specificities are p50/RelA, p50/c-Rel, p52/c-Rel, p65/c-Rel, RelA/RelA, p50/p50, p52/p52, RelB/p50 and RelB/p52<sup>[16]</sup>. The term NF- $\kappa$ B commonly refers specifically to a p50–RelA (p50/p65) heterodimer, which is the major Rel/NF- $\kappa$ B complex in most cells<sup>[14]</sup>.

NF- $\kappa$ B dimers are sequestered in the cytoplasm by a class of inhibitor proteins, called I $\kappa$ B. In mammalian cells, the major regulatory I $\kappa$ B proteins are I $\kappa$ B- $\alpha$ , I $\kappa$ B- $\beta$ , I $\kappa$ B- $\epsilon$ , and Bcl-3. The most common complex that is activated in mammalian cells appears to involve I $\kappa$ B- $\alpha$ , which binds to the p50/RelA heterodimer. I $\kappa$ B function as inhibitors through ankyrin repeats that interact with the RHD in NF- $\kappa$ B to mask the NLS and inhibit the nuclear translocation of NF- $\kappa$ B. The N-termini of these I $\kappa$ B proteins constitute a signal response domain, which is targeted for phosphorylation and ubiquitination by a variety of stimuli. The newly-synthesized I $\kappa$ B- $\alpha$  protein actively shuttles between the nucleus and the cytoplasm and both inhibit nuclear import and mediate the nuclear export of NF- $\kappa$ B/Rel proteins. In contrast, the I $\kappa$ B- $\beta$  protein can inhibit the nuclear import of NF- $\kappa$ B/Rel proteins, but does not remove NF- $\kappa$ B/Rel proteins from the nucleus<sup>[17,18]</sup>.

**NF- $\kappa$ B activation pathway** Signals that induce NF- $\kappa$ B activity cause the phosphorylation of I $\kappa$ B thereby activating the NF- $\kappa$ B complex. The activated NF- $\kappa$ B complex translocates into the nucleus and binds to DNA at the  $\kappa$ B binding motifs and alters gene expression. Most signals that lead to the activation of NF- $\kappa$ B act on a high molecular weight complex containing a serine-specific I $\kappa$ B kinase (IKK). IKK contains at least 3 distinct subunits: IKK- $\alpha$ , IKK- $\beta$ , and IKK- $\gamma$ . IKK- $\alpha$  and IKK- $\beta$  are catalytic kinase subunits, while IKK- $\gamma$  is a regulator for sensing and integrating upstream activating signals<sup>[18]</sup>. There are 2 NF- $\kappa$ B activation pathways: the classical or canonical pathway and the non-canonical pathway. In the canonical pathway, the activation of the IKK complex leads to the phosphorylation of 2 specific serines (Ser32 and Ser36) in I $\kappa$ B- $\alpha$ , which targets I $\kappa$ B- $\alpha$  for ubiquitination and degradation by the 26S proteasome. In the non-canonical pathway, the p100–RelB complex is acti-

vated through phosphorylation by an IKK- $\alpha$  homodimer (lacking IKK- $\gamma$ ) to generate p52–RelB. In either pathway, the unmasked NF- $\kappa$ B complex can then enter the nucleus to activate target gene expression. In the classical pathway, one of the target genes activated by NF- $\kappa$ B can encode I $\kappa$ B- $\alpha$ , and newly-synthesized I $\kappa$ B- $\alpha$  enters the nucleus recombined with NF- $\kappa$ B, which can remove NF- $\kappa$ B from DNA, and export the complex back to the cytoplasm to restore the original latent state. Thus, the NF- $\kappa$ B activating pathway is a transient process, generally lasting from 30 to 60 min in most cells<sup>[20,21]</sup>.

## Dual roles of NF- $\kappa$ B in cell death and survival

NF- $\kappa$ B targets many genes to activate their expression. These target genes include cytokines/chemokines and their modulators, immunoreceptors, proteins involved in antigen presentation, cell adhesion molecules, acute-phase proteins, early response genes, stress response genes, cell surface receptors, transcription factors and regulators, regulators of apoptosis, growth factors, and cell death receptor ligands and their modulators. In the central nervous system (CNS), NF- $\kappa$ B can play an anti-apoptotic or pro-apoptotic role in cell death<sup>[22]</sup>. This is not unexpected as NF- $\kappa$ B regulates the genes involved in neuronal death and survival.

**Anti-apoptotic activity of NF- $\kappa$ B** A large number of studies have demonstrated that NF- $\kappa$ B plays a prosurvival role in proliferating cells, including tumor cells. Two actions of NF- $\kappa$ B make it an important cell survival transcription factor in these cells: the regulation of the cell cycle and the inhibition of apoptosis. The best example for elucidating the prosurvival action of NF- $\kappa$ B in cells is finding an inhibitory role of NF- $\kappa$ B in death receptor-induced apoptosis<sup>[23]</sup>. Binding to death receptors by TNF- $\alpha$  activates caspase-8 through TRAF1 and NF- $\kappa$ B through TRAF2. The activation of caspase-8 leads to apoptosis. Blocking NF- $\kappa$ B activation potentiates TNF- $\alpha$ -induced apoptosis, indicating NF- $\kappa$ B exerts anti-apoptotic action. It has been found that nerve growth factor (NGF) promotes neuronal survival through activating NF- $\kappa$ B<sup>[24]</sup>. Yu *et al*<sup>[25]</sup> reported that mice lacking the p50 subunit of NF- $\kappa$ B exhibited increased damage to hippocampal pyramidal neurons after the administration of the excitotoxin kainate. In immortalized mouse hippocampal cell line HT22 cells, glutamate-induced apoptosis was inhibited by I $\kappa$ B inhibitor aspirin, while the NF- $\kappa$ B decoy oligonucleotide potentiated it<sup>[26]</sup>. The action of NF- $\kappa$ B on neuronal survival is mediated through the upregulation of several prosurvival genes.

**Superoxide dismutase** Manganese superoxide dis-

mutase (Mn-SOD) is an important antioxidant enzyme, which is a potent scavenger of superoxide anion and is likely to serve important cytoprotective roles against cellular damage. It has been reported that NF- $\kappa$ B is involved in the expression of Mn-SOD<sup>[27]</sup>. The incubation of human endometrial stromal cells with TNF- $\alpha$  or the phorbol 12-myristate 13-acetate (TPA), a protein kinase C activator, caused marked increases in nuclear NF- $\kappa$ B DNA binding activity and Mn-SOD mRNA and activity. These effects of TNF- $\alpha$  and TPA were completely inhibited by the proteasome inhibitor MG132 and the recombinant peptide capable of blocking NF- $\kappa$ B nuclear translocation, SN50<sup>[28]</sup>. Activities of Mn-SOD and SOD1 increased after spinal cord injury (SCI) and exposure to neurotoxins<sup>[29,30]</sup>. The increase in SOD appeared to be NF- $\kappa$ B-dependent, and overexpression significantly protected against the deleterious effect of reactive oxygen species, ceramide, or *N*-methyl-*D*-aspartate (NMDA). Several other studies have shown that the overexpression of copper-Zn SOD or the activation of Mn-SOD is neuroprotective against ischemia, excitotoxicity, and A $\beta$  toxicity<sup>[31-33]</sup>.

**Bcl-2** Bcl-2 and Bcl-X<sub>L</sub> are well-defined anti-apoptotic proteins. The NF- $\kappa$ B binding site is identified in the promoter of murine Bcl-x<sup>[34]</sup>. Bui *et al*<sup>[35]</sup> found that NGF increased the expression of Bcl-X<sub>L</sub>, possibly through the activation of NF- $\kappa$ B. Some studies indicate that TNF- $\alpha$  has neuroprotective effects. TNF- $\alpha$  increases the mRNA and protein levels of Bcl-2 and Bcl-x<sup>[36]</sup>. It has also been found that the exposure of cultured neurons to hypoxia/reoxygenation increases the levels of Bcl-2 and Bcl-X. The inhibition of NF- $\kappa$ B activation abolished the hypoxia-induced induction of Bcl-2 and Bcl-X, indicating that the induction of Bcl-2 and Bcl-X is mediated by NF- $\kappa$ B<sup>[37]</sup>. NF- $\kappa$ B is also reportedly involved in the activation of the Bcl-2 family member A1/Bfl-1<sup>[38]</sup>.

**Pro-apoptotic activity of NF- $\kappa$ B** A large number of studies have found that NF- $\kappa$ B activation participates in neuronal apoptosis. The mechanism by which NF- $\kappa$ B translocation induces apoptosis is not completely clear, but it is assumed that this mechanism involves the regulation of 1 or more genes known to play a pro-apoptotic role in apoptosis. Among the NF- $\kappa$ B-responsive genes possibly involved in the control of neuronal cell death, pro-apoptotic genes p53, c-Myc, cyclin D1, Bcl-Xs, and the Fas ligand and its receptor are activated by various pathological stimuli.

**p53** The p53 protein is a tumor suppressor and plays important roles in neuronal apoptosis via promoting the expression of the pro-apoptotic gene Bax and PUMA, but suppresses the expression of the cytoprotective gene Bcl-2. NF- $\kappa$ B may contribute to neuronal apoptosis through the induc-

tion of p53. In the study of glutamate receptor-mediated excitotoxicity, upon stimulation of glutamate receptors, a quick and robust induction in the levels of p53 mRNA and protein was observed. The induction of p53 was blocked by NF- $\kappa$ B inhibitors<sup>[39,40]</sup>. Qin *et al*<sup>[40]</sup> investigated the role of NF- $\kappa$ B in apoptosis induced by the NMDA receptor agonists in rat striatal medium spiny neurons. The administration of the excitotoxin quinolinic acid and NMDA induced apoptosis in the rat striatum. The inhibition of NF- $\kappa$ B nuclear translocation by the SN50, a recombinant cell permeable peptide containing the p50 nuclear localization sequence, reduced apoptotic death of striatal neurons and p53 expression. Uberti *et al*<sup>[41]</sup> pretreated the neuronal cultures with aspirin, which inhibits NF- $\kappa$ B activation, or with a specific p53 antisense oligonucleotide, which inhibits p53 transcription, resulting in a complete prevention of glutamate-induced p53 induction and apoptosis. The NF- $\kappa$ B-dependent induction of p53 was also found in response to DNA damage and oxidative stress. The induced p53 was apparently involved in cell death under these conditions as the synthetic p53 inhibitor pifithrin- $\alpha$  blocked neuronal apoptosis<sup>[42-45]</sup>. NF- $\kappa$ B not only regulates the levels of p53, but also increases the stability of the DNA binding of p53, providing an additional mechanism for promoting p53-mediated pro-apoptotic signaling<sup>[46]</sup>.

**Cyclin D1 and c-Myc** The cyclins are a family of proteins that are involved in cell cycle progression and apoptosis. The best explored link between NF- $\kappa$ B activation and cell cycle progression involves cyclin D1, a cyclin which is expressed relatively early in the cell cycle and is crucial to DNA synthesis<sup>[47]</sup>. The NF- $\kappa$ B regulation of cyclin D1 occurs at the transcriptional level and is mediated by the direct binding of NF- $\kappa$ B to multiple sites in the cyclin D1 promoter<sup>[48]</sup>. NF- $\kappa$ B promotes G<sub>1</sub> to S phase transition in mouse embryonic fibroblasts and in T47D mammary carcinoma cells<sup>[49]</sup>. The NF- $\kappa$ B-mediated induction of cyclin D1 was found in dorsal root ganglion neurons in response to ceramide-induced apoptosis. The inhibition of NF- $\kappa$ B blocked cyclin D1 induction and increased the viability of neurons<sup>[50]</sup>. Liang *et al*<sup>[44]</sup> reported that overstimulation NMDA receptors with quinolinic acid induced a NF- $\kappa$ B-dependent elevation in cyclin D1 mRNA and protein levels. The incorporation of BrdU was observed in some neurons undergoing apoptosis. NF- $\kappa$ B binding sites have also been identified in the c-Myc exon and upstream sites and positively regulate the expression of c-Myc<sup>[51]</sup>. In excitotoxic models, c-Myc was upregulated through NF- $\kappa$ B activation<sup>[39,44]</sup>. The NF- $\kappa$ B-dependent increase in c-Myc expression was also observed in 6-hydroxydopamine-induced Parkinson's dis-

ease<sup>[45,52]</sup>.

The involvement of cell cycle regulators in neuronal apoptosis has been shown by many investigators. The expression of certain cell cycle regulators, such as cyclin D1, cyclin G, c-Myc, and cdk4 have been found during neuronal apoptosis<sup>[53–56]</sup>. To support the role of cell cycle regulators in neuronal apoptosis, some studies showed that cdk inhibitors blocked neurotrophic factor withdrawal-induced apoptosis<sup>[57]</sup>. Cyclin D1 antisense or cell cycle inhibitors and cdk inhibitors partially blocked the excitotoxin-induced apoptosis of striatal neurons<sup>[44,58]</sup>, suggesting cycle regulators play an important role in neuronal apoptosis.

**Bcl-Xs and BAX** The *bcl-x* gene functions to regulate cell death. *Bcl-x* transcripts are alternatively spliced into a long and short form or the form lacking the transmembrane domain. The long form (*bcl-x<sub>L</sub>*) represses cell death, while the short form (*bcl-x<sub>s</sub>*) favors apoptosis. NF- $\kappa$ B binding sites have been identified in the *Bcl-x* promoter<sup>[59]</sup>. The NF- $\kappa$ B-dependent induction in *Bcl-Xs* has been reported. Dixon *et al*<sup>[60]</sup> showed that following ischemia and NF- $\kappa$ B activation, *Bcl-x<sub>s</sub>* messenger RNA levels increase in the CA1 hippocampal region. In cultured endothelial cells, hypoxia decreased *Bcl-2* mRNA levels, whereas the transfection of the NF- $\kappa$ B decoy significantly attenuated a decrease in *Bcl-2* mRNA, increased *Bcl-2/BAX* ratio, and inhibited hypoxia-induced cell death<sup>[61]</sup>. The prolonged activation of NMDA receptors results in NF- $\kappa$ B nuclear translocation, release of LDH, increases in the *BAX/Bcl-X<sub>L</sub>* ratio, and DNA fragmentation. SN50 blocked the NMDA-induced increase in the *Bax/Bcl-X<sub>L</sub>* ratio and cell death<sup>[62]</sup>. Glutamate also reportedly increased the expression of BAX, which was inhibited with BAY 11-7082, a selective inhibitor of I $\kappa$ B- $\alpha$  phosphorylation<sup>[63]</sup>. In cyanide-induced apoptosis, the expression levels of 2 anti-apoptotic *Bcl-2* proteins, *Bcl-2* and *Bcl-X<sub>L</sub>*, remained unchanged after cyanide treatment, whereas the mRNA levels of *Bcl-Xs* and *Bax* began to increase within 2 h, and their protein levels increased 6 h after treatment. Both NF- $\kappa$ B SN50 and the NF- $\kappa$ B decoy blocked the upregulation of *Bcl-Xs* and BAX<sup>[64]</sup>. In low potassium-induced apoptosis of cortical neurons, NF- $\kappa$ B DNA binding increased, and this was accompanied by an elevation in *Bcl-Xs* transcription. The latter was abolished by the inhibition of NF- $\kappa$ B or the restoration of potassium levels<sup>[65]</sup>.

**Nitric oxide** The role of nitric oxide (NO) in apoptosis is complex, as it may exert proapoptotic or antiapoptotic effects depending on experimental conditions. NF- $\kappa$ B plays a role in regulating the expression of NO synthase (NOS). Xie *et al*<sup>[66]</sup> defined a NF- $\kappa$ B binding domain in murine inducible NOS (iNOS). NF- $\kappa$ B stimulated the expression of iNOS. The


NF- $\kappa$ B inhibitor pyrrolidine dithiolidin inhibited the activation of NF- $\kappa$ B and the production of NO in lipopolysaccharide (LPS)-treated macrophages, suggesting that the activation of NF- $\kappa$ B/Rel is critical in the induction of iNOS by LPS.

NOS has been demonstrated to play a proapoptotic role in several *in vitro* and *in vivo* studies. The incubation of human breast cancer cell line MCF-7 cells and differentiated neuronal PC12 cells with TNF- $\alpha$  increased the expression and activity of iNOS. In addition to NOS inhibitors, iNOS antisense oligonucleotides effectively prevented NO<sub>2</sub> generation and apoptosis, suggesting that the TNF- $\alpha$ -induced cell death is mediated by iNOS-derived NO<sup>[67,68]</sup>. Employing the intrastriatal injection of autologous blood in rats to model intracerebral hemorrhage, Zhao *et al*<sup>[69]</sup> demonstrated a robust and prolonged NF- $\kappa$ B activation and a robust induction of iNOS at both the mRNA and protein levels. In SCI models, iNOS was also found to be increased<sup>[70]</sup>, and the drugs inhibiting NOS offered protective effects<sup>[71–73]</sup>.

## NF- $\kappa$ B inhibitors and neuroprotective therapy

Many human nervous system diseases have an association with NF- $\kappa$ B activation. These conditions, including aging<sup>[74]</sup>, headache<sup>[75]</sup>, pain<sup>[76]</sup>, stroke<sup>[77]</sup>, traumatic brain injury<sup>[78]</sup>, SCI<sup>[79]</sup>, Parkinson's disease<sup>[80,81]</sup>, multiple sclerosis<sup>[82]</sup>, Alzheimer's disease<sup>[83,84]</sup>, amyotrophic lateral sclerosis<sup>[85]</sup>, Huntington's disease<sup>[86]</sup>, and brain tumors<sup>[87–91]</sup>, have been associated with the NF- $\kappa$ B pathway. As NF- $\kappa$ B plays important roles in regulating cell survival and death in a broad array of physiological and pathological conditions, it is an attractive proposal to manipulate NF- $\kappa$ B functions to obtain its beneficial effects or abolish its harmful actions when it is required<sup>[92]</sup>. Multiple signaling events are involved in NF- $\kappa$ B activation, including the phosphorylation and degradation of I $\kappa$ B, NF- $\kappa$ B nuclear translocation, and DNA interaction, thus making it a relatively easy target for drug actions (Figure 1). There are a large number of compounds have been reported to inhibit NF- $\kappa$ B functions. These compounds mainly include antioxidants, non-steroidal anti-inflammatory drugs (NSAID), flavonoids, protease inhibitors. A few of these compounds have been used in the clinical setting.

**Antioxidants** Oxidative stress is one of the common pathogenic mechanisms in neurodegenerative disorders. Thus, antioxidants are frequently employed in the treatment of several neurodegenerative diseases, and are the most valuable therapeutic strategy for fighting neurodegeneration. Although it is hard to attribute a single mechanism to any antioxidants' neuroprotective effects, the inhibition of



**Figure 1.** Schematic elustration of NF-κB activation and possible sites where drugs interfere with the NF-κB signaling pathway. NF-κB is sequestered in the cytoplasm by IκB family proteins. Upon stimulation, the IκB is phosphorylated and ubiquitinated and subsequently degraded by the 26S proteasome. Degradation of IκB releases NF-κB dimers that translocate into the nucleus, where NF-κB binds to the consensus DNA sequence and regulates gene expression. ⊗: site of drug action.

NF-κB activation is a prominent feature of antioxidants. The antioxidants include *N*-acetyl-*L*-cysteine (NAC), α-lipoic acid, glutathione monoester, pyrrolidine dithiocarbamate (PDTC), tepoxalin, and flavonoids.

Free radicals are important mediators for NF-κB activation. NAC, a well-characterized antioxidant, is found to exert neuroprotective effects against free radical-related neuronal injury<sup>[93,94]</sup>. NAC influences many cellular signaling pathways, including c-Jun N-terminal kinase, p38 mitogen-activated protein kinase, and redox-sensitive activating protein-1. NAC can also prevent apoptosis and promote cell survival by activating the extracellular signal-regulated kinase pathway. NAC directly modifies the activity of several proteins by its reducing activity<sup>[95]</sup>, and is demonstrated to inhibit the degradation of IκB-α and the activation of NF-κB<sup>[96-98]</sup>. In animal models of global ischemia, pretreatment with NAC (300 mg/kg) or another antioxidant PDTC (200 mg/kg) significantly reduced the infarct volume. NAC has also been reported to increase the survival of dopaminergic neurons. The local or systemic administration of NAC protected dopamine neurons against 6-hydroxydopamine-induced oxidative damage<sup>[99]</sup>.

PDTC is an antioxidant that has been studied for many years. Using cell cultures, Schreck *et al*<sup>[100]</sup> found that micromolar concentrations of PDTC reversibly suppressed

NF-κB activation. PDTC specifically prevented the NF-κB-dependent transactivation of reporter genes under the control of the HIV-1 long terminal repeat and simian virus 40 enhancer. In other studies, PDTC inhibited NF-κB activation while enhancing the binding activity of activator protein-1<sup>[101]</sup>. In addition, PDTC can inhibit the NF-κB-mediated production of TNF-α, hypoxia-induced dephosphorylation of Akt, and inflammatory responses<sup>[102-104]</sup>. It has been shown that treatment with PDTC significantly attenuates reperfusion-induced lung injury<sup>[105]</sup>, glycerol-induced renal injury<sup>[106]</sup>, cholestatic liver injury<sup>[107]</sup>, and adriamycin-induced myocardial apoptosis<sup>[108]</sup>. Crack *et al*<sup>[109]</sup> observed that knockout glutathione peroxidase-1 (Gpx1) in mice increased the ischemia-induced activation of NF-κB. PDTC was able to afford partial neuroprotection in the Gpx1-null mice. In Wistar rats, PDTC prevented NF-κB activation in the ischemic brain, as determined by the reduced DNA binding and nuclear translocation of NF-κB in neurons. PDTC treatment reduced the infarction volume by 48% when given 6 h after MCAO<sup>[110]</sup>.

Flavonoids are potent antioxidants found in many natural products. They are widely used as food supplements and as anti-inflammation and antitumor drugs. Recently, many studies have demonstrated that flavonoids have neuroprotective effects in animal models<sup>[46]</sup>. Among them, *Ginkgo biloba* has received particular attention. Chen *et al*<sup>[111]</sup> found that *Ginkgo biloba* extract significantly reduced intracellular reactive oxygen species formation and NF-κB activation induced by TNF-α. Tea extracts have been previously reported to possess radical scavenger, iron chelating, and anti-inflammatory properties in a variety of tissues. Recent studies found that green tea extracts were capable of inhibiting NF-κB<sup>[112]</sup>. Studies demonstrated that green tea extracts inhibited iron-induced lipid peroxidation, NF-κB activation, and 6-hydroxydopamine (6-OHDA)-induced neuronal death. 6-OHDA-induced apoptosis of catecholaminergic PC12 cells was inhibited by green tea polyphenols and their major effective component epigallocatechin-3-gallate at a concentration of 200 mmol/L<sup>[113]</sup>. Green tea polyphenols have been shown to reduce the toxic effects of β-amyloid, ischemia/reperfusion-induced apoptosis, and the infarct volume<sup>[114,115]</sup>. Given by brain penetrating property of polyphenols, these compounds may be utilized as a class of drugs for the treatment of neurodegenerative diseases.

**IκB phosphorylation and degradation inhibitors** Phosphorylation and the subsequent proteasomal degradation of IκB are key steps for NF-κB activation. NSAID and cyclopentone prostaglandins are now found to be IκB inhibitors. In 1994, Kopp and Ghosh<sup>[116]</sup> reported that sodium salicylate and aspirin inhibited the activation of NF-

$\kappa$ B through blocking the degradation of the I $\kappa$ B. IKK- $\alpha$  and IKK- $\beta$  phosphorylate I $\kappa$ B. Aspirin and sodium salicylate can inhibit IKK- $\beta$  activity *in vitro* and *in vivo*. The mechanism by which aspirin and sodium salicylate inhibit NF- $\kappa$ B is the binding of these agents to IKK- $\beta$  to reduce ATP binding<sup>[117]</sup>. These studies not only further explain the new mechanism of actions, but also suggest new implications of these drugs. Grilli *et al*<sup>[118]</sup> found that acetylsalicylic acid and its metabolite sodium salicylate protected neurons against neurotoxicity elicited by the excitatory amino acid glutamate in rat primary neuronal cultures and hippocampal slices. This inhibitory effect may be involved in the inhibition of NF- $\kappa$ B activation, protein kinase C zeta activity, superoxide anion generation, and lipid peroxidation<sup>[118–120]</sup>. Recent studies suggest that aspirin and other NSAID protected cultured mesencephalic cells against 6-OHDA, 1-methyl-4 phenylpyridinium, and glutamate-induced toxicity<sup>[121,122]</sup>. Using NSAID in Parkinson's disease (PD) has also been proposed<sup>[123]</sup>. It has been found that arthritis patients taking aspirin have a lower incident and later onset of Alzheimer's disease (AD). Emerging evidence shows that aspirin and other NSAID have multiple influences on the AD pathogenic process, including inhibiting the formation of fibrillar A $\beta$ , destabilizing preformed fibrillar  $\beta$ -amyloid (A $\beta$ ), and preventing the aggregation of A $\beta$  and attenuating its toxicity<sup>[124–126]</sup>. Recent studies also indicate that acetylsalicylic acid inhibits tau phosphorylation<sup>[127]</sup>. Other potentially interesting anti-inflammatory drugs have also been reported to exert neuroprotective effects and inhibit NF- $\kappa$ B, including cannabinoid dexamibinol and caffeic acid<sup>[128–130]</sup>.

Recent studies have identified that cyclopentenone prostaglandins (cPG), including prostaglandin (PG)A1 and PGJ2 are ligands for peroxisome-proliferation activator receptor- $\gamma$  and inhibitors of NF- $\kappa$ B. Rossi *et al*<sup>[131]</sup> reported that PGA1 could block NF- $\kappa$ B activation by inhibiting I $\kappa$ B phosphorylation. In a subsequent study, they further identified that PGA1 could directly inhibit I $\kappa$ B kinase- $\beta$ <sup>[132]</sup>. PGA1 also increases the expression of I $\kappa$ B- $\alpha$ <sup>[133]</sup>. Similar inhibitory effects of PGJ2 and PGE1 on NF- $\kappa$ B were observed<sup>[134–136]</sup>. In addition, PGJ2 was found to interfere with DNA binding through covalently modifying the NF- $\kappa$ B p50 subunit<sup>[137]</sup>. Thus, cPG inhibit NF- $\kappa$ B by interfering in multiple sites in the NF- $\kappa$ B signaling pathway from I $\kappa$ B synthesis to DNA binding<sup>[138]</sup>. Studies have shown that cPG have neuroprotective effects under certain pathological conditions. Qin *et al*<sup>[139]</sup> first reported that PGA1 protected striatal neurons against NMDA receptor agonist quinolinic acid-induced apoptosis through inhibiting NF- $\kappa$ B activation. PGA1 also inhibited the mitochondrial toxin rotenone-induced death of

dopaminergic cell line SH-SY5Y cells<sup>[140]</sup>. In recent studies, PGA1 and PGJ2 were reported to reduce ischemic brain damage through inducing heat shock proteins, inhibiting NF- $\kappa$ B activation, and inflammation<sup>[141–143]</sup>. These studies revealed that the administration of PGA1 and PGJ2 2–3 h post-ischemia was still effective. Other studies have also reported that PGJ2 reduced ischemic myocardial infarction<sup>[144]</sup>.

Other drugs, such as estrogen, curcumin, and quercetin have been reported to inhibit I $\kappa$ B degradation<sup>[145–148]</sup>. All these compounds are reported to have neuroprotective effects under certain experimental conditions.

**NF- $\kappa$ B nuclear translocation and DNA binding inhibitors** NF- $\kappa$ B has to translocate into the nucleus in order to regulate gene expression. Drugs acting on NF- $\kappa$ B nuclear transport and DNA binding have received considerable attention. Lin *et al*<sup>[149]</sup> synthesized a membrane-permeable recombinant peptide NF- $\kappa$ B SN50. This peptide contains a signal peptide, which confers the cell membrane permeability of SN50, and a nuclear localization signal, which competes with NF- $\kappa$ B for nuclear entry. The peptide inhibits the nuclear translocation of NF- $\kappa$ B in cultured endothelial and monocytic cells stimulated with LPS or TNF- $\alpha$  in a concentration-dependent manner. This peptide has been widely used for dissecting cellular functions of NF- $\kappa$ B<sup>[150,151]</sup>. We, along with others, have successfully used SN50 for inhibiting NF- $\kappa$ B nuclear import *in vivo* and have found that it is very effective in blocking NF- $\kappa$ B nuclear entry and NF- $\kappa$ B-mediated target gene transcription in response to various stimuli<sup>[39,44,45,150,152–156]</sup>. An intrastriatal or nigral injection of SN50 substantially inhibited the nuclear translocation of NF- $\kappa$ B, inhibiting the expression of NF- $\kappa$ B target genes p53, c-Myc and cyclin D1, and attenuating excitotoxicity<sup>[39,44,45,153]</sup>. One study has successfully blocked cholecystokinin-octapeptide-induced pancreatitis with an intraperitoneal injection of SN50<sup>[155]</sup>. These studies suggest that nuclear import inhibitors represent an important class of NF- $\kappa$ B inhibitors.

Many studies have focused on NF- $\kappa$ B DNA decoys. The NF- $\kappa$ B DNA decoys are double-stranded DNA oligonucleotides (ODN) containing the NF- $\kappa$ B-binding motif. When delivered to cells, it competes with NF- $\kappa$ B for DNA binding sites, and thus inhibits NF- $\kappa$ B function. Some studies have demonstrated that ODN that are delivered locally or systemically are effective in blocking NF- $\kappa$ B transactivation activity<sup>[157–160]</sup>. Some *in vivo* therapeutic studies with ODN have generated promising results. The systemic administration of ODN suppressed NF- $\kappa$ B activity, and the expression of cytokines protected liver grafts against ischemia/reperfusion-induced injury in rats<sup>[161]</sup>. The animal studies showed that NF- $\kappa$ B decoys reduced lung vascular perme-

ability in septic mice, improved lung function<sup>[162]</sup>, and inhibited hepatic metastasis in the mice loaded with murine reticulosarcoma M5076<sup>[163]</sup>. The clinical usefulness of ODN has been tested in 2 patients with percutaneous coronary intervention. The initial results showed the suppression of restenosis with no observed adverse effect<sup>[164]</sup>. NF- $\kappa$ B decoys may be a potential therapeutic strategy for certain types of diseases<sup>[165,166]</sup>.

Drugs that directly inhibit NF- $\kappa$ B DNA binding have been found, but have not been well characterized. For example, it has been reported that the metal-chelating drug aurine tri-carboxylic acid inhibited NF- $\kappa$ B DNA interaction<sup>[167]</sup>.

### Neurological disorders may benefit from NF- $\kappa$ B inhibitors

**Ischemic brain injury** The distribution of NF- $\kappa$ B was investigated immunohistochemically in post-mortem brains of stroke patients. An enhanced immunoreactivity of NF- $\kappa$ B was observed in glial cells of infarcted areas, particularly in the penumbra or border zone between the ischemic and non-ischemic areas<sup>[168]</sup>. In animal studies, early activation of NF- $\kappa$ B has been found to precede DNA damage after ischemic attack<sup>[169,170]</sup>. It was reported that ischemia induced a TNF-like weak inducer of apoptosis (TWEAK) and its membrane receptor Fn14. TWEAK promotes neuronal cell death and activates NF- $\kappa$ B through the upstream kinase IKK<sup>[171]</sup>. The deletion of the neuronal IKK2 subunit or inhibition of IKK activity reduced the infarct size and neuronal cell loss. The role of NF- $\kappa$ B in neuronal death was further suggested, as several neuroprotective agents, such as antioxidant LY231617, PDTC, and PGA1, have been shown to inhibit NF- $\kappa$ B activation, reduce infarct volume, and improve behavior deficits<sup>[110,173,174]</sup>. The contribution of NF- $\kappa$ B activation to ischemic neuronal damage has also been assessed with either the expression of mutant I $\kappa$ B- $\alpha$  in neurons and glial, or NF- $\kappa$ B p50 knockout mice and transgenic mice. The results indicated that the neuronal expression of the NF- $\kappa$ B inhibitor reduced both the infarct size and cell death<sup>[174]</sup>. Mice lacking the p50 subunit of NF- $\kappa$ B develop significantly smaller infarcts after transient focal ischemia<sup>[173,175]</sup>. These studies may have established the rationale for use of NF- $\kappa$ B inhibitors in ischemic brain injury.

**PD** Recently, the role of the neuron-glia interaction and the inflammatory process in PD has been the focus of intense study by the research community. The increase in NF- $\kappa$ B has been found in the post-mortem brains of PD patients. In PD patients, the proportion of nigral dopaminergic neurons with immunoreactive NF- $\kappa$ B and interferon- $\gamma$  was sig-

nificantly increased in comparison with control patients<sup>[81,176]</sup>. A possible relationship between the nuclear localization of NF- $\kappa$ B in the mesencephalic neurons of PD patients and oxidative stress in such neurons has been shown *in vitro* with primary cultures of rat mesencephalon, where the translocation of NF- $\kappa$ B is preceded by a transient production of free radicals during apoptosis induced by the activation of the sphingomyelin-dependent signaling pathway with C2-ceramide. The data suggest that this oxidant-mediated apoptogenic transduction pathway may play a role in the mechanism of neuronal death in PD<sup>[176]</sup>. In animal models of PD, the inhibition of NF- $\kappa$ B achieves neuroprotection against the 6-OHDA- and MPTP-induced degeneration of dopaminergic neurons<sup>[44,45,112,177]</sup>, suggesting that NF- $\kappa$ B inhibitors could be beneficial in PD.

**AD** The distribution of NF- $\kappa$ B was investigated immunohistochemically in the post-mortem cases of AD. In the AD cases, increased staining for NF- $\kappa$ B p65 was seen in neurons and their processes, neurofibrillary tangles, and dystrophic neurites. The neuronal staining observed in AD was strongest in the hippocampal formation and entorhinal cortex<sup>[178]</sup>. Boissière *et al*<sup>[179]</sup> studied the cellular distribution of NF- $\kappa$ B in the nucleus basalis of Meynert of AD and control patients. The proportion of large cholinergic neurons with elevated nuclear immunostaining of NF- $\kappa$ B was significantly increased in AD, suggesting an association between NF- $\kappa$ B functions and the process of cholinergic degeneration in AD. In another report, NF- $\kappa$ B immunoreactivity was found in the neutrophil of diffuse A $\beta$  deposits. In addition, NF- $\kappa$ B immunoreactivity was found in the nuclei of neurons, but not in the nuclei of reactive astrocytes, in the vicinity of diffuse plaques<sup>[180]</sup>. The discovery of NF- $\kappa$ B activation in AD has been confirmed by other investigators<sup>[181-183]</sup>. Since inflammation is a prominent feature in AD, NF- $\kappa$ B may participate in the inflammatory response. A more direct connection of AD pathogenesis and NF- $\kappa$ B was observed as A $\beta$  activates NF- $\kappa$ B<sup>[184-186]</sup>. Although an early *in vitro* study found that the activation of NF- $\kappa$ B by a low dose of A $\beta$  may have neuroprotective effects<sup>[187]</sup>, other studies have been observed that NF- $\kappa$ B inhibitors inhibit the production of A $\beta$ <sup>[188]</sup>. In rat primary neurons and human post-mitotic neuronal cells, the A $\beta$  peptide induced dose-dependent neuronal death, the nuclear translocation of the p65 and p50 subunits, and an apoptotic profile of gene expression. The anti-inflammatory drug aspirin and the selective I $\kappa$ B kinase 2 inhibitor AS602868 completely inhibited p50/p65 nuclear translocation and neuronal damage<sup>[189]</sup>. The clinical trials with some NSAID did not generate encouraging outcomes, as noted by Valerio *et al*<sup>[190]</sup>, since not all NSAID can inhibit A $\beta$  production. Better

compounds with the ability to reduce A $\beta$  should be selected in future studies.

**Excitotoxicity** Excitotoxicity has been implicated in several neurodegenerative diseases. Kaltschmidt *et al*<sup>[191]</sup> reported that the ionotropic glutamate receptor agonist kainic acid (KA) activates NF- $\kappa$ B. Later studies defined a proapoptotic role of NF- $\kappa$ B activation and nuclear translocation mediated by AMPA/KA receptors<sup>[153,192]</sup>. Similarly, the stimulation of glutamate NMDA receptors robustly activates NF- $\kappa$ B through the degradation of I $\kappa$ B- $\alpha$ <sup>[139,152]</sup>. In other studies, the pharmacological upregulation of NF- $\kappa$ B increased glutamate-induced excitotoxicity, while the upregulation of CREB decreased excitotoxicity<sup>[193]</sup>. Grilli *et al*<sup>[118]</sup> reported a neuroprotective role of aspirin on the glutamate-induced death of hippocampal neurons, opening a new avenue for the study of excitotoxicity. Since then, several studies have reported that the inhibition of NF- $\kappa$ B has neuroprotective effects. In studies conducted by Casper *et al*<sup>[121]</sup>, neuroprotection against glutamate-mediated excitotoxicity was also found with ibuprofen. The inhibition of NF- $\kappa$ B with a herbal active component glycyrrhiza acid<sup>[193]</sup>, free radical scavenger OCT14117<sup>[194]</sup>, and glutamate metabotropic receptor agonists (2*S*, 1'*S*, 2'*S*)-(carboxy-cyclopropyl) glycine and amino-4-phosphonobutyric acid<sup>[195]</sup> was associated with a neuroprotective effect. These results suggest that NF- $\kappa$ B inhibitors could be suitable drugs for blocking excitotoxicity.

## Summary

The signaling pathway and the role of NF- $\kappa$ B have been studied for more than 2 decades. However, we still have limited knowledge on the role of NF- $\kappa$ B in CNS neurons and the molecular mechanisms underlying its actions. Many controversial findings need to be consolidated. In particular, its dual roles in neuronal death and survival and underlying molecular mechanisms need to be carefully evaluated in relation to human neurological diseases<sup>[197–200]</sup>. The involvement of NF- $\kappa$ B in human diseases certainly establishes it as a potential target for therapy. Many common synthetic (eg aspirin) and traditional remedies target, at least in part, the NF- $\kappa$ B signaling. Our knowledge of the molecular details of the NF- $\kappa$ B pathway will enable us to develop more specific NF- $\kappa$ B inhibitors to treat neurological diseases.

## References

- Baldwin AS Jr. The NF-kappa B and I kappa B proteins: new discoveries and insights. *Annu Rev Immunol* 1996; 14: 649–83.
- Shimada M, Satoh N, Yokosawa H. Involvement of Rel/NF-kappaB in regulation of ascidian notochord formation. *Dev Growth Differ* 2001; 43: 145–54.
- Weih F, Caamaño J. Regulation of secondary lymphoid organ development by the nuclear factor-kappaB signal transduction pathway. *Immunol Rev* 2003; 195: 91–105.
- Sen R, Baltimore D. Inducibility of k immunoglobulin enhancer-binding protein NF-kB by a post-translational mechanism. *Cell* 1986; 47: 921–8.
- O'Neill LA, Kaltschmidt C. NF-kappa B: a crucial transcription factor for glial and neuronal cell function. *Trends Neurosci* 1997; 20: 252–8.
- O'Mahony A, Raber J, Montano M, Foehr E, Han V, Lu SM, *et al*. NF-kappaB/Rel regulates inhibitory and excitatory neuronal function and synaptic plasticity. *Mol Cell Biol* 2006; 26: 7283–98.
- Meffert MK, Chang JM, Wiltgen BJ, Fanselow MS, Baltimore D. NF-kappa B functions in synaptic signaling and behavior. *Nat Neurosci* 2003; 6: 1072–8.
- Mémet S. NF-kappaB functions in the nervous system: from development to disease. *Biochem Pharmacol* 2006; 72: 1180–95.
- Xiao G, Rabson AB, Young W, Qing G, Qu Z. Alternative pathways of NF-kappaB activation: a double-edged sword in health and disease. *Cytokine Growth Factor Rev* 2006; 17: 281–93.
- Denk A, Wirth T, Baumann B. NF-kappaB transcription factors: critical regulators of hematopoiesis and neuronal survival. *Cytokine Growth Factor Rev* 2000; 11: 303–20.
- Grilli M, Memo M. Nuclear factor-kappaB/Rel proteins: a point of convergence of signalling pathways relevant in neuronal function and dysfunction. *Biochem Pharmacol* 1999; 57: 1–7.
- Mattson MP, Meffert MK. Roles for NF-kappaB in nerve cell survival, plasticity, and disease. *Cell Death Differ* 2006; 13: 852–60.
- Cornwell WD, Kirkpatrick RB. Cactus-independent nuclear translocation of Drosophila RELISH. *J Cell Biochem* 2001; 82: 22–37.
- Lätzer J, Papoian GA, Prentiss MC, Komives EA, Wolynes PG. Induced fit, folding, and recognition of the NF-kappaB-nuclear localization signals by IkappaBalpha and IkappaBbeta. *J Mol Biol* 2007; 367: 262–74.
- Heissmeyer V, Krappmann D, Hatada EN, Scheidereit C. Shared pathways of I $\kappa$ B $\alpha$  kinase-induced SCF TrCP-mediated ubiquitination and degradation for the NF- $\kappa$ B precursor p105 and I $\kappa$ B $\alpha$ . *Mol Cell Biol* 2001; 21: 1024–35.
- Dejardin E. The alternative NF-kappaB pathway from biochemistry to biology: pitfalls and promises for future drug development. *Biochem Pharmacol* 2006; 72: 1161–79.
- Baeuerle PA. I-kappa-B — NF-kappa-B structures: at the interface of inflammation control. *Cell* 1998; 95: 729–31.
- Lee SH, Hannink M. Characterization of the nuclear import and export functions of IkappaBalpha. *J Biol Chem* 2002; 26: 23358–66.
- Piette J, Piret B, Bonizzi G, Schoonbroodt S, Merville MP, Legrand-Poels S, *et al*. Multiple redox regulation in NF-kappaB transcription factor activation. *Biol Chem* 1997; 378: 1237–45.
- Huynh QK, Kishore N, Mathialagan S. Kinetic mechanisms of IkappaB-related kinases (IKK) inducible IKK and TBK-1 differ from IKK-1/IKK-2 heterodimer. *J Biol Chem* 2002; 277: 12550–8.



- 21 Gilmore TD. Introduction to NF- $\kappa$ B: players, pathways, perspectives. *Oncogene* 2006; 25: 6680–4.
- 22 Kaltschmidt B, Widera D, Kaltschmidt C. Signaling via NF-kappaB in the nervous system. *Biochim Biophys Acta* 2005; 1745: 287–99.
- 23 Wang CY, Mayo MW, Korneluk RG, Goeddel DV, Baldwin AS Jr. NF-kappaB antiapoptosis: induction of TRAF1 and TRAF2 and c-IAP1 and c-IAP2 to suppress caspase-8 activation. *Science* 1998; 281: 1680–3.
- 24 Maggirwar SB, Sarmiere PD, Dewhurst S, Freeman RS. Nerve growth factor-dependent activation of NF-kappaB contributes to survival of sympathetic neurons. *J Neurosci* 1998; 18: 10356–65.
- 25 Yu Z, Zhou D, Bruce-Keller AJ, Kindy MS, Mattson MP. Lack of the p50 subunit of nuclear factor-kappaB increases the vulnerability of hippocampal neurons to excitotoxic injury. *J Neurosci* 1999; 19: 8856–65.
- 26 Ishige K, Tanaka M, Arakawa M, Saito H, Ito Y. Distinct nuclear factor-kappaB/Rel proteins have opposing modulatory effects in glutamate-induced cell death in HT22 cells. *Neurochem Int* 2005; 47: 545–55.
- 27 Maehara K, Hasegawa T, Isobe KI. A NF-kappaB p65 subunit is indispensable for activating manganese superoxide dismutase gene transcription mediated by tumor necrosis factor-alpha. *J Cell Biochem* 2000; 77: 474–86.
- 28 Sugino N, Karube-Harada A, Sakata A, Takiguchi S, Kato H. Nuclear factor-kappa B is required for tumor necrosis factor-alpha-induced manganese superoxide dismutase expression in human endometrial stromal cells. *J Clin Endocrinol Metab* 2002; 87: 3845–50.
- 29 Borg J, London J. Copper/zinc superoxide dismutase overexpression promotes survival of cortical neurons exposed to neurotoxins *in vitro*. *J Neurosci Res* 2002; 70: 180–9.
- 30 Yune TY, Lee SM, Kim SJ, Park HK, Oh YJ, Kim YC, *et al*. Manganese superoxide dismutase induced by TNF-beta is regulated transcriptionally by NF-kappaB after spinal cord injury in rats. *J Neurotrauma* 2004; 21: 1778–94.
- 31 Mattson MP, Goodman Y, Luo H, Fu W, Furukawa K. Activation of NF-kappaB protects hippocampal neurons against oxidative stress-induced apoptosis: evidence for induction of manganese superoxide dismutase and suppression of peroxynitrite production and protein tyrosine nitration. *J Neurosci Res* 1997; 49: 681–97.
- 32 Saito A, Hayashi T, Okuno S, Ferrand-Drake M, Chan PH. Overexpression of copper/zinc superoxide dismutase in transgenic mice protects against neuronal cell death after transient focal ischemia by blocking activation of the Bad cell death signaling pathway. *J Neurosci* 2003; 23: 1710–8.
- 33 Sompol P, Xu Y, Ittarat W, Daosukho C, St Clair D. NF-kappaB-associated MnSOD induction protects against beta-amyloid-induced neuronal apoptosis. *J Mol Neurosci* 2006; 29: 279–88.
- 34 Glasgow JN, Wood T, Perez-Polo JR. Identification and characterization of nuclear factor kappaB binding sites in the murine bcl-x promoter. *J Neurochem* 2000; 75: 1377–89.
- 35 Bui NT, Livolsi A, Peyron JF, Prehn JH. Activation of nuclear factor kappaB and Bcl-x survival gene expression by nerve growth factor requires tyrosine phosphorylation of IkappaBalpha. *J Cell Biol* 2001; 152: 753–64.
- 36 Tamatani M, Che YH, Matsuzaki H, Ogawa S, Okado H, Miyake S, *et al*. Tumor necrosis factor induces Bcl-2 and Bcl-x expression through NFkappaB activation in primary hippocampal neurons. *J Biol Chem* 1999; 274: 8531–8.
- 37 Tamatani M, Mitsuda N, Matsuzaki H, Okado H, Miyake S, Vitek MP, *et al*. A pathway of neuronal apoptosis induced by hypoxia/reoxygenation: roles of nuclear factor-kappaB and Bcl-2. *J Neurochem* 2000; 75: 683–93.
- 38 Wang CY, Guttridge DC, Mayo MW, Baldwin AS Jr. NF-kappaB induces expression of the Bcl-2 homologue A1/Bfl-1 to preferentially suppress chemotherapy-induced apoptosis. *Mol Cell Biol* 1999; 19: 5923–9.
- 39 Qin ZH, Chen RW, Wang Y, Nakai M, Chuang DM, Chase TN. Nuclear factor kappaB nuclear translocation upregulates c-Myc and p53 expression during NMDA receptor-mediated apoptosis in rat striatum. *J Neurosci* 1999; 19: 4023–33.
- 40 Grilli M, Memo M. Possible role of NF-kappaB and p53 in the glutamate-induced pro-apoptotic neuronal pathway. *Cell Death Differ* 1999; 6: 22–7.
- 41 Uberti D, Grilli M, Memo M. Contribution of NF-kappaB and p53 in the glutamate-induced apoptosis. *Int J Dev Neurosci* 2000; 18: 447–54.
- 42 Culmsee C, Zhu X, Yu QS, Chan SL, Camandola S, Guo Z, *et al*. A synthetic inhibitor of p53 protects neurons against death induced by ischemic and excitotoxic insults, and amyloid beta-peptide. *J Neurochem* 2001; 77: 220–8.
- 43 Aleyasin H, Cregan SP, Iyirhario G, O'Hare MJ, Callaghan SM, Slack RS, *et al*. Nuclear factor-(kappa)B modulates the p53 response in neurons exposed to DNA damage. *J Neurosci* 2004; 24: 2963–73.
- 44 Liang ZQ, Wang X, Li LY, Wang Y, Chen RW, Chuang DM, *et al*. Nuclear factor-kappaB-dependent cyclin D1 induction and DNA replication associated with N-methyl-D-aspartate receptor-mediated apoptosis in rat striatum. *J Neurosci Res* 2007; 85: 1295–309.
- 45 Liang ZQ, Li YL, Zhao XL, Han R, Wang XX, Wang Y, *et al*. NF-kappaB contributes to 6-hydroxydopamine-induced apoptosis of nigral dopaminergic neurons through p53. *Brain Res* 2007; 1145: 190–203.
- 46 Fujioka S, Schmidt C, Scwabas GM, Li Z, Pelicano H, Peng B, *et al*. Stabilization of p53 is a novel mechanism for proapoptotic function of NF-kappaB. *J Biol Chem* 2004; 279: 27549–59.
- 47 Joyce D, Albanese C, Steer J, Fu M, Bouzahzah B, Pestell RG. NF-kappaB and cell-cycle regulation: the cyclin connection. *Cytokine Growth Factor Rev* 2001; 12: 73–90.
- 48 Guttridge DC, Albanese C, Reuther JY, Pestell RG, Baldwin AS Jr. NF-kappaB controls cell growth and differentiation through transcriptional regulation of cyclin D1. *Mol Cell Biol* 1999; 19: 5785–99.
- 49 Hinz M, Krappmann D, Eichten A, Heder A, Scheidereit C, Strauss M. NF-kappaB function in growth control: regulation of cyclin D1 expression and G0/G1-to-S-phase transition. *Mol Cell Biol* 1999; 19: 2690–8.
- 50 Gill JS, Windebank AJ. Ceramide initiates NFkappaB-mediated caspase activation in neuronal apoptosis. *Neurobiol Dis* 2000; 7: 448–61.
- 51 Ji L, Arcinas M, Boxer LM. NF-kappa B sites function as positive regulators of expression of the translocated c-myc allele in Burkitt's lymphoma. *Mol Cell Biol* 1994; 14: 7967–74.

- 52 Tarabin V, Schwaninger M. The role of NF-kappaB in 6-hydroxydopamine- and TNFalpha-induced apoptosis of PC12 cells. *Naunyn Schmiedebergs Arch Pharmacol* 2004; 369: 563–9.
- 53 Timsit S, Rivera S, Ouaghi P, Guisard F, Tremblay E, Ben-Ari Y, *et al*. Increased cyclin D1 in vulnerable neurons in the hippocampus after ischaemia and epilepsy: a modulator of *in vivo* programmed cell death? *Eur J Neurosci* 1999; 11: 263–78.
- 54 Di Giovanni S, Knoblach SM, Brandoli C, Aden SA, Hoffman EP, Faden AI. Gene profiling in spinal cord injury shows role of cell cycle in neuronal death. *Ann Neurol* 2003; 53: 454–68.
- 55 Wen Y, Yang S, Liu R, Simpkins JW. Cell-cycle regulators are involved in transient cerebral ischemia induced neuronal apoptosis in female rats. *FEBS Lett* 2005; 579: 4591–9.
- 56 Rao HV, Thirumangalakudi L, Desmond P, Grammas P. Cyclin D1, cdk4, and Bim are involved in thrombin-induced apoptosis in cultured cortical neurons. *J Neurochem* 2007; 101: 498–505.
- 57 Appert-Collin A, Hugel B, Levy R, Niederhoffer N, Coupin G, Lombard Y, *et al*. Cyclin dependent kinase inhibitors prevent apoptosis of postmitotic mouse motoneurons. *Life Sci* 2006; 79: 484–90.
- 58 Ino H, Chiba T. Cyclin-dependent kinase 4 and cyclin D1 are required for excitotoxin-induced neuronal cell death *in vivo*. *J Neurosci* 2001; 21: 6086–94.
- 59 Glasgow JN, Qiu J, Rassin D, Grafe M, Wood T, Perez-Pol JR. Transcriptional regulation of the BCL-X gene by NF-kappaB is an element of hypoxic responses in the rat brain. *Neurochem Res* 2001; 26: 647–59.
- 60 Dixon EP, Stephenson DT, Clemens JA, Little SP. Bcl-Xshort is elevated following severe global ischemia in rat brains. *Brain Res* 1997; 776: 222–9.
- 61 Matsushita H, Morishita R, Nata T, Aoki M, Nakagami H, Taniyama Y, *et al*. Hypoxia-induced endothelial apoptosis through nuclear factor-kappaB (NF-kappaB)-mediated bcl-2 suppression: *in vivo* evidence of the importance of NF-kappaB in endothelial cell regulation. *Circ Res* 2000; 86: 974–81.
- 62 McInnis J, Wang C, Anastasio N, Hultman M, Ye Y, Salvemini D, *et al*. The role of superoxide and nuclear factor-kappaB signaling in *N*-methyl-*D*-aspartate-induced necrosis and apoptosis. *J Pharmacol Exp Ther* 2002; 301: 478–87.
- 63 Pizzi M, Sarnico I, Boroni F, Benetti A, Benarese M, Spano PF. Inhibition of IkappaBalpha phosphorylation prevents glutamate-induced NF-kappaB activation and neuronal cell death. *Acta Neurochir Suppl* 2005; 93: 59–63.
- 64 Shou Y, Li N, Li L, Borowitz JL, Isom GE. NF-kappaB-mediated up-regulation of Bcl-X(S) and Bax contributes to cytochrome c release in cyanide-induced apoptosis. *J Neurochem* 2002; 81: 842–52.
- 65 Tao Y, Yan D, Yang Q, Zeng R, Wang Y. Low K<sup>+</sup> promotes NF-kappaB/DNA binding in neuronal apoptosis induced by K<sup>+</sup> loss. *Mol Cell Biol* 2006; 26: 1038–50.
- 66 Xie QW, Kashiwabara Y, Nathan C. Role of transcription factor NF-kappa B/Rel in induction of nitric oxide synthase. *J Biol Chem* 1994; 269: 4705–8.
- 67 Heneka MT, Löschmann PA, Gleichmann M, Weller M, Schulz JB, Wüllner U, *et al*. Induction of nitric oxide synthase and nitric oxide-mediated apoptosis in neuronal PC12 cells after stimulation with tumor necrosis factor-alpha/lipopolysaccharide. *J Neurochem* 1998; 71: 88–94.
- 68 Binder C, Schulz M, Hiddemann W, Oellerich M. Induction of inducible nitric oxide synthase is an essential part of tumor necrosis factor-alpha-induced apoptosis in MCF-7 and other epithelial tumor cells. *Lab Invest* 1999; 79: 1703–12.
- 69 Zhao X, Zhang Y, Strong R, Zhang J, Grotta JC, Aronowski J. Distinct patterns of intracerebral hemorrhage-induced alterations in NF-kappaB subunit, iNOS, and COX-2 expression. *J Neurochem* 2007; 101: 652–63.
- 70 Miscusi M, Ebner F, Ceccariglia S, Menegazzi M, Mariotto S, Berra L, *et al*. Early nuclear factor-kappaB activation and inducible nitric oxide synthase expression in injured spinal cord neurons correlating with a diffuse reduction of constitutive nitric oxide synthase activity. *J Neurosurg Spine* 2006; 4: 485–93.
- 71 Hecker M, Preiss C, Klemm P, Busse R. Inhibition by antioxidants of nitric oxide synthase expression in murine macrophages: role of nuclear factor kappa B and interferon regulatory factor 1. *Br J Pharmacol* 1996; 118: 2178–84.
- 72 Liu X, Buffington JA, Tjalkens RB. NF-kappaB-dependent production of nitric oxide by astrocytes mediates apoptosis in differentiated PC12 neurons following exposure to manganese and cytokines. *Brain Res Mol Brain Res* 2005; 141: 39–47.
- 73 Pokharel YR, Liu QH, Oh JW, Woo ER, Kang KW. 4-Hydroxykobusin inhibits the induction of nitric oxide synthase by inhibiting NF-kappaB and AP-1 activation. *Biol Pharm Bull* 2007; 30: 1097–101.
- 74 Chung HY, Kim HJ, Kim KW, Choi JS, Yu BP. Molecular inflammation hypothesis of aging based on the anti-aging mechanism of calorie restriction. *Microsc Res Tech* 2002; 59: 264–72.
- 75 Reuter U, Chiarugi A, Bolay H, Moskowitz MA. Nuclear factor-kappaB as a molecular target for migraine therapy. *Ann Neurol* 2002; 51: 507–16.
- 76 Tegeder I, Niederberger E, Schmidt R, Kunz S, Gühring H, Ritzeler O, *et al*. Specific Inhibition of IkappaB kinase reduces hyperalgesia in inflammatory and neuropathic pain models in rats. *J Neurosci* 2004; 24: 1637–45.
- 77 Herrmann O, Baumann B, de Lorenzi R, Muhammad S, Zhang W, Kleesiek J, *et al*. IKK mediates ischemia-induced neuronal death. *Nat Med* 2005; 11: 1322–9.
- 78 Hang CH, Shi JX, Li JS, Wu W, Yin HX. Concomitant upregulation of nuclear factor-kappaB activity, proinflammatory cytokines and ICAM-1 in the injured brain after cortical contusion trauma in a rat model. *Neurol India* 2005; 53: 312–7.
- 79 Brambilla R, Bracchi-Ricard V, Hu WH, Frydel B, Bramwell A, Karmally S, *et al*. Inhibition of astroglial nuclear factor kappaB reduces inflammation and improves functional recovery after spinal cord injury. *J Exp Med* 2005; 202: 145–56.
- 80 Soós J, Engelhardt JI, Siklós L, Havas L, Majtényi K. The expression of PARP, NF-kappa B and parvalbumin is increased in Parkinson disease. *Neuroreport* 2004; 15: 1715–8.
- 81 Mogi M, Kondo T, Mizuno Y, Nagatsu T. p53 protein, interferon-gamma, and NF-kappaB levels are elevated in the parkinsonian brain. *Neurosci Lett* 2007; 414: 94–7. Epub 2006 Dec 29.
- 82 Satoh J, Illes Z, Peterfalvi A, Tabunoki H, Rozsa C, Yamamura T. Aberrant transcriptional regulatory network in T cells of multiple sclerosis. *Neurosci Lett* 2007; 422: 30–3.
- 83 Mattson MP, Camandola S. NF-kappaB in neuronal plasticity and neurodegenerative disorders. *J Clin Invest* 2001; 107: 247–

- 54.
- 84 Collister KA, Albeni BC. Potential therapeutic targets in the NF-kappaB pathway for Alzheimer's disease. *Drug News Perspect* 2005; 18: 623-9.
- 85 Xu Z, Chen S, Li X, Luo G, Li L, Le W. Neuroprotective effects of (-)-epigallocatechin-3-gallate in a transgenic mouse model of amyotrophic lateral sclerosis. *Neurochem Res* 2006; 31: 1263-9.
- 86 Khoshnan A, Ko J, Watkin EE, Paige LA, Reinhart PH, Patterson PH. Activation of the IkappaB kinase complex and nuclear factor-kappaB contributes to mutant huntingtin neurotoxicity. *J Neurosci* 2004; 24: 7999-8008.
- 87 Hayashi S, Yamamoto M, Ueno Y, Ikeda K, Ohshima K, Soma G, *et al.* Expression of nuclear factor-kappa B, tumor necrosis factor receptor type 1, and c-Myc in human astrocytomas. *Neurol Med Chir (Tokyo)* 2001; 41: 187-95.
- 88 Bian X, Oipari AW Jr, Ratanaproeksa AB, Boitano AE, Lucas PC, Castle VP. Constitutively active NFkappa B is required for the survival of S-type neuroblastoma. *J Biol Chem* 2002; 277: 42144-50.
- 89 Garkavtsev I, Kozin SV, Chernova O, Xu L, Winkler F, Brown E, *et al.* The candidate tumour suppressor protein ING4 regulates brain tumour growth and angiogenesis. *Nature* 2004; 428: 328-32.
- 90 Brown RE, Tan D, Taylor JS, Miller M, Prichard JW, Kott MM. Morphoproteomic confirmation of constitutively activated mTOR, ERK, and NF-kappaB pathways in high risk neuroblastoma, with cell cycle and protein analyte correlates. *Ann Clin Lab Sci* 2007; 37: 141-7.
- 91 Raychaudhuri B, Han Y, Lu T, Vogelbaum MA. Aberrant constitutive activation of nuclear factor kappaB in glioblastoma multiforme drives invasive phenotype. *J Neurooncol* 2007; [Epub ahead of print]
- 92 Camandola S, Mattson MP. NF-kappa B as a therapeutic target in neurodegenerative diseases. *Expert Opin Ther Targets* 2007; 11: 123-32.
- 93 Zhang CG, Welin D, Novikov L, Kellerth JO, Wiberg M, Hart AM. Motorneuron protection by N-acetyl-cysteine after ventral root avulsion and ventral rhizotomy. *Br J Plast Surg* 2005; 58: 765-73.
- 94 Arakawa M, Ishimura A, Arai Y, Kawabe K, Suzuki S, Ishige K, *et al.* N-Acetylcysteine and ebselen but not nifedipine protected cerebellar granule neurons against 4-hydroxynonenal-induced neuronal death. *Neurosci Res* 2007; 57: 220-9.
- 95 Zafarullah M, Li WQ, Sylvester J, Ahmad M. Molecular mechanisms of N-acetylcysteine actions. *Cell Mol Life Sci* 2003; 60: 6-20.
- 96 Mihm S, Ennen J, Pessara U, Kurth R, Dröge W. Inhibition of HIV-1 replication and NF-kappa B activity by cysteine and cysteine derivatives. *AIDS* 1991; 5: 497-503.
- 97 Schreck R, Rieber P, Baeuerle PA. Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kappa B transcription factor and HIV-1. *EMBO J* 1991; 10: 2247-58.
- 98 Shen WH, Zhang CY, Zhang GY. Antioxidants attenuate reperfusion injury after global brain ischemia through inhibiting nuclear factor-kappa B activity in rats. *Acta Pharmacol Sin* 2003; 24: 1125-30.
- 99 Muñoz AM, Rey P, Soto-Otero R, Guerra MJ, Labandeira-Garcia JL. Systemic administration of N-acetylcysteine protects dopaminergic neurons against 6-hydroxydopamine-induced degeneration. *J Neurosci Res* 2004; 76: 551-62.
- 100 Schreck R, Meier B, Männel DN, Dröge W, Baeuerle PA. Dithiocarbamates as potent inhibitors of nuclear factor kappa B activation in intact cells. *J Exp Med* 1992; 175: 1181-94.
- 101 Schenk H, Klein M, Erdbrügger W, Dröge W, Schulze-Osthoff K. Distinct effects of thioredoxin and antioxidants on the activation of transcription factors NF-kappa B and AP-1. *Proc Natl Acad Sci USA* 1994; 91: 1672-6.
- 102 Ziegler-Heitbrock HW, Sternsdorf T, Liese J, Belohradsky B, Weber C, Wedel A, *et al.* Pyrrolidine dithiocarbamate inhibits NF-kappa B mobilization and TNF production in human monocytes. *J Immunol* 1993; 151: 6986-93.
- 103 Cuzzocrea S, Chatterjee PK, Mazzon E, Dugo L, Serrano I, Britti D, *et al.* Pyrrolidine dithiocarbamate attenuates the development of acute and chronic inflammation. *Br J Pharmacol* 2002; 135: 496-510.
- 104 Nurmi A, Goldsteins G, Närväinen J, Pihlaja R, Ahtoniemi T, Gröhn O, *et al.* Antioxidant pyrrolidine dithiocarbamate activates Akt-GSK signaling and is neuroprotective in neonatal hypoxia-ischemia. *Free Radic Biol Med* 2006; 40: 1776-84.
- 105 Kabay B, Teke Z, Aytakin FO, Yenisey C, Bir F, Sacar M, *et al.* Pyrrolidine dithiocarbamate reduces lung injury caused by mesenteric ischemia/reperfusion in a rat model. *World J Surg* 2007; 31: 1707-15.
- 106 de Jesus Soares T, Costa RS, Balbi AP, Coimbra TM. Inhibition of nuclear factor-kappa B activation reduces glycerol-induced renal injury. *J Nephrol* 2006; 19: 439-48.
- 107 Demirbilek S, Akin M, Gürünlöödü K, Aydın NE, Emre MH, Tas E, *et al.* The NF-kappaB inhibitors attenuate hepatic injury in bile duct ligated rats. *Pediatr Surg Int* 2006; 22: 655-63.
- 108 Li H, Gu H, Sun B. Protective effects of pyrrolidine dithiocarbamate on myocardium apoptosis induced by adriamycin in rats. *Int J Cardiol* 2007; 114: 159-65.
- 109 Crack PJ, Taylor JM, Ali U, Mansell A, Hertzog PJ. Potential contribution of NF-kappaB in neuronal cell death in the glutathione peroxidase-1 knockout mouse in response to ischemia-reperfusion injury. *Stroke* 2006; 37: 1533-8.
- 110 Nurmi A, Vartiainen N, Pihlaja R, Goldsteins G, Yrjänheikki J, Koistinaho J. Pyrrolidine dithiocarbamate inhibits translocation of nuclear factor kappa-B in neurons and protects against brain ischaemia with a wide therapeutic time window. *J Neurochem* 2004b; 91: 755-65.
- 111 Chen JW, Chen YH, Lin FY, Chen YL, Lin SJ. *Ginkgo biloba* extract inhibits tumor necrosis factor-alpha-induced reactive oxygen species generation, transcription factor activation, and cell adhesion molecule expression in human aortic endothelial cells. *Arterioscler Thromb Vasc Biol* 2003; 23: 1559-66.
- 112 Levites Y, Youdim MB, Maor G, Mandel S. Attenuation of 6-hydroxydopamine (6-OHDA)-induced nuclear factor-kappaB (NF-kappaB) activation and cell death by tea extracts in neuronal cultures. *Biochem Pharmacol* 2002; 63: 21-9.
- 113 Nie G, Cao Y, Zhao B. Protective effects of green tea polyphenols and their major component, (-)-epigallocatechin-3-gallate (EGCG), on 6-hydroxydopamine-induced apoptosis in PC12 cells. *Redox Rep* 2002; 7: 171-7.
- 114 Hong JT, Ryu SR, Kim HJ, Lee JK, Lee SH, Yun YP, *et al.*

- Protective effect of green tea extract on ischemia/reperfusion-induced brain injury in Mongolian gerbils. *Brain Res* 2001; 888: 11–8.
- 115 Lee SY, Lee JW, Lee H, Yoo HS, Yun YP, Oh KW, *et al*. Inhibitory effect of green tea extract on beta-amyloid-induced PC12 cell death by inhibition of the activation of NF-kappaB and ERK/p38 MAP kinase pathway through antioxidant mechanisms. *Brain Res Mol Brain Res* 2005; 140: 45–54.
- 116 Kopp E, Ghosh S. Inhibition of NF-kappa B by sodium salicylate and aspirin. *Science* 1994; 265: 956–9.
- 117 Yin MJ, Yamamoto Y, Gaynor RB. The anti-inflammatory agents aspirin and salicylate inhibit the activity of I(kappa)B kinase-beta. *Nature* 1998; 396: 77–80.
- 118 Grilli M, Pizzi M, Memo M, Spano P. Neuroprotection by aspirin and sodium salicylate through blockade of NF-kappaB activation. *Science* 1996; 274: 1383–5.
- 119 Crisanti P, Leon A, Lim DM, Omri B. Aspirin prevention of NMDA-induced neuronal death by direct protein kinase Czeta inhibition. *J Neurochem* 2005; 93: 1587–93.
- 120 Maharaj H, Maharaj DS, Daya S. Acetylsalicylic acid and acetaminophen protect against oxidative neurotoxicity. *Metab Brain Dis* 2006; 21: 189–99.
- 121 Casper D, Yaparalvi U, Rempel N, Werner P. Ibuprofen protects dopaminergic neurons against glutamate toxicity *in vitro*. *Neurosci Lett* 2000; 289: 201–4.
- 122 Carrasco E, Werner P. Selective destruction of dopaminergic neurons by low concentrations of 6-OHDA and MPP+: protection by acetylsalicylic acid aspirin. *Parkinsonism Relat Disord* 2002; 8: 407–11.
- 123 Esposito E, Di Matteo V, Benigno A, Pierucci M, Crescimanno G, Di Giovanni G. Non-steroidal anti-inflammatory drugs in Parkinson's disease. *Exp Neurol* 2007; 205: 295–312.
- 124 Thomas T, Nadackal TG, Thomas K. Aspirin and non-steroidal anti-inflammatory drugs inhibit amyloid-beta aggregation. *Neuroreport* 2001; 12: 3263–7.
- 125 Bisaglia M, Venezia V, Piccioli P, Stanzione S, Porcile C, Russo C, *et al*. Acetaminophen protects hippocampal neurons and PC12 cultures from amyloid beta-peptides induced oxidative stress and reduces NF-kappaB activation. *Neurochem Int* 2002; 41: 43–54.
- 126 Hirohata M, Ono K, Naiki H, Yamada M. Non-steroidal anti-inflammatory drugs have anti-amyloidogenic effects for Alzheimer's beta-amyloid fibrils *in vitro*. *Neuropharmacology* 2005; 49: 1088–99.
- 127 Tortosa E, Avila J, Pérez M. Acetylsalicylic acid decreases tau phosphorylation at serine 422. *Neurosci Lett* 2006; 396: 77–80.
- 128 Natarajan K, Singh S, Burke TR Jr, Grunberger D, Aggarwal BB. Caffeic acid phenethyl ester is a potent and specific inhibitor of activation of nuclear transcription factor NF-kappa B. *Proc Natl Acad Sci USA* 1996; 93: 9090–5.
- 129 Amodio R, De Ruvo C, Sacchetti A, Di Santo A, Martelli N, Di Matteo V, *et al*. Caffeic acid phenethyl ester blocks apoptosis induced by low potassium in cerebellar granule cells. *Int J Dev Neurosci* 2003; 21: 379–89.
- 130 Jüttler E, Potrovita I, Tarabin V, Prinz S, Dong-Si T, Fink G, *et al*. The cannabinoid dexamabinol is an inhibitor of the nuclear factor-kappa B (NF-kappa B). *Neuropharmacology* 2004; 47: 580–92.
- 131 Rossi A, Elia G, Santoro MG. Inhibition of nuclear factor kappa B by prostaglandin A1: an effect associated with heat shock transcription factor activation. *Proc Natl Acad Sci USA* 1997; 94: 746–50.
- 132 Rossi A, Kapahi P, Natoli G, Takahashi T, Chen Y, Karin M, *et al*. Anti-inflammatory cyclopentenone prostaglandins are direct inhibitors of I(kappa)B kinase. *Nature* 2000; 403: 103–8.
- 133 Thomas SC, Ryan MA, Shanley TP, Wong HR. Induction of the stress response with prostaglandin A1 increases I-kappaBalpha gene expression. *FASEB J* 1998; 12: 1371–8.
- 134 Castrillo A, Díaz-Guerra MJ, Hortelano S, Martín-Sanz P, Boscá L. Inhibition of I(kappa)B kinase and I(kappa)B phosphorylation by 15-deoxy-delta(12,14)-prostaglandin J(2) in activated murine macrophages. *Mol Cell Biol* 2000; 20: 1692–8.
- 135 Rovin BH, Lu L, Cosio A. Cyclopentenone prostaglandins inhibit cytokine-induced NF-kappaB activation and chemokine production by human mesangial cells. *J Am Soc Nephrol* 2001; 12: 1659–67.
- 136 Siendones E, Fouad D, Díaz-Guerra MJ, de la Mata M, Boscá L, Muntané J. PGE1-induced NO reduces apoptosis by D-galactosamine through attenuation of NF-kappaB and NOS-2 expression in rat hepatocytes. *Hepatology* 2004; 40: 1295–303.
- 137 Cernuda-Morollón E, Pineda-Molina E, Cañada FJ, Pérez-Sala D. 15-Deoxy-delta 12,14-prostaglandin J2 inhibition of NF-kappaB-DNA binding through covalent modification of the p50 subunit. *J Biol Chem* 2001; 276: 35 530–6.
- 138 Straus DS, Pascual G, Li M, Welch JS, Ricote M, Hsiang CH, *et al*. 15-deoxy-delta 12,14-prostaglandin J2 inhibits multiple steps in the NF-kappa B signaling pathway. *Proc Natl Acad Sci USA* 2000; 97: 4844–9.
- 139 Qin ZH, Wang Y, Chen RW, Wang X, Ren M, Chuang DM, *et al*. Prostaglandin A(1) protects striatal neurons against excitotoxic injury in rat striatum. *J Pharmacol Exp Ther* 2001; 297: 78–87.
- 140 Wang X, Qin ZH, Leng Y, Wang Y, Jin X, Chase TN, *et al*. Prostaglandin A1 inhibits rotenone-induced apoptosis in SH-SY5Y cells. *J Neurochem* 2002; 83: 1094–102.
- 141 Zhao X, Zhang Y, Strong R, Grotta JC, Aronowski J. 15d-Prostaglandin J2 activates peroxisome proliferator-activated receptor-gamma, promotes expression of catalase, and reduces inflammation, behavioral dysfunction, and neuronal loss after intracerebral hemorrhage in rats. *J Cereb Blood Flow Metab* 2006; 26: 811–20.
- 142 Pereira MP, Hurtado O, Cárdenas A, Boscá L, Castillo J, Dávalos A, *et al*. Rosiglitazone and 15-deoxy-delta 12,14-prostaglandin J2 cause potent neuroprotection after experimental stroke through noncompletely overlapping mechanisms. *J Cereb Blood Flow Metab* 2006; 26: 218–29.
- 143 Xu XH, Zhang HL, Han R, Gu ZL, Qin ZH. Enhancement of neuroprotection and heat shock protein induction by combined prostaglandin A1 and lithium in rodent models of focal ischemia. *Brain Res* 2006; 1102: 154–62.
- 144 Wayman NS, Hattori Y, McDonald MC, Mota-Filipe H, Cuzzocrea S, Pisano B, *et al*. Ligands of the peroxisome proliferator-activated receptors (PPAR-gamma and PPAR-alpha) reduce myocardial infarct size. *FASEB J* 2002; 16: 1027–40.
- 145 Sun WH, Keller ET, Stebler BS, Ershler WB. Estrogen inhibits

- phorbol ester-induced IkappaBalpha transcription and protein degradation. *Biochem Biophys Res Commun* 1998; 244: 691–5.
- 146 Peet GW, Li J. IkappaB kinases alpha and beta show a random sequential kinetic mechanism and are inhibited by staurosporine and quercetin. *J Biol Chem* 1999; 274: 32 655–61.
- 147 Kalaitzidis D, Gilmore TD. Transcription factor cross-talk: the estrogen receptor and NF-kappaB. *Trends Endocrinol Metab* 2005; 16: 46–52.
- 148 Dikshit P, Goswami A, Mishra A, Chatterjee M, Jana NR. Curcumin induces stress response, neurite outgrowth and prevents NF-kappaB activation by inhibiting the proteasome function. *Neurotox Res* 2006; 9: 29–37.
- 149 Lin YZ, Yao SY, Veach RA, Torgerson TR, Hawiger J. Inhibition of nuclear translocation of transcription factor NF-kappa B by a synthetic peptide containing a cell membrane-permeable motif and nuclear localization sequence. *J Biol Chem* 1995; 270: 14 255–8.
- 150 Liu D, Liu XY, Robinson D, Burnett C, Jackson C, Seele L, *et al*. Suppression of staphylococcal enterotoxin B-induced toxicity by a nuclear Import inhibitor. *J Biol Chem* 2004; 279: 19 239–46.
- 151 Chen YL, Law PY, Loh HH. Sustained activation of phosphatidylinositol 3-kinase/Akt/nuclear factor kappaB signaling mediates G protein-coupled delta-opioid receptor gene expression. *J Biol Chem* 2006; 281: 3067–74.
- 152 Qin ZH, Wang Y, Nakai M, Chase TN. Nuclear factor-kappa B contributes to excitotoxin-induced apoptosis in rat striatum. *Mol Pharmacol* 1998; 53: 33–42.
- 153 Nakai M, Qin ZH, Chen JF, Wang Y, Chase TN. Kainic acid-induced apoptosis in rat striatum is associated with nuclear factor-kappaB activation. *J Neurochem* 2000; 74: 647–58.
- 154 Shou Y, Gunasekar PG, Borowitz JL, Isom GE. Cyanide-induced apoptosis involves oxidative-stress-activated NF-kappaB in cortical neurons. *Toxicol Appl Pharmacol* 2000; 164: 196–205.
- 155 Letoha T, Somlai C, Takacs T, Szabolcs A, Jarmay K, Rakonczay Z Jr, *et al*. A nuclear import inhibitory peptide ameliorates the severity of cholecystokinin-induced acute pancreatitis. *World J Gastroenterol* 2005; 11: 990–9.
- 156 Saika S, Miyamoto T, Yamanaka O, Kato T, Ohnishi Y, Flanders KC, *et al*. Therapeutic effect of topical administration of SN50, an inhibitor of nuclear factor-kappaB, in treatment of corneal alkali burns in mice. *Am J Pathol*; 166: 1393–403.
- 157 Ono S, Date I, Onoda K, Shiota T, Ohmoto T, Ninomiya Y, *et al*. Decoy administration of NF-kappaB into the subarachnoid space for cerebral angiopathy. *Hum Gene Ther* 1998; 9: 1003–11.
- 158 De Vry CG, Prasad S, Komuves L, Lorenzana C, Parham C, Le T, *et al*. Non-viral delivery of nuclear factor-kappaB decoy ameliorates murine inflammatory bowel disease and restores tissue homeostasis. *Gut* 2007; 56: 524–33.
- 159 Laguillier C, Hbib AT, Baran-Marszak F, Metelev V, Cao A, Cymbalista F, *et al*. Cell death in NF-kappaB-dependent tumour cell lines as a result of NF-kappaB trapping by linker-modified hairpin decoy oligonucleotide. *FEBS Lett* 2007; 581: 1143–50.
- 160 Takeuchi K, Itoh H, Yonemitsu Y, Matsumoto T, Kume M, Komori K, *et al*. *In vivo* reduction of the nuclear factor-kappaB activity using synthetic *cis*-element decoy oligonucleotides suppresses intimal hyperplasia in the injured carotid arteries in rabbits. *Surg Today* 2007; 37: 575–83.
- 161 Xu MQ, Shuai XR, Yan ML, Zhang MM, Yan LN. Nuclear factor-kappaB decoy oligodeoxynucleotides attenuates ischemia/reperfusion injury in rat liver graft. *World J Gastroenterol* 2005; 11: 6960–7.
- 162 Matsuda N, Hattori Y, Takahashi Y, Nishihira J, Jesmin S, Kobayashi M, *et al*. Therapeutic effect of *in vivo* transfection of transcription factor decoy to NF-kappaB on septic lung in mice. *Am J Physiol Lung Cell Mol Physiol* 2004; 287: L1248–55.
- 163 Kawamura I, Morishita R, Tsujimoto S, Manda T, Tomoi M, Tomita N, *et al*. Intravenous injection of oligodeoxynucleotides to the NF-kappaB binding site inhibits hepatic metastasis of M5076 reticulosarcoma in mice. *Gene Ther* 2001; 8: 905–12.
- 164 Jun-Ichi S, Hiroshi I, Ryo G, Ryuichi M, Kensuke E, Mitsuaki I. Initial clinical cases of the use of a NF-kappaB decoy at the site of coronary stenting for the prevention of restenosis. *Circ J* 2004; 68: 270–1.
- 165 Isomura I, Morita A. Regulation of NF-kappaB signaling by decoy oligodeoxynucleotides. *Microbiol Immunol* 2006; 50: 559–63.
- 166 Nakagami H, Tomita N, Kaneda Y, Ogihara T, Morishita R. Anti-oxidant gene therapy by NF kappa B decoy oligodeoxynucleotide. *Curr Pharm Biotechnol* 2006; 7: 95–100.
- 167 Sharma RK, Garg BS, Kurosaki H, Goto M, Otsuka M, Yamamoto T, *et al*. Aurine tricarboxylic acid, a potent metal-chelating inhibitor of NFkappaB-DNA binding. *Bioorg Med Chem* 2000; 8: 1819–23.
- 168 Terai K, Matsuo A, McGeer EG, McGeer PL. Enhancement of immunoreactivity for NF-kappa B in human cerebral infarctions. *Brain Res* 1996; 739: 343–9.
- 169 Clemens JA, Stephenson DT, Dixon EP, Smalstig EB, Mincy RE, Rash KS, *et al*. Global cerebral ischemia activates nuclear factor-kappa B prior to evidence of DNA fragmentation. *Brain Res Mol Brain Res* 1997; 48: 187–96.
- 170 Stephenson D, Yin T, Smalstig EB, Hsu MA, Panetta J, Little S, *et al*. Transcription factor nuclear factor-kappa B is activated in neurons after focal cerebral ischemia. *J Cereb Blood Flow Metab* 2000; 20: 592–603.
- 171 Schwaninger M, Inta I, Herrmann O. NF-kappaB signalling in cerebral ischaemia. *Biochem Soc Trans* 2006; 34: 1291–4.
- 172 Clemens JA, Stephenson DT, Yin T, Smalstig EB, Panetta JA, Little SP. Drug-induced neuroprotection from global ischemia is associated with prevention of persistent but not transient activation of nuclear factor-kappaB in rats. *Stroke* 1998; 29: 677–82.
- 173 Nurmi A, Lindsberg PJ, Koistinaho M, Zhang W, Juettler E, Karjalainen-Lindsberg ML, *et al*. Nuclear factor-kappaB contributes to infarction after permanent focal ischemia. *Stroke* 2004a; 35: 987–91.
- 174 Zhang HL, Huang ZH, Zhu Y, Liang ZQ, Han R, Wang XX, *et al*. Neuroprotective effects of prostaglandin A1 in animal models of focal ischemia. *Brain Res* 2005; 1039: 203–6.
- 175 Schneider A, Martin-Villalba A, Weih F, Vogel J, Wirth T, Schwaninger M. NF-kappaB is activated and promotes cell death in focal cerebral ischemia. *Nat Med* 1999; 5: 554–9.
- 176 Hunot S, Brugg B, Ricard D, Michel PP, Muriel MP, Ruberg M, *et al*. Nuclear translocation of NF-kappaB is increased in dopam-

- nergic neurons of patients with parkinson disease. *Proc Natl Acad Sci USA* 1997; 94: 7531–6.
- 177 Dehmer, T, Heneka, MT, Sastre M, Dichgans J. Protection by pioglitazone in the MPTP model of Parkinson's disease correlates with I $\kappa$ B $\alpha$  induction and block of NF kappa B and iNOS activation. *J Neurochem* 2004; 88: 494–501.
- 178 Terai K, Matsuo A, McGeer PL. Enhancement of immunoreactivity for NF-kappa B in the hippocampal formation and cerebral cortex of Alzheimer's disease. *Brain Res* 1996; 735: 159–68.
- 179 Boissière F, Hunot S, Faucheux B, Duyckaerts C, Hauw JJ, Agid Y, *et al*. Nuclear translocation of NF-kappaB in cholinergic neurons of patients with Alzheimer's disease. *Neuroreport* 1997; 8: 2849–52.
- 180 Ferrer I, Martí E, López E, Tortosa A. NF- $\kappa$ B immunoreactivity is observed in association with beta A4 diffuse plaques in patients with Alzheimer's disease. *Neuropathol Appl Neurobiol* 1998; 24: 271–7.
- 181 Kitamura Y, Shimohama S, Ota T, Matsuoka Y, Nomura Y, Taniguchi T. Alteration of transcription factors NF-kappaB and STAT1 in Alzheimer's disease brains. *Neurosci Lett* 1997; 237: 17–20.
- 182 Lukiw WJ, Bazan NG. Strong nuclear factor-kappaB-DNA binding parallels cyclooxygenase-2 gene transcription in aging and in sporadic Alzheimer's disease superior temporal lobe neocortex. *J Neurosci Res* 1998; 53: 583–92.
- 183 Huang Y, Liu F, Grundke-Iqbal I, Iqbal K, Gong CX. NF-kappaB precursor, p105, and NF-kappaB inhibitor, I $\kappa$ B $\gamma$ , are both elevated in Alzheimer disease brain. *Neurosci Lett* 2005; 373: 115–8.
- 184 Kaltschmidt B, Uherek M, Volk B, Baeuerle PA, Kaltschmidt C. Transcription factor NF-kappaB is activated in primary neurons by amyloid beta peptides and in neurons surrounding early plaques from patients with Alzheimer disease. *Proc Natl Acad Sci USA* 1997; 94: 2642–7.
- 185 Bales KR, Du Y, Dodel RC, Yan GM, Hamilton-Byrd E, Paul SM. The NF-kappaB/Rel family of proteins mediates Abeta-induced neurotoxicity and glial activation. *Brain Res Mol Brain Res* 1998; 57: 63–72.
- 186 Samuelsson M, Fisher L, Iverfeldt K. beta-Amyloid and interleukin-1beta induce persistent NF-kappaB activation in rat primary glial cells. *Int J Mol Med* 2005; 16: 449–53.
- 187 Kaltschmidt B, Uherek M, Wellmann H, Volk B, Kaltschmidt C. Inhibition of NF-kappaB potentiates amyloid beta-mediated neuronal apoptosis. *Proc Natl Acad Sci USA* 1999; 96: 9409–14.
- 188 Paris D, Patel N, Quadros A, Linan M, Bakshi P, Ait-Ghezala G, *et al*. Inhibition of Abeta production by NF-kappaB inhibitors. *Neurosci Lett* 2007; 415: 11–6.
- 189 Valerio A, Boroni F, Benarese M, Sarnico I, Ghisi V, Bresciani LG, *et al*. NF-kappaB pathway: a target for preventing beta-amyloid (Abeta)-induced neuronal damage and Abeta42 production. *Eur J Neurosci* 2006; 23: 1711–20.
- 190 Imbimbo BP. The potential role of non-steroidal anti-inflammatory drugs in treating Alzheimer's disease. *Expert Opin Investig Drugs* 2004; 13: 1469–81.
- 191 Kaltschmidt C, Kaltschmidt B, Baeuerle PA. Stimulation of ionotropic glutamate receptors activates transcription factor NF-kappa B in primary neurons. *Proc Natl Acad Sci USA* 1995; 92: 9618–22.
- 192 de Erasquin GA, Hyrc K, Dorsey DA, Mamah D, Dokucu M, Mascó DH, *et al*. Nuclear translocation of nuclear transcription factor-kappa B by alpha-amino-3-hydroxy-5-methyl-4-isoxazole-propionic acid receptors leads to transcription of p53 and cell death in dopaminergic neurons. *Mol Pharmacol* 2003; 63: 784–90.
- 193 Zou J, Crews F. CREB and NF-kappaB transcription factors regulate sensitivity to excitotoxic and oxidative stress induced neuronal cell death. *Cell Mol Neurobiol* 2006; 26: 385–405.
- 194 Cherg JM, Lin HJ, Hung MS, Lin YR, Chan MH, Lin JC. Inhibition of nuclear factor kappaB is associated with neuroprotective effects of glycyrrhizic acid on glutamate-induced excitotoxicity in primary neurons. *Eur J Pharmacol* 2006; 547: 10–21.
- 195 Nakai M, Qin ZH, Wang Y, Chase TN. Free radical scavenger OPC-14117 attenuates quinolinic acid-induced NF-kappaB activation and apoptosis in rat striatum. *Brain Res Mol Brain Res* 1999; 64: 59–68.
- 196 Wang Y, Qin ZH, Nakai M, Chen RW, Chuang DM, Chase TN. Co-stimulation of cyclic-AMP-linked metabotropic glutamate receptors in rat striatum attenuates excitotoxin-induced nuclear factor-kappaB activation and apoptosis. *Neuroscience* 1999; 94: 1153–62.
- 197 Lipton SA. Janus faces of NF-kappa B: neurodestruction versus neuroprotection. *Nat Med* 1997; 3: 20–2.
- 198 Barkett M, Gilmore TD. Control of apoptosis by Rel/NF-kappaB transcription factors. *Oncogene* 1999; 18: 6910–24.
- 199 Kaltschmidt B, Baeuerle PA, Kaltschmidt C. Potential involvement of the transcription factor NF-kappa B in neurological disorders. *Mol Aspects Med* 1993; 14: 171–90.
- 200 Pizzi M, Spano P. Distinct roles of diverse nuclear factor-kappaB complexes in neuropathological mechanisms. *Eur J Pharmacol* 2006; 545: 22–8.