

 Open access • Journal Article • DOI:10.1126/SCIENCE.273.5282.1686

## Dynamic control and amplification of molecular chirality by circular polarized light

— [Source link](#) 

Nina P. M. Huck, Wolter F. Jager, Ben de Lange, Ben L. Feringa

**Institutions:** University of Groningen

**Published on:** 20 Sep 1996 - Science (American Association for the Advancement of Science)

**Topics:** Chirality (chemistry), Planar chirality, Liquid crystal and Circular dichroism

Related papers:

- [Light-driven monodirectional molecular rotor](#)
- [Chiroptical switching between liquid crystalline phases](#)
- [Chiroptical Molecular Switches](#)
- [Absolute Asymmetric Synthesis: The Origin, Control, and Amplification of Chirality.](#)
- [Photoresolution of an Axially Chiral Bicyclo\[3.3.0\]octan-3-one: Phototriggers for a Liquid-Crystal-Based Optical Switch](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/dynamic-control-and-amplification-of-molecular-chirality-by-4hlde7ey3q>

University of Groningen

## Dynamic control and amplification of molecular chirality by circular polarized light

Huck, N.P.M.; Jager, W.F.; de Lange, B.; Feringa, B.L.

*Published in:*  
 Science

*DOI:*  
[10.1126/science.273.5282.1686](https://doi.org/10.1126/science.273.5282.1686)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
 Publisher's PDF, also known as Version of record

*Publication date:*  
 1996

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Huck, N. P. M., Jager, W. F., de Lange, B., & Feringa, B. L. (1996). Dynamic control and amplification of molecular chirality by circular polarized light. *Science*, 273(5282), 1686-1688.  
<https://doi.org/10.1126/science.273.5282.1686>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*



Scienceexpress

SCIENCE ONLINE | SCIENCE MAGAZINE HOME | SCIENCE NOW | NEXT WAVE | SCIENCE'S STKE | SCIENCE CAREERS | E-MARKETPLACE

Institution: University Library | Sign In as Individual | FAQ

Science magazine

HELP | SUBSCRIPTIONS | FEEDBACK | SIGN IN

SEARCH

BROWSE

▶ ORDER THIS ARTICLE

## Dynamic Control and Amplification of Molecular Chirality by Circular Polarized Light

Nina P. M. Huck, Wolter F. Jager, Ben de Lange,  
Ben L. Feringa\*

The enantiomers of a racemic photoresponsive material represent two distinct states that can be modulated with irradiation at a single wavelength by changing the handedness of the light. Dynamic control over molecular chirality was obtained by the interconversion of enantiomers of helically shaped molecules with either left or right circular polarized light (CPL). Photoresolution of the bistable compound as a dopant in a nematic liquid crystalline phase by CPL irradiation led to a chiral mesoscopic phase. The chiral information inherent to CPL is therefore transmitted to the bistable molecule, followed by amplification and macroscopic expression of the chirality.

Department of Organic and Molecular Inorganic Chemistry, University of Groningen, Nijenborgh 4, 9747 AG Groningen, Netherlands.

\* To whom correspondence should be addressed.

The development of organic materials for reversible optical data storage and photochemical switches requires components whose structures and physical properties can be modulated by light (1, 2, 3). Photochromic compounds exist in two distinct, interconvertible forms but often suffer from poor fatigue resistance and destructive read-out (3). Nondestructive read-out has been achieved by monitoring changes in the optical activity of chiral optical switches (4). This requires bistable, nonracemic molecules with sufficiently different optical spectra so that the individual diastereoisomers can be addressed (5). Switching between the enantiomers in a racemic photoresponsive material would constitute a novel approach to a molecular memory element in a binary logic system. Because of the identical optical spectra of enantiomers, distinct states cannot be reached with irradiation unless the unique interaction of enantiomers with CPL can be exploited (6, 7).

We show that photoresolution of helical alkene **1** (Fig. 1A) as well as photomodulation of the helicity is possible with CPL, and that the very small bias for one enantiomer upon CPL irradiation can be macroscopically expressed. Distinct from photochemical molecular switches reported so far, switching between the enantiomers of a racemic compound and modulation of the chiroptical properties and three-dimensional structure have been achieved exclusively by modulation of the chirality of the light. The photochemical modulation of mesophases and physical properties of liquid crystalline (LC) materials is a further challenge for information

▶ [Abstract of this Article](#)

▶ Similar articles found in: [SCIENCE Online](#)

▶ This Article has been cited by:

[other online articles](#)

▶ Search Medline for articles by:


[Huck, N. P. M.](#) || [Feringa, B. L.](#)

▶ Alert me when: [new articles cite this article](#)

▶ [Download to Citation Manager](#)

▶ Collections under which this article appears: [Chemistry](#)

physical properties of liquid crystalline (LC) materials is a further challenge for information technology (8). We show that amplification of the chirality from molecule to mesophase and switching between chiral and achiral mesophases are also affected by CPL irradiation of racemic **1**, added as a dopant to a nematic LC phase.




**Fig. 1.** (A) Photochemical interconversion of P (right-handed) and M (left-handed) helices of **1** upon irradiation with *l*- or *r*-CPL light. (B) Proposed optical data storage system based on the chiral optical switch **1**. Writing is with CPL, and reading and erasing are with LPL.

[\[View Larger Versions of these Images \(32K GIF file\)\]](#)

Of many sterically overcrowded chiral alkenes that we synthesized, compound **1** (Fig. 1A) meets the requirements for a successful switch: (i) The enantiomers are stable at ambient temperatures (free enthalpy of racemization,  $\Delta G_{\text{rac}} = 25.9 \pm 0.2 \text{ kcal mol}^{-1}$ ) and fatigue resistant. (ii) A stereospecific photochemical isomerization process takes place that reverses the helicity of the molecules. (iii) Large circular dichroism (CD) absorptions and optical rotations, essential for detection, are observed.

Racemic **1**, 12-(9<sup>H</sup>-thioxanthene-9<sup>ylidene</sup>-12H-benzo[a]xanthene, was resolved by chiral high-pressure liquid chromatography (HPLC) on a (+)-poly(triphenylmethylmethacrylate) column. The CD spectrum of M-**1** shows a negative Cotton effect at 314 nm (difference in extinction coefficient,  $\Delta\epsilon = -51.3 \text{ cm}^2 \text{ mol}^{-1}$ ) in *n*-hexane:isopropanol (9:1) (Fig. 2A). The anisotropy factor (*g*) is decisive for the selectivity of a CPL switch. The maximum enantiomeric excess that can be achieved upon irradiation with CPL is *g*/2, and for practical purposes the chiral alkene must exhibit sufficiently large *g* values at wavelengths above 300 nm. The experimental *g* values for alkene **1** at selected wavelengths are as follows:  $g \times 10^3 = 2.8$  ( $\lambda = 236 \text{ nm}$ ;  $\Delta\epsilon$  at maximum =  $128.6 \text{ cm}^2 \text{ mol}^{-1}$ ),  $-6.4$  (314 nm,  $-51.3 \text{ cm}^2 \text{ mol}^{-1}$ ), and  $0.4$  (356 nm,  $5.0 \text{ cm}^2 \text{ mol}^{-1}$ ) (units for  $\Delta\epsilon$  are omitted hereafter).



**Fig. 2.** (A) CD spectrum of M-**1** taken in hexane:isopropanol (9:1). (B) The difference in CD absorption at 313 and 400 nm ( $\Delta\epsilon_{313} - \Delta\epsilon_{400}$ ) for a solution of **1** ( $9 \times 10^{-5} \text{ mol liter}^{-1}$ ) in *n*-hexane upon alternating irradiation with *l*- and *r*-CPL. Irradiation was carried out with a 200-W high-pressure mercury lamp equipped with a water filter, a 10-nm bandwidth mercury line filter (313 or 340 nm), a polarization filter, and a  $\lambda/4$  plate (340 nm). The desired wavelength was selected by an interference filter. LPL was generated with a polarization filter, and CPL was generated from this LPL with a 340-nm  $\lambda/4$  plate. Irradiations of degassed alkene solutions were


generated from this LPL with a 570-nm AOT plate. Irradiations of degassed aprotic solutions were performed in 2-cm quartz CD cells. [\[View Larger Versions of these Images \(34K GIF file\)\]](#)

Irradiation of P-1 at 300 nm resulted in rapid photoisomerization into M-1 without notable degradation. The quantum yield for the photochemical racemization was  $\Phi = 0.40$  (pure n-hexane). Next, we examined photoisomerization of racemic **1** under the influence of *l*-CPL and *r*-CPL. CD spectroscopy was used as a detection technique to establish optical enrichment, that is, deracemization, upon irradiation with CPL. Irradiation of the racemate MP-1 ( $9 \times 10^{-5}$  mol liter $^{-1}$ , n-hexane) with *l*-CPL at 313 nm resulted in a negative CD absorption at 313 nm and spectral features comparable with the CD spectrum of M-1 ( $g = 4.2 \times 10^{-6}$ ) (Fig. 2A). Because weak CD effects were expected, we sampled both at 313 nm, where  $\Delta\epsilon$  reaches a maximum, and at 400 nm, where no CD absorption is found. Through use of this method, the effects of CPL irradiation on the difference in  $\Delta\epsilon$  at  $\lambda = 313$  nm and at  $\lambda = 400$  nm ( $\Delta\epsilon_{313} - \Delta\epsilon_{400}$ ) could be determined accurately. Upon successive irradiations with *l*- and *r*-CPL at 313 nm, a modulation of CD absorptions was observed (Fig. 2B). This modulation was entirely reproducible in independent experiments, and no deterioration of the modulated CD signal was seen during eight switching cycles.


For a solution of MP-1 in n-hexane of  $9 \times 10^{-5}$  mol liter $^{-1}$  ( $\Delta\epsilon_{313} = 0.7$ ), the magnitude of the modulation of  $\Delta\epsilon$  was  $1 \times 10^{-5}$  when an irradiation time of 30 min was used. These results indicate that switching occurs between photostationary states with excess P and M helices, with an enantiomeric excess (EE) of 0.07 and  $-0.07\%$ , respectively. The EE values are smaller than anticipated, but taking into account that the light is 90% circular polarized at best and that the bandwidth is 10 nm at most, EEs of no greater than 0.1 to 0.2% can be expected.

A potential data storage system based on racemic **1** can be envisioned (Fig. 1B). Irradiation of a racemate (MP) with *r*-CPL or *l*-CPL for the writing process generates P-enriched or M-enriched regions, respectively. Detection (read-out) is achieved through use of linearly polarized light (LPL) by measuring CD, or optical rotation in transmission, or reflection outside the absorption band. Written information can be erased by LPL (or unpolarized light) at the original wavelength, regenerating MP-1. An erasable direct read after write (EDRAW) process might be executed by changing the chirality of the light at a particular wavelength. In principle, this system constitutes a three-position switch of racemic, P-enriched, and M-enriched **1**, with the distinct advantage that all the processes can be performed at a single wavelength.

Once the switching process with CPL irradiation had been firmly established, we studied the photoisomerization in LC materials using racemic **1** as a dopant (10, 11). A stable nematic phase was obtained from 20 weight % MP-1 and 4'-*t*-(pentyloxy)-4-biphenylcarbonitrile **2**. Differential scanning calorimetry showed a solid-nematic transition ( $T_{S \rightarrow N}$ ) at 48.2°C [enthalpy change,  $\Delta H = 71$  J/g; for pure **2** the transition occurs at 52.4°C ( $\Delta H = 96$  J/g)]. Irradiation of a thin film of racemic **2**, doped with MP-1 at 313 nm for 90 min with *l*-CPL, analogous to the experiments in solution, resulted in a cholesteric phase (Fig. 3A), showing that the photostationary state P-1  $\rightleftharpoons$  M-1 with an optical enrichment in M-1 is also reached in the LC phase. The formation of the cholesteric phase was accompanied by a change in phase transition ( $T_{S \rightarrow N}$  at 48.7°C,  $\Delta H = 66$  to 69 J/g). Control experiments showed that a similar effect on the phase transition occurred with doping of **2** with 20 weight % of racemic **1**, 1.5% enriched in either P-1 or M-1 ( $T_{S \rightarrow N}$  at 48.7°C,  $\Delta H = 69$  J/g).



**Fig. 3.** (A) Cholesteric texture observed for 4'-*t*-(pentyloxy)-4-biphenylcarbonitrile (**2**) doped with 20 weight % racemic **1** after irradiation with 313-nm *r*-CPL light. Phase separation can be seen. (B) Cholesteric structure obtained for **2** doped with 20 weight % racemic



cholesteric structure obtained for **2** doped with 20 weight % racemic **1** enriched with 1.5 weight % M-**1**. A similar cholesteric structure as in (A) can be observed. Bar indicates 100  $\mu\text{m}$ .

[\[View Larger Version of this Image \(143K GIF file\)\]](#)

In a separate experiment on a thin film of **2** doped with racemic **1** in which *r*-CPL was used instead of *l*-CPL, photoderacemization was again observed. Irradiation of the film of the cholesteric phase reached after CPL irradiation, containing excess M-**1**, with unpolarized light at 313 nm resulted in a nematic phase. This is a consequence of the formation of a 50:50 ratio of opposite helices of **1**; that is, photoracemization occurred. A large pitch ( $p = 580 \mu\text{m}$  based on helical twisting power 0.1) is seen in the cholesteric phase as a result of low resolution of **1** by CPL irradiation. We performed control experiments to establish independently the optical enrichment of dopant **1** as the origin of the induced cholesteric phase. An LC phase was prepared from **2** and 20 weight % racemic **1** enriched with 1.5% M-**1**. A cholesteric texture (Fig. 3B) was obtained similar to that observed after CPL irradiation (Fig. 3A).

Because of the high amount of dopant, some phase separation (demixing) was observed after CPL irradiation (Fig. 3A). This phenomenon, reminiscent of conglomerate or racemate formation in the solid state (12), was observed in several experiments and points to differences in the stability of chiral and racemic phases at low enantiomeric excesses of the dopant. The photochemical switching processes of the LC phases are completely controlled by the changes in chirality of the light at a single (313 nm) wavelength: (i) switching between LPL and CPL results in a nematic to cholesteric modulation, and (ii) switching between *l*-CPL and *r*-CPL modulates the chirality of the cholesteric phase (left-handed and right-handed cholesteric phases, respectively). Great care was taken in all experiments to exclude chiral contamination or any other sources besides CPL irradiation of the induced chirality resulting in a cholesteric phase. It should be emphasized that rather large amounts of racemic dopant are required, compared with optically active dopants (10), to observe the cholesteric phase; on the basis of an EE of 0.07% in solution, 0.14% enrichment can be achieved when the LC phase is doped with 20 weight % of MP-**1**. An increase of the helical twisting power and anisotropy factor of the dopant, by structural modifications of the chromophore in **1**, may decrease the required dopant concentration.

## REFERENCES AND NOTES

1. J.-M. Lehn, *Supramolecular Chemistry* (VCH, Weinheim, 1995).
2. P. Ball and L. Garwin, *Nature* **355**, 761 (1992).
3. B. L. Feringa, W. F. Jager, B. de Lange, *Tetrahedron* **49**, 8267 (1993); I. Willner and B. Willner, *Adv. Mater.* **17**, 587 (1995). For a pertinent example of a stable molecular switch, see S. H. Kawai, S. L. Gilat, J.-M. Lehn, *J. Chem. Soc. Chem. Commun.* **8**, 1011 (1994).
4. For switching process of diastereoisomers with unpolarized light, see W. F. Jager *et al.*, *Angew. Chem. Int. Ed. Engl.* **34**, 348 (1995).
5. For a brief review of chiroptical molecular switches based on bistable nonracemic materials, see B. L. Feringa, N. P. M. Huck, A. M. Schoevaars, *Adv. Mater.*, in press.
6. Discriminative interactions of CPL with racemic materials have been proposed in the context of the origin of chirality [S. L. Milla and L. E. Orgel, *The Origin of Life on the Earth* (Prentice-Hall, Englewood Cliffs, NJ, 1974); W. A. Bonner, *Top. Stereochem.* **18**, 1 (1988); S. F. Mason, *Chem. Soc. Rev.* **17**, 347 (1988); M. Gardner, in *The Ambidextrous Universe*, C. Scribner, Ed. (Penguin, Harmondsworth, UK, 1982); R. A. Hegstrom and



- D. R. Kondepudi, *Sci. Am.* **262**, 108 (January 1990); S. F. Mason, *Molecular Optical Activity and the Chiral Discriminations* (Cambridge Univ. Press, Cambridge, 1982); A. Salam, *J. Mol. Evol.* **33**, 105 (1991) ; L. D. Barron, *New Developments in Molecular Chirality* (Kluwer, Dordrecht, Netherlands, 1991); D. K. Kondepudi and G. W. Nelson, *Nature* **314**, 438 (1985) ; B. S. Green, M. Lahav, D. Rabinovitch, *Acc. Chem. Res.* **12**, 191 (1979) ].
7. Photodestruction and photoresolution with CPL has been achieved [ K. L. Stevenson and J. K. Verdick, *J. Am. Chem. Soc.* **90**, 2974 (1968) ; A. Moradpour, J. F. Nicoud, G. Balavoine, H. Kagan, G. Tsoucaris, *ibid.* **93**, 2353 (1971) ; Y. Zhang and G. B. Schuster, *J. Org. Chem.* **60**, 7192 (1995) ].
  8. D. Kreyzig and J. Stumpe, in *Selected Topics in Liquid Crystal Research*, H. D. Koswig, Ed. (Verlag Chemie, Berlin, 1990).
  9. H. Rau, *Chem. Rev.* **83**, 535 (1983) .
  10. Small amounts of an optically active guest molecule added to a nematic host can induce a cholesteric phase, and the helical organization (pitch) in the mesoscopic system is sensitive to modifications in the chiral guest molecule [ G. Solladié and R. G. Zimmermann, *Angew. Chem. Int. Ed. Engl.* **23**, 348 (1984) ].
  11. Recently, photochemical switching between LC phases through use of diastereomeric photoresponsive guest molecules was shown [ B. L. **Feringa**, N. P. M. Huck, H. A. van Doren, *J. Am. Chem. Soc.* **117**, 9929 (1995) ].
  12. J. Jacques, A. Collet, S. Wilen, *Enantiomers, Racemates and Resolutions* (Wiley, New York, 1981).
  13. We thank R. M. Kellogg, H. A. van Doren, and J. van Esch for usefull comments.

31 May 1996; accepted 18 July 1996

---

- ▼ [Abstract of this Article](#)
- ▼ Similar articles found in:  
[SCIENCE Online](#)
- ▼ This Article has been cited by:
- ▼ Search Medline for articles by:  
[Huck, N. P. M.](#) || [Feringa, B. L.](#)
- ▼ Alert me when:  
[new articles cite this article](#)
- ▼ [Download to Citation Manager](#)
- ▼ Collections under which this article appears:  
[Chemistry](#)

## This article has been cited by other articles:

- Zahn, S., Canary, J. W. (2000). Electron-Induced Inversion of Helical Chirality in Copper Complexes of N,N-Dialkylmethionines. *Science* 288: 1404-1407 [[Abstract](#)] [[Full Text](#)]

Volume 273, Number 5282, Issue of 20 Sep 1996, pp. 1686-1688.

Copyright © 1996 by The American Association for the Advancement of Science.

**SIGNAL TRANSDUCTION  
KNOWLEDGE ENVIRONMENT**

Customize Your  
Job Search



---

▲ PAGE TOP