

Dynamics of the 2006/2007 dengue outbreak in Brazil

Giovanini E Coelho, Marcelo Nascimento Burattini^{1/+}, Maria da Glória Teixeira²,
Francisco Antonio Bezerra Coutinho¹, Eduardo Massad^{1,3}

Programa Nacional de Controle da Dengue, Ministério da Saúde, Brasília, Brasil ¹Faculdade de Medicina, Universidade de São Paulo, R. Teodoro Sampaio 115, 05405-000 São Paulo, SP, Brasil ²Instituto de Saúde Coletiva, Universidade Federal da Bahia, Salvador, BA, Brasil ³London School of Hygiene and Tropical Medicine, London, UK

We analyzed dengue incidence in the period between October 2006-July 2007 of 146 cities around the country where Larval Index Rapid Assay (LIRA) surveillance was carried out in October 2006. Of these, we chosen 61 cities that had 500 or more cases reported during this period. We calculated the incidence coefficient, the force of infection (λ) and the basic reproduction number (R_0) of dengue in those 61 cities and correlated those variables with the LIRA. We concluded that λ and R_0 are more associated with the number of cases than LIRA. In addition, the average R_0 for the 2006/2007 dengue season was almost as high as that calculated for the 2001/2002 season, the worst in Brazilian history.

Key words: dengue - larval index - basic reproduction number - force of infection

Dengue fever (DF), a Flaviridae infection transmitted by peridomestic *Aedes aegypti* (Gubler & Kuno 1997) re-appeared as a major urban epidemic in Brazil in 1986 (Marques et al. 1994, Massad et al. 2001, Luz et al. 2003) and is currently the most important arthropod-borne viral disease in Brazil (Luz et al. 2003). By the year 2000, all 26 states had reported DF cases. In the period 1986-2007, DF caused 4,559,812 officially reported cases (MS 2007a) with 493 deaths (MS 2007b).

The worst year in history was 2002, with the introduction of DEN-3, when 794,219 cases were reported. In the following years, DEN-3 spread itself throughout the country, causing several important outbreaks (MS 2007c). In 2006, 345,922 dengue cases were reported, with 682 dengue hemorrhagic fever (DHF) cases causing 76 deaths (MS 2007c). From January to July 2007, 438,949 classical DF cases were reported, with 926 cases of DHF, causing 98 deaths (MS 2007c). In this year DEN-3 virus caused 81% of the cases.

An important observation related to dengue transmission is that, in some subtropical areas like Brazil, the density of mosquitoes (Forattini et al. 1993a, b, 1995) and the number of DF cases starts to increase in number at the beginning of the rainy season (October) with the cases of dengue peaking more than three months later (Coutinho et al. 2006).

Long-term control of dengue and condition for outbreak occurrence has relied mainly on larval index (LI), that is, the proportion of houses with breeding places harboring aedes larvae which, paradoxically, has not

demonstrated a good correlation with the number of DF cases (Gubler & Kuno 1997, Burattini et al. 2007). One of the few studies that indirectly show some correlation between the house index and the number of dengue cases is the work by Cunha et al. (1997), in which it is suggested, although it is not demonstrated, a relationship between low number of cases and low house index in a community of Rio de Janeiro. In Brazil, an alternative approach to the so-called container indexes (German et al. 1980, Bang et al. 1981) for larval surveillance was proposed in order to optimize the logistics of its operation. It consists of random sampling a relatively small number of dwellings in which surveillance of aedes-positive breeding places is carried out simultaneously in several cities around the country. This strategy has been called Larval Index Rapid Assay (LIRA).

In this paper we analyze the period October 2006-July 2007 of 146 Brazilian cities that carried out LIRA surveillance in October 2006. Of these, we choose those 61 cities that experienced 500 or more dengue cases in the period. We calculate the incidence coefficient (IC), the force of infection (λ) and the basic reproduction number (R_0) of dengue in those 61 cities and cross-correlate those variables along with the LIRA, in order to estimate which parameter related to the intensity of transmission is the best predictor of the number of cases.

MATERIALS AND METHODS

LIRA - Consisted in the simultaneous visit of dwellings randomly chosen of 146 cities representing all the Brazilian regions (MS 2005). LIRA is a random sampling technique in which the sample unit corresponds to 9,000 to 12,000 dwellings, from which a maximum of 450 houses are randomly selected for inspection. The LIs obtained are the proportion of houses with *A. aegypti* larvae and the Breteau index. It is being used in Brazil since 2003.

Calculation of λ and R_0 - Of the 146 cities visited for the LIRA surveillance, we selected 61 that had 500 or

Financial support: CNPq, FAPESP, LIM01 HC-FMUSP

+ Corresponding author: edmassad@usp.br

Received 28 February 2008

Accepted 21 July 2008

more reported cases and experienced a dengue outbreak in the period. By outbreak we understand the sudden exponential rise in the number of cases in the beginning of the transmission season. The list of those 61 cities can be seen in Table.

The calculation of R_0 can be carried out from the initial exponential growing phase of the number of cases as detailed in Massad et al. (2001). However, we show in the appendix a summary of the derivation of R_0 and λ .

From equations (5) and (6) of the appendix (Supplementary data) we obtain the relationship between R_0 and λ for each one of the 61 cities, such as:

$$R_0 = 1 + \frac{\lambda^2 + \lambda(\mu + \gamma)}{\gamma \mu}$$

In this equation, λ is the growing rate of the number of dengue cases in the initial exponential phase, the incidence rate or λ of the disease in this phase. By fitting the proportion of new cases to an exponential curve, we can estimate R_0 . The values of the parameters were $\mu = 0.156 \text{ weeks}^{-1}$ and $\gamma = 1 \text{ week}^{-1}$ (Massad et al. 2001).

Statistical analysis - We fitted exponential curves by the method of the Minimum Squares to the number of cases in each week starting in October 2006 for each one of the 61 chosen cities. As mentioned above, the growing rate of the number of cases in the initial exponential phase, λ corresponds to the incidence rate or λ of the disease in this phase (equation 4 of the appendix).

The IC per 100,000 inhabitants, λ and R_0 of dengue in those 61 cities were correlated with their respective LIRA. As a first approximation to estimate the correlations between the variables, we carried out a linear regression analysis that correlated the LIRA versus λ ; LIRA versus R_0 ; LIRA versus IC; λ versus the IC; and R_0 versus IC. The strength of the above correlations was estimated by the calculation of the Pearson's correlation coefficients. Of course other functional associations between the variables could serve the same purpose. However, we chosen a linear regression because the correlations found were poor for all tested function and so we chosen the linear for simplicity. All the statistical analyses were carried out with the software SPSS 15.0.

RESULTS

Table shows the 61 cities analyzed with the respective values of the LIRA, λ , R_0 , the population size, IC and the goodness of fit of the exponential model.

In Fig. 1 we show an example of a city (Paranaí) that resulted in a particularly accurate fitting of dengue cases. Note the exponential shape of the fitted curve and the resulting incidence of 0.32 per capita new cases per week.

In Figs 2 and 3 we show the correlation between LIRA and λ and LIRA and the R_0 , respectively. Note that LIRA significantly correlates with both λ and R_0 .

Finally, we tested the relation between LIRA, λ and R_0 with dengue IC per 100,000 inhabitants for each city analyzed. Results are shown in Figs 4-6, respectively. It is noteworthy that LIRA correlates with both λ and R_0 but none of them correlated with IC. Note also the pres-




Fig. 1: example of the exponential fitting for the number of dengue cases for the city of Paranaí.




Fig. 2: correlation between LIRA and λ for the 61 cities analyzed.




Fig. 3: correlation between LIRA and R_0 for the 61 cities analyzed.




Fig. 4: correlation between LIRA and the IC per 100,000 inhabitants for the 61 cities analyzed.

TABLE
Results of the analysis with the 61 chosen Brazilian cities

City	<i>LIRA</i>	λ	r	p	R_0	dengue cases	incidence
Uberlândia	0.44	0.15	0.810	< 0.001	2.26	682	110.83
Parnamirim	0.36	0.16	0.875	< 0.001	2.35	1.175	664.19
Duque de Caxias	1.79	0.17	0.798	< 0.001	2.45	664	76.58
Belford Roxo	2.85	0.17	0.934	< 0.001	2.45	834	167.73
Nova Iguaçu	1.96	0.19	0.901	< 0.001	2.64	1.154	134.48
Serra	3.58	0.19	0.673	0.007	2.64	1.908	470.61
Juiz de Fora	0.31	0.2	0.949	< 0.001	2.74	509	98.45
Vitória	3.49	0.21	0.962	< 0.001	2.84	888	276.79
Sabará	1.34	0.22	0.765	< 0.001	2.94	654	476.88
Rio Verde	1.90	0.22	0.963	< 0.001	2.94	872	626.44
Londrina	0.54	0.23	0.957	< 0.001	3.04	3.891	773.50
Brasília	1.37	0.23	0.919	< 0.001	3.04	2.157	88.62
Belém	1.63	0.23	0.821	< 0.001	3.04	4.812	331.70
Araguaína	2.06	0.23	0.970	< 0.001	3.04	1.838	1385.42
Manaus	1.65	0.24	0.973	< 0.001	3.15	2.504	144.57
Teresina	0.32	0.25	0.898	< 0.001	3.25	4.957	608.18
Belo Horizonte	1.78	0.25	0.977	< 0.001	3.25	8.037	331.52
São José do Rio Preto	1.66	0.26	0.979	< 0.001	3.36	10.012	2360.69
Vila Velha	2.67	0.26	0.856	0.002	3.36	541	130.57
Maracanaú	0.93	0.27	0.983	< 0.001	3.47	751	377.50
Araçatuba	1.25	0.27	0.981	< 0.001	3.47	2.117	1153.90
Cambé	1.34	0.27	0.924	< 0.001	3.47	1.484	1478.26
Campos dos Goytacazes	2.18	0.27	0.818	< 0.001	3.47	2.103	485.58
Itaboraí	2.23	0.27	0.932	< 0.001	3.47	1.200	530.87
Niterói	2.30	0.27	0.887	< 0.001	3.47	7.423	1548.82
Contagem	1.10	0.28	0.979	< 0.001	3.58	619	100.94
Ribeirão das Neves	1.68	0.28	0.989	< 0.001	3.58	648	193.74
Barretos	2.86	0.28	0.969	< 0.001	3.58	501	450.76
Angra dos Reis	2.00	0.285	0.826	< 0.001	3.63	827	559.18
Aparecida de Goiânia	0.68	0.29	0.927	< 0.001	3.69	3.475	738.21
Betim	1.40	0.29	0.971	< 0.001	3.69	763	180.74
Parnaíba	0.92	0.3	0.965	< 0.001	3.80	1.183	813.64
Cabo Frio	2.72	0.3	0.941	< 0.001	3.80	708	412.95
Sobral	0.79	0.31	0.924	< 0.001	3.91	3.603	2013.79
Paranavaí	3.97	0.32	0.968	< 0.001	4.03	1.502	1883.46
São Luís	2.17	0.33	0.962	< 0.001	4.14	1.874	184.13
Resende	1.40	0.35	0.940	< 0.001	4.38	743	608.92
Goiânia	3.14	0.35	0.948	< 0.001	4.38	6.503	524.58
Guarulhos	0.65	0.38	0.919	0.002	4.74	864	65.70
Fortaleza	1.10	0.38	0.918	0.002	4.74	16.083	654.17
Macaé	2.01	0.38	0.906	< 0.001	4.74	1.784	1081.18
São Paulo	0.23	0.4	0.932	< 0.001	4.99	2.743	24.70
Rio de Janeiro	6.20	0.4	0.890	0.003	4.99	21.821	353.16
Governador Valadares	6.25	0.4	0.818	0.015	4.99	624	238.84
Montes Claros	4.22	0.41	0.970	< 0.001	5.12	1.052	296.05
Santos	0.89	0.42	0.673	0.03	5.24	830	198.36
Maringá	1.51	0.43	0.866	< 0.001	5.37	8.356	2533.66
Foz do Iguaçu	9.85	0.43	0.981	< 0.001	5.37	4.592	1449.71
Cascavel	1.09	0.45	0.872	< 0.001	5.63	543	187.29
Magé	2.72	0.47	0.918	< 0.001	5.90	1.149	475.37
Osasco	1.47	0.49	0.922	< 0.001	6.17	893	123.28
São Gonçalo	2.00	0.49	0.966	< 0.001	6.17	3.006	304.93
Campo Grande	1.71	0.52	0.980	< 0.001	6.59	48.574	6222.71
Macapá	1.06	0.53	0.580	0.04	6.73	4.544	1191.98
Piracicaba	1.85	0.54	0.937	< 0.001	6.87	2.608	700.94
Ribeirão Preto	3.92	0.56	0.964	< 0.001	7.16	2.867	504.83
Juazeiro do Norte	1.79	0.6	0.928	< 0.001	7.75	737	300.89
Coronel Fabriciano	7.42	0.65	0.947	< 0.001	8.53	543	512.40
Marília	1.13	0.67	0.929	< 0.001	8.84	978	428.69
Cáceres	6.86	0.8	0.908	0.008	11.03	877	956.23
Cacoal	2.74	0.82	0.967	0.004	11.39	726	944.65




Fig. 5: correlation between λ and IC per 100,000 inhabitants for the 61 cities analyzed.




Fig. 6: correlation between R_0 and IC per 100,000 inhabitants for the 61 cities analyzed.

ence of an outlier in Figs. However, its presence does not change significantly the correlation between the variables analyzed, and so we decided to keep it in the analysis.

DISCUSSION

The 2006/2007 dengue season in Brazil was the most severe in number of cases since the 2001/2002 season. This is reflected by the values of R_0 available so far. If we compare the average values of R_0 for the years of 1991 (2.03, Marques et al. 1994), 2001 (6.29, Massad et al. 2001) and 2007 (4.51, Burattini et al. 2007) we conclude that 2007 is closer to 2001 values than to 1991 as far as dengue intensity of transmission is concerned. We should stress here that the entire estimated LIRA used in this analysis refer to whole town averages, although dengue, as many of the vector-borne infections, have heterogeneous spatial distribution over the affected areas. This would theoretically underestimate the incidence, λ and R_0 estimates (Coutinho et al. 1999). However, our model is based on known cases and our equations system is scale-invariant, that is, it could be expressed in terms of proportions or absolute numbers in an approximately homogeneous area where the cases are happening. Moreover, dengue control measures as proposed by the Brazilian National Program of Dengue Control, in spite of being aimed to cover whole towns, should prioritize the focuses where transmission is likely to occur.

The regression analysis indicates that, despite the low values obtained for the correlation coefficients, they are significant for the associations between LIRA and λ and LIRA and R_0 . Since LIRA is an entomologically derived variable and λ and R_0 were estimated from human cases, it is noteworthy that there are weak, although significant correlations between them. To the best of our knowledge, this has not been demonstrated before. The correlations found between the entomological indexes and the estimators of the intensity of transmission are due to the fact that all indexes used are based on larval stages. These have to develop into the further stages until the adult mosquitoes that, in turn, have to bite infected individuals and the new susceptibles. It is, therefore, expected that the large number of unknown parameters governing each step of that chain, would vary from place to place and from moment to moment and, therefore, would weakened those correlation. In spite of this, LIRA could be used as a proxy for the velocity of the following dengue epidemic growth. In addition, those regions with the higher R_0 are those likely to experience major impact of future outbreaks, especially when a new serotype starts to circulate.

However, when we try to calculate the association of these parameters with the incidence of dengue, none demonstrated to be significantly correlated with it. Nevertheless, when we compare the significance of the association between LIRA and incidence rate of dengue with the associations between λ and R_0 and the incidence rate we observe that both λ and R_0 are more correlated with incidence than LIRA. If we consider the large number of variables associated with dengue transmission we should expect a poor association between each of the parameters analyzed and the incidence of the infection. However, in view of the p-values found for each of them, we may conclude that both λ and R_0 are better predictors of the total number of cases than the LI currently used. This is probably so because λ and R_0 are also related to human dengue epidemiological factors, such as previous dengue outbreaks and herd immunity effects, in addition to the entomological parameters.

We would like to emphasize the importance of the intensity of transmission parameters proposed in this work, namely λ and R_0 , both of which are more closely associated with the total number of cases than the LIs, although far from being reliable predictors of incidence. However, they are clear indicators of the intensity of outbreaks and relatively easy to calculate in the particular situation in which the number of cases start to increase after a relatively long period with very small number of cases.

As mentioned above, the current predominant strain in Brazil is DEN-3. It would be interesting to include a comment on potential differences in the outcomes now that DEN-2 started to predominate in several states of the country. Certainly, the fact that a great cohort of susceptibles to DEN-2 is available would increase all the estimators of intensity of transmission analyzed in this study.

Finally, it is noteworthy that LIRA performed in nearly October has some predictive power on the velocity of epidemic growth, as demonstrated by the association found between LIRA, λ and R_0 . Nevertheless, LIRA is not reliable as a predictor of dengue incidence, a fact already

known for other LIs. However, the relation between LIRA and the rate of growth of dengue epidemic demonstrated herein is a new result and it is one of the most important results of this paper, since the results of LIRA obtained almost two months before the beginning of dengue outbreaks could serve as a guide for early intervention.

REFERENCES

- Bang YH, Bown DN, Onwubiko AO 1981. Prevalence of larvae of potential yellow fever vectors in domestic water containers in south-east Nigeria. *Bull WHO* 59: 107-114.
- Burattini MN, Chen M, Chow A, Coutinho FAB, Goh KT, Lopez LF, Ma S, Massad E 2007. Modelling the control strategies against dengue in Singapore. *Epidemiol Infect* 136: 309-319.
- Coutinho FAB, Burattini MN, Lopez LF, Massad E 2006. Threshold conditions for a non-autonomous epidemic system describing the population dynamics of dengue. *Bull Math Biol* 68: 2263-2282.
- Coutinho FAB, Lopez LF, Massad E, Burattini MN, Struchiner CJ, Azevedo-Neto RS 1999. Modelling heterogeneities in individual frailties in epidemic models. *Math and Comput Modell* 30: 97-115.
- Cunha V, Maspero RC, Miagostovich MP 1997. Dengue infection in Paracambi, state of Rio de Janeiro, 1990-1995. *Rev Soc Bras Med Trop* 30: 379-383.
- Forattini OP, Kakitani I, Massad E, Marucci D 1993a. Studies on mosquitoes (Diptera: Culicidae) and anthropic environment. 2. Immature stages research at a rice irrigation system location in South-Eastern Brazil. *Rev Saúde Pública* 27: 227-236.
- Forattini OP, Kakitani I, Massad E, Marucci D 1993b. Studies on mosquitoes (Diptera: Culicidae) and anthropic environment. 4. Survey of resisting adults and synanthropic behaviour in South-Eastern Brazil. *Rev Saúde Pública* 27: 398-411.
- Forattini OP, Kakitani I, Massad E, Marucci D 1995. Studies mosquitoes (Diptera: Culicidae) and anthropic environment. 9. Synanthropic and epidemiological vector role of *Aedes scapularis* in South-Eastern Brazil. *Rev Saúde Pública* 29: 199-207.
- Germain M, Francy DB, Monath TP, Ferrara L, Bryan J, Salaun J-J, Heme G, Renaudet J, Adam C, Digoutte J-P 1980. Yellow fever in The Gambia, 1978-1979; entomological aspects and epidemiological correlations. *Am J Trop Med Hyg* 29: 929-940.
- Gubler DJ, Kuno G 1997. *Dengue and Dengue Hemorrhagic Fever*. CABI publishing, Wallingford, 478 pp.
- Luz PM, Codeço PT, Massad E, Struchiner CJ 2003. Uncertainties regarding dengue modeling in Rio de Janeiro, Brazil. *Mem Inst Oswaldo Cruz* 98: 871-878.
- Marques CA, Forattini OP, Massad E 1994. The basic reproduction number for dengue fever in São Paulo state, Brazil: 1990-1991 epidemics. *Trans R Soc Trop Med Hyg* 88: 58-59.
- Massad E, Coutinho FAB, Burattini MN, Lopez LF 2001. The risk of yellow fever in a dengue infested area. *Trans R Soc Trop Med Hyg* 95: 370-374.
- MS-Ministério da Saúde 2005. *Diagnóstico rápido nos municípios para vigilância entomológica de Aedes aegypti no Brasil – LIRAA, metodologia para avaliação dos índices de Breteau e predial (portuguese)*. Secretaria de Vigilância em Saúde, Programa Nacional de Controle da Dengue, Brasília, Brazil.
- MS-Ministério da Saúde 2007a. Graphic showing the historical series of reported dengue cases between 1980 and 2005. [cited 2007 Oct 13] Available from: http://portal.saude.gov.br/portal/arquivos/pdf/Graficos_DNC_dengue_verde.pdf.
- MS-Ministério da Saúde 2007b. Graphic showing the historical series of mortality due to dengue cases between 1980 and 2005 [cited 2007 Oct 13]. Available from: http://portal.saude.gov.br/portal/arquivos/pdf/Graficos_DNC_dengue_vermelho.pdf.
- MS-Ministério da Saúde 2007c. Epidemiological Bulletin reporting the 2007 dengue outbreak in Brazil. [cited 2007 Oct 13]. Available from: http://portal.saude.gov.br/portal/arquivos/pdf/dengue_0210.pdf.

Appendix - Let S_H and I_H be the number of susceptible and infectious individuals in a human population. Let also S_m and I_m be number of susceptible and infectious individuals, respectively, in a mosquito population. We can write (Massad et al. 2001):

$$(1) \quad \begin{aligned} \frac{dI_H}{dt} &= aN_m \frac{I_m}{N_m} b \frac{S_H}{N_H} - \gamma I_H \\ \frac{dI_m}{dt} &= \exp(-\mu\lambda) a S_m(t-\tau) \frac{I_H}{N_H} - \mu I_m \end{aligned}$$

In the first equation of system (1) a is the average number of bites a mosquito inflicts in the human population, so that aN_m is the total number of bites per unit of time and b is the fraction of those bites that are actually infective for humans. Humans recover from infection with rate γ . Finally, only a fraction S_H/N_H of the infective bites is in susceptible individuals, resulting in new infections.

In the second equation of system (1) the extrinsic incubation period is represented by τ . Therefore, $aS_m(t-\tau)$ is the number of bites per unit time given by susceptible mosquitoes in the instant $(t-\tau)$. A fraction I_H/N_H of those bites is in infected humans, producing infected mosquitoes, a fraction $\exp(-\mu\tau)$ of which survives the extrinsic incubation period τ to become infective for the rest of their lives. Mosquitoes are assumed to die by natural causes at a rate μ .

Dividing the first equation of system (1) by N_H (total human population) and the second by N_m (total mosquito population) we have:

$$(2) \quad \begin{aligned} \frac{di_H}{dt} &= mai_m b S_H - \gamma i_H \\ \frac{di_m}{dt} &= \exp(-\mu\lambda) a S_m(t-\tau) i_H - \mu i_m \end{aligned}$$

where $m = N_m/N_H$ and the lower case letters represent proportions.

At the beginning of an outbreak, we can assume that $S_H \cong 1$ and $S_m(t-\tau) \cong 1$ to get the linearized system

$$(3) \quad \begin{aligned} \frac{di_H}{dt} &= mai_m b - \gamma i_H \\ \frac{di_m}{dt} &= \exp(-\mu\lambda) ai_H - \mu i_m \end{aligned}$$

whose general solution is

$$(4) \quad \begin{aligned} i_H &= c_H \exp(\lambda t) \\ i_m &= c_m \exp(\lambda t) \end{aligned}$$

Taking the derivative of equations (4) and substituting the result in equations (3) we get:

$$(5) \quad \lambda = \frac{1}{2} [-(\mu + \lambda) \pm \sqrt{((\mu + \lambda)^2 - 4\mu\gamma + 4ma^2 b \exp(-\mu\tau))}]$$

Remembering that R_0 is

$$(6) \quad R_0 = \frac{ma^2 b \exp(\mu\lambda)}{\gamma\mu}$$

Therefore, from equations (5) and (6) we obtain (Massad et al. 2001):

$$(7) \quad R_0 = 1 + \frac{\lambda^2 + \lambda(\mu + \gamma)}{\gamma\mu}$$