

E_8 instantons on type-A ALE spaces and supersymmetric field theories

Noppadol Mekareeya,^a Kantaro Ohmori,^b Yuji Tachikawa^c and Gabi Zafrir^c

^a*Dipartimento di Fisica, Università di Milano-Bicocca, and INFN, sezione di Milano-Bicocca, Piazza della Scienza 3, I-20126 Milano, Italy*

^b*School of Natural Sciences, Institute for Advanced Study, Princeton, NJ 08540, U.S.A.*

^c*Kavli Institute for the Physics and Mathematics of the Universe, University of Tokyo, Kashiwa, Chiba 277-8583, Japan*

E-mail: n.mekareeya@gmail.com, kantaro@ias.edu,
yuji.tachikawa@ipmu.jp, gabi.zafrir@ipmu.jp

ABSTRACT: We consider the 6d superconformal field theory realized on M5-branes probing the E_8 end-of-the-world brane on the deformed and resolved $\mathbb{C}^2/\mathbb{Z}_k$ singularity. We give an explicit algorithm which determines, for arbitrary holonomy at infinity, the 6d quiver gauge theory on the tensor branch, the type-A class S description of the T^2 compactification, and the star-shaped quiver obtained as the mirror of the T^3 compactification.

KEYWORDS: Supersymmetry and Duality, Field Theories in Higher Dimensions, Field Theories in Lower Dimensions

ARXIV EPRINT: [1707.04370](https://arxiv.org/abs/1707.04370)

Contents

1	Introduction and summary	1
2	Geometric preliminaries	4
2.1	Topological data of the instanton configuration	4
2.2	Dimension of the instanton moduli space	5
3	Six-dimensional description	7
3.1	The general structure of the quiver	7
3.2	The algorithm	8
3.3	A subtlety concerning the 6d θ angle	9
3.4	Anomalies and the inflow	10
4	Lower dimensional incarnations	12
4.1	Five-dimensional brane-web description	12
4.2	Four-dimensional class S description	14
4.3	Three-dimensional star-shaped quiver description	15
5	Examples	17
5.1	The case of $k = 2$	17
5.2	Enhanced flavor symmetries from 3d quivers	20
5.3	The case of $k = 4$	23
5.4	Theories differing by the 6d θ angle	27
5.5	Massive E-string theories	28
5.6	Higgsing the $SU(k)$ flavour symmetry	29

1 Introduction and summary

One of the many surprises during the second superstring revolution was the realization that the construction of $SU(N)$ instantons on \mathbb{R}^4 by Aiyah-Drinfeld-Hitchin-Manin [1] and on asymptotically locally Euclidean (ALE) spaces by Kronheimer-Nakajima [2, 3] have a physical realization in terms of Dp -branes probing $D(p+4)$ -brane on \mathbb{R}^4 [4] and/or ALE spaces [5]. There, we have a gauge theory with eight supercharges on Dp -branes such that its Higgs branch is given by the corresponding instanton moduli spaces: the equations of the ADHM and Kronheimer-Nakajima construction are the F-term and D-term conditions of the supersymmetric gauge theory.

As a variation of this construction, we can consider M5-branes probing the E_8 end-of-the-world brane of the M-theory, either on \mathbb{R}^4 or on ALE spaces. The low-energy worldvolume theory on these M5-branes is a 6d $\mathcal{N} = 1$ supersymmetric theory whose Higgs

branch is intimately related to the E_8 instanton moduli spaces. These theories were studied already in the heyday of the second revolution, see e.g. [6]. We did not, however, have many methods to understand the properties of these theories back then, since these theories are intrinsically strongly-coupled. Therefore we could not say anything new regarding the mathematics of the E_8 instanton moduli spaces on \mathbb{R}^4 or ALE spaces, for which no constructions analogous to ADHM or Kronheimer-Nakajima are known even today.

The situation has changed drastically since then, thanks to our improved understanding of strongly-coupled supersymmetric theories. Among others, we can count the class S construction in four dimensions initiated by [7], the determination of the chiral ring of the Coulomb branch in three dimensions starting with [8], and a new method to study $\mathcal{N}=1$ theories in six dimensions pioneered by [9]. Combining these developments, we believe there might be a chance that the physics might shed new lights on the mathematics of the structure of the E_8 instanton moduli spaces on ALE spaces.

The main target of our study in this paper is the 6d $\mathcal{N}=1$ theory on M5-branes probing the E_8 9-brane on the A-type ALE singularity $\mathbb{C}^2/\mathbb{Z}_k$. Such a system can be labeled by the M5-brane charge Q and the asymptotic holonomy $\rho : \mathbb{Z}_k \rightarrow E_8$. For some simple choices of ρ , the structure of the 6d theory on generic points on its tensor branch was already determined in [10, 11], which was further extended in [12–14].

Our first aim is to determine the tensor branch structure for an *arbitrary* choice of the asymptotic holonomy ρ . We give a complete algorithm determining the gauge group and the matter content in terms of ρ . Along the way, we encounter a subtle feature that there are two distinct ways to gauge $\mathfrak{su}(2n+8)$ symmetry of $\mathfrak{so}(4n+16)$ flavor symmetry of an $\mathfrak{usp}(2n)$ gauge theory with $N_f = 2n+8$ flavors, due to the fact that the outer automorphism of $\mathfrak{so}(4n+16)$ is *not* a symmetry of the latter gauge theory.

The Higgs branch $\mathcal{M}_{Q,\rho}$ of our theory $\mathcal{T}_{Q,\rho}^{6d}$ is not directly the instanton moduli space. In particular, $\mathcal{M}_{Q,\rho}$ has an action of $SU(k)$, which we do not expect for the instanton moduli space. Rather, by a small generalization of the argument in [15], we see that the E_8 instanton moduli space $\mathcal{M}_{Q,\rho,\xi}^{\text{inst}}$ of charge Q and asymptotic holonomy ρ on the ALE space $\widetilde{\mathbb{C}^2/\mathbb{Z}_k}$ is given by

$$\mathcal{M}_{Q,\rho,\xi}^{\text{inst}} = (\mathcal{M}_{Q,\rho} \times \mathcal{O}_\xi) // SU(k) \tag{1.1}$$

where $\xi = (\xi_{\mathbb{C}}, \xi_{\mathbb{R}}) \in \mathfrak{su}(k) \otimes (\mathbb{C} \oplus \mathbb{R})$ is an element in the Cartan of $\mathfrak{su}(k)$ tensored by \mathbb{R}^3 specifying the hyperkähler deformation parameter of the ALE space, \mathcal{O}_ξ is the orbit of $\xi_{\mathbb{C}}$ in $\mathfrak{su}(k)_{\mathbb{C}}$ with the hyperkähler metric specified by $\xi_{\mathbb{R}}$ as in [16], and the symbol $///$ denotes the hyperkähler quotient construction. This means that the space $\mathcal{M}_{Q,\rho}$ knows the structure of the instanton moduli on the ALE space for arbitrary deformation parameter ξ . The existence of such a generating space was conjectured by one of the authors in [17], based on a study of $SO(8)$ instantons on the ALE spaces.

We then study the 4d theory which arises from the T^2 compactification of the 6d theory as in [15]. We find that they always correspond to a class S theory of type A, given by a sphere with three punctures. The 3d mirror of its S^1 compactification is a star-shaped quiver, whose structure can be deduced from the class S description by the methods of [18]. We find that they have the form of an *over-extended* E_8 quiver. In 3d,

the relation (1.1) can be physically implemented by realizing \mathcal{O}_ξ as the Coulomb branch of the $T[\text{SU}(k)]$ theory. Using this, we will find that $\mathcal{M}_{Q,\rho,\xi}^{\text{inst}}$ is the Higgs branch of an affine E_8 quiver where ξ is now the mass parameter of an $\text{SU}(k)$ flavor symmetry. For $\xi = 0$ this was already conjectured by mathematicians [19, 20] and by physicists [21, 22].

Organization of the paper. The rest of the paper is organized as follows. We start by recalling the geometric data characterizing our system in section 2. Then in section 3, we provide the algorithm determining the 6d quiver theory in terms of the asymptotic holonomy. In section 4, we discuss its dimensional reduction to 5d, 4d and 3d in turn. In 5d and 4d, we translate the Kac labels to the three Young diagrams characterizing the brane web and the class S description. In 3d, we give the star-shaped quiver. Finally in section 5, we provide many examples illustrating our discussions.

Accompanying Mathematica file. The paper comes with a Mathematica file which implements the algorithm to produce the 6d quiver given the asymptotic E_8 holonomy. In addition, it allows the user to determine the 4d class S theory, and compute the anomalies from three different methods, namely the 6d field theory, the M-theoretic inflow, and the 4d class S technique.

Summary of notations.

- The asymptotic holonomy $\rho : \mathbb{Z}_k \rightarrow E_8$ is given by an element $\mathbf{w} \in \mathfrak{e}_8$ in the Cartan subalgebra, or equivalently in terms of the Kac label

$$\underline{n} := \begin{matrix} & & & & n_{3'} & & & & \\ & & & & & & & & \\ n_1 & n_2 & n_3 & n_4 & n_5 & n_6 & n_{4'} & n_{2'} & \end{matrix}, \tag{1.2}$$

a set of non-negative integers arranged on the affine E_8 Dynkin diagram. For more details, see section 2.1.

- We have closely related quantities N_{inst} , N_3 , N_S , N_6 , and Q , which are all essentially the number of M5-branes or equivalently the instanton charge on the ALE space. They all increase by one when we add one M5-brane to the system. Their constant parts are however different. We could have used just one out of them, but any choice would make at least one of the formulas quite unseemly. We therefore decided to keep them and provide a summary here.

- The integer N_{inst} is defined in terms of the instanton number as

$$\int_{\widetilde{\mathbb{C}^2/\Gamma}} \text{tr } F \wedge F \propto N_{\text{inst}} - \frac{\langle \mathbf{w}, \mathbf{w} \rangle}{2k}. \tag{1.3}$$

See (2.6) for details.

- Another integer N_3 satisfies $N_3 = N_{\text{inst}} - k$, see (3.14), (4.18). This is useful to parameterize the ranks of groups in the 3d quiver, see (4.3).
- Another integer N_S defined by $N_S = N_3 + n_1 + \dots + n_6$ is useful to parameterize the class S data, see (3.5).

- The integer N_6 is the number of tensors of the 6d quiver. The difference between N_S and N_6 is determined by the Kac label and is described in the algorithm in section 3.2.
- A rational number Q is the M5-charge which appears in the inflow computation, and satisfies

$$Q = N_{\text{inst}} - \frac{\langle \mathbf{w}, \mathbf{w} \rangle}{2k} - \frac{1}{2} \left(k - \frac{1}{k} \right), \tag{1.4}$$

see (3.12).

2 Geometric preliminaries

2.1 Topological data of the instanton configuration

Here we recall the topological data necessary to specify a G -instanton on \mathbb{C}^2/Γ or its resolution $\widetilde{\mathbb{C}^2}/\Gamma$, where $\Gamma \in \text{SU}(2)$.

On \mathbb{C}^2/Γ , we first need to specify the holonomy at the origin and at the infinity. They determine the representation $\rho_{0,\infty} : \Gamma \rightarrow G$, which we consider as a linear action on the complexified adjoint representation \mathfrak{g} .

On $\widetilde{\mathbb{C}^2}/\Gamma$, we specify the holonomy at infinity ρ_∞ . In addition, we need to specify the class in $H^2(\widetilde{\mathbb{C}^2}/\Gamma, \pi_1(G))$. This is the first Chern class when $G = \text{U}(N)$ and the second Stiefel-Whitney class when $G = \text{SO}(N)$.

Finally we need to specify the instanton number, defined as the integral of $\text{tr } F \wedge F$ over the ALE space. Unless otherwise mentioned, we normalize the trace so that the instanton on \mathbb{R}^4 of the smallest positive instanton number satisfies

$$\int \text{tr } F \wedge F = 1. \tag{2.1}$$

On the ALE space, the instanton number is in general fractional.

Our main interest lies in the case $G = E_8$ and $\Gamma = \mathbb{Z}_k$. Since $\pi_1(E_8)$ is trivial, we do not have to specify the class in H^2 .

A holonomy $\rho : \mathbb{Z}_k \rightarrow E_8$ can be nicely encoded by its Kac label

$$\underline{n} := \begin{matrix} & & & & n_{3'} & & & & \\ & & & & & & & & \\ n_1 & n_2 & n_3 & n_4 & n_5 & n_6 & n_{4'} & n_{2'} & \end{matrix}. \tag{2.2}$$

introduced in §8.6 of Kac’s textbook [23]. Let us quickly recall how it works. Let the image g of the generator of \mathbb{Z}_k in E_8 be

$$g = e^{2\pi i \mathbf{w}/k} \in E_8 \tag{2.3}$$

where

$$\mathbf{w} = \sum_{i \neq 0} n_i \mathbf{w}_i \in \mathfrak{e}_8. \tag{2.4}$$

where \mathbf{w}_i are the fundamental weights of E_8 . Since g is of order k , n_i are integers. We define n_0 so that $\sum d_i n_i = k$, where the Dynkin marks \underline{d} are given by

$$\underline{d} = \begin{array}{cccccccc} & & & & 3 & & & \\ & & & & | & & & \\ & & & & 1 & 2 & 3 & 4 & 5 & 6 & 4 & 2 \end{array}. \quad (2.5)$$

It is known that by the Weyl reflections and the shifts, we can arrange $n_i \geq 0$ for all i and then the result is unique. This is the Kac label of the holonomy.

The subalgebra of \mathfrak{e}_8 left unbroken by the holonomy ρ can be easily read off from its Kac label. Namely, it is given by the subalgebra corresponding to the nodes i of the Dynkin diagram where $n_i = 0$, together with an Abelian subalgebra making the total rank 8.

On $\widetilde{\mathbb{C}^2/\mathbb{Z}_k}$, the instanton number modulo one is given by the classical Chern-Simons invariant evaluated on S^3/\mathbb{Z}_k at infinity. One way to compute it is to introduce coordinates on S^3/\mathbb{Z}_k using polar coordinates θ, ϕ on S^2 and the angle ψ along the S^1 fiber. The connection itself is $\propto \mathbf{w}(d\psi + \dots)$. One finds that

$$\int \text{tr } F \wedge F = N_{\text{inst}} - \frac{\langle \mathbf{w}, \mathbf{w} \rangle}{2k} \quad (2.6)$$

where N_{inst} is an integer. The hyperkähler dimension of the moduli space is given by the formula

$$\dim_{\mathbb{H}} \mathcal{M}_{\widetilde{\mathbb{C}^2/\Gamma, \rho_\infty}} = 30N_{\text{inst}} - \langle \mathbf{w}, \boldsymbol{\rho} \rangle. \quad (2.7)$$

2.2 Dimension of the instanton moduli space

In this subsection we derive the formula (2.7) of the dimension of the moduli space. Those readers who trust the authors can skip this subsection. This computation is of course not new. It is provided here to make this paper more self-contained.

The basic tool is the Atiyah-Patodi-Singer index theorem. Its explicit form on the orbifold of \mathbb{C}^2 was worked out e.g. in [24] for $\Gamma \subset \text{U}(2)$. Here we quote the form used in Kronheimer-Nakajima [2] for $\Gamma \subset \text{SU}(2)$. The formula for the orbifold is:

$$\dim_{\mathbb{H}} \mathcal{M}_{\mathbb{C}^2/\Gamma, \rho_\infty, \rho_0} = h^\vee(G) \left(\int \text{tr } F \wedge F \right) + \frac{1}{2|\Gamma|} \sum_{\gamma \neq e} \frac{\chi_{\rho_\infty}(\gamma)}{2 - \chi_Q(\gamma)} - \frac{1}{2|\Gamma|} \sum_{\gamma \neq e} \frac{\chi_{\rho_0}(\gamma)}{2 - \chi_Q(\gamma)}. \quad (2.8)$$

Here, $h^\vee(G)$ is the dual Coxeter number of G , and the second and the third terms are the contributions from the η invariant of S^3/Γ at the asymptotic infinity and at the origin, respectively, and Q is the standard two-dimensional representation of Γ from the defining embedding $\Gamma \subset \text{SU}(2)$,

On the ALE space $\widetilde{\mathbb{C}^2/\Gamma}$, we have:

$$\dim_{\mathbb{H}} \mathcal{M}_{\widetilde{\mathbb{C}^2/\Gamma, \rho_\infty}} = h^\vee(G) \left(\int \text{tr } F \wedge F \right) + \frac{1}{2|\Gamma|} \sum_{\gamma \neq e} \frac{\chi_{\rho_\infty}(\gamma)}{2 - \chi_Q(\gamma)} - \frac{1}{24} \dim G \chi_\Gamma \quad (2.9)$$

where the quantity

$$\chi_\Gamma := r_\Gamma + 1 - \frac{1}{|\Gamma|} \tag{2.10}$$

is the Euler number of $\widetilde{\mathbb{C}^2/\Gamma}$ as defined by the integral of the Pontrjagin density. Furthermore,

$$\frac{1}{24} \left(r_\Gamma + 1 - \frac{1}{|\Gamma|} \right) = \frac{1}{2|\Gamma|} \sum_{\gamma \neq e} \frac{1}{2 - \chi_Q(\gamma)}, \tag{2.11}$$

reflecting the fact that if the holonomy at the origin of an instanton on \mathbb{C}^2/Γ is trivial, we can resolve/deform the instanton and the ALE at the same time to be on $\widetilde{\mathbb{C}^2/\Gamma}$. In the end, we find the formula

$$\dim_{\mathbb{H}} \mathcal{M}_{\widetilde{\mathbb{C}^2/\Gamma, \rho_\infty} = h^\vee(G) \left(\int \text{tr } F \wedge F \right) + \Delta\eta, \tag{2.12}$$

where

$$\Delta\eta := \frac{1}{2|\Gamma|} \sum_{\gamma \neq e} \frac{(\chi_{\rho_\infty}(\gamma) - \dim \mathfrak{g})}{2 - \chi_Q(\gamma)} \tag{2.13}$$

Let us evaluate this formula when $G = E_8$ with the holonomy ρ_∞ specified by $g = e^{2\pi i \mathbf{w}/k}$ with the Kac label \underline{n} . The eta invariant is

$$\Delta\eta = \frac{1}{2|\Gamma|} \sum_{\gamma \neq e} \frac{(\chi_{\rho_\infty}(\gamma) - \dim \mathfrak{g})}{2 - \chi_Q(\gamma)} = \frac{1}{2k} \sum_{\alpha: \text{all roots}} \sum_{\gamma \neq 1} \frac{\chi_{\langle \mathbf{w}, \alpha \rangle}(\gamma) - 1}{2 - \chi_Q(\gamma)} \tag{2.14}$$

where

$$\chi_a(g^j) = e^{2\pi i a j/k}, \quad \chi_Q(g^j) = 2 \cos 2\pi j/k. \tag{2.15}$$

Now, we note

$$\frac{1}{2k} \sum_{j=1}^{k-1} \frac{\chi_a(g^j) - 1}{2 - \chi_Q(g^j)} = -\frac{a(a-k)}{4k} \tag{2.16}$$

for $a = 0, 1, \dots, k$. Then

$$\Delta\eta = 2 \sum_{\alpha: \text{positive roots}} -\frac{\langle \alpha, \mathbf{w} \rangle (\langle \alpha, \mathbf{w} \rangle - k)}{4k} \tag{2.17}$$

since $0 \leq \langle \alpha, \mathbf{w} \rangle \leq k$ for positive roots α . Now we use

$$\sum_{\alpha: \text{positive roots}} \alpha = 2\rho, \quad \sum_{\alpha: \text{positive roots}} \langle \mathbf{v}_1, \alpha \rangle \langle \alpha, \mathbf{v}_2 \rangle = h^\vee \langle \mathbf{v}_1, \mathbf{v}_2 \rangle \tag{2.18}$$

and find

$$\Delta\eta = \frac{h^\vee}{2k} \langle \mathbf{w}, \mathbf{w} \rangle - \langle \mathbf{w}, \rho \rangle. \tag{2.19}$$

To compute the dimension, we now plug in to (2.12) the formula for $\Delta\eta$ found just above and the formula for the instanton number (2.6). The term proportional to $\langle \mathbf{w}, \mathbf{w} \rangle$ cancels out, and we indeed have the desired result (2.7).

3 Six-dimensional description

After these geometrical preliminaries, we move on to the field theoretical analysis. We start with the six-dimensional quiver descriptions. As already mentioned in the introduction, for various simple choices of ρ , the six-dimensional quivers were already determined in [10–14]. By a series of trials and errors, and following the principle that the quiver should be determined in terms of the Kac label, the authors found the following algorithm.

3.1 The general structure of the quiver

Our 6d SCFT on the generic points on its tensor branch consists of a collection of N_6 tensors, corresponding to a linear quiver of the form

$$G_1 \times \text{SU}(m_2) \times \text{SU}(m_3) \times \cdots \times \text{SU}(m_{N_6}) \times [\text{SU}(k)] \tag{3.1}$$

where G_1 is on the -1 curve, the rest is on -2 curves, and the final $\text{SU}(k)$ is a flavor symmetry. In the notation of [11], we have

$$\begin{array}{ccccccc} G_1 & \text{su}(m_2) & \text{su}(m_3) & \cdots & \text{su}(m_{N_6}) & & \\ 1 & 2 & 2 & \cdots & 2 & & [\text{SU}(k)]. \end{array} \tag{3.2}$$

Below, we slightly abuse the notation and refer by G_1 the combination of the group and the non-fundamental hypermultiplets on the -1 curve. The choices are:

- $G_1 = \text{USp}(m_1)$,
- $G_1 = \text{SU}(m_1)$ with an antisymmetric hyper, or
- $G_1 = \text{SU}(m_1 = 6)$ with a rank 3 antisymmetric half-hyper.

We consider the rank 1 E-string theory as $\text{USp}(0)$, and furthermore, the rank 2 E-string theory is considered as a $\text{USp}(0)$ connecting to an $\text{SU}(1)$ group.

We have $m_1 \leq m_2 \leq \cdots \leq m_{N_6}$, and we define a_1, \dots, a_9 by

$$a_s = \#\{i \mid m_{i+1} - m_i = s\}. \tag{3.3}$$

We can reconstruct the whole of m_i from m_1, N_6 and a_1, \dots, a_9 . For example, when the quiver is

$$\text{SU}(3) \times \text{SU}(9) \times \text{SU}(13) \times \text{SU}(17) \times \text{SU}(18) \times \text{SU}(19) \times \text{SU}(20) \tag{3.4}$$

we have $a_8 = a_7 = a_5 = a_3 = a_2 = 0, a_6 = 1, a_4 = 2, a_1 = 3$.

There are bifundamentals between two consecutive groups in the quiver, and finally fundamental hypers are added such that each group is anomaly free, that is:

- $N_f = 2N$ for $\text{SU}(N)$,
- $N_f = N + 8$ for $\text{USp}(N)$ or $\text{SU}(N)$ with an antisymmetric hyper, and
- $N_f = 15$ for $\text{SU}(6)$ with a rank 3 antisymmetric half-hyper.

3.2 The algorithm

Now we present the algorithm to determine the structure of the quiver given the Kac label \underline{n} and the number N_6 of the groups. Along the way, we also define the quantity N_S which will be used in the following. We will also need the quantity

$$N_3 = N_S - n_1 - n_2 - n_3 - n_4 - n_5 - n_6. \tag{3.5}$$

The algorithm is implemented in the accompanying Mathematica file, so that the reader can easily try it around.

In general we have

$$a_i = n_i \quad \text{for } i = 1, 2, 3, 4, 5, 6. \tag{3.6}$$

To specify $a_{7,8,9}$, we need to consider various cases as summarized below:

$$\left\{ \begin{array}{l} n'_4 \geq n'_3 \quad \longrightarrow \left\{ \begin{array}{l} n'_4 - n'_3 = \text{even} \quad \longrightarrow \text{Case 1,} \\ n'_4 - n'_3 = \text{odd} \quad \longrightarrow \text{Case 2,} \end{array} \right. \\ \\ n'_3 \geq n'_4 \quad \longrightarrow \left\{ \begin{array}{l} n'_2 < (n'_3 - n'_4)/2 \quad \longrightarrow \text{Case 4,} \\ n'_2 \geq (n'_3 - n'_4)/2 \quad \longrightarrow \left\{ \begin{array}{l} n'_3 - n'_4 = \text{odd} \quad \longrightarrow \text{Case 3,} \\ n'_3 - n'_4 = \text{even} \quad \longrightarrow \text{Case 5.} \end{array} \right. \end{array} \right. \end{array} \right.$$

For each case, the output of the algorithm is $(a_{7,8,9}, G_1, N_S)$ as shown below:¹

1. $n'_4 \geq n'_3, n'_4 - n'_3 = \text{even}$:
 - $a_7 = n'_3, a_8 = \frac{n'_4 - n'_3}{2}, a_9 = 0$.
 - $G_1 = \text{USp}(2n'_2)$.
 - $N_S = N_6 - \frac{n'_4 + n'_3}{2}$.
2. $n'_4 \geq n'_3 + 1, n'_4 - n'_3 = \text{odd}$:
 - $a_7 = n'_3, a_8 = \frac{n'_4 - n'_3 - 1}{2}, a_9 = 0$.
 - $G_1 = \text{SU}(2n'_2 + 4)$ group with an antisymmetric hyper.
 - $N_S = N_6 - \frac{n'_4 + n'_3 - 1}{2}$.
3. $n'_3 \geq n'_4 + 1, n'_3 - n'_4 = \text{odd}, n'_2 \geq \frac{n'_3 - n'_4 - 1}{2}$:
 - $a_7 = n'_4, a_8 = \frac{n'_3 - n'_4 - 1}{2}, a_9 = 0$.
 - $G_1 = \text{SU}(2n'_2 + n'_4 - n'_3 + 4)$ group with an antisymmetric hyper.
 - $N_S = N_6 - \frac{n'_4 + n'_3 - 1}{2}$.

¹When more than two cases apply to the same Kac label, they produce the same quiver.

4. $n'_3 > n'_4 + 2n'_2 + \ell, n'_3 - n'_4 - 2n'_2 = 3x + \ell, x \in \mathbb{Z}, \ell = 0, 1, 2$:
 - $a_7 = n'_4, a_8 = n'_2, a_9 = \frac{n'_3 - n'_4 - 2n'_2 - \ell}{3}$.
 - G_1 is
 - empty for $\ell = 0$, in which case this node corresponds to a rank-1 E-string,
 - SU(3) for $\ell = 1$,
 - SU(6) with a half-hyper in the rank 3 antisymmetric for $\ell = 2$.
 - $N_S = N_6 - \frac{n'_3 + 2n'_4 + n'_2 - \ell}{3}$.

5. $n'_3 \geq n'_4, n'_3 - n'_4 = \text{even}, n'_2 \geq \frac{n'_3 - n'_4}{2}$:
 - $a_7 = n'_4, a_8 = \frac{n'_3 - n'_4}{2}, a_9 = 0$.
 - $G_1 = \text{USp}(2n'_2 + n'_4 - n'_3)$.
 - $N_S = N_6 - \frac{n'_4 + n'_3}{2}$.

3.3 A subtlety concerning the 6d θ angle

Note that the quivers produced in Case 5 are the same ones as the ones produced by Case 1, as far as the data we described so far are concerned. This is perfectly fine when $n'_4 = n'_3$, since in this case we are just applying the different cases to the same Kac label. However, when $n'_4 \neq n'_3$, or equivalently when $a_8 \neq 0$, the resulting quivers should however be subtly different, since e.g. they reduce to different 4d class S theories and 3d star-shaped quivers. We argue that the difference between them is how one embeds the SU(2N + 8) group into the SO(4N + 16) global symmetry group of USp(2N).

A relatively simple case is the following. Let us first consider the cases when $n'_4 = 2, n'_3 = 0, n'_2 = 0$ versus $n'_4 = 0, n'_3 = 2, n'_2 = 1$, with the rest of labels being zero $n_{1,\dots,6} = 0$. Both theories have the form of a long SU(8) quiver gauging an SU(8) subgroup of the rank 1 E_8 theory. The two differ by the embedding of SU(8) inside E_8 and in fact have different global symmetries. To see this, consider embedding SU(8) inside $\text{SO}(16) \subset E_8$. The adjoint of E_8 decomposes under its SO(16) maximal subgroup as **248** \rightarrow **120** + **128**. Now consider decomposing SO(16) to its U(1) \times SU(8) maximal subgroup. Under this embedding the spinors of SO(16) decompose to the rank x antisymmetric tensors of SU(8) for $x = 0, 2, 4, 6, 8$ for one spinor and $x = 1, 3, 5, 7$ for the other. However only one spinor appears in the adjoint of E_8 , and therefore there are two different embedding of SU(8) inside E_8 . In one of them the **128** contains gauge invariant contributions leading to the larger global symmetry.

The general case corresponds to the situation where SU(2N + 8) is embedded in SO(4N + 16). There is no distinction in the perturbative sector of the theory. However the theory possesses instanton strings. The ones for USp groups will be in a chiral spinor of the SO group and so will decompose differently depending on the embedding. This then leads to theories with distinct spectrum of string excitations. Also note that this only occurs if the entire SO symmetry is gauged leaving only a U(1) commutant. If

we gauge an $SU(x) \subset SO(2x) \subset SO(4N + 16)$ with $x < 2N + 8$, then the chiral spinor of $SO(4N + 16)$ decomposes to non-chiral spinors of $SO(2x)$ and therefore there is a single embedding. This agrees with the fact that the cases coincide when $a_8 = 0$.

We can understand this distinction from the existence of the discrete θ angle in 6d, due to the fact that $\pi_5(\text{USp}(2N))_5 = \mathbb{Z}_2$. Suppose now that the USp group has $2n$ half-hypermultiplets in the fundamental. Classically it has an $O(2n)$ flavor symmetry, but the parity part flips the discrete theta angle. Therefore the flavor symmetry is actually $\mathfrak{so}(2n)$. The two embeddings of $\mathfrak{su}(n)$ into $\mathfrak{so}(2n)$ are related exactly by the parity part of $O(2n)$, and therefore are inequivalent. The F-theoretical interpretation of these two inequivalent embeddings seems to be unknown. It would be interesting to work it out.²

Note that an analogous phenomenon exists in $5d$, where given a pure USp group there are two distinct $5d$ SCFTs associated with this theory differing by the instanton spectrum of the $5d$ gauge theory. This is related to the existence of a \mathbb{Z}_2 valued θ angle originating from the fact that $\pi_4(\text{USp}(2N))_4 = \mathbb{Z}_2$.

3.4 Anomalies and the inflow

The anomaly of these 6d SCFTs can be computed from their quiver description using the technique of [25, 26]. We should be able to match it to the anomaly computed from the inflow using the M-theory description.

The inflow computations of M5-branes probing the E_8 end-of-the-world brane and of M5-branes probing the $\mathbb{C}^2/\mathbb{Z}_k$ singularity was given in [27] and in an appendix of [25], respectively. We can combine the two computations into one and one finds the following contribution to the anomaly, excluding the most subtle contribution from the codimension-5 singularity where the $\mathbb{C}^2/\mathbb{Z}_k$ singularity hits the end-of-the-world brane:

$$I_{\text{inflow}}^{\text{naive}}(Q) = \frac{Q^3 k^2}{6} c_2(R)^2 - \frac{Q^2 k}{2} c_2(R) I_4 + Q \left(\frac{1}{2} I_4^2 - I_8 \right) + (I_4 - Qk c_2(R)) J_4 - \frac{1}{2} I^{\text{vec}}(\text{SU}(k)) \quad (3.7)$$

where Q is the M5-charge of the configuration,

$$I_8 = \frac{1}{48} (p_2(N) + p_2(T) - \frac{1}{4} (p_1(N) - p_1(T))^2), \quad (3.8)$$

$$I_4 = \frac{1}{4} (p_1(T) - 2c_2(R)), \quad (3.9)$$

$$J_4 = \frac{1}{48} (k - \frac{1}{k}) (4c_2(R) + p_1(T)) + \frac{1}{4} \text{tr} F_{\text{SU}(k)}^2. \quad (3.10)$$

Here I_8 comes from the M-theory interaction $\int C \wedge I_8$, I_4 appears in the boundary condition $G = I_4$ at the E_8 wall, and J_4 is the interaction on the $\mathbb{C}^2/\mathbb{Z}_k$ singular locus $\int C \wedge J_4$. In this section the normalization of tr is as in [27].

Let \underline{n} be the Kac label, and let $\mathbf{w} = \sum \mathbf{w}_i n_i$ be the corresponding weight vector. By performing computations for many choices of \underline{n} , we find that

$$I_{\text{quiver}}(\underline{n}, N_3) = I_{\text{inflow}}^{\text{naive}}(Q) + c(\underline{n}) \quad (3.11)$$

²The authors thank D. R. Morrison for the correspondence on this point.

where $c(\underline{n})$ is a constant depending on the Kac label \underline{n} but independent of N_3 and

$$Q = N_3 + \frac{1}{2} \left(k + \frac{1}{k} - \frac{\langle \mathbf{w}, \mathbf{w} \rangle}{k} \right). \quad (3.12)$$

Recall that the instanton number as defined by the integral of $\text{tr } F \wedge F$ was given by

$$\int \text{tr } F \wedge F = N_{\text{inst}} - \frac{\langle \mathbf{w}, \mathbf{w} \rangle}{2k}, \quad (3.13)$$

see (2.6), and that $k - 1/k$ is the Euler number of $\widetilde{\mathbb{C}^2/\Gamma}$, or equivalently of the integral of $-p_1/4$ there, see (2.10). Then, assuming that

$$N_{\text{inst}} = N_3 + k, \quad (3.14)$$

we can rewrite the effective M5-brane charge Q as

$$Q = \int \text{tr } F \wedge F + \int_{\widetilde{\mathbb{C}^2/\mathbb{Z}_k}} \frac{p_1}{4} \quad (3.15)$$

which is what we expect from the curvature coupling on the E8 end-of-the-world brane.

The authors made a guess of the formula for $c(\underline{n})$ by trial and error. It has the form

$$c(\underline{n}) = \frac{1}{k} (P_0(\underline{n}) + P_2(\underline{n}) + P_4(\underline{n}) + P_6(\underline{n})) + \frac{1}{2} I_{\text{free vector}} \quad (3.16)$$

where

$$I_{\text{free vector}} = \frac{1}{5760} (-240c_2(R)^2 - 120c_2(R)p_1(T) - 7p_1(T)^2 + 4p_2(T)) \quad (3.17)$$

is the anomaly polynomial of a free vector multiplet and $P_i(\underline{n})$ is a homogeneous polynomial of n_i 's of degree i . Those polynomials are identified as

$$P_0 = \frac{1}{384} (-88c_2(R)^2 + 32c_2(R)p_1(T) - 5p_1(T)^2 + 4p_2(T)) \quad (3.18)$$

$$P_2 = \frac{1}{11520} k^2 (2512c_2(R)^2 - 760c_2(R)p_1(T) + 157p_1(T)^2 - 124p_2(T)) \\ + \frac{1}{5760} (15\langle \mathbf{w}, \mathbf{w} \rangle - k\langle \mathbf{w}, \boldsymbol{\rho} \rangle) (112c_2(R)^2 - 40c_2(R)p_1(T) + 7p_1(T)^2 - 4p_2(T)) \quad (3.19)$$

$$P_4 = -\frac{1}{288} \left(9\langle \mathbf{w}, \mathbf{w} \rangle^2 + 15k^2\langle \mathbf{w}, \mathbf{w} \rangle - 2k^4 - k \sum_{\alpha \in \Delta^+} \langle \mathbf{w}, \boldsymbol{\alpha} \rangle^3 \right) (4c_2(R)^2 - c_2(R)p_1(T)) \quad (3.20)$$

$$P_6 = \frac{1}{240} \left(5\langle \mathbf{w}, \mathbf{w} \rangle^3 + 15k^2\langle \mathbf{w}, \mathbf{w} \rangle^2 - 5k^4\langle \mathbf{w}, \mathbf{w} \rangle + k^6 - k \sum_{\alpha \in \Delta^+} \langle \mathbf{w}, \boldsymbol{\alpha} \rangle^5 \right) c_2(R)^2, \quad (3.21)$$

where Δ^+ is the set of positive roots of E_8 . The authors have not been able to determine how this formula come from the correct anomaly inflow calculation. It would be interesting to understand it.

4 Lower dimensional incarnations

4.1 Five-dimensional brane-web description

We can reduce the 6d theory on a circle to 5d. Roughly speaking, there are two different types of reductions. For example, starting from the E-string theory, one can obtain $SU(2)$ theory with eight flavors in one way, or the 5d SCFT with E_8 flavor symmetry in the other way.

First reduction. Keeping the radius of the circle non-zero the low-energy 5d theory is sometimes a 5d gauge theory. Specifically, the class of 6d theories we are considering can be realized by a brane construction involving a system of NS5-branes and D6-branes in the presence of an $O8^-$ -plane [28, 29]. Performing T-duality on this system results in a brane configuration involving NS5-branes and D6-branes in the presence of an $O8^-$ -plane. Alternatively, the system can also be described as D4-branes immersed in an $O8^-$ -plane and D8-branes, in the presence of a $\mathbb{C}^2/\mathbb{Z}_k$ singularity [30].

Either way, the system can sometimes be deformed so as to describe a 5d gauge theory. Specifically, when compactifying we have a choice of the value of the radius as well as the freedom to turn on holonomies for the global symmetries. These then become mass parameters in the 5d theory. In specific ranges of these parameters the 6d theory may flow at low-energy to a 5d quiver gauge theory with the coupling constants of the gauge theory identified with the mass deformations. In general, a given 6d SCFT may have several different low-energy 5d gauge theory descriptions depending on the specific deformations used. Various 5d descriptions of 6d theories, including the type we are interested in, were studied in [12, 14, 31, 32]. We will not consider this problem here.

Second reduction. Instead we shall take the limit of zero radius. In this case we argue that the 6d theory flows in the IR to a 5d SCFT. Furthermore, we claim that the 5d SCFT can be readily described in terms of the integer N and the Kac label \underline{n} . To find the 5d theory, we first write down the 6d quiver following the algorithm presented in the last section.³ We realize this 6d quiver in type IIA using $O8$ -planes, D8-branes, D6-branes and NS5-branes as in [28, 29]. We then compactify it on S^1 , T-dualize it to type IIB, and manipulate the branes. We will detail the procedure in slightly more detail below.

The result can be conveniently represented by a brane web, which has a star shape form with a group of $(1,0)$, $(0,1)$ and $(1,1)$ 5-branes all intersecting at a point. The 5-branes end on the appropriate 7-branes where some collection of 5-branes end on the same 7-brane. Specifying the configuration then is done by giving the distribution of 5-branes on the 7-branes. This is conveniently done by a Young diagram where each column represents a 7-brane, and the number of boxes in it represents the number of 5-branes ending on it.

³Purely field theoretically, the 6d quiver only contains the information on the low energy limit on the generic points on the tensor branch of a given 6d SCFT. Therefore further manipulations of the quiver such as dimensional reductions are not guaranteed to tell every detail of the original 6d SCFT. What we do is, instead, to realize the quiver using branes, and apply string dualities. This way we can keep all the ultraviolet information required in the process.

The three Young diagrams for the SCFTs we are considering are given by:

$$\begin{aligned}
 Y_1 = & (N_S - n_6, \\
 & N_S - n_6 - n_5, \\
 & N_S - n_6 - n_5 - n_4, \\
 & N_S - n_6 - n_5 - n_4 - n_3, \\
 & N_S - n_6 - n_5 - n_4 - n_3 - n_2, \\
 & N_S - n_6 - n_5 - n_4 - n_3 - n_2 - n_1, \\
 & 1^k), \\
 Y_2 = & (2N_S + 2n_{4'} + n_{2'} + n_{3'}, \\
 & 2N_S + n_{4'} + n_{2'} + n_{3'}, \\
 & 2N_S + n_{4'} + n_{3'}), \\
 Y_3 = & (3N_S + 2n_{4'} + n_{2'} + 2n_{3'}, \\
 & 3N_S + 2n_{4'} + n_{2'} + n_{3'}).
 \end{aligned}
 \tag{4.1}$$

More detail of the second reduction. For cases 1, 2 and 3, these results can be derived using the standard techniques. But there are some issues for cases 4 and 5. Case 4 naively does not have a brane construction of the type considered in [29] so this procedure appears to be inapplicable in this case. However, a conjecture for the 5d theories that lift to these types of 6d SCFTs was given in [12, 14], and we can use this conjecture to fill in this step for case 4.

This leaves case 5. We can ask how does the 6d θ angle appears in the brane construction. In fact a similar issue arises in the analogue 5d system: D5-branes suspended between NS5-branes in the presence of an $O7^-$ -plane. In that case it was observed by [33] that accounting for the 5d θ angle seems to necessitate the introduction of two variants of the $O7^-$ -plane, where one is an $SL(2, \mathbb{Z})$ T-transform of the other. This in particular means that they differ by their decomposition into a pair of 7-branes. Note that the distinction between the two cases vanishes when there are D7-branes on the $O7^-$ -plane. This becomes clear after we decompose the $O7^-$ -plane into 7-branes which can be moved through the monodromy lines of the 7-branes which will change them by a T-transformation. This of course agrees with the unphysical nature of the 5d θ angle once flavors are present. There should be a similar distinction for the $O8^-$ -plane, and so can account for the apparent 6d θ angle we observe. We will not pursue this here.

However once we perform T-duality we end with a system with two $O7^-$ -planes, and we expect that we can accommodate this in the observed difference in $O7^-$ -planes. We have a discrete choice for each $O7^-$ -plane leading to four possibilities. However we are free to perform a global T-transformation. Since all the external branes are D7-branes, this will lead us to the same system, save for changing the types of both orientifolds. Thus we conclude that there are only two distinct choices: the same or differing types. These cases are expected to differ only when there are no 7-branes on the $O7^-$ -planes, and thus no D8-brane on the original $O8^-$ -plane. This exactly agrees with the two cases, which

coincide once $a_8 = 0$. We indeed find different 5d theories for these two choices, where the former is identified with case 1 while the latter with case 5. In this manner we can apply this procedure also to case 5.

4.2 Four-dimensional class S description

We can compactify on an additional circle to 4d. Using the results of [34], it is straightforward to write the 4d theory. It is just an A type class S theory given by the same set of Young diagrams as the 5d description, given above in (4.1).

In fact it is also possible to motivate this class S description with the Young diagrams (4.1) directly from the 4d description, and then use the preceding discussion to connect the 6d quiver data to the Kac labels. We start from the observation that the class S theory whose Young diagrams are (4.1) can be thought of as generated by modifying the Young diagrams of the rank N E_8 theory, which is given by a class S theory of type $SU(6N)$ with Young diagrams $Y_1 = (N^6)$, $Y_2 = (2N^3)$, $Y_3 = (3N^2)$.

First the 4d theory needs to have the $SU(k)$ global symmetry, coming from the $\mathbb{C}^2/\mathbb{Z}_k$ singularity. This is given by the k boxes attached to the Young diagram Y_1 of the E_8 theory. That this is the correct way to account for it can be seen by comparing anomalies. For the type of 6d theories we are considering, there is a result due to [15] that allows for the computations of the central charges of the 4d result of the compactification of the 6d theory from the anomaly polynomial of the latter. Furthermore the anomaly polynomial of the 6d theories of the type we considered was studied in [12]. When applied to our case we find that $k_{SU(k)}^{4d} = 2k + 12$ independent of the details of the Kac label. This agrees with the anomaly of the class S theory.

In addition to the $SU(k)$ we also have the commutant of the orbifold in E_8 as a global symmetry, which depends on the Kac labels. The E_8 global symmetry is accommodated by the Young diagram structure of the starting E_8 SCFT so it is natural to expect that modifying this will give the required global symmetry and take into account the Kac labels. The global symmetry which is manifest in the class S construction is $SU(2) \times SU(3) \times SU(6)$ which can be identified with the three legs of the affine Dynkin diagram. This becomes more apparent once we compactify to 3d and consider the mirror dual, which we consider more extensively in the next subsection.

The point is that we can associate a node in the legs of the affine E_8 Dynkin diagram roughly with the difference between neighboring columns. The central node can be associated with the difference between the sum of the first columns of the three Young diagrams and the total number of boxes in any of them. When that difference is zero, we get the E_8 theory. It is now natural to associate that difference to the Kac label of the corresponding node. By the Kac prescription, this ensures that we get the correct global symmetry. This leads to the conjectured form. There is one ambiguity in determining the total number of boxes which is related to the rank of the initial E_8 theory. This should be related to the number of tensors in 6d, but we need to determine the exact mapping. For this we use the relation outlined in the previous sections between the 6d and 4d theories.

We can perform various consistency checks of this proposal. One check is to compare anomalies. We already mentioned that these can be computed from the 6d anomaly poly-

The dimension of the Coulomb branch $\hat{\mathcal{M}}$ is then

$$\dim_{\mathbb{H}} \hat{\mathcal{M}} = 30(N_3 + k) - \langle \mathbf{w}, \boldsymbol{\rho} \rangle + \frac{k(k+1)}{2} - 1 \tag{4.9}$$

where $\mathbf{w} = \sum n_i \mathbf{w}_i$ is the Kac label as a weight vector and $\boldsymbol{\rho} = \sum_i \mathbf{w}_i$ is the Weyl vector.

The Coulomb branch $\hat{\mathcal{M}}$ of this system \hat{X} is closely related to the instanton moduli space $\mathcal{M}^{\text{inst}}$ on the ALE space $\widetilde{\mathbb{C}^2/\mathbb{Z}_k}$. To explain the relation, let us first recall that the resolution and deformation parameters of the ALE space can be specified by a parameter

$$\xi = (\xi_{\mathbb{C}}, \xi_{\mathbb{R}}) \in \mathfrak{su}(k) \otimes (\mathbb{C} \oplus \mathbb{R}) \tag{4.10}$$

which takes values in the Cartan of $\mathfrak{su}(k)$ tensored by \mathbb{R}^3 . We now need an auxiliary hyperkähler space \mathcal{O}_{ξ} , which is the $SU(k)_{\mathbb{C}}$ orbit of $\xi_{\mathbb{C}}$ in $\mathfrak{su}(k)$ with the hyperkähler metric specified by $\xi_{\mathbb{R}}$. Equivalently, \mathcal{O}_{ξ} is the Coulomb/Higgs branch of the $T[SU(k)]$ theory whose quiver realization is given by

$$T[SU(k)] = \bullet_1 - \bullet_2 - \cdots - \bullet_{k-1} - \blacksquare_k \tag{4.11}$$

where the rightmost square node is a flavor symmetry and ξ is the $SU(2)_R$ triplet of mass parameters associated to it.

We can now state the relation between $\hat{\mathcal{M}}$ and $\mathcal{M}^{\text{inst}}$ by slightly modifying an argument given in [15]:

$$\mathcal{M}^{\text{inst}} = (\hat{\mathcal{M}} \times \mathcal{O}_{\xi}) // SU(k). \tag{4.12}$$

This relation can be understood as follows. The resolution/deformation parameter ξ of the ALE space can be identified with the scalar vacuum expectation values of the 7d super $SU(k)$ Yang-Mills theory supported on the M-theory singularity $\mathbb{C}^2/\mathbb{Z}_k$. The 6d SCFT on the M5-branes at the intersection of the E_8 wall and the $\mathbb{C}^2/\mathbb{Z}_k$ singularity couples to this 7d super Yang-Mills, via the standard coupling where the triplet moment map field of the 6d theory is identified with the limiting value of the triplet of scalars of the 7d bulk. The resulting hyperkähler manifold is then given by the hyperkähler reduction as in (4.12).

Now, our system \hat{X} can also be written using the theory \tilde{X}

$$\tilde{X} := \blacksquare_k - \bullet_{\tilde{N}_1} - \bullet_{\tilde{N}_2} - \bullet_{\tilde{N}_3} - \bullet_{\tilde{N}_4} - \bullet_{\tilde{N}_5} - \bullet_{\tilde{N}_6} - \bullet_{\tilde{N}_4'} - \bullet_{\tilde{N}_2'} \tag{4.13}$$

Indeed,

$$\hat{X} = (T[SU(k)] \times \tilde{X}) // SU(k) \tag{4.14}$$

where the symbol $T//G$ means that we gauge the flavor symmetry G of the theory T .

So the theory X whose Coulomb branch is $\mathcal{M}^{\text{inst}}$ in (4.12) is given by

$$X = (T[SU(k)] \times T[SU(k)] \times \tilde{X}) // (SU(k) \times SU(k)) \tag{4.15}$$

But two $T[SU(k)]$ gauged by a diagonal $SU(k)$ is known to disappear, since it is the domain wall of 4d $\mathcal{N}=4$ SYM implementing the S-duality [37]. So we have, in fact,

$$X = \tilde{X} \tag{4.16}$$

and the ALE deformation parameter ξ is now the mass parameter of the $SU(k)$ flavor symmetry. We have

$$\dim_{\mathbb{H}} \mathcal{M}^{\text{inst}} = 30(N_3 + k) - \langle \mathbf{w}, \boldsymbol{\rho} \rangle. \tag{4.17}$$

This nicely agrees with the computation from the geometry (2.7) by the identification

$$N_{\text{inst}} = N_3 + k. \tag{4.18}$$

This relation between N_3 and N_{inst} is also consistent with what we found from the inflow, see (3.14).

We note that the theory $X = \tilde{X}$ is the theory whose Higgs branch is the $U(k)$ instanton moduli on $\mathbb{C}^2/\Gamma_{E_8}$ [2, 5]. From this reason, the Coulomb branch, at least when the mass parameter is zero, has been conjectured to be the E_8 instanton moduli space on the singular space $\mathbb{C}^2/\mathbb{Z}_k$ by various people. This follows, at least in a rough form, from the string duality: consider the theory on M2-branes on $\mathbb{C}^2/\mathbb{Z}_k \times \mathbb{C}^2/\Gamma_{E_8}$. It has two supersymmetric branches of vacua, one describing E_8 instantons on $\mathbb{C}^2/\mathbb{Z}_k$ and another describing $U(k)$ instantons on $\mathbb{C}^2/\Gamma_{E_8}$. If the former is the Coulomb branch, then the latter is the Higgs branch.

Note that we arrived at the quiver gauge theory $X = \tilde{X}$ from a totally different method, by first studying the 6d quiver and then by reducing on successively on circles. Therefore, this agreement can be thought of as an overall consistency check of our construction.

Now, applying [2] and [5] in our case, we see that the $U(k)$ holonomy at infinity of $\mathbb{C}^2/\Gamma_{E_8}$ is trivial, and the first Chern class c_1 satisfies $\int_{E_i} c_1 = n_i$ which can be read off from (4.8). It would be interesting to understand from M-theory point of view why the first Chern class on the $\mathbb{C}^2/\Gamma_{E_8}$ side is given by the asymptotic E_8 holonomy on the $\mathbb{C}^2/\mathbb{Z}_k$. It seems important for the full story to consider a more general case where $\mathbb{C}^2/\mathbb{Z}_k$ is replaced by the multicenter Taub-NUT space, see e.g. [20, 38].

5 Examples

Let us demonstrate the above general statement in various examples. In this section, we take N to be the number of tensor multiplets in the 6d theory, which was denoted by N_6 in the other sections.

5.1 The case of $k = 2$

There are three possibilities. We label the cases with the Kac label \underline{n} and the group $H \subset E_8$ left unbroken by the Kac label. The choice $k = 2$ is somewhat special, since the ALE space $\widetilde{\mathbb{C}^2/\mathbb{Z}_2}$, also known as the Eguchi-Hanson space, has an exceptional isometry $SU(2)$. Then the generic flavor symmetry of the 6d SCFT should be $H \times SU(2)^2$, where one $SU(2)$ comes from the 7d gauge field on the singularity and another $SU(2)$ comes from the isometry.

1. The first case is

$$\underline{n} = \begin{matrix} 0 \\ 20000000 \end{matrix}, \quad H = E_8. \tag{5.1}$$

The corresponding 6d theory is

$$[E_8] \quad 1 \quad \begin{array}{c} \mathfrak{su}(1) \\ 2 \end{array} \quad \begin{array}{c} \mathfrak{su}(2) \\ 2 \end{array} \quad \underbrace{\begin{array}{c} \mathfrak{su}(2) \\ 2 \end{array} \quad \cdots \quad \begin{array}{c} \mathfrak{su}(2) \\ 2 \end{array}}_{N-3} \quad [\mathrm{SU}(2)]. \quad (5.2)$$

The T^3 reduction of this theory gives the following 3d $\mathcal{N} = 4$ theory:

$$\bullet_1 - \bullet_2 - \bullet_N - \bullet_{2N} - \bullet_{3N} - \bullet_{4N} - \bullet_{5N} - \begin{array}{c} \bullet_{3N} \\ | \\ \bullet_{6N} \end{array} - \bullet_{4N} - \bullet_{2N}. \quad (5.3)$$

2. The second case is

$$\underline{n} = \begin{array}{c} 0 \\ 01000000 \end{array}, \quad H = E_7 \times \mathrm{SU}(2). \quad (5.4)$$

The corresponding 6d theory is

$$[E_7] \quad 1 \quad \begin{array}{c} \mathfrak{su}(2) \\ 2 \end{array} \quad \begin{array}{c} \mathfrak{su}(2) \\ 2 \end{array} \quad \cdots \quad \begin{array}{c} \mathfrak{su}(2) \\ 2 \end{array} \quad [\mathrm{SU}(2)] \quad (5.5)$$

$[N_f=2] \quad \underbrace{\hspace{10em}}_{N-2}$

where the number of $\mathfrak{su}(2)$ gauge groups in the quiver is $N - 1$. The Higgs branch dimension of the UV fixed point of this theory is $29N + 4 + 4(N - 1) - 3(N - 1) = 30N + 3$. The mirror of the T^3 compactification of this theory is

$$\bullet_1 - \bullet_2 - \bullet_{N+1} - \bullet_{2N} - \bullet_{3N} - \bullet_{4N} - \bullet_{5N} - \begin{array}{c} \bullet_{3N} \\ | \\ \bullet_{6N} \end{array} - \bullet_{4N} - \bullet_{2N}, \quad (5.6)$$

The Coulomb branch dimension, which is the sum of the rank of the gauge groups minus one, is indeed $30N + 3$.

3. The third case is

$$\underline{n} = \begin{array}{c} 0 \\ 00000001 \end{array}, \quad H = \mathrm{SO}(16). \quad (5.7)$$

The corresponding 6d theory is

$$[\mathrm{SO}(16)] \quad \begin{array}{c} \mathfrak{sp}(1) \\ 1 \end{array} \quad \begin{array}{c} \mathfrak{su}(2) \\ 2 \end{array} \quad \begin{array}{c} \mathfrak{su}(2) \\ 2 \end{array} \quad \cdots \quad \begin{array}{c} \mathfrak{su}(2) \\ 2 \end{array} \quad [\mathrm{SU}(2)] \quad (5.8)$$

$\underbrace{\hspace{10em}}_{N-1}$

where the number of $\mathrm{SU}(2)$ gauge groups associated with the (-2) curves is $N - 1$. The Higgs branch dimension of the UV fixed point of this theory is $29N + 16 + 4 + 4(N - 1) - 3 - 3(N - 1) = 30N + 16$. The mirror of the T^3 compactification of this theory is

$$\bullet_1 - \bullet_2 - \bullet_{N+2} - \bullet_{2N+2} - \bullet_{3N+2} - \bullet_{4N+2} - \bullet_{5N+2} - \begin{array}{c} \bullet_{3N+1} \\ | \\ \bullet_{6N+2} \end{array} - \bullet_{4N+1} - \bullet_{2N}. \quad (5.9)$$

The Coulomb branch dimension, which is the sum of the rank of the gauge groups minus one, is indeed $30N + 16$. This is consistent with figure 45 of [12], namely the T^2 compactification of (5.8) yields the class \mathcal{S} theory whose Gaiotto curve is a sphere with punctures:

$$[(3N + 1)^2], \quad [(2N + 1)^2, 2N], \quad [N^6, 1^2]. \quad (5.10)$$

Now let us comment on the flavor symmetry from the point of view of the 6d quiver. Since an $SU(2)$ - $SU(2)$ bifundamental has an $SU(2)$ flavor symmetry, the three 6d quivers presented above have order N copies of $SU(2)$ symmetries on the generic points of the tensor branch. In fact the same issue already appears in the case of N M5-branes probing the $\mathbb{C}^2/\mathbb{Z}_2$ singularity, which has the quiver

$$[SU(2)] \underbrace{\begin{matrix} \text{su}(2) & & \text{su}(2) \\ 2 & \cdots & 2 \end{matrix}}_N [SU(2)] \tag{5.11}$$

which naively has too many $SU(2)$ flavor symmetries.

The issue can be resolved by recalling the fact derived in appendix A of [39] that the basic 6d SCFT whose quiver on the tensor branch is given by $SU(2)$ with $N_f = 4$ with a naive $SO(8)$ symmetry, only has an $SO(7)$ symmetry under which the flavors transform in the spin representation. In the quiver representation of the same theory as

$$[SU(2)_1] \begin{matrix} \text{su}(2) \\ 2 \end{matrix} [SU(2)_2], \tag{5.12}$$

this means the following: regard the bifundamental hypermultiplets on the left and on the right of the gauge group as the trifundamental half-hypermultiplets. At the quiver level there are therefore the flavor symmetry $SU(2)_1 \times SU(2)'_1 \times SU(2)_2 \times SU(2)'_2 \subset SO(8)$. Under the $SO(7)$ symmetry which is the flavor symmetry of the SCFT, only the diagonal subgroup of $SU(2)'_1$ and $SU(2)'_2$ survives. Applying this argument at every $\mathfrak{su}(2)$ node in (5.2), (5.5), (5.8), and (5.12), we see that the number of $SU(2)$ flavor symmetries is reduced appropriately.

There are also some interesting special cases with enhanced flavour symmetries when N is small:

1. $N = 2, H = E_8$. In this case the quiver (5.2) degenerates to

$$[E_8] \begin{matrix} 1 & \text{su}(1) \\ & 2 \\ & [SU(2)] \\ & [N_f=1] \end{matrix} \tag{5.13}$$

which is just the rank-2 E-string theory with three decoupled hypermultiplets. The 3d quiver in this case is (5.3) for $N = 2$:

$$\begin{matrix} & & & & & & & \bullet 6 \\ & & & & & & & | \\ \bullet 1 & - & \bullet 2 & - & \bullet 2 & - & \bullet 4 & - & \bullet 6 & - & \bullet 8 & - & \bullet 10 & - & \bullet 12 & - & \bullet 8 & - & \bullet 4 \end{matrix} \tag{5.14}$$

Its Coulomb branch is

$$\mathbb{H}^3 \times (\text{the reduced moduli space of } 2 E_8 \text{ instantons on } \mathbb{C}^2) \tag{5.15}$$

and we indeed see the same decoupled structure. The explanation from the perspective of the Coulomb branch operators will be described below.

2. $N = 3$ with $H = E_8$. The 6d quiver is

$$[E_8] \quad 1 \quad \begin{array}{c} \text{su}(1) \\ 2 \end{array} \quad \begin{array}{c} \text{su}(2) \\ 2 \\ [N_f=1] \end{array} \quad [\text{SU}(2)]. \quad (5.16)$$

The 3d quiver is

$$\begin{array}{cccccccccccccccc} \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet \\ 1 & & 2 & & 3 & & 6 & & 9 & & 12 & & 15 & & 18 & & 12 & & 6 & & \bullet \\ & & & & & & & & & & & & & & \uparrow & & & & & & & 9 \end{array} \quad (5.17)$$

The flavour symmetry is enhanced to $G_2 \times E_8$. The explanation from the perspective of the Coulomb branch operators will also be described below.

3. $N = 2$, $H = E_7 \times \text{SU}(2)$. The 6d quiver for this case reduces to

$$[E_7] \quad 1 \quad \begin{array}{c} \text{su}(2) \\ 2 \\ [N_f=2] \end{array} \quad [\text{SU}(2)]. \quad (5.18)$$

On the tensor branch, there is an $\text{SO}(8)$ symmetry acting on the four flavors of $\text{SU}(2)$ gauge group. In the SCFT it is known that there is only $\text{SO}(7)$. The total symmetry is then $\text{SO}(7) \times E_7$. In fact this 6d theory is the $(E_7, \text{SO}(7))$ minimal conformal matter [10], which describes “half M5-branes” on the E_7 singularity.

The 3d quiver in this case is (5.6) for $N = 2$:

$$\begin{array}{cccccccccccccccc} \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet \\ 1 & & 2 & & 3 & & 4 & & 6 & & 8 & & 10 & & 12 & & 8 & & 4 & & \bullet \\ & & & & & & & & & & & & & & \uparrow & & & & & & & 6 \end{array} \quad (5.19)$$

This theory is the mirror of the S^1 reduction of the class \mathcal{S} theory whose Gaiotto curve is a sphere with punctures

$$[2^4, 1^4], \quad [6^2], \quad [4^3]. \quad (5.20)$$

In [12, 15] the T^2 compactification was also identified with a class \mathcal{S} theory of the E_6 type associated with the sphere with punctures $0, 2A_1$ and $E_6(a_1)$. For consistency, these two class \mathcal{S} theories should in fact be the same. Let us compute the central charges of (5.20). We find that the effective numbers of vector multiplets and hypermultiplets are $n_H = 112$ and $n_V = 49$, respectively. Thus,

$$a = \frac{1}{24}(5n_V + n_H) = \frac{119}{8}, \quad c = \frac{1}{12}(2n_V + n_H) = \frac{35}{2}. \quad (5.21)$$

This agrees with a and c of the aforementioned class \mathcal{S} theory of the E_6 type; see (7.1) of [15].

5.2 Enhanced flavor symmetries from 3d quivers

In fact the symmetry enhancement of each of the three cases above can be generalized to other over-extended Dynkin quivers in 3d, namely:

1. For the quiver consisting of a tail $\bullet - \bullet$ attached to the affine Dynkin diagram of type \mathfrak{g} with gauge groups being unitary groups of the ranks given by 2 times the dual Coxeter labels, the Coulomb branch moduli space is $\mathbb{H}^3 \times \widetilde{\mathcal{M}}_{2,\mathfrak{g}}$, where $\widetilde{\mathcal{M}}_{2,\mathfrak{g}}$ denotes the reduced two-instanton moduli space of group \mathfrak{g} on \mathbb{C}^2 . For example, the Coulomb branch of the quiver

$$\bullet - \bullet - \bullet = \bullet \tag{5.22}$$

$\begin{matrix} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{matrix}$

is $\mathbb{H}^3 \times \widetilde{\mathcal{M}}_{2, \mathfrak{su}(2)}$, and the Coulomb branch of the quiver

$$\begin{matrix} & & & \bullet & & \\ & & & | & & \\ \bullet & - & \bullet & - & \bullet & - & \bullet \\ & & & | & & & \\ & & & \bullet & & & \end{matrix} \tag{5.23}$$

is $\mathbb{H}^3 \times \widetilde{\mathcal{M}}_{2, \mathfrak{so}(8)}$.

2. For the quiver consisting of a tail $\bullet - \bullet$ attached to the affine Dynkin diagram of type \mathfrak{g} with gauge groups being unitary groups of the ranks given by 3 times the dual Coxeter labels, the Coulomb branch moduli space has a symmetry $G_2 \times \mathfrak{g}$. For example, the Coulomb branch of the quiver

$$\bullet - \bullet - \bullet = \bullet \tag{5.24}$$

$\begin{matrix} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{matrix}$

has a symmetry $G_2 \times \text{SU}(2)$, and the Coulomb branch of the quiver

$$\begin{matrix} & & & \bullet & & \\ & & & | & & \\ \bullet & - & \bullet & - & \bullet & - & \bullet \\ & & & | & & & \\ & & & \bullet & & & \end{matrix} \tag{5.25}$$

has a symmetry $G_2 \times \text{SO}(8)$.

3. For the quiver consisting of a tail $\bullet - \bullet$ attached to the affine Dynkin diagram of type \mathfrak{g} with the affine node being $\text{U}(3)$ and other gauge groups being unitary groups of the ranks given by 2 times the dual Coxeter labels, the Coulomb branch has a symmetry $\text{SO}(7) \times \tilde{\mathfrak{g}}$, where $\tilde{\mathfrak{g}}$ is the commutant of $\mathfrak{su}(2)$ in \mathfrak{g} . For example, the Coulomb branch of the following quiver

$$\begin{matrix} & & & & \bullet & & \\ & & & & | & & \\ & & & & \bullet & & \\ & & & & | & & \\ \bullet & - & \bullet & - & \bullet & - & \bullet \\ & & & & | & & \\ & & & & \bullet & & \\ & & & & | & & \\ & & & & \bullet & & \\ & & & & | & & \\ \bullet & - & \bullet & - & \bullet & - & \bullet \\ & & & & | & & \\ & & & & \bullet & & \end{matrix} \tag{5.26}$$

has a symmetry $\text{SO}(7) \times \text{SU}(6)$, where $\text{SU}(6)$ is the commutant of $\text{SU}(2)$ in E_6 .

In each of the above examples, the quiver contains of a balanced affine Dynkin quiver diagram as a subquiver. If we consider only this subquiver, the R-charges of the monopole operators in this theory vanish, and hence this subquiver is indeed a bad theory. By attaching a quiver tail $\bullet - \bullet - \dots - \bullet$ to such a subquiver, the total quiver becomes good

or ugly.⁴ We would like to consider the contribution of this quiver tail to the Coulomb branch of the total quiver.

1. For this case, the node \bullet_2 , which is the affine node in the affine Dynkin diagram, is over-balanced in the sense of [37]. Following [37], we can split the quiver into two parts, namely $\bullet_1 - \bullet_2 - \bullet_2$ and the rest of the Dynkin diagram. The R-charge of the monopole operators from the subquiver $\bullet_1 - \bullet_2 - \bullet_2$ receives the contribution from the hypermultiplets and vector multiplets in the way described in [37], except that there is no contribution from the vector multiplet of the rightmost node \bullet_2 , since this was cancelled inside the affine Dynkin quiver. The contribution from the subquiver is therefore the same as that of the quiver $\bullet_1 - \bullet_2 - \overset{\cap}{\bullet_2}$, where \cap denotes an adjoint hypermultiplet of the U(2) rightmost node. The Coulomb branch of $\bullet_1 - \bullet_2 - \overset{\cap}{\bullet_2}$ contains 3 free hypermultiplets, which can be seen from the monopole operators with $SU(2)_R$ -spin 1/2. This explains the \mathbb{H}^3 factor in (5.15). The reduced moduli space of two E_8 instantons on \mathbb{C}^2 can be realised as in [21].
2. Similarly, for this case, the total quiver can be split into $\bullet_1 - \bullet_2 - \bullet_3$ and the rest of the Dynkin diagram. The contribution to the R-charge of the monopole operators from the subquiver $\bullet_1 - \bullet_2 - \bullet_3$ can be realised from the quiver $\bullet_1 - \bullet_2 - \overset{\cap}{\bullet_3}$, where \cap denotes an adjoint hypermultiplet of the U(3) rightmost node.⁵ Indeed, it was pointed out in section 3.3.2 of [41] that the Coulomb branch of the latter model has a G_2 symmetry. (Note that the corresponding 4d class S theory had been studied in [40]. The G_2 symmetry on the Higgs branch of such a theory had also been pointed out in that reference.) This therefore explains the G_2 symmetry in case 3. The E_8 symmetry follows from the Dynkin subquiver.
3. Finally, for this case, \bullet_4 is the unbalanced node in the quiver. There are two contributions to the Coulomb branch operators with $SU(2)_R$ -spin 1. One contribution can be realised using the quiver $\bullet_1 - \bullet_2 - \bullet_3 - \overset{\cap}{\bullet_4}$ in a similar fashion to the above discussion. This quiver has a Coulomb branch symmetry $SU(4)$ and thus gives 15 operators with $SU(2)_R$ -spin 1 in the adjoint representation of $SU(4)$. The other contribution can be seen as follows. Since the node \bullet_3 , which was originally a part of the affine Dynkin subquiver, now belongs to the tail $\bullet_1 - \bullet_2 - \bullet_3 - \bullet_4$, we also need to take into account the contribution that arises from the removal of this node from such an affine Dynkin diagram. The second contribution thus comes from considering $\bullet_1 - \bullet_2 - \overset{\cap}{\bullet_4}$. There are 6 Coulomb branch operators with $SU(2)_R$ -spin 1 in the latter. Therefore, we have in total $15+6=21$ operators with $SU(2)_R$ -spin 1; this explains the enhancement to the $SO(7)$ symmetry. The remaining symmetry is thus the commutant of $SU(2)$, which arises from node \bullet_3 , in the original symmetry associated with the affine Dynkin diagram.

⁴See also [40] for a related consideration from the 4d point of view.

⁵The authors thank S. Cremonesi for this argument.

5.3 The case of $k = 4$

There are ten possibilities. The F-theory quiver for the $6d$ theories are listed on page 73 of [11]. Here are the mirrors of the T^3 compactification of them.

1. The first case is

$$\underline{n} = \begin{matrix} 0 \\ 40000000 \end{matrix}, \quad H = E_8. \quad (5.27)$$

The $6d$ quiver is

$$[E_8] \quad 1 \quad \begin{matrix} \text{su}(1) \\ 2 \\ [N_f=1] \end{matrix} \quad \begin{matrix} \text{su}(2) \\ 2 \\ [N_f=1] \end{matrix} \quad \begin{matrix} \text{su}(3) \\ 2 \\ [N_f=1] \end{matrix} \quad \overbrace{\begin{matrix} \text{su}(4) & \dots & \text{su}(4) \\ 2 & & 2 \end{matrix}}^{N-4} \quad [SU(4)] \quad (5.28)$$

and the $3d$ quiver is

$$\bullet_1 - \bullet_2 - \bullet_3 - \bullet_4 - \bullet_N - \bullet_{2N} - \bullet_{3N} - \bullet_{4N} - \bullet_{5N} - \overset{\bullet_{3N}}{\downarrow} \bullet_{6N} - \bullet_{4N} - \bullet_{2N}. \quad (5.29)$$

2. The second case is

$$\underline{n} = \begin{matrix} 0 \\ 21000000 \end{matrix}, \quad H = E_7 \times U(1) \quad (5.30)$$

with the $6d$ quiver

$$[E_7] \quad 1 \quad \begin{matrix} \text{su}(2) \\ 2 \\ [N_f=1] \end{matrix} \quad \begin{matrix} \text{su}(3) \\ 2 \\ [N_f=1] \end{matrix} \quad \overbrace{\begin{matrix} \text{su}(4) & \dots & \text{su}(4) \\ 2 & & 2 \end{matrix}}^{N-3} \quad [SU(4)]. \quad (5.31)$$

The dimension of the SCFT Higgs branch is

$$29N + 2 + 6 + 12 + 4 + 16(N - 3) - 3 - 8 - 15(N - 3) = 30N + 10. \quad (5.32)$$

The $3d$ quiver is

$$\bullet_1 - \bullet_2 - \bullet_3 - \bullet_4 - \bullet_{N+1} - \bullet_{2N} - \bullet_{3N} - \bullet_{4N} - \bullet_{5N} - \overset{\bullet_{3N}}{\downarrow} \bullet_{6N} - \bullet_{4N} - \bullet_{2N} \quad (5.33)$$

and the dimension of the Coulomb branch is $30N + 10$.

3. The third case is

$$\underline{n} = \begin{matrix} 0 \\ 20000001 \end{matrix}, \quad H = SO(14) \times U(1) \quad (5.34)$$

with the $6d$ quiver

$$[SO(14)] \quad \begin{matrix} \text{sp}(1) \\ 1 \\ [N_f=1] \end{matrix} \quad \begin{matrix} \text{su}(3) \\ 2 \\ [N_f=1] \end{matrix} \quad \overbrace{\begin{matrix} \text{su}(4) & \dots & \text{su}(4) \\ 2 & & 2 \end{matrix}}^{N-2} \quad [SU(4)]. \quad (5.35)$$

The dimension of the SCFT Higgs branch is

$$29N + 14 + 6 + 12 + 4 + 16(N - 2) - 3 - 8 - 15(N - 2) = 30N + 23 . \quad (5.36)$$

The 3d quiver is

$$\begin{array}{cccccccccccc}
 \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet \\
 1 & & 2 & & 3 & & 4 & & N+2 & & 2N+2 & & 3N+2 & & 4N+2 & & 5N+2 & & 6N+2 & & 4N+1 & & 2N \\
 & & & & & & & & & & & & & & & & & & \uparrow & & & & & \\
 & & & & & & & & & & & & & & & & & & 3N+1 & & & & & \\
 \end{array} \quad (5.37)$$

and the Coulomb branch dimension is $30N + 23$.

4. The fourth case is

$$\underline{n} = \begin{array}{c} 0 \\ 02000000 \end{array}, \quad H = E_7 \times \text{SU}(2). \quad (5.38)$$

with the 6d quiver

$$[E_7] \begin{array}{c} 1 \\ \text{su}(2) \\ \text{[SU}(2)] \end{array} \begin{array}{c} 2 \\ \text{su}(4) \\ 2 \end{array} \overbrace{\begin{array}{c} \text{su}(4) \\ \dots \\ \text{su}(4) \end{array}}^{N-2} \begin{array}{c} 2 \\ \text{[SU}(4)] \end{array}. \quad (5.39)$$

The dimension of the SCFT Higgs branch is

$$29N + 8 + 8 + 16(N - 2) - 3 - 15(N - 2) = 30N + 11 . \quad (5.40)$$

The 3d mirror is

$$\begin{array}{cccccccccccc}
 \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet \\
 1 & & 2 & & 3 & & 4 & & N+2 & & 2N & & 3N & & 4N & & 5N & & 6N & & 4N & & 2N \\
 & & & & & & & & & & & & & & & & & & \uparrow & & & & & \\
 & & & & & & & & & & & & & & & & & & 3N & & & & & \\
 \end{array} \quad (5.41)$$

The Coulomb branch dimension is $30N + 11$.

5. The fifth case is

$$\underline{n} = \begin{array}{c} 0 \\ 00000002 \end{array}, \quad H = \text{SO}(16) \quad (5.42)$$

with the 6d quiver

$$[\text{SO}(16)] \begin{array}{c} 1 \\ \text{sp}(2) \\ \text{[SU}(4)] \end{array} \begin{array}{c} 2 \\ \text{su}(4) \\ 2 \end{array} \overbrace{\begin{array}{c} \text{su}(4) \\ \dots \\ \text{su}(4) \end{array}}^{N-1} \begin{array}{c} 2 \\ \text{[SU}(4)] \end{array} \quad (5.43)$$

The dimension of the SCFT Higgs branch is

$$29N + 32 + 16 + 16(N - 1) - 10 - 15(N - 1) = 30N + 37 . \quad (5.44)$$

The 3d quiver is

$$\begin{array}{cccccccccccc}
 \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet \\
 1 & & 2 & & 3 & & 4 & & N+4 & & 2N+4 & & 3N+4 & & 4N+4 & & 5N+4 & & 6N+4 & & 4N+2 & & 2N \\
 & & & & & & & & & & & & & & & & & & \uparrow & & & & & \\
 & & & & & & & & & & & & & & & & & & 3N+2 & & & & & \\
 \end{array} \quad (5.45)$$

and the Coulomb branch dimension is $30N + 37$.

6. The sixth case is

$$\underline{n} = \begin{matrix} 0 \\ 01000001 \end{matrix}, \quad H = \text{SO}(12) \times \text{SU}(2) \times \text{U}(1) \quad (5.46)$$

with the 6d quiver

$$[\text{SO}(12)] \overset{\text{sp}(1)}{1} \overbrace{\begin{matrix} \text{su}(4) & \text{su}(4) \\ 2 & 2 \end{matrix}}^{N-1} \text{[SU}(4)] \quad (5.47)$$

The dimension of the SCFT Higgs branch is

$$29N + 12 + 8 + 8 + 16(N - 1) - 3 - 15(N - 1) = 30N + 24. \quad (5.48)$$

The 3d quiver is

$$\begin{matrix} \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet \\ 1 & & 2 & & 3 & & 4 & & N+3 & & 2N+2 & & 3N+2 & & 4N+2 & & 5N+2 & & 6N+2 & & 4N+1 & & 2N \end{matrix} \quad (5.49)$$

The Coulomb branch dimension is $30N + 24$.

7. The seventh case is

$$\underline{n} = \begin{matrix} 0 \\ 10100000 \end{matrix}, \quad H = E_6 \times \text{SU}(2) \times \text{U}(1). \quad (5.50)$$

with the 6d quiver

$$[E_6] \overset{\text{su}(3)}{1} \overbrace{\begin{matrix} \text{su}(4) & \text{su}(4) \\ 2 & 2 \end{matrix}}^{N-2} \text{[SU}(4)]. \quad (5.51)$$

The dimension of the SCFT Higgs branch is

$$29N + 6 + 12 + 4 + 16(N - 2) - 8 - 15(N - 2) = 30N + 12. \quad (5.52)$$

The 3d mirror is

$$\begin{matrix} \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet \\ 1 & & 2 & & 3 & & 4 & & N+2 & & 2N+1 & & 3N & & 4N & & 5N & & 6N & & 4N & & 2N \end{matrix} \quad (5.53)$$

and the Coulomb branch dimension is $30N + 12$.

8. The eighth case is

$$\underline{n} = \begin{matrix} 1 \\ 10000000 \end{matrix}, \quad H = \text{SU}(8) \times \text{U}(1) \quad (5.54)$$

with the 6d quiver

$$[\text{SU}(8)] \overset{\text{su}(3)}{1} \overbrace{\begin{matrix} \text{su}(4) & \text{su}(4) \\ 2 & 2 \end{matrix}}^{N-1} \text{[SU}(4)]. \quad (5.55)$$

We can observe this from the $3d$ quivers. We get a tower of monopole operators from every node. The basic monopole operator from the balanced nodes leads to enhancement of symmetry. We also have a basic monopole operator from the unbalanced nodes. These provide operators with higher R-charges, and we can read of their R-charges and non-abelian global symmetry charges from the quiver.

We have three unbalanced nodes. Two of them give the same contribution in both theories: one operator of $SU(2)_R$ spin $\frac{N}{2}$ in the bifundamental of the $SU(2l+8) \times SU(8)$ global symmetry, and one operator of $SU(2)_R$ spin 2 in the **28** of the $SU(8)$ global symmetry. These can be readily identified with gauge invariants in the $6d$ quiver, where the former is the one made from $N-2$ $SU(2l+8) \times SU(2l+8)$ bifundamentals and the flavors, and the later is made from two $SU(8)$ flavors and the $USp(2l) \times SU(2l+8)$ bifundamental. The last one differ slightly between the two theories.

In the $\theta = 0$ case it is a flavor singlet with $SU(2)_R$ spin $\frac{l+2}{2}$. Particularly for $l = 0$ this gives the conserved current enhancing the $U(1)$ to $SU(2)$. In the $\theta = \pi$ case, however, it is in the **8** of $SU(8)$ with $SU(2)_R$ spin $\frac{l+3}{2}$. We can interpret these states as coming from the USp gauge group instanton strings wrapped on the circle. These are in the spinor of $SO(4l+16)$, and depending on the θ angle decompose to all the even or odd rank antisymmetric tensor representations of the gauge $SU(2l+8)$ connected to the USp gauge group. In the $\theta = 0$ case we get the even rank representations, which contain a gauge invariant part which is a flavor symmetry singlet. In the $\theta = \pi$ case we get the odd rank representations, which do not contain any gauge invariants. However we can combine it with one of the $SU(2l+8)$ flavors to form an invariant. This should contribute a state in the **8** of $SU(8)$ with $SU(2)_R$ spin which is greater by $\frac{1}{2}$ from that of the singlet. This agrees with what we observe. It might be interesting to study more accurately the spectrum, particularly, the Higgs branch chiral ring, and compare against the $6d$ expectations. We will not pursue this here.

5.5 Massive E-string theories

In this subsection, we consider the following $6d$ theory

$$\mathcal{T}_E^{6d}(\varpi, m_0, N) : [E_{9-m_0}] \overset{su_{m_0}}{2} \overset{su_{2m_0}}{2} \dots \overset{su_{(\varpi-1)m_0}}{2} \overset{su_{\varpi m_0}}{2} \underbrace{\overset{su_{\varpi m_0}}{2} \dots \overset{su_{\varpi m_0}}{2}}_{N-\varpi-1} [SU(\varpi m_0)]. \tag{5.71}$$

These theories were studied in [10, 12, 13, 42]. They can be called the “massive E-string theories” as in the last reference, since they correspond to NS5-branes probing the O8-D8 combination in the presence of the Romans mass.

The mirror of the T^3 compactification of (5.71) is

$$\bullet_1 - \bullet_2 - \dots - \bullet_{\varpi m_0} - \bullet_{N+r_1} - \bullet_{2N+r_2} - \bullet_{3N+r_3} - \bullet_{4N+r_4} - \bullet_{5N+r_5} - \overset{\bullet_{3N+r_{3'}}}{\downarrow} \bullet_{6N+r_6} - \bullet_{4N+r_{4'}} - \bullet_{2N+r_{2'}} \tag{5.72}$$

where the values of r_i and the Kac labels for each m_0 are given in table 1. Note that

$$\sum_i r_i = \frac{1}{2} \varpi m_0 (m_0 - 1) . \tag{5.73}$$

m_0	E_{9-m_0}	Kac label \underline{n}/ϖ	r_i/ϖ
1	E_8	0 10000000	0 00000000
2	E_7	0 01000000	0 10000000
3	E_6	0 00100000	0 21000000
4	SO(10)	0 00010000	0 32100000
5	SU(5)	0 00001000	0 43210000
6	SU(3) × SU(2)	0 00000100	0 54321000
7	SU(2) × U(1)	1 00000010	0 65432100
8	SU(2)	0 00000020	1 76543200

Table 1. The values of r_i in (5.72) and the Kac label for each m_0 .


The SCFT Higgs branch dimension of (5.71) is

$$\dim_{\mathbb{H}}^{\text{SCFT}} \text{Higgs of } \mathcal{T}_E^{6d}(\varpi, m_0, N) = 30N + \frac{1}{2}\varpi m_0^2(\varpi + 1) - 1; \quad (5.74)$$

this is equal to the Coulomb branch dimension of (5.72).

5.6 Higgsing the SU(k) flavour symmetry

In the theories we have discussed so far, there is always an SU(k) flavour symmetry which came from the gauge symmetry on the $\mathbb{C}^2/\mathbb{Z}_k$ singularity. From the 3d quiver perspective, this symmetry arises from the topological symmetry associated with the nodes in the tail



We can obtain another class of models by on nilpotent VEVs that Higgs the flavour symmetry SU(k).⁶ Suppose that such VEVs are in the nilpotent orbit of SU(k) given by $\bigoplus_i J_{s_i}$ where J_s is a $s \times s$ Jordan block so that $Y = [s_1, s_2, \dots, s_\ell]$ is a corresponding partition of k .

Assuming that the 6d quiver theory before the Higgsing has a sufficiently long plateau of SU(k) gauge groups, this Higgsing can be performed exactly as in 4d class S theory e.g. as described in section 12.5 of [43]. Its effect in 6d quiver was studied in [44, 45]. In the end, we see that the tail on the right-hand side of the quiver to have the form

$$\dots \begin{array}{c} \text{su}(k) \\ 2 \\ [N_f=u_{\ell'}] \end{array} \begin{array}{c} \text{su}(k) \\ 2 \\ [N_f=(u_{\ell'-1}-u_{\ell'})] \end{array} \dots \begin{array}{c} \text{su}(u_2+u_1) \\ 2 \\ [N_f=(u_2-u_3)] \end{array} \begin{array}{c} \text{su}(u_1) \\ 2 \\ [N_f=(u_1-u_2)] \end{array}, \quad (5.75)$$

⁶The authors thank Alessandro Tomasiello for the discussion about this class of theories.

where u_i are the elements of the transpose $Y^T = [u_1, u_2, \dots, u_{\ell'}]$, and we define $u_i = 0$ for $i > \ell'$.

The SCFT Higgs branch dimension of (5.75) is

$$\dim_{\mathbb{H}}^{\text{SCFT}} \text{Higgs of (5.75)} = \left[30(N_3 + k) - \langle \mathbf{w}, \boldsymbol{\rho} \rangle + \frac{1}{2}k(k+1) - 1 \right] - \dim_{\mathbb{H}} \mathcal{O}_Y \quad (5.76)$$

where \mathcal{O}_Y is the nilpotent orbit labeled by Y .

The mirror of the T^3 compactification of (5.75) is

$$\frac{T_Y(\text{SU}(k)) \times \begin{array}{ccccccccccc} \blacksquare & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ & k & \tilde{N}_1 & \tilde{N}_2 & \tilde{N}_3 & \tilde{N}_4 & \tilde{N}_5 & \tilde{N}_6 & \tilde{N}_{4'} & \tilde{N}_{2'} & \end{array}}{\text{U}(k)/\text{U}(1)} \quad (5.77)$$

In other words, we simply replace the tail $\bullet - \bullet - \dots - \blacksquare$ for the theories discussed in the preceding sections by $T_Y(\text{SU}(k))$, where the latter is defined as in [37]. The Coulomb branch dimension of (5.77) is

$$\begin{aligned} \dim_{\mathbb{H}} \text{Coulomb of (5.77)} &= [30(N_3 + k) - \langle \mathbf{w}, \boldsymbol{\rho} \rangle] + \left[\frac{1}{2}\{(k^2 - 1) - (k - 1)\} - \dim_{\mathbb{H}} \mathcal{O}_Y \right] + (k - 1) \\ &= 30(N_3 + k) + \frac{1}{2}k(k+1) - 1 - \dim_{\mathbb{H}} \mathcal{O}_Y - \langle \mathbf{w}, \boldsymbol{\rho} \rangle, \end{aligned} \quad (5.78)$$

where the terms in the second square brackets in the second line denote the Coulomb branch dimension of $T_Y(\text{SU}(k))$. This result is indeed in agreement with (5.76).

As an example, let us consider $\mathcal{T}_E^{6d}(k, m_0 = 1, N)$ of the previous section and perform the Higgsing with $Y = [k - 1, 1]$. The resulting $6d$ theory is

$$[E_8] \begin{array}{cccccccccccc} \text{su}(1) & \text{su}(2) & & \text{su}(k-1) & \text{su}(k) & \text{su}(k)^{N-2k} & \text{su}(k) & \text{su}(k-1) & & \text{su}(2) & \text{su}(1) \\ 1 & 2 & 2 & \dots & 2 & 2 & 2 & 2 & \dots & 2 & 2 \\ & & & & [N_f=1] & & [N_f=1] & & & & \end{array}, \quad (5.79)$$

where the number of tensor multiplets is N . This theory is similar to that discussed in (36) of [6], (5.2) of [29], except that we have only one (-1) -curve in the quiver, instead of two. The mirror of the T^3 compactification of this theory is

$$\begin{array}{ccccccccccc} & & & & & & & \bullet & & & \\ & & & & & & & | & & & \\ \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet & - & \bullet \\ 1 & & k & & N & & 2N & & 3N & & 4N & & 5N & & 6N & & 4N & & 2N \end{array}, \quad (5.80)$$

This quiver is a “good” theory in the sense of [37] if $N + 1 \geq 2k$ and $k \geq 2$. In this case, this quiver is the $3d$ mirror theory of the S^1 reduction of the class \mathcal{S} theory of type $\text{SU}(6N)$ associated a sphere with the punctures

$$[N^5, N - k, k - 1, 1], \quad [(3N)^2], \quad [(2N)^3]. \quad (5.81)$$

Acknowledgments

The authors thank Hiroyuki Shimizu for the collaboration at the early stages. NM sincerely thanks Stefano Cremonesi, Amihay Hanany and Alessandro Tomasiello for a close collaboration, invaluable insights, and several useful discussions. He is also grateful to the hospitality of the organisers of the Pollica Summer Workshop 2017, including Fernando Alday, Philip Argyres, Madalena Lemos and Mario Martone. He is supported in part by the INFN, the ERC Starting Grant 637844-HBQFTNCER, as well as the ERC STG grant 306260 through the Pollica Summer Workshop. KO gratefully acknowledges support from the Institute for Advanced Study. YT is partially supported in part by JSPS KAKENHI Grant-in-Aid (Wakate-A), No.17H04837 and JSPS KAKENHI Grant-in-Aid (Kiban-S), No.16H06335. YT and GZ are partially supported by WPI Initiative, MEXT, Japan at IPMU, the University of Tokyo.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] M.F. Atiyah, N.J. Hitchin, V.G. Drinfeld and Yu. I. Manin, *Construction of Instantons*, *Phys. Lett. A* **65** (1978) 185 [[INSPIRE](#)].
- [2] P.B. Kronheimer and H. Nakajima, *Yang-Mills instantons on ALE gravitational instantons*, *Math. Ann.* **288** (1990) 263.
- [3] M. Bianchi, F. Fucito, G. Rossi and M. Martellini, *Explicit construction of Yang-Mills instantons on ALE spaces*, *Nucl. Phys. B* **473** (1996) 367 [[hep-th/9601162](#)] [[INSPIRE](#)].
- [4] E. Witten, *Small instantons in string theory*, *Nucl. Phys. B* **460** (1996) 541 [[hep-th/9511030](#)] [[INSPIRE](#)].
- [5] M.R. Douglas and G.W. Moore, *D-branes, quivers and ALE instantons*, [hep-th/9603167](#) [[INSPIRE](#)].
- [6] P.S. Aspinwall and D.R. Morrison, *Point-like instantons on K3 orbifolds*, *Nucl. Phys. B* **503** (1997) 533 [[hep-th/9705104](#)] [[INSPIRE](#)].
- [7] D. Gaiotto, *$\mathcal{N} = 2$ dualities*, *JHEP* **08** (2012) 034 [[arXiv:0904.2715](#)] [[INSPIRE](#)].
- [8] S. Cremonesi, A. Hanany and A. Zaffaroni, *Monopole operators and Hilbert series of Coulomb branches of 3d $\mathcal{N} = 4$ gauge theories*, *JHEP* **01** (2014) 005 [[arXiv:1309.2657](#)] [[INSPIRE](#)].
- [9] J.J. Heckman, D.R. Morrison and C. Vafa, *On the Classification of 6D SCFTs and Generalized ADE Orbifolds*, *JHEP* **05** (2014) 028 [*Erratum ibid.* **1506** (2015) 017] [[arXiv:1312.5746](#)] [[INSPIRE](#)].
- [10] M. Del Zotto, J.J. Heckman, A. Tomasiello and C. Vafa, *6d Conformal Matter*, *JHEP* **02** (2015) 054 [[arXiv:1407.6359](#)] [[INSPIRE](#)].
- [11] J.J. Heckman, D.R. Morrison, T. Rudelius and C. Vafa, *Atomic Classification of 6D SCFTs*, *Fortsch. Phys.* **63** (2015) 468 [[arXiv:1502.05405](#)] [[INSPIRE](#)].

- [12] G. Zafrir, *Brane webs, 5d gauge theories and 6d $\mathcal{N} = (1, 0)$ SCFT's*, *JHEP* **12** (2015) 157 [[arXiv:1509.02016](#)] [[INSPIRE](#)].
- [13] K. Ohmori and H. Shimizu, *S^1/T^2 compactifications of 6d $\mathcal{N} = (1, 0)$ theories and brane webs*, *JHEP* **03** (2016) 024 [[arXiv:1509.03195](#)] [[INSPIRE](#)].
- [14] H. Hayashi, S.-S. Kim, K. Lee and F. Yagi, *6d SCFTs, 5d Dualities and Tao Web Diagrams*, [arXiv:1509.03300](#) [[INSPIRE](#)].
- [15] K. Ohmori, H. Shimizu, Y. Tachikawa and K. Yonekura, *6d $\mathcal{N} = (1, 0)$ theories on T^2 and class S theories: Part I*, *JHEP* **07** (2015) 014 [[arXiv:1503.06217](#)] [[INSPIRE](#)].
- [16] P.B. Kronheimer, *Instantons and the geometry of the nilpotent variety*, *J. Diff. Geom.* **32** (1990) 473 [[INSPIRE](#)].
- [17] Y. Tachikawa, *Moduli spaces of SO(8) instantons on smooth ALE spaces as Higgs branches of 4d $\mathcal{N} = 2$ supersymmetric theories*, *JHEP* **06** (2014) 056 [[arXiv:1402.4200](#)] [[INSPIRE](#)].
- [18] F. Benini, Y. Tachikawa and D. Xie, *Mirrors of 3d Sicilian theories*, *JHEP* **09** (2010) 063 [[arXiv:1007.0992](#)] [[INSPIRE](#)].
- [19] H. Nakajima, *Towards a mathematical definition of Coulomb branches of 3-dimensional $\mathcal{N} = 4$ gauge theories, I*, *Adv. Theor. Math. Phys.* **20** (2016) 595 [[arXiv:1503.03676](#)] [[INSPIRE](#)].
- [20] H. Nakajima, *Questions on provisional Coulomb branches of 3-dimensional $\mathcal{N} = 4$ gauge theories*, [arXiv:1510.03908](#) [[INSPIRE](#)].
- [21] S. Cremonesi, G. Ferlito, A. Hanany and N. Mekareeya, *Coulomb Branch and The Moduli Space of Instantons*, *JHEP* **12** (2014) 103 [[arXiv:1408.6835](#)] [[INSPIRE](#)].
- [22] N. Mekareeya, *The moduli space of instantons on an ALE space from 3d $\mathcal{N} = 4$ field theories*, *JHEP* **12** (2015) 174 [[arXiv:1508.06813](#)] [[INSPIRE](#)].
- [23] V.G. Kac, *Infinite Dimensional Lie Algebras*, Cambridge University Press, Cambridge U.K. (1994).
- [24] H. Nakajima, *Moduli spaces of anti-self-dual connections on ALE gravitational instantons*, *Invent. Math.* **102** (1990) 267.
- [25] K. Ohmori, H. Shimizu, Y. Tachikawa and K. Yonekura, *Anomaly polynomial of general 6d SCFTs*, *PTEP* **2014** (2014) 103B07 [[arXiv:1408.5572](#)] [[INSPIRE](#)].
- [26] K. Intriligator, *6d, $\mathcal{N} = (1, 0)$ Coulomb branch anomaly matching*, *JHEP* **10** (2014) 162 [[arXiv:1408.6745](#)] [[INSPIRE](#)].
- [27] K. Ohmori, H. Shimizu and Y. Tachikawa, *Anomaly polynomial of E-string theories*, *JHEP* **08** (2014) 002 [[arXiv:1404.3887](#)] [[INSPIRE](#)].
- [28] I. Brunner and A. Karch, *Branes at orbifolds versus Hanany Witten in six-dimensions*, *JHEP* **03** (1998) 003 [[hep-th/9712143](#)] [[INSPIRE](#)].
- [29] A. Hanany and A. Zaffaroni, *Branes and six-dimensional supersymmetric theories*, *Nucl. Phys. B* **529** (1998) 180 [[hep-th/9712145](#)] [[INSPIRE](#)].
- [30] O. Bergman and D. Rodriguez-Gomez, *5d quivers and their AdS_6 duals*, *JHEP* **07** (2012) 171 [[arXiv:1206.3503](#)] [[INSPIRE](#)].
- [31] H. Hayashi, S.-S. Kim, K. Lee, M. Taki and F. Yagi, *A new 5d description of 6d D-type minimal conformal matter*, *JHEP* **08** (2015) 097 [[arXiv:1505.04439](#)] [[INSPIRE](#)].

- [32] H. Hayashi, S.-S. Kim, K. Lee, M. Taki and F. Yagi, *More on 5d descriptions of 6d SCFTs*, *JHEP* **10** (2016) 126 [[arXiv:1512.08239](#)] [[INSPIRE](#)].
- [33] O. Bergman and G. Zafrir, *5d fixed points from brane webs and O7-planes*, *JHEP* **12** (2015) 163 [[arXiv:1507.03860](#)] [[INSPIRE](#)].
- [34] F. Benini, S. Benvenuti and Y. Tachikawa, *Webs of five-branes and $N = 2$ superconformal field theories*, *JHEP* **09** (2009) 052 [[arXiv:0906.0359](#)] [[INSPIRE](#)].
- [35] O. Chacaltana and J. Distler, *Tinkertoys for Gaiotto Duality*, *JHEP* **11** (2010) 099 [[arXiv:1008.5203](#)] [[INSPIRE](#)].
- [36] Y. Tachikawa, *A review of the T_N theory and its cousins*, *PTEP* **2015** (2015) 11B102 [[arXiv:1504.01481](#)] [[INSPIRE](#)].
- [37] D. Gaiotto and E. Witten, *S-duality of Boundary Conditions In $N = 4$ Super Yang-Mills Theory*, *Adv. Theor. Math. Phys.* **13** (2009) 721 [[arXiv:0807.3720](#)] [[INSPIRE](#)].
- [38] M. Bullimore, T. Dimofte and D. Gaiotto, *The Coulomb Branch of 3d $\mathcal{N} = 4$ Theories*, *Commun. Math. Phys.* **354** (2017) 671 [[arXiv:1503.04817](#)] [[INSPIRE](#)].
- [39] K. Ohmori, H. Shimizu, Y. Tachikawa and K. Yonekura, *6d $\mathcal{N} = (1, 0)$ theories on S^1/T^2 and class S theories: part II*, *JHEP* **12** (2015) 131 [[arXiv:1508.00915](#)] [[INSPIRE](#)].
- [40] D. Gaiotto and S.S. Razamat, *Exceptional Indices*, *JHEP* **05** (2012) 145 [[arXiv:1203.5517](#)] [[INSPIRE](#)].
- [41] S. Cremonesi, A. Hanany, N. Mekareeya and A. Zaffaroni, *Coulomb branch Hilbert series and Three Dimensional Sicilian Theories*, *JHEP* **09** (2014) 185 [[arXiv:1403.2384](#)] [[INSPIRE](#)].
- [42] I. Bah, A. Passias and A. Tomasiello, *AdS5 compactifications with punctures in massive IIA supergravity*, [arXiv:1704.07389](#) [[INSPIRE](#)].
- [43] Y. Tachikawa, *$\mathcal{N} = 2$ Supersymmetric Dynamics for Pedestrians*, *Lect. Notes Phys.* **890** (2013) 2014 [[arXiv:1312.2684](#)] [[INSPIRE](#)].
- [44] J.J. Heckman, T. Rudelius and A. Tomasiello, *6D RG Flows and Nilpotent Hierarchies*, *JHEP* **07** (2016) 082 [[arXiv:1601.04078](#)] [[INSPIRE](#)].
- [45] N. Mekareeya, T. Rudelius and A. Tomasiello, *T-branes, Anomalies and Moduli Spaces in 6D SCFTs*, [arXiv:1612.06399](#) [[INSPIRE](#)].