



# HHS Public Access

Author manuscript

*Prog Brain Res.* Author manuscript; available in PMC 2018 March 16.

Published in final edited form as:

*Prog Brain Res.* 2008 ; 167: 137–149. doi:10.1016/S0079-6123(07)67010-1.

## Early care experiences and HPA axis regulation in children: a mechanism for later trauma vulnerability

Megan R. Gunnar\* and Karina M. Quevedo

Institute of Child Development, University of Minnesota, Minneapolis, MN 55455, USA

### Abstract

Post-traumatic stress disorder (PTSD) is associated with functional abnormalities of the hypothalamic- pituitary-adrenocortical (HPA) axis. Emerging evidence suggests that failures in social regulation of the HPA axis in young children manifested as neglectful or abusive care may play a role in shaping cortico-limbic circuits involved in processing experiences threatening experiences encountered later in life. Low cortisol levels, particularly near the peak of the diurnal rhythm, have been reported in abused, neglected and deprived children. Thus early imprinting effects of parenting quality on the HPA system regulation may be one of the mechanisms causing heightened risk of PTSD in responses to later trauma. However there is also evidence that the altered patterns of cortisol production seen in the context of early adverse care are not permanent, and remit once the care children receive improves. What awaits study is whether periods of atypical cortisol levels and altered HPA function early in life, even if transient, impact brain development in ways that heighten vulnerability to PTSD in response to traumas experienced later.

### Keywords

stress; early experience; cortisol; children

---

Post-traumatic stress disorder (PTSD) develops in only a subset of individuals who experience trauma. Hyper-arousal of threat-responsive neurobiological systems, including the sympathetic nervous system, and hypo-arousal of the hypothalamic- pituitary- adrenocortical (HPA) system at the time of trauma exposure are hypothesized to represent pre-existing risk factors for the development of PTSD (for review, see Delahanty and Nugent, 2006). Hypo-activity of the HPA axis is believed to enhance risk of PTSD perhaps through failing to counter-regulate hyper-arousal in other trauma-sensitive neurobiological systems (Yehuda, 2001). In adults, prior trauma increases both the risk of PTSD and the risk of blunted cortisol responses to trauma (Delahanty and Nugent, 2006). Exposure to trauma during childhood is believed to be particularly critical in increasing the risk of PTSD in response to later traumatic experiences (Teicher et al., 2002). In studies predicting PTSD from early childhood experiences, however, measures of childhood trauma include a range of experiences from sexual abuse to more subtle forms of emotional abuse (Yehuda, 2002). Furthermore, the animal models used to support arguments that childhood abuse shapes vulnerability to developing PTSD do not clearly point to childhood trauma as a necessary

---

\*Corresponding author. Tel.: +1-612-624-2846; Fax: +1-612-624-6373; gunnar@umn.edu.

antecedent (Heim et al., 1997). Rather, they point to lack or loss of expectable parental stimulation as critical in altering the development of the HPA axis and neurobiological systems involved in regulating reactions to stressful and traumatic events (Gunnar and Fisher, 2006).

In the following review we will examine the impact of variations in parental care on activity of the HPA axis in animal models and young children. Our focus on the HPA axis is based on the putative role of this neuroendocrine system in the etiology of PTSD (Yehuda, 2001). We will argue that parental care provides a social regulator of activity of the HPA axis in infants and young children (Gunnar and Donzella, 2002). Disturbances in parental care, thus, disrupt normal regulation of this neuroendocrine system. However, the effects of these disruptions are varied. In some instances, disturbances in the typical circadian rhythm of the axis are observed, in others the presence of the caregiver fails to buffer activation of the axis in response to events that produce fearful behavior, while in still others the parent's presence and interaction with the child stimulates elevations in cortisol. Whether these different effects on HPA axis activity reflect common or varied elements in the parent-offspring relationships are not yet known. However, there is now substantial evidence that parental care impacts activity of the HPA axis in the young of many species, including our own. As such, this evidence supports arguments that failures in the caregiving system may increase the risk of PTSD through altering the development of stress- and threat-responsive neurobiological systems. This review assumes the reader's knowledge of the neuroanatomy and physiology of the HPA axis (for review, see de Kloet, 1991; Herman et al., 2005) and mechanisms through which altered activity of this axis may impact neurodevelopment (for review, see Gunnar and Vazquez, 2006).

### **Animal models: parental care and HPA activity**

Much of the animal work has been conducted on rats. During infancy (postnatal days 4–14) rats exhibit low basal levels of corticosterone (the predominant steroid hormone produced by the HPA axis in rodents) and hypo-responsiveness of the HPA axis to many stressors (Rosenfeld et al., 1992). This hypo-responsive period in rats is maintained by specific components of maternal care: licking and grooming and the delivery of milk into the gut (Suchecki et al., 1993a, b, 1995). Both manipulations that alter licking and grooming and normal variations in this aspect of maternal care produce long-term alterations in the HPA axis and in emotional behavior. Short (3–5 min) repeated daily separations in infancy result in the development of attenuated HPA reactivity and fearfulness to stressors encountered later in life. In contrast, longer separations (e.g., 180 min daily) produce the development of a more hyper-responsive HPA axis and a more fearful animal (for review, see Sanchez et al., 2001). Repeated, prolonged separations also produce numerous other alterations that support vulnerability to stressors throughout life (Cirulli et al., 2000, 2003; Roceri et al., 2004). Early, repeated separations are believed to produce their effects, in part, through altering maternal care: increasing maternal licking and grooming in the case of brief, daily separations and reducing maternal licking and grooming in the case of prolonged daily separations (Heim and Nemeroff, 2001; Denenberg, 1999; Tang et al., 2006).

The argument that parental care mediates the impact of early separations is enhanced by evidence that normal variations in maternal licking and grooming produce the same effects as early separations. When the offspring of high licking and grooming dams are followed, they exhibit better regulation of the HPA axis and reduced fearful behavior in adulthood (Caldji et al., 1998). These effects are not due to common maternal and offspring genetics, as they can be produced in pups cross-fostered from low-to-high licking and grooming dams (Francis and Meaney, 1999). Furthermore, there is now good evidence that maternal care in rats produces long-term impacts on stress and emotional behavior through affecting methylation of the glucocorticoid receptor (GR) gene in the hippocampus. This gene produces the GR, the receptor that is critically involved in negative feedback of the HPA axis in response to psychosocial stressors (De Kloet, 1991). In response to low maternal care (i.e., low licking and grooming), there is greater methylation of the GR gene which results in fewer glucocorticoid receptors in the hippocampus and poorer negative feedback regulation of the HPA axis (Meaney and Szyf, 2005). Pharmacological intervention can correct these impacts on GR methylation. Thus offspring of low licking and grooming dams who were centrally infused with methionine showed a reversal of the effects of maternal behavior on the fearful behavior, HPA axis regulation and GR methylation (Weaver et al., 2005). Postnatal experiences can also alter some, but not all of these effects. Thus exposure of juvenile rats to an enriched environment prevents expression of the heightened fearfulness and elevated corticosterone in response to stressors, but it does not alter patterns of GR methylation, suggesting that these animals may maintain a vulnerability to traumatic stimulation later in life (Francis et al., 2002).

In non-human primates, the work on early parental care has progressed through the lens of attachment theory. Through this lens, the mother–infant relationship is viewed as a stress buffer (Bowlby, 1969; Suomi, 1995). The natural ecology of most primate species is social. Mother and infant live in troupes that include other mothers and infants, adult males and the infant’s older siblings. Species differ in how often the infant is parented by other members of the troupe — a phenomenon termed alloparenting or aunting. Although the availability of alloparents does not prevent elevations in cortisol to maternal separation, it does result in a more rapid termination of the HPA response and a reduced likelihood that despair behavior will develop during prolonged separations (Levine and Wiener, 1988). Thus, in primate infants maternal buffering is a function that can be provided to some extent by other nurturing conspecifics.

While repeated prolonged separations in rats may produce hyper-cortisolism, in monkeys they may reduce HPA activity (Levine et al., 1997; Dettling et al., 2002; Sanchez et al., 2005). In rhesus macaques, patterns of low basal output of cortisol following repeated separations have been accompanied by evidence of heightened startle responses (Sanchez et al., 2005), a pattern reminiscent of PTSD. Similarly, manipulations like variable foraging that produced disturbed mother–infant relations do not necessarily produce hyper-activation of the HPA axis in adulthood. Nevertheless, animals reared in these paradigms do exhibit elevated corticotropin-releasing hormone (CRH) in cerebrospinal fluid, altered serotonergic and noradrenergic activity and fearful behavior (Rosenblum and Andrews, 1994). Likewise, animals reared under conditions of parental deprivation as adults do not exhibit heightened HPA responses to stressors or neurochemical and structural changes in the HPA system (for

review, see Sanchez et al., 2001), but do exhibit changes in monoamine systems (Kraemer et al., 1989), hippocampal development (Siegel et al., 1993) and in CRH receptors in the prefrontal cortex and amygdala (Sanchez et al., 1999). These differences from the rodent model likely reflect the greater maturity of the HPA axis at birth relative to its maturity in the rat (Gunnar and Vazquez, 2006). To summarize, in non-human primates parental care is a potent regulator of activity of the infant's HPA axis; however, hyper-reactivity of the axis as a consequence of adverse early care has typically not been observed. Despite a general failure to observe hyper-reactivity of the axis following adverse early care experiences, in non-human primates many of the anxiety-associated behavioral and neurobiological sequelae of disturbances in early parental care have been observed.

## Human development and the hypothalamic-pituitary axis

With these animal data in mind, we turn now to data on parental care and activity of the HPA axis in human infants and children. These data, however, must be viewed in the context of developmental changes in HPA axis activity. We will describe these changes first.

### Development of HPA activity during childhood

The HPA axis is highly responsive to stressors at birth; indeed, HPA stress responses are observed by as early as 18–20 weeks of gestation (Giannakoulopoulos et al., 1999). In adults, approximately 80% of circulating cortisol is usually bound to corticosteroid-binding globulin (CBG) and is thus rendered biologically inactive. However, CBG levels are low in the human newborn and do not achieve adult levels until about 6 months (Hadjian et al., 1972). As a result, plasma levels of total cortisol are low in the newborn and increase over the first months of life; while the free fraction of the hormone is as high or higher than levels observed in older infants and children (Gunnar et al., 1988). In addition, when salivary measures of cortisol are used which reflect only the free fraction of the hormone, newborns show marked elevations in cortisol to even very minor perturbations (e.g., undressing, weighing and measuring; Gunnar et al., 1992). Therefore, by 6 months of age, the HPA system of the human child is relatively mature.

Over the first year of life, the human HPA system becomes progressively less responsive to stressors (Gunnar et al., 1996). Indeed, by one year of age it becomes difficult to elevate cortisol to a variety of events, which nonetheless, elicit significant behavioral distress (Ramsay and Lewis, 1994; Gunnar and Donzella, 1999). As will be discussed below, this apparent hypo-responsivity of the HPA axis reflects the fact that during the first year of life the human HPA axis comes under strong social regulation or parental buffering (Gunnar and Donzella, 2002). During childhood, small increases in cortisol can be observed to the experimentally induced stressors (Blair et al., 2005), although in most cases only some children exhibit these elevations (Talge et al., 2007). Likewise, stressors such as the beginning kindergarten or a new school year and important school exams produce cortisol increases in some children (Boyce et al., 1995; Spangler, 1995; Davis et al., 1999; Bruce et al., 2002).

Epidemiological data indicate that onset of stress-related emotional disorders increases during adolescence (Angold and Rutter, 1992). It has been suggested that this heightened

vulnerability to stressors may reflect increased reactivity of the HPA axis and other neurobiological changes over the transition to adolescence (Spear, 2000). There is now good evidence from both cross-sectional and longitudinal studies that basal cortisol levels increase as children progress into adolescence, perhaps related to pubertal changes (Gunnar and Vazquez, 2006). Indeed, a recent study found that adolescents at higher stages of pubertal development had more elevated overall diurnal cortisol curves reflected in cortisol levels sampled throughout the day (Adam, 2006). Several studies now suggest that over the pubertal transition the HPA system becomes more adult-like in its responses to stressors (Gunnar, in preparation; Stroud, in preparation). It thus seems likely that the impact of early care experiences on stress reactivity and regulation may not be fully evident until after the pubertal transition.

### **Normative variations in parental care**

As in research on non-human primates, regulation of the human infant's HPA system has been viewed through the lens of attachment theory. By the end of the first year, infants develop strong attachment relationships with one or a few consistent caregivers (Bowlby, 1969). The attachment figure provides a source of comfort and stress modulation. Patterns of behavior in response to mildly stressful provocation (stranger approach and brief separations) have been described which presumably reflect differences in the child's sense of security in the attachment relationship (Ainsworth et al., 1978). Secure relationships are characterized by caregivers that are consistently responsive and sensitive to their children's needs for both exploration and reassurance (Ainsworth et al., 1978). Thus, both parental sensitivity/ responsiveness and patterns of infant attachment behavior have been examined as a means of examining the parental buffering hypothesis in humans.

In general, both parental sensitivity/responsiveness and secure attachment relationships buffer the child's HPA axis in response to stressors. For example, Nachmias et al. (1996) exposed 18-month-olds to a series of events that elicited wary or fearful reactions in approximately half of the participants. Elevations in cortisol, however, were only observed in fearful toddlers who were insecurely attached to the parent who was with them during testing. Ahnert et al. (2004) examined cortisol responses of toddlers when they visited a new child care center in the presence of their mothers. In their mother's presence, but not during the child's initial days without mother at child care, securely attached infants exhibited markedly lower cortisol elevations compared to insecurely attached toddlers. In several studies, Spangler has demonstrated elevations in cortisol only among insecurely but not securely attached infants (for review, see Spangler and Grossmann, 1999).

Insensitive patterns of parental care early in the first year of life also predict larger cortisol responses to stressors (Gunnar et al., 1996). Indeed, when mothers were asked to play with their infants, infants with more sensitive mothers exhibited decreases in cortisol over the play period, while those with less sensitive mothers exhibited small increases in cortisol (Spangler et al., 1994). Harsh parenting (spanking) has also been shown to predict larger cortisol responses to mildly stressful events in infants (Bugental et al., 2003). Furthermore, when provided (randomly) with a sensitive and responsive surrogate caregiver, even 9-month-old infants with negative emotional temperaments did not show a cortisol elevation to

30 min of maternal separation; while those given an insensitive, unresponsive surrogate caregiver did (Gunnar et al., 1992). Similar findings have been obtained for toddlers and preschoolers in full-day out-of-home childcare (for review, see Gunnar and Donzella, 2002). These data support the hypothesis that children who experience less supportive and responsive care may frequently experience stressor-induced elevations in cortisol that might, over time, impact neurobehavioral development (see also De Kloet et al., 1996).

Studies of “at risk” children also support the argument that early care may have long-term impacts on vulnerability to stressful or traumatic events. A number of studies have been conducted on children whose mothers experience postpartum and/or major depression in the child’s first years of life. During bouts of clinical depression, mothers are typically less responsive and more rejecting with their infants and young children (Ashman et al., 2002). Because bouts of clinical depression are associated with disturbances in parenting, associations between when the mother was depressed during the child’s development and child outcomes provide a window into the timing of disturbances in parental care and neurobehavioral development.

Preschoolers whose mothers have a history of clinical depression or who report high number of depressive symptoms during the child’s first 2 years, exhibit elevated home cortisol levels, particularly in the context of on-going stress in the family (Dawson and Ashman, 2000; Essex et al., 2002). School-aged children exposed to more prolonged periods of maternal depression in infancy and early childhood have been noted to have higher and more variable early morning cortisol levels even after controlling for maternal depression later in the child’s life (Halligan et al., 2004; Lupien et al., 2000). It is not clear, however, whether maternal depression during a child’s early development increases reactivity of the HPA axis, as this has been noted only for girls and then only for girls with concurrent internalizing behavior problems (Ashman et al., 2002). Several of these studies implicate elevated cortisol levels in the etiology of internalizing problems in childhood (Smider et al., 2002) and the emergence of depression in response to stressful events during adolescence (Halligan et al., 2007) for the offspring of depressed mothers. Unfortunately, although maternal depression has often been shown to be correlated with less sensitive and responsive care (Dawson and Ashman, 2000), in none of these studies was parental care examined as a mediator of the impact of maternal depressive symptoms on HPA axis activity. Furthermore, they do not allow us to disentangle genetic vulnerabilities in children of depressed mothers from the impact of postnatal care. That is, we do not know whether bouts of disturbed parental care would have the same impact on children who did not have a high genetic load for depression.

To summarize, studies of variations in parental care within the normal range in humans strongly indicate that less sensitive and responsive caregiving is associated with poorer concurrent regulation of the HPA axis in children. Sensitive and responsive care and associated secure attachment relationships appear to provide a powerful buffer for the HPA axis during early development. Children exposed to less sensitive/responsive care and those of mothers who use more harsh (though not necessarily abusive) discipline practices exhibit higher basal cortisol levels and their mother’s presence does not prevent elevations in cortisol to the type of everyday stressors experienced by young children. Studies of the

offspring of depressed mothers appear to indicate that these early care experiences may produce permanent alteration in HPA axis functioning. However, because the children of depressed mothers may have inherited genes that increase their vulnerability to variations in maternal care, it is difficult to know whether we can extrapolate these findings to children with different genetic heritages.

### **Severe deprivation, neglect and abuse**

Thus far we have reviewed human studies examining care experiences, which though not perhaps optimal, would not reach criteria for maltreatment. Unfortunately for comparisons with animal studies, HPA reactivity to stressors has not yet been studied in children exposed to severe deprivation, neglect and abuse (see Heim et al., 2004, for such studies of adults exposed to abuse during childhood). Instead, the focus in the child research has been on basal activity of the axis. Evidence for both elevated and extremely low cortisol levels have been obtained. When extremely low levels have been noted, these have most often been seen in early morning, near the peak of the circadian rhythm (Gunnar and Vazquez, 2001). Because the diurnal slope of cortisol is largely determined by the height of the early morning peak, children with the extremely low morning levels have a flat daytime cortisol rhythm. Whether suppressed or elevated basal cortisol levels are noted, however, may depend on the age of the child, the type of maltreatment, whether the maltreatment is on-going and how long it has been since the child was placed in more supportive care arrangements.

There are now several studies of preschool-aged children in foster care that reveal quite different patterns of HPA axis functioning than those noted in studies of children of depressed mothers. Dozier et al. (2006) found that approximately 40% of their sample of foster preschoolers exhibited extremely low early morning cortisol levels, while about 20% exhibited markedly high levels relative to children living in non-maltreating families. Bruce et al. (under review) noted similar findings and also found that prior neglect, rather than abuse, predicted low early morning levels. In a study following preschoolers in foster care, children in regular foster care increasingly exhibited low early morning levels over time (Fisher et al., 2007). These children also were more likely than children of the same social class living with their non-maltreating parents to develop insecure patterns of attachment over this time period (Fisher et al., in press). These results are consistent with data on toddlers living in institutions (i.e., orphanages). Orphanage-reared children also have very low early morning cortisol levels and a relatively flat pattern of cortisol production over the day (Carlson and Earls, 1997). Institutional care tends to be characterized by little individualized care, low levels of adult-child interaction, but especially among infants and toddlers, little frank physical or sexual abuse (Zeanah et al., 2006). Benign neglect is a term used to typify institutional care of infants and young children, and hence these data appear remarkably similar to the results reported for preschool-aged children in foster care.

The findings for children exposed to significant neglect and deprivation are also similar to reports of hypo-cortisolism among adults who experience chronic stress (Heim et al., 2000). One mechanism invoked to explain the development of hypocortisolism is down-regulation of the HPA axis at the level of the pituitary in response to chronic CRH drive (Fries et al., 2005). Accordingly, one might expect that prior to appearance of very low early morning

cortisol levels, there might be a period of exceptionally high cortisol (Fries et al., 2005). Unfortunately, most of the children examined in both the foster care and institutional care studies experienced adverse care from birth. Examining such children earlier in their care histories also means examining them when they are infants. We have already noted the marked changes in HPA axis activity over the course of the first year or two of life. Developmental change and duration of exposure to maltreatment are difficult to disentangle. Notably, however, the one foster care study examining infants, as opposed to preschoolers, reported exceptionally high cortisol levels for infants in regular foster care relative to infants of the same social class living with their non-maltreating parents (Dozier et al., in press).

There is little reason to believe that this hypocortisolism in basal activity of the HPA axis observed early in life in the context of neglect and deprivation is permanent. At least in work with adults, low cortisol output can be reversed by alleviating chronic stress conditions (for review, see Fries et al., 2005). Among infants in foster care, training foster parents to be more sensitive to the needs of previously neglected and abused infants results in normalization of basal cortisol levels with only 10 weeks of parent training (Dozier et al., in press). What is not clear is whether alterations in HPA axis activity persist once children have time to recover in more supportive care arrangements. In addressing this question we will not cover studies of children with chronic PTSD (Carrion et al., 2001, 2002; De Bellis et al., 1999a, b), although we will consider whether current psychiatric morbidity might explain associations between early care experiences and later HPA activity.

Two studies have now examined basal cortisol levels in children adopted from institutions. For children adopted from the type of global deprivation that characterized Romanian institutions in the early 1990s, elevated rather than suppressed cortisol levels were noted 6–7 years post-adoption (Gunnar et al., 2001). Most of these children were severely growth delayed at adoption due to poor health, nutrition and social stimulation in the orphanage (Rutter, 1998). In a subsequent study, growth delay at adoption rather than duration of institutional care or growth parameters at assessment predicted elevated early morning cortisol levels (Kertes et al., 2007). In this latter study, very few of the children who were growth delayed at adoption met criteria for internalizing problems at assessment; thus, it was unlikely that concurrent psychopathology moderated the relations between early care, growth delay and basal cortisol. Studies of maltreated children suggest that only those who experience the most severe and prolonged maltreatment may have elevated cortisol levels (Cicchetti and Rogosch, 2001a; De Bellis et al., 1999a). However, these factors also increase the risk of both affective and conduct disorder, and thus cortisol levels and psychopathology are difficult to disentangle (Cicchetti and Rogosch, 2001b). Even after their maltreatment has been exposed and child protective services are involved, on-going stress in the family, rather than maltreatment history, may explain increased HPA activity in the children (Kaufman et al., 1997).

Taken together, the studies of post-institutionalized children and children with maltreatment histories do not suggest that for most children, periods of adverse early care increases the basal set point of the HPA axis. Lack of a persistent increase in basal cortisol levels pursuant to early life stress would be consistent with findings in the animal literature where HPA reactivity but not basal levels appear to be affected. When early life stress results in long-



term increases in basal HPA levels even after the conditions producing the stress have been improved, the child either appears to have experienced growth failure due to the stressful conditions or, as in the data on children of mothers with depression, may carry a high genetic load for depression or other affective disorders. However, because there are no studies on HPA stress reactivity in children exposed as infants to adverse care, we cannot conclude that their early experiences failed to impact how the axis responds to stressors later in development. This, rather than long-term changes in basal activity is what has been noted in the early experience animal literature and this is what remains to be studied in children exposed to significantly adverse care early in life. Obviously, studies of HPA axis stress reactivity pursuant to early deprivation and maltreatment are crucial to fully understanding how early parental care may impact vulnerability to PTSD. However, because a fully adult-like patterns of HPA responses to stressors may not emerge until sometime in adolescence, studies of the impact of adverse care on subsequent HPA responses to stressors will need to take into account the child's developmental level.

## Conclusions

Pre-clinical human studies, animal models and recent studies with children deprived of adequate parenting provide evidence that the HPA axis is under strong social regulation early in life. Accordingly, variations in parental care have potent effects on concurrent regulation of the axis. Unlike in studies of rats, however, in neither non-human primates nor human children do early adverse care experiences appear to produce permanent alterations activity of the HPA axis. The exceptions to this conclusion can be found in studies of children of depressed mothers, children whose early care produced severe physical growth delays, and in children who develop chronic PTSD. Few studies of human children exposed to adverse early care have examined HPA responses to laboratory stressors or to real-life traumatic events. These studies are needed. Failure to observe long-term alterations in HPA axis activity in humans does not mean that failure of the caregiving system to provide regulation of this axis early in development is of no consequence. Rather, as in nonhuman primates, it may suggest that the HPA axis, being relatively mature at birth, is less susceptible to the long-term alterations noted in studies of infant rats. Nonetheless, failures in social regulation of the HPA axis in young children because of inadequate and/or abusive care may play a role in shaping cortico-limbic circuits involved in processing traumatic experiences. Evidence for this hypothesis, however, will require finding disturbances in HPA activity in early childhood that are reversible by the provision of more supportive care, but that nevertheless predict the development of individual differences in neurobiological systems associated with risk for PTSD. Longitudinal studies of this sort are just beginning to be conducted (Dozier et al., in press; Fisher et al., under review).

## Acknowledgments

Preparation of this manuscript was supported by a National Institute of Mental Health pre-doctoral fellowship (T32 MH15755) to the second author, and a National Institute of Mental Health Senior Scientist Award (K05 MH66208) to the first author.

## Abbreviations

<b>CBG</b>	corticosteroid-binding globulin
<b>CRH</b>	corticotropin-releasing hormone
<b>GR</b>	glucocorticoid receptor
<b>HPA</b>	hypothalamic-pituitary-adrenocortical
<b>PTSD</b>	post-traumatic stress disorder

## References

- Adam EK. Transactions among adolescent trait and state emotion and diurnal and momentary cortisol activity in naturalistic settings. *Psychoneuroendocrinology*. 2006; 31(5):664–679. [PubMed: 16584847]
- Ahnert L, Gunnar MR, Lamb ME, Barthel M. Transition to child care: associations with infant–mother attachment, infant negative emotion, and cortisol elevations. *Child Dev*. 2004; 75(3):639–650. [PubMed: 15144478]
- Ainsworth, MS., Blehar, MC., Waters, E., Wall, S. *Patterns of Attachment: A Psychological Study of the Strange Situation*. Lawrence Erlbaum; Oxford, England: 1978. p. 391
- Angold A, Rutter M. Effects of age and pubertal status on depression in a large clinical sample. *Dev. Psychopathol*. 1992; 4(1):5–28.
- Ashman SB, Dawson G, Panagiotides H, Yamada E, Wilkinson CW. Stress hormone levels of children of depressed mothers. *Dev. Psychopathol*. 2002; 14(2):333–349. [PubMed: 12030695]
- Blair C, Granger D, Peters R. Cortisol reactivity is positively related to executive function in preschool children attending head start. *Child Dev*. 2005; 76(3):554–567. [PubMed: 15892778]
- Bowlby, J. *Volume 1 of Attachment and Loss*. 2. Basic Books; New York, NY: 1969. Attachment; p. 425
- Boyce WT, Adams S, Tschann JM, Cohen F, Wara D, Gunnar MR. Adrenocortical and behavioral predictors of immune responses to starting school. *Pediatr. Res*. 1995; 38(6):1009–1017. [PubMed: 8618776]
- Bruce J, Davis EP, Gunnar MR. Individual differences in children’s cortisol response to the beginning of a new school year. *Psychoneuroendocrinology*. 2002; 27(6):635–650. [PubMed: 12084657]
- Bruce, J., Pears, KC., Levine, S., Fisher, PA. *Morning cortisol levels in preschool-aged foster children: differential effects of maltreatment type*. Oregon Social Learning Center; Eugene, Oregon: (under review)
- Bugental DB, Martorell GA, Barraza V. The hormonal costs of subtle forms of infant maltreatment. *Horm. Behav*. 2003; 43(1):237–244. [PubMed: 12614655]
- Caldji C, Tannenbaum B, Sharma S, Francis D, Plotsky PM, Meaney MJ. Maternal care during infancy regulates the development of neural systems mediating the expression of fearfulness in the rat. *Proc. Natl. Acad. Sci. U.S.A.* 1998; 95(9):5335–5340. [PubMed: 9560276]
- Carlson M, Earls F. Psychological and neuroendocrinological sequelae of early social deprivation in institutionalized children in Romania. *Ann. N.Y. Acad. Sci.* 1997; 807:419–428. [PubMed: 9071367]
- Carrion VG, Weems CF, Eliez S, Patwardhan A, Brown W, Ray RD, et al. Attenuation of frontal asymmetry in pediatric posttraumatic stress disorder. *Biol. Psychiatry*. 2001; 50(12):943–951. [PubMed: 11750890]
- Carrion VG, Weems CF, Ray RD, Glaser B, Hessel D, Reiss AL. Diurnal salivary cortisol in pediatric posttraumatic stress disorder. *Biol. Psychiatry*. 2002; 51(7):575–582. [PubMed: 11950459]
- Cicchetti D, Rogosch FA. Diverse patterns of neuroendocrine activity in maltreated children. *Dev. Psychopathol*. 2001a; 13(3):677–693. [PubMed: 11523854]

- Cicchetti D, Rogosch FA. The impact of child maltreatment and psychopathology on neuroendocrine functioning. *Dev. Psychopathol.* 2001b; 13(4):783–804. [PubMed: 11771908]
- Cirulli F, Alleva E, Antonelli A, Aloe L. NGF expression in the developing rat brain: effects of maternal separation. *Dev. Brain Res.* 2000; 123:129–134. [PubMed: 11042341]
- Cirulli F, Berry A, Alleva E. Early disruption of the mother–infant relationship: effects on brain plasticity and implications for psychopathology. *Neurosci. Biobehav. Rev.* 2003; 27(1–2):73–82. [PubMed: 12732224]
- Davis EP, Donzella B, Krueger WK, Gunnar MR. The start of a new school year: Individual differences in salivary cortisol response in relation to child temperament. *Dev. Psychobiol.* 1999; 35(3):188–196. [PubMed: 10531531]
- Dawson, G., Ashman, S. On the origins of a vulnerability to depression: the influence of early social environment on the development of psychobiological systems related to risk for affective disorder. In: Nelson, CA., editor. *The Effects of Adversity on Neurobehavioral Development*. Minnesota Symposia on Child Psychology. Vol. 31. Erlbaum Associates; New York: 2000. p. 245–278.
- De Bellis MD, Baum AS, Birmaher B, Keshavan MS, Eccard CH, Boring AM, et al. Developmental traumatology. Part I: biological stress systems. *Biol. Psychiatry.* 1999a; 45(10):1259–1270. [PubMed: 10349032]
- De Bellis MD, Keshavan MS, Clark DB, Casey BJ, Giedd JN, Boring AM, et al. Developmental traumatology. Part II: brain development. *Biol. Psychiatry.* 1999b; 45(10):1271–1284. [PubMed: 10349033]
- Delahanty DL, Nugent NR. Predicting PTSD prospectively based on prior trauma history and immediate biological responses. *Ann. N.Y. Acad. Sci.* 2006; 1071:27–40. [PubMed: 16891559]
- Denenberg VH. Commentary: is maternal stimulation the mediator of the handling effects in infancy? *Dev. Psychobiol.* 1999; 34(1):1–3. [PubMed: 9919428]
- Dettling AC, Feldom J, Pryce CR. Repeated parental deprivation in the infant common marmoset (*Callithrix jacchus*, primates) and analysis of its effects on early development. *Biol. Psychiatry.* 2002; 52(11):1037–1046. [PubMed: 12460687]
- Dozier M, Manni M, Gordon MK, Peloso E, Gunnar MR, Stovall-McClough KC, et al. Foster children's diurnal production of cortisol: an exploratory study. *Child Maltreat.* 2006; 11(2):189–197. [PubMed: 16595852]
- Dozier M, Peloso E, Sepulveda S, Manni M, Lindhiem O, Ackerman J, et al. Preliminary evidence from a randomized clinical trial: intervention effects on behavioral and biobehavioral regulation of young foster children. *J. Soc. Sci.* (in press).
- Essex MJ, Klein MH, Cho E, Kalin NH. Maternal stress beginning in infancy may sensitize children to later stress exposure: effects on cortisol and behavior. *Biol. Psychiatry.* 2002; 52(8):776–784. [PubMed: 12372649]
- Fisher P, Gunnar M, Dozier M, Bruce J, Pears K. Effects of therapeutic interventions for foster children on behavioral problems, caregiver attachment and stress regulatory neural systems. *Ann. N.Y. Acad. Sci.* (in press).
- Fisher PA, Stoolmiller M, Gunnar MR. Effects of a therapeutic intervention for foster preschoolers on diurnal cortisol activity. *Psychoneuroendocrinology.* 2007 Jul 24. [Epub ahead of print].
- Francis DD, Diorio J, Plotsky PM, Meaney MJ. Environmental enrichment reverses the effects of maternal separation on stress reactivity. *J. Neurosci.* 2002; 22(18):7840–7843. [PubMed: 12223535]
- Francis DD, Meaney MJ. Maternal care and the development of stress responses. *Curr. Opin. Neurobiol.* 1999; 9(1):128–134. [PubMed: 10072372]
- Fries E, Hesse J, Hellhammer J, Hellhammer DH. A new view on hypocortisolism. *Psychoneuroendocrinology.* 2005; 30(10):1010–1016. [PubMed: 15950390]
- Giannakouloupoulos X, Teixeira J, Fisk N, Glover V. Human fetal and maternal noradrenaline responses to invasive procedures. *Pediatr. Res.* 1999; 45(4):494–499. [PubMed: 10203140]
- Gunnar MR, Fisher P. Bringing basic research on early experience and stress neurobiology to bear on preventive interventions for neglected and maltreated children. *Dev. Psychopathol.* 2006; 18:651–677. [PubMed: 17152395]

- Gunnar, M., Marvinney, D., Isensee, J., Fisch, RO. Coping with uncertainty: new models of the relations between hormonal behavioral and cognitive processes. In: Palermo, D., editor. *Copying with Uncertainty: Biological, Behavioral, and Development Perspectives*. Erlbaum; Hillsdale, NJ: 1988. p. 101-130.
- Gunnar MR, Brodersen L, Krueger K, Rigatuso J. Dampening of adrenocortical responses during infancy: normative changes and individual differences. *Child Dev.* 1996; 67(3):877–889. [PubMed: 8706532]
- Gunnar, MR., Donzella, B. Looking for the Rosetta Stone: an essay on crying, soothing, and stress. In: Lewis, M., Ramsay, D., editors. *Soothing and Stress*. Lawrence Erlbaum Associates; Mahwah, NJ: 1999. p. 39-56.
- Gunnar MR, Donzella B. Social regulation of the cortisol levels in early human development. *Psychoneuroendocrinology.* 2002; 27(1–2):199–220. [PubMed: 11750779]
- Gunnar MR, Larson MC, Hertsgaard L, Harris ML, et al. The stressfulness of separation among nine-month-old infants: effects of social context variables and infant temperament. *Child Dev.* 1992; 63(2):290–303. [PubMed: 1611934]
- Gunnar MR, Morison SJ, Chisholm K, Schuder M. Salivary cortisol levels in children adopted from Romanian orphanages. *Dev. Psychopathol.* 2001; 13(3):611–628. [PubMed: 11523851]
- Gunnar, MR., Vazquez, D. Stress neurobiology and developmental psychopathology. In: Cicchetti, D., Cohen, D., editors. *Developmental Psychopathology: Developmental Neuroscience*. 2. Vol. 2. Wiley; New York: 2006. p. 533-577.
- Gunnar, MR., Wewerka, S., Frenn, K., Long, JD., Griggs, C. Developmental Changes in HPA Activity over the Transition to Adolescence: Normative Changes and Associations with Puberty. University of Minnesota; (in preparation)
- Gunnar MR, Vazquez DM. Low cortisol and a flattening of expected daytime rhythm: potential indices of risk in human development. *Dev. Psychopathol.* 2001; 13(3):515–538. [PubMed: 11523846]
- Hadjian AJ, Chedin M, Chambaz EM. Characteristics of cortisol binding to plasma proteins: peculiar aspects of the perinatal period. *Arch. Fr. Pediatr.* 1972; 29(6):669–670. [PubMed: 5078368]
- Halligan SL, Herbert J, Goodyer L, Murray L. Disturbances in morning cortisol secretion in association with maternal postnatal depression predict subsequent depressive symptomatology in adolescents. *Biol. Psychiatry.* 2007; 62:40–46. [PubMed: 17188253]
- Halligan SL, Herbert J, Goodyer IM, Murray L. Exposure to postnatal depression predicts elevated cortisol in adolescent offspring. *Biol. Psychiatry.* 2004; 55:376–381. [PubMed: 14960290]
- Heim C, Ehlert U, Hellhammer DH. The potential role of hypocortisolism in the pathophysiology of stress-related bodily disorders. *Psychoneuroendocrinology.* 2000; 25(1):1–35. [PubMed: 10633533]
- Heim C, Nemeroff CB. The role of childhood trauma in the neurobiology of mood and anxiety disorders: preclinical and clinical studies. *Biol. Psychiatry.* 2001; 49:1023–1039. [PubMed: 11430844]
- Heim, C., Owens, MJ., Plotsky, PM., Nemeroff, CB. The role of early adverse life events in the etiology of depression and posttraumatic stress disorder. Focus on corticotropin-releasing factor. In: Yehuda, RM., Alexander, C., editors. *Psychobiology of Posttraumatic Stress Disorder*. Vol. 821. New York Academy of Sciences; New York: 1997. p. 194-207.
- Heim C, Plotsky P, Nemeroff CB. The importance of studying the contributions of early adverse experiences to the neurobiological findings in depression. *Neuropsychopharmacology.* 2004; 29:641–648. [PubMed: 15034558]
- Herman JP, Ostrander MM, Mueller NK, Figueiredo H. Limbic system mechanisms of stress regulation: hypothalamo-pituitary-adrenocortical axis. *Prog. Neuropsychopharmacol. Biol. Psychiatry.* 2005; 29(8):1201–1213. [PubMed: 16271821]
- Kaufman J, Birmaher B, Perel J, Dahl RE, Moreci P, Nelson B, et al. The corticotropin-releasing hormone challenge in depressed abused, depressed nonabused, and normal control children. *Biol. Psychiatry.* 1997; 42(8):669–679. [PubMed: 9325560]
- Kertes DA, Gunnar MR, Madsen NJ, Long J. Early deprivation and home basal cortisol levels: a study of internationally-adopted children. *Dev. Psychopathol.* 2007 (in press).

- de Kloet ER. Brain corticosteroid receptor balance and homeostatic control. *Front. Neuroendocrinol.* 1991; 12(2):95–164.
- de Kloet ER, Rots NY, Cools AR. Brain-corticosteroid hormone dialogue: slow and persistent. *Cell. Mol. Neurobiol.* 1996; 16(3):345–356. [PubMed: 8818401]
- Kraemer GW, Ebert MH, Schmidt DE, McKinney WT. A longitudinal study of the effect of different social rearing conditions on cerebrospinal fluid norepinephrine and biogenic amine metabolites in rhesus monkeys. *Neuropsychopharmacology.* 1989; 2(3):175–189. [PubMed: 2477005]
- Levine S, Lyons DM, Schatzberg AF. Psychobiological consequences of social relationships. *Ann. N.Y. Acad. Sci.* 1997; 807:210–218. [PubMed: 9071353]
- Levine S, Wiener SG. Psychoendocrine aspects of mother–infant relationships in nonhuman primates. *Psychoneuroendocrinology.* 1988; 13(1–2):143–154. [PubMed: 3287413]
- Lupien S, King S, Meaney MJ, McEwen B. Child’s stress hormone levels correlate with mother’s socioeconomic status and depressive state. *Biol. Psychiatry.* 2000; 48:976–980. [PubMed: 11082471]
- Meaney MJ, Szyf M. Environmental programming of stress responses through DNA methylation: life at the interface between a dynamic environment and a fixed genome. *Dialogues Clin. Neurosci.* 2005; 7(2):103–123. [PubMed: 16262207]
- Nachmias M, Gunnar M, Mangelsdorf S, Parritz RH, Buss K. Behavioral inhibition and stress reactivity: the moderating role of attachment security. *Child Dev.* 1996; 67(2):508–522. [PubMed: 8625725]
- Ramsay DS, Lewis M. Developmental change in infant cortisol and behavioral response to inoculation. *Child Dev.* 1994; 65(5):1491–1502. [PubMed: 7982364]
- Roceri M, Cirulli F, Pessina C, Peretto P, Racagni G, Riva MA. Postnatal repeated maternal deprivation produces age-dependent changes of brain-derived neurotrophic factor expression in selected rat brain regions. *Biol. Psychiatry.* 2004; 55(7):708–714. [PubMed: 15038999]
- Rosenblum LA, Andrews MW. Influences of environmental demand on maternal behavior and infant development. *Acta Paediatr. Suppl.* 1994; 397:57–63. [PubMed: 7981475]
- Rosenfeld P, Suchecki D, Levine S. Multifactorial regulation of the hypothalamic–pituitary–adrenal axis during development. *Neurosci. Biobehav. Rev.* 1992; 16(4):553–568. [PubMed: 1480351]
- Rutter M. Developmental catch-up, and deficit, following adoption after severe global early privation, English and Romanian Adoptees (ERA) Study Team. *J. Child Psychol. Psychiatry.* 1998; 39(4):465–476. [PubMed: 9599775]
- Sanchez MM, Ladd CO, Plotsky PM. Early adverse experience as a developmental risk factor for later psychopathology: evidence from rodent and primate models. *Dev. Psychopathol.* 2001; 13(3):419–449. [PubMed: 11523842]
- Sanchez MM, Noble PM, Lyon CK, Plotsky PM, Davis M, Nemeroff CB, et al. Alterations in diurnal cortisol rhythm and acoustic startle response in nonhuman primates with adverse rearing. *Biol. Psychiatry.* 2005; 57(4):373–381. [PubMed: 15705353]
- Sanchez, MM., Young, LJ., Plotsky, P., Insel, TR. Different rearing conditions affect the development of corticotropin-releasing hormone (CRF) and vasopressin (AVP) systems in the non-human primate; Paper presented at the Society for Neuroscience Abstracts; October 23–28; Miami Beach, Florida. 1999.
- Siegel SJ, Ginsberg SD, Hof PR, Foote SL, Young WG, Kraemer GW, et al. Effects of social deprivation in prepubescent rhesus monkeys: immunohistochemical analysis of the neurofilament protein triplet in the hippocampal formation. *Brain Res.* 1993; 619(1–2):299–305. [PubMed: 8374785]
- Smider NA, Essex MJ, Kalin NH, Buss KA, Klein MH, Davidson RJ, et al. Salivary cortisol as a predictor of socioemotional adjustment during kindergarten: a prospective study. *Child Dev.* 2002; 73(1):75–92. [PubMed: 14717245]
- Spangler G. School performance, type A behavior and adrenocortical activity in primary school children. *Anxiety, Stress Coping Int. J.* 1995; 8(4):299–310.
- Spangler, G., Grossmann, K. Individual and physiological correlates of attachment disorganization in infancy. In: Salomon, J., George, C., editors. *Attachment Disorganization.* Guilford; New York, NY: 1999. p. 95-124.

- Spangler G, Schieche M, Ilg U, Maier U, Ackermann C. Maternal sensitivity as an external organizer for biobehavioral regulation in infancy. *Dev. Psychobiol.* 1994; 27(7):425–437. [PubMed: 7843497]
- Spear LP. The adolescent brain and age-related behavioral manifestations. *Neurosci. Biobehav. Rev.* 2000; 24(4):417–463. [PubMed: 10817843]
- Stroud, LR., Foster, E., Handwerker, K., Papandonatos, GD., Granger, DA., Kivlighan, KT., Niaura, R. Stress Response and the Adolescent Transition: Performance versus Peer Rejection Stress. Brown University Medical School; (in preparation)
- Suchecki D, Mazzafarian D, Gross G, Rosenfeld P, Levine S. Effects of maternal deprivation on the ACTH stress response in the infant rat. *Neuroendocrinology.* 1993a; 57(2):204–212. [PubMed: 8389994]
- Suchecki D, Nelson DY, Van Oers H, Levine S. Activation and inhibition of the hypothalamic-pituitary-adrenal axis of the neonatal rat: effects of maternal deprivation. *Psychoneuroendocrinology.* 1995; 20(2):169–182. [PubMed: 7899536]
- Suchecki D, Rosenfeld P, Levine S. Maternal regulation of the hypothalamic-pituitary-adrenal axis in the infant rat: the roles of feeding and stroking. *Dev. Brain Res.* 1993b; 75(2):185–192. [PubMed: 8261610]
- Suomi, SJ. Influence of attachment theory on ethological studies of biobehavioral development in nonhuman primates. In: Goldberg, MS., Kerr, RJ., editors. *Attachment Theory: Social, Developmental, and Clinical Perspectives.* The Analytic Press; Hillsdale, NJ: 1995. p. 185-201.
- Talge NM, Donzella B, Gunnar MR. Fearful temperament and stress reactivity among preschool-aged children. *Infant and Child Development.* 2007 (in press).
- Tang AC, Akers K, Reeb B, Romeo RD, McEwen B. Programming social, cognitive, and neuroendocrine development by early exposure to novelty. *Proc. Nat. Acad. Sci.* 2006; 103:15716–15721. [PubMed: 17030787]
- Teicher MH, Andersen SL, Polcarri A, Anderson CM, Navalta CP. Developmental neurobiology of childhood stress and trauma. *Psychiatric Clinics of North America.* 2002; 25:397–426. [PubMed: 12136507]
- Weaver IC, Champagne FA, Brown SE, Dymov S, Sharma S, Meaney MJ, et al. Reversal of maternal programming of stress responses in adult offspring through methyl supplementation: altering epigenetic marking later in life. *J. Neurosci. Biology of posttraumatic stress disorder.* 2005; 25(47):11045–11054.
- Yehuda R. Biology of posttraumatic stress disorder. *J. Clin. Psychiatry.* 2001; 62:41–46.
- Yehuda R. Current status of cortisol findings in posttraumatic stress disorder. *Psychiatr. Clin. North Am.* 2002; 25(2):341–368. [PubMed: 12136504]
- Zeanah, CH., Smyke, AT., Settles, L. Children in orphanages. In: McCartney, K., Phillips, DA., editors. *Handbook of Early Childhood Development.* Blackwell Publishing; Malden, MA: 2006. p. 224-254.