



## UvA-DARE (Digital Academic Repository)

### Early micronutrient supplementation protects against early stress-induced cognitive impairments

Naninck, E.F.G.; Oosterink, J.E.; Yam, K.-Y.; de Vries, L.P.; Schierbeek, H.; van Goudoever, J.B.; Verkaik-Schakel, R.N.; Plantinga, J.A.; Plosch, T.; Lucassen, P.J.; Korosi, A.

**DOI**

[10.1096/fj.201600834R](https://doi.org/10.1096/fj.201600834R)

**Publication date**

2017

**Document Version**

Final published version

**Published in**

The FASEB Journal

**License**

Article 25fa Dutch Copyright Act

[Link to publication](#)

**Citation for published version (APA):**

Naninck, E. F. G., Oosterink, J. E., Yam, K.-Y., de Vries, L. P., Schierbeek, H., van Goudoever, J. B., Verkaik-Schakel, R. N., Plantinga, J. A., Plosch, T., Lucassen, P. J., & Korosi, A. (2017). Early micronutrient supplementation protects against early stress-induced cognitive impairments. *The FASEB Journal*, 31(2), 505-518.  
<https://doi.org/10.1096/fj.201600834R>

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

*UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)*

## Early micronutrient supplementation protects against early stress–induced cognitive impairments

Eva F. G. Naninck,\* J. Efraim Oosterink,<sup>†</sup> Kit-Yi Yam,\* Lennart P. de Vries,\* Henk Schierbeek,<sup>†</sup> Johannes B. van Goudoever,<sup>†</sup> Rikst-Nynke Verkaik-Schakel,<sup>‡</sup> Josèe A. Plantinga,<sup>‡</sup> Torsten Plosch,<sup>‡</sup> Paul J. Lucassen,\* and Aniko Korosi\*<sup>1</sup>

\*Swammerdam Institute for Life Sciences, Center for Neuroscience, University of Amsterdam, Amsterdam, The Netherlands; <sup>†</sup>Department of Mother and Child, Emma Children’s Hospital, Academic Medical Center, Amsterdam, The Netherlands; and <sup>‡</sup>Department of Obstetrics and Gynecology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

**ABSTRACT:** Early-life stress (ES) impairs cognition later in life. Because ES prevention is problematic, intervention is needed, yet the mechanisms that underlie ES remain largely unknown. So far, the role of early nutrition in brain programming has been largely ignored. Here, we demonstrate that essential 1-carbon metabolism–associated micronutrients (1-CMAMs; *i.e.*, methionine and B vitamins) early in life are crucial in programming later cognition by ES. ES was induced in male C57Bl/6 mice from postnatal d (P)2–9. 1-CMAM levels were measured centrally and peripherally by using liquid chromatography-mass spectroscopy. Next, we supplemented the maternal diet with 1-CMAM only during the ES period and studied cognitive, neuroendocrine, neurogenic, transcriptional, and epigenetic changes in adult offspring. We demonstrate that ES specifically reduces methionine in offspring plasma and brain. Of note, dietary 1-CMAM enrichment during P2–9 restored methionine levels and rescued ES-induced adult cognitive impairments. Beneficial effects of this early dietary enrichment were associated with prevention of the ES-induced rise in corticosterone and adrenal gland hypertrophy did not involve changes in maternal care, hippocampal volume, neurogenesis, or global/*Nr3c1*-specific DNA methylation. In summary, nutrition is important in brain programming by ES. A short, early supplementation with essential micronutrients can already prevent lasting effects of ES. This concept opens new avenues for nutritional intervention.—Naninck, E. F. G., Oosterink, J. E., Yam, K.-Y., de Vries, L. P., Schierbeek, H., van Goudoever, J. B., Verkaik-Schakel, R.-N., Plantinga, J. A., Plosch, T., Lucassen, P. J., Korosi, A. Early micronutrient supplementation protects against early stress–induced cognitive impairments. *FASEB J.* 31, 505–518 (2017). www.fasebj.org

**KEY WORDS:** 1-carbon metabolism · HPA-axis · hippocampus · limited nesting material · cognition

Adversities early in life are strongly associated with decreased cognitive performance in adulthood, an increased vulnerability to developing psychopathology, and impaired hippocampal integrity and function, as established in several clinical (1, 2) and preclinical studies (3). As a result of a lack of insight into the key factors that contribute to these detrimental effects of early-life stress (ES), there are currently no strategies available to solve this problem. The past 50 yr have already identified some key processes that are involved in brain programming, including maternal care (4, 5)

and neuroendocrine factors [*e.g.*, stress-related hormones/neuropeptides (6–9)], as well as some molecular mechanisms [*e.g.*, those involved in neuronal plasticity (10, 11) and epigenetic regulation (12–14)]. Recently, inflammation (15, 16) and microbiota (17) have also attracted considerable attention, but the role of early nutrition in brain programming has so far been poorly studied.

Nutrition, however, is potentially very important in early programming, as: 1) brain development comes with an extremely high demand for nutrients (18), 2) there is an intense cross-talk between the stress and metabolic pathways (19, 20), and 3) ES and early malnutrition generally lead to similar cognitive deficits (21). Thus, a better understanding of the role of nutrition in ES effects on brain programming could create novel opportunities for early intervention in vulnerable populations.

Here, we focus on a group of essential micronutrients (as their availability fully depends on dietary intake), including methionine, homocysteine, vitamins B<sub>6</sub>, B<sub>12</sub>, B<sub>9</sub> (folic acid), and their metabolites. These micronutrients are notably critical in the 1-carbon (1-C) metabolism (22),

**ABBREVIATIONS:** 1-CMAM, 1-carbon metabolism–associated micronutrient; BrdU, 5-bromo-2'-deoxyuridine; CpG, 5'-C-phosphate-G-3'; Ctl, control; DG, dentate gyrus; DNMT, DNA-methyl transferase; EPM, elevated plus maze; ES, early-life stress; GR, glucocorticoid receptor; HPA, hypothalamo-pituitary-adrenal; LINE-1, long interspersed nuclear element-1; MWM, Morris water maze; *Nr3c1*, nuclear receptor subfamily 3, group C, member 1; OLT, object location task; ORT, object recognition task; P, postnatal day

<sup>1</sup> Correspondence: Swammerdam Institute for Life Sciences, Center for Neuroscience, Brain Plasticity Group, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands. E-mail: a.korosi@uva.nl

doi: 10.1096/fj.201600834R

which is required for methylation and for synthesis of proteins, phospholipids, and neurotransmitters. In this paper, we further refer to them as 1-C metabolism-associated micronutrients (1-CMAMs).

1-CMAMs are vital for brain function and development as they are involved in many biological processes, including neurogenesis (23, 24) and epigenetic mechanisms (25). Accordingly, impaired 1-C metabolism has been associated with neurological pathologies (26) and developmental anomalies [*e.g.*, neural-tube defects (27), neurodegeneration (28), and depression (29)]. In addition, a deficit in 1-CMAMs during critical developmental periods leads to brain dysfunction and lasting cognitive impairments, which suggests the involvement of 1-CMAMs in brain programming (30–33). Finally, 1-CMAMs are modulated by stress in adulthood (34–36), and polymorphisms in the methylene-tetrahydrofolate reductase gene relate to depression when studied in the context of early adversity (37).

This led us to hypothesize that 1-CMAMs are critical for programming cognition by ES. To this end, we assessed in mice: 1) whether peripheral and central levels of 1-CMAMs are altered by ES, 2) if restoring early micronutrient availability ameliorates ES-induced cognitive deficits in adulthood, and 3) which processes mediate the effects of this nutritional intervention. We took a multilevel approach at 3 different ages, that is, postnatal day (P)9, P120, and P265, and studied possible changes in maternal behavior, hypothalamic-pituitary-adrenal (HPA) axis activity of the offspring, neurogenesis in and volume of the hippocampal dentate gyrus (DG), hippocampal DNA methylation [globally and glucocorticoid receptor (GR)-specific], and hippocampal expression of key epigenetic regulators [DNA methyltransferases (DNMTs), the enzymes required for *de novo* DNA methylation (DNMT3a/3b) or its maintenance (DNMT1)].

We demonstrate that: 1) 1-CMAMs are critical in brain programming by ES, 2) maternal nutritional supplementation with 1-CMAMs ameliorates several lasting ES-induced cognitive impairments in offspring, and 3) mechanistically, the beneficial effects of the diet are mediated by preventing the ES-induced rise in corticosterone and adrenal gland hypertrophy.

## MATERIALS AND METHODS

### Animals

A total of 48 litters [13 control (Ctl), 13 ES, 10 Ctl-supplemented, 12 ES-supplemented] of 5–6 C57Bl/6j mice each were generated; only male offspring were used in these experiments. All procedures were conducted under European Union directives on animal experiments and approved by the animal welfare committee of the University of Amsterdam.

### ES paradigm

The ES procedure consisted of the limited nesting/bedding material model from P2 to P9 as previously described (38, 39). On the morning of P2, litters were randomly assigned to one of the following conditions: control condition with standard diet (Ctl), ES condition with standard diet (ES), control condition with

1-CMAM supplementation (Ctl-supplemented), or ES condition with 1-CMAM supplementation (ES-supplemented). On P9, pups were killed for short-term studies or moved to standard cages for long-term studies and weaned at P21.

### Micronutrient supplementation

Considering that offspring nutrition during early postnatal life depends on breastmilk, the composition of which is sensitive to maternal diet, specifically regarding water-soluble B vitamins (40), we enriched the maternal diet with 1-CMAMs during ES exposure to achieve supplementation of the offspring diet.

Throughout their lifetime, all animals were fed Teklad global rodent diet 2018 (Harlan Laboratories, Venray, The Netherlands), which contained 1200 mg/kg choline, 4 mg/kg folic acid, 0.08 mg/kg vitamin B<sub>12</sub> (cyanocobalamin), 0.4% methionine, 70 mg/kg zinc, and 18 mg/kg vitamin B<sub>6</sub> (pyridoxine). From P2–9, Ctl-supplemented and ES-supplemented dams received a 1-CMAM-supplemented diet (Ssniff, Soest, Germany) that was supplemented with 15g/kg Choline, 15g/kg betaine, 15 mg/kg folic acid, 1.5 mg/kg vitamin B<sub>12</sub>, 7.5 mg/kg L-methionine, and 150 mg/kg zinc [from ZnSO<sub>4</sub>·7H<sub>2</sub>O; according to Wolff *et al.* (41)], as well as drinking water that was supplemented with 15.3 µg/ml vitamin B<sub>6</sub> (pyridoxine-HCL P9755; Sigma-Aldrich, Zwijndrecht, The Netherlands). Both standard and 1-CMAM diets had an energy density of 3.1 kcal/g (24% of calories from protein, 18% from fat, 58% from carbohydrate).

### Behavioral analyses

Four-month-old mice (Ctl *n* = 14; ES *n* = 13; Ctl-supplemented *n* = 9; ES-supplemented *n* = 9) were tested in the elevated plus maze (EPM), object recognition task (ORT), object location task (OLT), Morris water maze (MWM), and T maze, as previously described (39). In the EPM, velocity, exploration time, and open/closed arm entries were scored to assess anxiety-like behavior.

In OLT and ORT, the ratio of novel *vs.* familiar object exploration time (on d 2) was used as an index of memory; a ratio >1 indicates a preference for the novel object/location and is thus associated with good memory of the object/location. Mice that spent <10 s exploring the objects were excluded from analysis. All necessary assumptions for the ORT and OLT were met: during acquisition, mice had no object preference and there was no difference in the total exploration time between groups.

MWM was used to assess spatial learning and memory, a 6-d acquisition phase (two 60-s trials/d) was followed by a single probe trial, as previously described (39). For MWM, one Ctl-supplemented animal was excluded from analysis, as it did not obtain an active search strategy.

Reference memory was tested by assessment of spontaneous choice alternations during 2 consecutive testing days (3 trials/d with an interval of 90 min) in a T-maze apparatus (dimensions arms 30 L × 10 W × 20 H), as previously described (42). Animals with latency to choice of >120 s were excluded from analyses.

Eight weeks after the final behavioral test, 100 mg/kg 5-bromo-2'-deoxyuridine (BrdU; Sigma-Aldrich; dissolved in sterile saline + 0.007 M NaOH) was injected 3 × /d, *i.p.* at P234 and P235 to assess survival of adult-born neurons. Animals were killed at P265.

### Tissue collection

For short-term studies (assessment of nutritional status, neurogenesis, gene expression, and DNA methylation), animals were



were considered statistically significant at  $P < 0.05$ . *Post hoc* analyses were performed by using Bonferroni multiple comparisons tests. Animals from multiple litters were included in each experiment and nested under the condition factor (3–7 litters/group), when appropriate. Relative gene expression data were logarithmically transformed to meet assumptions of parametric statistics.

## RESULTS

### ES reduces methionine levels, which are restored by 1-CMAM supplementation of the maternal diet


At P9, ES reduced methionine status both peripherally (30.1% reduction in plasma;  $P < 0.05$ ) and centrally (17.6% reduction in hippocampal tissue;  $P < 0.05$ ; **Table 1**), but did not affect levels of the other 1-CMAM nutrients ( $P > 0.05$ ).

During P2–9, ES-exposed dams ate more than control dams did (condition:  $F_{1,42} = 8.855$ ;  $P = 0.005$ ), and supplementation of the dam diet with 1-CMAM further increased food intake (diet:  $F_{1,42} = 11.774$ ;  $P = 0.001$ ), which confirmed that the diet was eaten by the dams. In addition, 1-CMAM supplementation reduced body-weight gain in pups (diet:  $F_{1,144} = 37.996$ ;  $P < 0.001$ ), but without affecting body weight in adulthood (diet:  $F_{1,29} = 0.8287$ ;  $P = 0.370$ ). Supplementation of the dam diet with 1-CMAM restored methionine levels in her pup's plasma and brain (**Table 1**). Plasma methionine was reduced by ES (univariable  $F$  test  $F_{1,51} = 26.25$ ;  $P < 0.0001$ ), but was increased by 1-CMAM diet ( $F_{1,51} = 5.95$ ;  $P = 0.018$ , no interaction  $F_{1,51} = 0.353$ ;  $P = 0.555$ ; **Fig. 1A**). Similarly, whereas ES reduced hippocampal methionine ( $F_{1,35} = 7.73$ ;  $P = 0.009$ ), 1-CMAM supplementation increased hippocampal methionine levels ( $F_{1,35} = 7.77$ ;  $P = 0.009$ , no interaction; **Fig. 1B**) and restored them

TABLE 1. Nutrient levels in stomach milk, plasma, and brain of P9 pups

Nutrient	Ctl	ES	Ctl-supplemented	ES-supplemented	Effect
<b>In milk</b>					
Folic acid (pmol/g)	167.4 ± 9.4	148.1 ± 9.4	323.2 ± 13.2 <sup>a</sup>	277.4 ± 14.2 <sup>a</sup>	Increase by diet: $F_{1,43} = 7.7$
5-MTHF (pmol/g)	295.7 ± 29.8	287.0 ± 29.8	193.7 ± 42.1	177.9 ± 45.0	Decrease by diet: $F_{1,43} = 8.0$
Vitamin B <sub>6</sub> (pmol/g)					
Pyridoxal	4675 ± 484	4755 ± 484	7779 ± 684 <sup>a</sup>	8973 ± 731 <sup>a</sup>	Increase by diet: $F_{1,43} = 36.5$
P5P	4089 ± 207	4400 ± 207	3693 ± 293	3894 ± 313	n.s.
Pyridoxamine	179.7 ± 21.7	195.3 ± 21.7	159.4 ± 30.7	174.9 ± 32.9	n.s.
Vitamin B <sub>12</sub> (pmol/g)	Undetectable	Undetectable	Undetectable	Undetectable	
Homocysteine (nmol/g)	2.38 ± 0.50	1.47 ± 0.50	8.15 ± 0.71 <sup>a</sup>	6.94 ± 0.76 <sup>a</sup>	Increase by diet: $F_{1,43} = 79.3$
Methionine (nmol/g)	25.2 ± 8.2	51.2 ± 8.2	17.4 ± 11.6	19.79 ± 12.4	n.s.
<b>In plasma</b>					
Folic acid (μM)	Undetectable	Undetectable	Undetectable	Undetectable	
5-MTHF (μM)	40.71 ± 4.17	39.64 ± 3.93	36.40 ± 5.03	49.14 ± 5.27	n.s.
Vitamin B <sub>6</sub> (μM)					
Pyridoxal	663.4 ± 95.5	605.3 ± 90.1	850.3 ± 101.4	577.7 ± 101.4	n.s.
P5P	126.3 ± 14.9	97.1 ± 14.0	92.5 ± 18.0	101.4 ± 18.8	n.s.
Pyridoxamine	<10	<10	<10	<10	—
Vitamin B <sub>12</sub> (μM)	Undetectable	Undetectable	Undetectable	Undetectable	
Homocysteine (mM)	5.64 ± 0.36	5.51 ± 0.33	5.25 ± 0.38	5.02 ± 0.38	n.s.
Methionine (mM)	125.9 ± 6.4 <sup>b</sup>	88.0 ± 5.9 <sup>b</sup>	143.7 ± 6.8 <sup>b</sup>	112.5 ± 6.8 <sup>b</sup>	Increase by diet: $F_{1,51} = 26.25$ ; decrease by ES: $F_{1,51} = 0.353$
<b>In brain</b>					
Folic acid (pmol/g)	Undetectable	Undetectable	Undetectable	Undetectable	
5-MTHF (pmol/g)	Undetectable	Undetectable	Undetectable	Undetectable	
Vitamin B <sub>6</sub> (pmol/g)					
Pyridoxal	268.7 ± 16.5	305.7 ± 16.5	323.1 ± 22.5	277.1 ± 16.5	
P5P	1803 ± 79	1725 ± 79	1640 ± 107	1450 ± 107	n.s.
Pyridoxamine	Undetectable	Undetectable	Undetectable	Undetectable	n.s.
Vitamin B <sub>12</sub> (pmol/g)	Undetectable	Undetectable	Undetectable	Undetectable	
Homocysteine (nmol/g)	2.59 ± 0.14	2.43 ± 0.14	2.88 ± 0.19	2.78 ± 0.19	n.s.
Methionine (nmol/g)	134.7 ± 5.5 <sup>b</sup>	110.9 ± 5.5 <sup>b</sup>	162.7 ± 7.5 <sup>b</sup>	147.3 ± 7.5 <sup>b</sup>	Increase by diet: $F_{1,35} = 7.73$ ; decrease by ES: $F_{1,35} = 7.77$

Data (represented as means ± SEM) were analyzed by a 2-factor multivariable ANOVA (MANOVA), with condition and diet as between-subject factors. If significant effects were detected, univariable  $F$  tests were used to identify the variables (nutrients) that contributed to these effects. Nutrient composition of the stomach milk of pups at P9 was altered by diet (2-factor MANOVA, HT = 7.22;  $F_{7,37} = 38.16$ ;  $P < 0.0001$ ) for folic acid, 5-methyltetrahydrofolate (5-MTHF), pyridoxal, and homocysteine (univariable  $F$  tests;  $P < 0.05$ ). Conversely, nutrient content of stomach milk was not altered by ES exposure (MANOVA revealed no multivariable effect of condition HT = 0.34;  $F_{7,37} = 1.79$ ;  $P = 0.118$ ). In plasma samples of ES pups, 1-CMAM supplementation restored nutrient levels in the pup's plasma. A 2-factor MANOVA for plasma nutrient levels revealed a significant effect of condition (HT = 0.814;  $F_{5,47} = 7.65$ ;  $P < 0.0001$ ) and diet (HT = 0.544;  $F_{5,47} = 512.16$ ;  $P = 0.001$ ) without interaction effect (HT = 0.210;  $F_{5,47} = 1.98$ ;  $P = 0.099$ ). In line with plasma data, nutrient levels in hippocampal tissue revealed significant multivariable effects for condition (HT = 0.345;  $F_{4,33} = 2.84$ ;  $P = 0.0039$ ) and diet (HT = 0.746;  $F_{4,33} = 6.156$ ;  $P = 0.001$ ) on micronutrient content. Hippocampal methionine levels were 17.6% lower in ES animals than in Ctl animals (*post hoc* analysis Ctl vs. ES;  $P < 0.05$ ). Levels of the other 1-CMAM nutrients detected in hippocampal tissue [pyridoxal, pyridoxal-5-phosphate (P5P), and homocysteine] were not affected by ES;  $P > 0.05$ . n.s., not significant. <sup>a</sup>Main effect for either diet or condition. <sup>b</sup>Significant effect for both condition and diet.



**Figure 1.** ES reduces central and peripheral methionine levels at P9. *A*) Methionine levels (mM) in pup plasma at P9 are reduced by ES ( $^{\#}F_{1,51} = 26.25$ ;  $P < 0.0001$ ) and increased by 1-CMAM diet ( $^*F_{1,51} = 5.95$ ;  $P = 0.018$ ), with no interaction of stress  $\times$  diet ( $F_{1,51} = 0.353$ ;  $P = 0.555$ ). *B*) Hippocampal methionine levels (nmol/g tissue) at P9 are reduced by ES ( $^{\#}F_{1,35} = 7.73$ ;  $P = 0.009$ ) and increased by 1-CMAM supplementation ( $^*F_{1,35} = 7.77$ ;  $P = 0.009$ ). Data are expressed as means  $\pm$  SEM.

to control levels [Student's *t* test, Ctl ( $n = 13$ ) *vs.* ES-supplemented ( $n = 7$ );  $P = 0.187$ ], whereas ES still affected them.

### 1-CMAM supplementation partly prevents ES-induced cognitive impairments

ES impaired performance in ORT, OLT, and MWM, which confirmed previous findings (38, 39). 1-CMAM supplementation prevented this in ORT and MWM, but not in OLT.

ES-induced impairments in ORT were fully prevented by the 1-CMAM diet. Two-way ANOVA revealed an interaction effect ( $F_{1,41} = 5.432$ ;  $P = 0.0248$ ; **Fig. 2A**), and *post hoc* testing showed impaired object recognition in ES animals compared with Ctl ( $t = 4164$ ;  $P < 0.001$ ), without difference between Ctl and Ctl-supplemented or ES-supplemented mice ( $P > 0.05$ ). Similarly, 1-CMAM diet improved acquisition in MWM. Acquisition consisted of 2 trials/d for 6 consecutive days. Whereas performance in the first trial on the day represents long-term (24 h) memory, the second trial of the day reflects properties of short-term (15 min) memory function.

As expected, latencies were shorter during the second (27.0  $\pm$  2.1 s) compared with the first trial of the day (33.0  $\pm$  1.8 s;  $P = 0.031$ ). A separate analysis of all first and second trials of the acquisition days revealed no differences in latencies during the first trial of the day (condition:  $F_{1,40} = 0.620$ ;  $P = 0.44$ ; diet:  $F_{1,40} = 1.1713$ ;  $P = 0.20$ , interaction:  $F_{1,40} = 1.161$ ;  $P = 0.690$ ), but when all second trials were compared, there was a strong trend toward an effect of diet ( $F_{1,40} = 3.88$ ;  $P = 0.056$ ; no condition effect:  $F_{1,40} = 0.766$ ;  $P = 0.387$ ; no interaction:  $F_{1,40} = 0.896$ ;  $P = 0.350$ ), which was largely attributable to shorter latencies of the ES-supplemented group (**Fig. 2C, D**). This suggests that 1-CMAM supplementation improved short-term memory in ES mice during task acquisition, but did not prevent ES-induced impairment in the testing phase of MWM (probe trial, **Fig. 2E, F**). Finally, ES-induced impairments in OLT persisted despite 1-CMAM supplementation (condition:  $F_{1,38} = 14.24$ ;  $P = 0.0005$ ; diet:  $F_{1,38} = 0.02$ ;  $P = 0.8853$ ; interaction:  $F_{1,38} = 0.22$ ;  $P = 0.645$ ; **Fig. 2B**).

Anxiety-like behavior reflected by the relative amount of open arm entries in the EPM was not affected by ES ( $F_{1,39} = 0.0875$ ;  $P = 0.78$ ), as found before (39), nor by diet ( $F_{1,39} = 0.9710$ ;  $P = 0.331$ ) or interaction ( $F_{1,39} = 3.115$ ;  $P = 0.0854$ ; data not shown).

Reference memory in the T maze (on average 75.4  $\pm$  2.6% of spontaneous alternations) was unaffected by condition ( $F_{1,18} = 0.2965$ ;  $P = 0.59$ ), diet ( $F_{1,18} = 1.747$ ;  $P = 0.20$ ), or interaction ( $F_{1,18} = 0.017$ ;  $P = 0.90$ ; data not shown).


### 1-CMAM supplementation of the dam prevents ES-induced HPA axis hyperactivity in her offspring

Basal plasma corticosterone levels at P9 were elevated by ES, which confirmed previous results (39). This was entirely prevented by 1-CMAM supplementation (interaction:  $F_{1,45} = 4.186$ ;  $P = 0.047$ ; *post hoc* ES *vs.* ES-supplemented;  $P < 0.01$ ; **Fig. 3A**). Similarly, 1-CMAM supplementation prevented ES-induced adrenal gland hypertrophy (interaction:  $F_{1,28} = 7.408$ ;  $P = 0.011$ ; *post hoc* Ctl *vs.* ES;  $P < 0.001$ ; **Fig. 3B**), which indicated an overall repressive effect of 1-CMAM supplementation on HPA axis activity and corticosterone production.


Neither ES, nor 1-CMAM supplementation led to lasting alterations in HPA axis activity (corticosterone at P90, condition:  $F_{1,14} = 0.087$ ;  $P = 0.772$ ; diet:  $F_{1,14} = 0.277$ ;  $P = 0.607$ ; interaction:  $F_{1,14} = 0.070$ ;  $P = 0.985$ ). 1-CMAM supplementation did not prevent the ES-induced reduction in pup thymus weight and body-weight gain (**Fig. 3C, D**).

### Beneficial effects of 1-CMAM supplementation are not mediated by normalizing maternal care, nor by preventing ES-induced alterations in offspring DG volume, neurogenesis, GR expression, or DNA methylation

Exposure to limited nesting and bedding material from P2 to P9 resulted in fragmented maternal care, which confirmed previous findings (38, 39), independent of diet



**Figure 2.** ES-induced cognitive impairments are prevented, in part, by 1-CMAM supplementation. *A*) In ORT, the ratio of novel to familiar was  $>1$  for all but the ES condition (1-sample Student's *t* tests, Ctl:  $1.96 \pm 0.16$ ;  $n = 14$ ;  $P < 0.0001$ ; Ctl-supplemented:  $2.03 \pm 0.07$ ;  $n = 9$ ;  $P < 0.00001$ ; ES-supplemented:  $1.94 \pm 0.18$ ;  $n = 9$ ;  $P = 0.0007$ ; ES:  $1.21 \pm 0.11$ ;  $n = 13$ ;  $P = 0.09$ ). ES reduces the ratio of novel to familiar exploration time in the standard diet condition (interaction effect of stress  $\times$  diet:  $F_{1,41} = 5.432$ ;  $P = 0.0248$ ; *post hoc* Ctl vs. ES:  $t = 4164$ ;  $*P < 0.001$ ). This impairment was prevented by 1-CMAM supplementation (Ctl-supplemented vs. ES-supplemented mice;  $P > 0.05$ ). *B*) In OLT, ES impairs performance ( $^{\#}F_{1,38} = 14.24$ ;  $P = 0.0005$ ), and this is not prevented by 1-CMAM supplementation (no effect of diet,  $F_{1,38} = 0.02$ ;  $P = 0.8853$ ; no interaction of stress  $\times$  diet ( $F_{1,38} = 0.22$ ;  $P = 0.645$ ). In fact, whereas both Ctl and Ctl-supplemented mice acquired the task (1-sample Student's *t* tests, Ctl:  $1.90 \pm 0.19$ ;  $n = 12$ ;  $P = 0.0005$ ; Ctl-supplemented:  $1.80 \pm 0.19$ ;  $n = 9$ ;  $P = 0.0029$ ), neither ES, nor ES-supplemented mice were able to discriminate between the objects (ES:  $1.21 \pm 0.12$ ;  $n = 12$ ;  $P = 0.1096$ ; ES-supplemented:  $1.26 \pm 0.131$ ;  $n = 9$ ;  $P = 0.0748$ ). *C*) Spatial acquisition in MWM. All animals were able to learn the task within 6 d as evident by the significantly reduced latency to find the platform between the first and the last training day in all 4 groups (paired Student's *t* test d 1 vs. d 6,  $P < 0.05$  in all groups). In Ctl animals, short-term spatial memory that was reflected by a short latency to find the hidden platform in each second training session of the day was not affected by diet (1-way repeated measures ANOVA,  $F_{1,21} = 0.907$ ;  $P = 0.352$ ). *D*) In ES animals, latency to find the platform during each second training session of the day was reduced by 1-CMAM supplementation (1-way repeated measures ANOVA,  $F_{1,20} = 4.434$ ;  $P = 0.048$ ). *E*) Performance in the probe trial of MWM was impaired by ES ( $^{\#}F_{1,40} = 9.523$ ;  $P = 0.0037$ ) but not restored by diet. ES and ES-supplemented mice showed no preference over chance (25%) for the target quadrant [1-sample Student's *t* tests compared with chance level, ES ( $n = 13$ )  $P = 0.34$ ; ES-supplemented ( $n = 9$ );  $P = 0.16$ ] and were significantly different from Ctl and Ctl-supplemented mice that spent significantly more time in the target quadrant [1-sample Student's *t* tests, Ctl ( $n = 14$ )  $P \leq 0.001$ ; Ctl-supplemented ( $n = 8$ )  $P \leq 0.0010$ ]. Differences in the performance of this task were not a result of differences in swimming ability or motivation, as swim speed was not different between conditions (data not shown). Data are expressed as means  $\pm$  SEM.



**Figure 3.** 1-CMAM supplementation dampens ES-induced HPA axis hyperactivation. *A*) ES-induced increase in basal plasma corticosterone (CORT) levels in pups (at P9) is prevented by 1-CMAM supplementation (2-way ANOVA, main effect of condition:  $F_{1,45} = 4.516$ ;  $P = 0.039$ ; main effect of diet:  $F_{1,45} = 13.287$ ;  $P = 0.001$ ; interaction effect:  $F_{1,45} = 4.186$ ;  $P = 0.047$ ). *Post hoc* analyses revealed that basal corticosterone levels were elevated after ES in the standard diet condition but not in the 1-CMAM-supplemented condition (*post hoc* Ctl vs. ES significant increase,  $*P < 0.01$ ). *B*) ES-induced increase in relative weight of the adrenal glands at P9 is prevented by 1-CMAM supplementation (significant stress  $\times$  diet interaction effect:  $F_{1,28} = 7.408$ ;  $P = 0.011$ ; *post hoc* Ctl vs. ES significant increase;  $*P < 0.001$ ). *C*) ES-induced reduction in relative thymus weight is not prevented by 1-CMAM supplementation (main effect of condition:  $^{\#}F_{1,40} = 10.875$ ;  $P = 0.002$ ; no effect of diet:  $F_{1,40} = 2.882$ ;  $P = 0.097$ ; no interaction effect:  $F_{1,40} = 2.745$ ;  $P = 0.105$ ). *D*) Body-weight gain from P2 to -9 is reduced by ES (main effect of condition:  $^{\#}F_{1,144} = 28.095$ ;  $P < 0.001$ ). 1-CMAM supplementation further reduced body-weight gain in both Ctl and ES pups (main effect of diet:  $*F_{1,144} = 37.996$ ;  $P < 0.001$ , no interaction effect). Data are expressed as means  $\pm$  SEM.

(Fig. 4); therefore, 1-CMAM supplementation did not exert its beneficial effects by normalizing maternal care.


We next investigated whether 1-CMAM supplementation exerted its effect *via* modulating neurobiologic processes in the offspring's hippocampus. 1-CMAM supplementation did not prevent ES-induced alterations in DG volume (Fig. 5A) and neurogenesis at P9 and P120 (Fig. 5B–E). Indeed, ES reduced the volume of the granular zone, defined as granular cell layer + subgranular zone, by approximately 20% at P9 (condition:  $F_{1,22} = 31.70$ ;  $P < 0.0001$ ; Fig. 5A), and this reduction was not prevented by 1-CMAM supplementation (diet:  $F_{1,22} = 1.147$ ;  $P = 0.2959$ ; no interaction:  $F_{1,22} = 1.319$ ;  $P = 0.2631$ ).

Similarly, at P9, the number of proliferating (Ki-67<sup>+</sup>) cells in the granular cell layer + subgranular zone was increased by ES (condition:  $F_{1,22} = 20.85$ ;  $P = 0.002$ ), which is in agreement with previous findings (39), and this was not prevented by 1-CMAM supplementation (diet:  $F_{1,22} = 2.733$ ;  $P = 0.1125$ ; no interaction:  $F_{1,22} = 3.346$ ;  $P = 0.0809$ ;

Fig. 5B). In adulthood, proliferation was not affected by ES and/or 1-CMAM supplementation (condition:  $F_{1,29} = 2.43$ ;  $P = 0.130$ ; diet:  $F_{1,29} = 0.55$ ;  $P = 0.461$ ; no interaction; Fig. 5C), but the 4-wk survival of adult-born (BrdU<sup>+</sup>) neurons was reduced by ES (condition:  $F_{1,41} = 4.658$ ;  $P = 0.037$ ; Fig. 5D, E), which was in line with prior results (39). This ES-induced reduction in survival of adult-born cells was not prevented by 1-CMAM supplementation ( $F_{1,41} = 0.008$ ;  $P = 0.930$ ; no interaction:  $F_{1,41} = 1.587$ ;  $P = 0.215$ ). None of the above mentioned alterations was specific for only the supra-/infrapyramidal blade or the rostral/caudal parts of the DG (data not shown).

We then investigated whether 1-CMAM supplementation exerted its effect *via* modulation of the epigenetic machinery in the hippocampus. Neither ES exposure, nor 1-CMAM supplementation—and the associated alterations in methionine status—were accompanied by clear indications of epigenetic dysregulation. In fact, even though global hippocampal DNA methylation levels at





**Figure 4.** Effects of ES and of micronutrient supplementation on maternal behavior. *A*) Exposure to the limited nesting and bedding material cage from P2 to P9 resulted in fragmented maternal care in both the standard and 1-CMAM-supplemented conditions. ES increases the number of nest exits in both diet groups (2-way ANOVA, main effect of condition:  $F_{1,22} = 11.99$ ;  $P = 0.002$ ). *B*) ES drastically increases the time during which  $\geq 1$  pups are outside the nest area (2-way ANOVA, main effect of condition:  $F_{1,22} = 244.76$ ;  $P < 0.0001$ ), but this is not affected by the diet. *C*) Dams from all 4 groups spent, on average, equal amounts of time on nursing behavior. Data are expressed as means  $\pm$  SEM.

P9, which were determined by a LINE1 pyrosequencing assay, were increased after ES by  $< 1\%$  in the standard condition and reduced by  $< 1\%$  after 1-CMAM supplementation (interaction:  $F_{1,41} = 4.730$ ;  $P = 0.035$ ; *post hoc* ES *vs.* ES-supplemented  $P < 0.05$ ; Fig. 6A), given the small effect size, we considered them biologically negligible. These effects were not lasting in adulthood (condition:  $F_{1,34} = 0.021$ ;  $P = 0.885$ ; diet:  $F_{1,34} = 0.013$ ;  $P = 0.911$ ; interaction:  $F_{1,34} = 0.072$ ;  $P = 0.789$ ; Fig. 6B). In addition, gene expression levels of Dnmts were not altered at P9 and P120 by ES or diet ( $P > 0.05$ ; Fig. 6C, D).

Finally, we investigated whether ES and/or diet affected expression and methylation status of GR gene (*Nr3c1*). At P9, hippocampal GR expression was increased by 1-CMAM diet in Ctl, but not ES, animals (interaction:  $F_{1,19} = 12.707$ ;  $P = 0.02$ ; *post hoc* Ctl *vs.* Ctl-supplemented,  $P < 0.05$ ; Fig. 6E). These alterations were not attributable to altered methylation patterns on exon1 of the *Nr3c1* gene. Two-way multivariable ANOVA of 9 CpGs within the *Nr3c1* promoter revealed no effects for condition (HT = 0.329;  $F_{9,31} = 1.134$ ;  $P = 0.370$ ) and diet (HT = 0.170;  $F_{9,31} = 1.585$ ;  $P = 0.799$ ), even though a trend toward an interaction effect was present (HT = 0.579;  $F_{9,31} = 1.996$ ;  $P = 0.074$ ; Fig. 6F). Separate analysis of individual CpGs revealed that this trend was largely attributable to changes in CpG8 methylation (univariable *F* tests, interaction:  $F_{1,39} = 2.207$ ;  $P = 0.020$ ; but *post hoc*  $P > 0.05$ ; Fig. 6G).

These alterations in hippocampal GR expression at P9 did not persist into adulthood (condition:  $F_{1,28} = 0.123$ ;  $P = 0.73$ ; diet:  $F_{1,28} = 0.505$ ;  $P = 0.48$ ; interaction:  $F_{1,28} = 0.205$ ;  $P = 0.65$ ; Fig. 6H). Similarly, levels of *Nr3c1* methylation in adulthood were not affected by ES (HT = 0.238;  $F_{9,25} = 0.662$ ;  $P = 0.734$ ) or diet (HT = 0.217;  $F_{9,25} = 0.602$ ;  $P = 0.783$ ), and no interaction was present (HT = 0.646;  $F_{9,25} = 1.795$ ;  $P = 0.120$ ; Fig. 6I).


## DISCUSSION

ES exposure has a lifelong impact on cognition. Here, we present evidence to suggest that restoring the ES-induced reduction in methionine levels, peripherally and centrally,

*via* supplementation of the dam's diet, and selectively during the ES period, has lasting benefits for her offspring. In fact, this short dietary intervention counteracted negative effects of ES in object recognition performance and improved acquisition in MWM. This indicates that the ES-induced lack of methionine during this critical phase might be a determinant factor in cognitive impairments observed in adult ES offspring. This supports our hypothesis (19) that essential micronutrients that have been implicated in 1-C metabolism might contribute to the programming effects of ES on later cognitive function. Furthermore, we established that the diet does not involve alterations in maternal care, hippocampal volume, neurogenesis, or epigenetic modifications, but prevents ES-induced hyperactivation of the HPA axis, which suggests that this might be implicated in its beneficial effects on later cognitive function.

## Early nutritional intervention ameliorates ES-induced cognitive impairments

Our finding that ES reduces levels of the essential amino acid methionine in the plasma and hippocampus of P9 pups highlights the stress sensitivity of this micronutrient during critical developmental periods. Methionine, other than being an important substrate of 1-C metabolism, is also a necessary amino acid for protein synthesis (45). Hence, a lack of methionine could impair these processes and thereby affect brain development and function. As such, it could be a critical factor for specific cognitive functions later in life. ES-induced alterations in nutrient content could not be detected in the (ingested) stomach milk. This might be a result of the fact that stomach milk is an accumulated amount and its content after ingestion may have been further altered by metabolic processes; therefore, it may not be entirely representative of exact breast milk composition). Hence, measurement in ingested stomach milk reflects not only nutritional intake of the offspring but, at least to some extent, also ongoing metabolic processes. ES-induced reduction in methionine can thus be either a result of a reduction of intake of maternal milk and/or impairing methionine uptake and bioavailability, for




**Figure 5.** ES-induced alterations in neurogenesis and DG volume reductions are not prevented by micronutrient supplementation. **A)** ES exposure reduces volume of the granular zone (GZ; granular cell layer + subgranular zone) at P9 with  $\pm 20\%$  (2-way ANOVA, main effect of condition:  $^{\#}F_{1,22} = 31.70$ ;  $P < 0.0001$ ). This reduction was not prevented by 1-CMAM supplementation (no effect of diet:  $F_{1,22} = 1.147$ ;  $P = 0.2959$ ; no interaction effect:  $F_{1,22} = 1.319$ ;  $P = 0.2631$ ). **B)** At P9, ES increases the number of proliferating (Ki-67<sup>+</sup>) cells in the GZ of the DG (expressed as numeric densities; 2-way ANOVA, main effect of condition:  $^{\#}F_{1,22} = 20.85$ ;  $P = 0.002$ ). This was not prevented by 1-CMAM supplementation (no effect of diet:  $F_{1,22} = 2.733$ ;  $P = 0.1125$ ; no interaction effect:  $F_{1,22} = 3.346$ ;  $P = 0.0809$ ). **C)** In adulthood, the number of proliferating (Ki-67<sup>+</sup>) cells was not affected by ES exposure and/or maternal 1-CMAM supplementation (2-way ANOVA, no effect of condition:  $F_{1,29} = 2.43$ ;  $P = 0.130$ ; no effect of diet:  $F_{1,29} = 0.55$ ;  $P = 0.461$ ; no interaction effect). **D)** Experimental timeline: to study survival of newborn cells in adulthood, animals were injected 3 wk after the final behavioral test with 100 mg/kg BrdU 3 times/d for 2 consecutive days and were killed by transcardial perfusion 4 wk after the last injection. **E)** ES lastingly reduces the survival of adult-born (BrdU<sup>+</sup>) neurons (2-way ANOVA, main effect of condition:  $^{\#}F_{1,41} = 4.658$ ;  $P = 0.037$ ). This effect on adult neurogenesis is not prevented by 1-CMAM supplementation ( $F_{1,41} = 0.008$ ;  $P = 0.930$ ; no interaction effect:  $F_{1,41} = 1.587$ ;  $P = 0.215$ ). None of the above-mentioned alterations was subregion specific, neither between the supra- vs. infrapyramidal blade, nor between the rostro vs. caudal parts of the DG. Data are expressed as means  $\pm$  SEM.

instance, by affecting gastrointestinal tract functioning. As stress is known to affect various physiologic functions of the gastrointestinal tract and the gut is an important site of methionine metabolism itself, this warrants further investigation.

Of importance, supplementation of the maternal diet during the ES period restored methionine to Ctl levels and ameliorated some of the negative effects of ES on cognition, fully restoring performance in ORT. Because the prefrontal cortex is also involved in ORT performance, our behavioral data indicate that, in addition to the hippocampus, other brain regions (e.g., prefrontal cortex) are likely involved in the beneficial effects of 1-CMAM supplementation on cognitive functioning.

In line with our findings, several preclinical studies in which fully 1-CMAM-deficient diets were administered during early life (30, 32) or adolescence (33, 46) reported lasting behavioral impairments, which further supports a critical role for 1-CMAM during early life for later cognitive function. This indicates that methionine levels need to reach a certain threshold to contribute to optimal adult brain functioning and points to early nutritional intervention as a promising tool to prevent lasting ES effects.

Here, we show that a short and early nutritional intervention with 1-CMAM can prevent lasting cognitive impairments induced by ES. Of interest, effects of methionine supplementation during adulthood have been reported before, although these were administered for a prolonged period of time under depletion conditions—for instance, severe adult folate deficiency induced cognitive impairments that could be prevented by a prolonged dietary methionine supplementation (47). In addition, also highlighting the powerful properties of these nutrients under specific conditions, acute or prolonged administration of methionine or the universal methyl donor S-adenosyl methionine during adulthood rescued cognitive impairments that were a result of earlier epilepsy or lead or cocaine exposure (12, 48–52) and improved drug-seeking behaviors (53). Our results show the sensitivity of specific nutrients to ES exposure and highlight a key role for them during early life in long-term brain programming, as well as the efficacy of early nutritional interventions to prevent later ES-induced effects on cognition. Because prevention of ES exposure (e.g., during conditions of hospitalization, abuse, neglect) itself is often not feasible, such adequate intervention strategies are



**Figure 6.** Global hippocampal DNA methylation and DNMT expression. *A*) At P9, average LINE-1 methylation is slightly higher in ES compared with ES-supplemented (no effect of condition:  $F_{1,41} = 0.803$ ;  $P = 0.375$ ; no effect of diet:  $F_{1,41} = 3.284$ ;  $P = 0.077$ ; interaction effect:  $F_{1,41} = 4.730$ ;  $P = 0.035$ ; \**post hoc* test ES vs. ES-supplemented,  $P < 0.05$ ). *B*) In adulthood, there was no difference in levels of LINE-1 methylation (no effect of condition:  $F_{1,34} = 0.021$ ;  $P = 0.885$ ; no effect of diet:  $F_{1,34} = 0.013$ ;  $P = 0.911$ ; no interaction effect:  $F_{1,34} = 0.072$ ;  $P = 0.789$ ). *C*) Relative DNMT mRNA expression at P9 is unaffected by ES and/or diet [determined by quantitative PCR, normalized to TATA-box binding protein (TBP),  $\alpha$ -tubulin, and hypoxanthine phosphoribosyltransferase (HPRT)]; for P9: DNMT1: no effect of condition:  $F_{1,19} = 0.031$ ;  $P = 0.86$ ; no effect of diet:  $F_{1,19} = 0.736$ ;  $P = 0.40$ ; no interaction effect:  $F_{1,19} = 0.001$ ;  $P = 0.99$ ; DNMT3a: no effect of condition:  $F_{1,19} = 0.496$ ;  $P = 0.49$ ; no effect of diet:  $F_{1,19} = 0.033$ ;  $P = 0.86$ ; no interaction effect:  $F_{1,19} = 0.339$ ;  $P = 0.57$ ; DNMT3b: no effect of condition:  $F_{1,19} = 2.109$ ;  $P = 0.17$ ; no effect of diet:  $F_{1,19} = 1.360$ ;  $P = 0.26$ ; no interaction effect:  $F_{1,19} = 1.057$ ;  $P = 0.32$ ]. *D*) At P120, relative DNMT expression levels were not altered by ES or diet (determined by quantitative PCR, normalized to TBP,  $\alpha$ -tubulin, and HPRT; DNMT1: no effect of condition:  $F_{1,28} = 0.686$ ;  $P = 0.42$ ; no effect of diet:  $F_{1,28} = 0.021$ ;  $P = 0.887$ ; no interaction effect:  $F_{1,28} = 0.056$ ;  $P = 0.82$ ; DNMT3a: no effect of condition:  $F_{1,28} = 0.123$ ;  $P = 0.73$ ; no effect of diet:  $F_{1,28} = 0.505$ ;  $P = 0.48$ ; no interaction effect:  $F_{1,28} = 0.205$ ;  $P = 0.65$ ; DNMT3b: no effect of condition:  $F_{1,19} = 0.839$ ;  $P = 0.37$ ; no effect of diet:  $F_{1,19} = 0.142$ ;  $P = 0.71$ ; no interaction effect:  $F_{1,19} = 1.994$ ;  $P = 0.17$ ). *E*) At P9, hippocampal GR expression is affected by ES exposure in a diet-dependent manner (2-way ANOVA, (continued on next page)

needed. As nutritional intervention may not be easily administered in all stressful environments, future studies that address additional windows of opportunity will aid in the development of practical intervention strategies.

### Which processes are involved?

Early 1-CMAM supplementation can exert its effects on later brain function either *via* a direct effect at the neuronal level (e.g., by influencing brain structure and plasticity) or by indirect modulation of other processes. Here, we investigated maternal care (4), the neuroendocrine stress system (54), hippocampal volume (2, 55, 56), neurogenesis (57, 58), and epigenetic modifications (13, 59, 60). 1-CMAM supplementation abolished ES-induced chronic HPA axis hyperactivation without affecting any of the other biologic processes we investigated. Hyperactivation of the stress axis is a likely candidate in determining long-term consequences of ES, as it is not only a common phenotype of ES exposure, but it has also been associated with later cognitive impairments. Indeed, a direct modulation of the HPA axis [e.g., *via* GR antagonists or adrenalectomy during ES (61, 62) or by exposure to stress at adulthood (63, 64)] can prevent ES-induced changes in several hippocampal parameters.

Our data demonstrate that 1-CMAM supplementation exerts a direct suppressive effect on ES-induced increase in basal plasma corticosterone and associated adrenal gland hypertrophy. We did not detect ES-induced changes in hippocampal GR expression, which was consistent with recent observations that hippocampal GR expression was unaffected by maternal separation in mice (65). Thus, in mice, chronic corticosterone elevation in early life does not seem to affect hippocampal GR, which is in contrast with well-established, lasting alterations in hippocampal GR that have been observed after ES exposure and limited maternal care in rats (4, 6, 66), which indicates a species-specific effect.

Furthermore, because methionine is a key factor in 1-C metabolism, which is crucial for epigenetic machinery, ES-induced reduction in methionine levels could potentially hamper epigenetic processes, which have been implicated in brain programming by early life experiences (6, 13, 67) and learning and memory (68). However, in contrast to our expectations, we did not find drastic alterations in global DNA methylation, which suggested that this process is tightly regulated and that global methylation levels remain relatively stable, even when

methionine input varies (69). In fact, the subtle differences in global hippocampal (LINE1) DNA methylation between ES and ES-supplemented animals comprised <1% (on average 81.2% methylation), and, therefore, its biologic relevance might be limited. Our finding of no alterations in global DNA methylation levels might be a result of the fact that central levels of the other measured 1-CMAM are unaffected by ES. These are further supported by evidence that both maternal care and acute methionine administration in a model of epilepsy failed to alter global DNA methylation in the hippocampus (12, 52).

In addition, ES exposure in mice did not alter any hippocampal DNMT levels, although others have recently described altered DNMT1 levels in neuronal progenitors after maternal separation—but this was in rats, which again points to a possible species-specific effect (70). However, clearly, methodologic differences could have also contributed to these discrepancies.

As the GR had been identified as being specifically vulnerable to epigenetic modifications in early life, both in human (71, 72) and rodents (6, 65), it was of considerable interest to study methylation of the *Nr3c1* promoter. We observed a subtle ES-induced hypermethylation of CpG8, in particular, in ES, which was not prevented in our ES-supplemented pups. These changes, however, were not sufficient to modulate GR gene expression levels. Similarly, maternal deprivation in mice failed to alter hippocampal *Nr3c1* methylation (65). In addition, central methionine infusion in adult rats left the majority of genes unaffected and modified only specific genes (48). Thus, effects of ES and methionine administration seem brain region and/or gene specific, and further research is needed to investigate if and how ES in mice leads to selective vulnerability of specific brain regions and genes.

### Early nutritional intervention as a clinically relevant tool

The current maternal dietary supplementation prevented several ES-induced cognitive deficits in offspring, at least part, because 1-CMAM supplementation repressed the ES-induced rise in corticosterone. 1-CAM supplementation did not have any detectable adverse effects, which was in agreement with previous studies that administered this diet (41). We have observed reduced body-weight gain in 1-CMAM-supplemented pups—probably as a result of the high intake of B vitamins, which also plays a role in energy expenditure (73)—that did not, however, lead to any lasting alteration in body weight (data not shown).

---

interaction of condition  $\times$  diet,  $F_{1,19} = 12.707$ ;  $P = 0.02$ ). 1-CMAM supplementation increased GR expression in control animals (*post hoc*:  $\#P < 0.05$ ), whereas in ES-animals, 1-CMAM supplementation had no effect on GR expression (*post hoc*:  $P > 0.05$ ). *F*) At P9, average methylation status of 9 CpGs within the *Nr3c1* promoter was not significantly different between the 4 groups (no effect for condition: HT = 0.329;  $F_{9,31} = 1.134$ ;  $P = 0.370$ ; no effect for diet: HT = 0.170;  $F_{9,31} = 1.585$ ;  $P = 0.799$ ), but a trend toward an interaction effect of condition  $\times$  diet (HT = 0.579;  $F_{9,31} = 1.996$ ;  $P = 0.074$ ) was observed. *G*) Separate analysis of the individual CpGs revealed that the trend toward an interaction is largely attributable to changes in methylation of CpG8 (interaction effect condition  $\times$  diet:  $F_{1,39} = 2.207$ ;  $P = 0.020$ ), but *post hoc* analysis revealed no significant differences between the groups. *H*) At P120, alterations in hippocampal GR expression were no longer present (no effect of condition:  $F_{1,28} = 0.123$ ;  $P = 0.73$ ; no effect of diet:  $F_{1,28} = 0.505$ ;  $P = 0.48$ ; no interaction effect:  $F_{1,28} = 0.205$ ;  $P = 0.65$ ). *I*) At P120, differences in levels of *Nr3c1* methylation were not detected (no effect of condition: HT = 0.238;  $F_{9,25} = 0.662$ ;  $P = 0.734$ ; no effect of diet: HT = 0.217;  $F_{9,25} = 0.602$ ;  $P = 0.783$ ; no interaction: HT = 0.646;  $F_{9,25} = 1.795$ ;  $P = 0.120$ ). Data are expressed as means  $\pm$  SEM.

Of interest, this supplementation did not modulate ES-induced changes in maternal care, which was in line with our hypothesis that ES-induced phenotype is likely the result of synergistic actions of multiple processes, including sensory stimuli, hormone levels, and nutritional status (19). A lack of effect on maternal care could be a possible explanation for why not all cognitive readouts were rescued. In support of this, stroking rat pups to mimic maternal sensory stimulation was previously shown to prevent ES-induced HPA axis derangements only when it was combined with feeding (74).

Our finding that supplementation of maternal diet with 1-CMAM did not prevent ES-induced changes in neurogenesis is consistent with the observed DG volumes, which were reduced by ES but not altered by 1-CMAM supplementation. These results, however, contrast with earlier described alterations in neurogenesis that were observed upon prolonged deficiencies in maternal folic acid (75), vitamin B<sub>12</sub> (30), or choline (76, 77) and upon prolonged protein and 1-CMAM supplementation throughout gestation and lactation (78). This is likely explained by the shorter duration of our intervention compared with previous studies.

In summary, until now, maternal care and glucocorticoids have often been held responsible for many ES-related lasting effects on brain structure and function. Here, we provide evidence for an additional, critical role of early nutrition in modulating ES-induced programming of adult brain function. Our findings open new avenues of early nutritional intervention to prevent lasting effects of ES in offspring, an approach that may have important implications for clinical practice. FJ

## ACKNOWLEDGMENTS

A.K. is supported by grants from the Netherlands Organization for Scientific Research (NWO; Meervoud; Food, Cognition and Brain). P.J.L. is supported by HersenStichting, Internationale Stichting Alzheimer Onderzoek (ISAO), and Alzheimer Nederland. T.P. is supported by a ZonMW TOP grant.

## AUTHOR CONTRIBUTIONS

E. F. G. Naninck and A. Korosi designed the research; E. F. G. Naninck, J. E. Oosterink, K.-Y. Yam, L. P. de Vries, J. A. Platinga, and R. N. Verkaik-Schakel performed research; H. Schierbeek and J. B. van Goudoever contributed reagents/analytic tools; E. F. G. Naninck and T. Plosch analyzed data; and E. F. G. Naninck, P. J. Lucassen, and A. Korosi wrote the paper.

## REFERENCES

1. Pesonen, A.-K., Eriksson, J. G., Heinonen, K., Kajantie, E., Tuovinen, S., Alastalo, H., Henriksson, M., Leskinen, J., Osmond, C., Barker, D. J. P., and Räikkönen, K. (2013) Cognitive ability and decline after early life stress exposure. *Neurobiol. Aging* **34**, 1674–1679
2. Hanson, J. L., Nacewicz, B. M., Sutterer, M. J., Cayo, A. A., Schaefer, S. M., Rudolph, K. D., Shirtcliff, E. A., Pollak, S. D., and Davidson, R. J. (2015) Behavioral problems after early life stress: contributions of the hippocampus and amygdala. *Biol. Psychiatry* **77**, 314–323

3. Molet, J., Maras, P. M., Avishai-Eliner, S., and Baram, T. Z. (2014) Naturalistic rodent models of chronic early-life stress. *Dev. Psychobiol.* **56**, 1675–1688
4. Liu, D., Diorio, J., Tannenbaum, B., Caldji, C., Francis, D., Freedman, A., Sharma, S., Pearson, D., Plotsky, P. M., and Meaney, M. J. (1997) Maternal care, hippocampal glucocorticoid receptors, and hypothalamic-pituitary-adrenal responses to stress. *Science* **277**, 1659–1662
5. Feldman, R., Singer, M., and Zagoory, O. (2010) Touch attenuates infants' physiological reactivity to stress. *Dev. Sci.* **13**, 271–278
6. Weaver, I. C. G., Cervoni, N., Champagne, F. A., D'Alessio, A. C., Sharma, S., Seckl, J. R., Dymov, S., Szyf, M., and Meaney, M. J. (2004) Epigenetic programming by maternal behavior. *Nat. Neurosci.* **7**, 847–854
7. Murgatroyd, C. A., Taliefar, M., Bradburn, S., Carini, L. M., Babb, J. A., and Nephew, B. C. (2015) Social stress during lactation, depressed maternal care, and neuropeptidergic gene expression. *Behav. Pharmacol.* **26**, 642–653
8. Ivy, A. S., Rex, C. S., Chen, Y., Dubé, C., Maras, P. M., Grigoriadis, D. E., Gall, C. M., Lynch, G., and Baram, T. Z. (2010) Hippocampal dysfunction and cognitive impairments provoked by chronic early-life stress involve excessive activation of CRH receptors. *J. Neurosci.* **30**, 13005–13015
9. Wang, X.-D., Rammes, G., Kraev, I., Wolf, M., Liebl, C., Scharf, S. H., Rice, C. J., Wurst, W., Holsboer, F., Deussing, J. M., Baram, T. Z., Stewart, M. G., Müller, M. B., and Schmidt, M. V. (2011) Forebrain CRF<sub>1</sub> modulates early-life stress-programmed cognitive deficits. *J. Neurosci.* **31**, 13625–13634
10. Kohl, C., Wang, X.-D., Grosse, J., Fournier, C., Harbich, D., Westerholz, S., Li, J.-T., Bacq, A., Sippel, C., Hausch, F., Sandi, C., and Schmidt, M. V. (2015) Hippocampal neurotrophin-2 links early-life stress with impaired social recognition and increased aggression in adult mice. *Psychoneuroendocrinology* **55**, 128–143
11. Oomen, C. A., Soeters, H., Audureau, N., Vermunt, L., van Hasselt, F. N., Manders, E. M. M., Joëls, M., Krugers, H., and Lucassen, P. J. (2011) Early maternal deprivation affects dentate gyrus structure and emotional learning in adult female rats. *Psychopharmacology (Berl.)* **214**, 249–260
12. Weaver, I. C. G., Champagne, F. A., Brown, S. E., Dymov, S., Sharma, S., Meaney, M. J., and Szyf, M. (2005) Reversal of maternal programming of stress responses in adult offspring through methyl supplementation: altering epigenetic marking later in life. *J. Neurosci.* **25**, 11045–11054
13. Murgatroyd, C., Patchev, A. V., Wu, Y., Micalé, V., Bockmühl, Y., Fischer, D., Holsboer, F., Wotjak, C. T., Almeida, O. F. X., and Spengler, D. (2009) Dynamic DNA methylation programs persistent adverse effects of early-life stress. *Nat. Neurosci.* **12**, 1559–1566
14. Bale, T. L., Baram, T. Z., Brown, A. S., Goldstein, J. M., Insel, T. R., McCarthy, M. M., Nemeroff, C. B., Reyes, T. M., Simerly, R. B., Susser, E. S., and Nestler, E. J. (2010) Early life programming and neurodevelopmental disorders. *Biol. Psychiatry* **68**, 314–319
15. Hoeijmakers, L., Lucassen, P. J., and Korosi, A. (2015) The interplay of early-life stress, nutrition, and immune activation programs adult hippocampal structure and function. *Front. Mol. Neurosci.* **7**, 103
16. Cattaneo, A., Macchi, F., Plazzotta, G., Veronica, B., Bocchio-Chiavetto, L., Riva, M. A., and Pariante, C. M. (2015) Inflammation and neuronal plasticity: a link between childhood trauma and depression pathogenesis. *Front. Cell. Neurosci.* **9**, 40
17. Borre, Y. E., O'Keefe, G. W., Clarke, G., Stanton, C., Dinan, T. G., and Cryan, J. F. (2014) Microbiota and neurodevelopmental windows: implications for brain disorders. *Trends Mol. Med.* **20**, 509–518
18. Benton, D. (2010) The influence of dietary status on the cognitive performance of children. *Mol. Nutr. Food Res.* **54**, 457–470
19. Lucassen, P. J., Naninck, E. F. G., van Goudoever, J. B., Fitzsimons, C., Joels, M., and Korosi, A. (2013) Perinatal programming of adult hippocampal structure and function; emerging roles of stress, nutrition and epigenetics. *Trends Neurosci.* **36**, 621–631
20. Spencer, S. J. (2013) Perinatal programming of neuroendocrine mechanisms connecting feeding behavior and stress. *Front. Neurosci.* **7**, 109
21. Naninck, E. F. G., Lucassen, P. J., and Korosi, A. (2013) Consequences of early-life experiences on cognition and emotion: a role for nutrition and epigenetic mechanisms. In *Oxford Handbook of Molecular Psychology* (Canli, T., ed.), pp. 311–337, Oxford University Press, New York
22. Feil, R., and Fraga, M. F. (2012) Epigenetics and the environment: emerging patterns and implications. *Nat. Rev. Genet.* **13**, 97–109

23. Kruman, I. I., Mouton, P. R., Emokpae, R., Jr., Cutler, R. G., and Mattson, M. P. (2005) Folate deficiency inhibits proliferation of adult hippocampal progenitors. *Neuroreport* **16**, 1055–1059
24. Rabaneda, L. G., Carrasco, M., López-Toledano, M. A., Murillo-Carretero, M., Ruiz, F. A., Estrada, C., and Castro, C. (2008) Homocysteine inhibits proliferation of neuronal precursors in the mouse adult brain by impairing the basic fibroblast growth factor signaling cascade and reducing extracellular regulated kinase 1/2-dependent cyclin E expression. *FASEB J.* **22**, 3823–3835
25. Friso, S., and Choi, S.-W. (2002) Gene-nutrient interactions and DNA methylation. *J. Nutr.* **132**, 2382S–2387S
26. Sugden, C. (2006) One-carbon metabolism in psychiatric illness. *Nutr. Res. Rev.* **19**, 117–136
27. Mitchell, L. E. (2005) Epidemiology of neural tube defects. *Am. J. Med. Genet. C. Semin. Med. Genet.* **135C**, 88–94
28. Mattson, M. P., and Shea, T. B. (2003) Folate and homocysteine metabolism in neural plasticity and neurodegenerative disorders. *Trends Neurosci.* **26**, 137–146
29. Bjelland, I., Tell, G. S., Vollset, S. E., Refsum, H., and Ueland, P. M. (2003) Folate, vitamin B12, homocysteine, and the MTHFR 677C>T polymorphism in anxiety and depression: the Hordaland Homocysteine Study. *Arch. Gen. Psychiatry* **60**, 618–626
30. Blaise, S. A., Nédélec, E., Schroeder, H., Alberto, J.-M., Bossenmeyer-Pouricé, C., Guéant, J.-L., and Daval, J.-L. (2007) Gestational vitamin B deficiency leads to homocysteine-associated brain apoptosis and alters neurobehavioral development in rats. *Am. J. Pathol.* **170**, 667–679
31. Ramaekers, V. T., and Blau, N. (2004) Cerebral folate deficiency. *Dev. Med. Child Neurol.* **46**, 843–851
32. Chehadeh, S. E. H., Pouricé, G., Martin, N., Alberto, J.-M., Daval, J.-L., Guéant, J.-L., and Leininger-Muller, B. (2013) Gestational methyl donor deficiency alters key proteins involved in neurosteroidogenesis in the olfactory bulbs of newborn female rats and is associated with impaired olfactory performance. *Br. J. Nutr.* **111**, 1021–1031
33. Ishii, D., Matsuzawa, D., Matsuda, S., Tomizawa, H., Sutoh, C., and Shimizu, E. (2014) Methyl donor-deficient diet during development can affect fear and anxiety in adulthood in C57BL/6j mice. *PLoS One* **9**, e105750
34. Stoney, C. M., and Engebretson, T. O. (2000) Plasma homocysteine concentrations are positively associated with hostility and anger. *Life Sci.* **66**, 2267–2275
35. de Oliveira, A. C., Suchecki, D., Cohen, S., and D'Almeida, V. (2004) Acute stressor-selective effect on total plasma homocysteine concentration in rats. *Pharmacol. Biochem. Behav.* **77**, 269–273
36. De Souza, F. G., Rodrigues, M. D., Tufik, S., Nobrega, J. N., and D'Almeida, V. (2006) Acute stressor-selective effects on homocysteine metabolism and oxidative stress parameters in female rats. *Pharmacol. Biochem. Behav.* **85**, 400–407
37. Lok, A., Bockting, C. L. H., Koeter, M. W. J., Snieder, H., Assies, J., Mocking, R. J. T., Vinkers, C. H., Kahn, R. S., Boks, M. P., and Schene, A. H. (2013) Interaction between the MTHFR C677T polymorphism and traumatic childhood events predicts depression. *Transl. Psychiatry* **3**, e288
38. Rice, C. J., Sandman, C. A., Lenjavi, M. R., and Baram, T. Z. (2008) A novel mouse model for acute and long-lasting consequences of early life stress. *Endocrinology* **149**, 4892–4900
39. Naninck, E. F. G., Hoeymakers, L., Kakava-Georgiadou, N., Meesters, A., Lazic, S. E., Lucassen, P. J., and Korosi, A. (2015) Chronic early life stress alters developmental and adult neurogenesis and impairs cognitive function in mice. *Hippocampus* **25**, 309–328
40. Allen, L. H. (2005) Multiple micronutrients in pregnancy and lactation: an overview. *Am. J. Clin. Nutr.* **81**, 1206S–1212S
41. Wolff, G. L., Kodell, R. L., Moore, S. R., and Cooney, C. A. (1998) Maternal epigenetics and methyl supplements affect agouti gene expression in *Ay/a* mice. *FASEB J.* **12**, 949–957
42. Deacon, R. M. J., and Rawlins, J. N. P. (2006) T-maze alternation in the rodent. *Nat. Protoc.* **1**, 7–12
43. Oosterink, J. E., Naninck, E. F. G., Korosi, A., Lucassen, P. J., van Goudoever, J. B., and Schierbeek, H. (2015) Accurate measurement of the essential micronutrients methionine, homocysteine, vitamins B<sub>6</sub>, B<sub>12</sub>, B<sub>9</sub> and their metabolites in plasma, brain and maternal milk of mice using LC/MS ion trap analysis. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* **998-999**, 106–113
44. Freitag, N., Zwier, M. V., Barrientos, G., Tirado-González, I., Conrad, M. L., Rose, M., Scherjon, S. A., Plösch, T., and Blois, S. M. (2014) Influence of relative NK-DC abundance on placentation and its relation to epigenetic programming in the offspring. *Cell Death Dis.* **5**, e1392
45. Kozak, M. (1983) Comparison of initiation of protein synthesis in procaryotes, eucaryotes, and organelles. *Microbiol. Rev.* **47**, 1–45
46. Tomizawa, H., Matsuzawa, D., Ishii, D., Matsuda, S., Kawai, K., Mashimo, Y., Sutoh, C., and Shimizu, E. (2015) Methyl-donor deficiency in adolescence affects memory and epigenetic status in the mouse hippocampus. *Genes Brain Behav.* **14**, 301–309
47. Troen, A. M., Chao, W.-H., Crivello, N. A., D'Anci, K. E., Shukitt-Hale, B., Smith, D. E., Selhub, J., and Rosenberg, I. H. (2008) Cognitive impairment in folate-deficient rats corresponds to depleted brain phosphatidylcholine and is prevented by dietary methionine without lowering plasma homocysteine. *J. Nutr.* **138**, 2502–2509
48. Weaver, I. C. G., Meaney, M. J., and Szyf, M. (2006) Maternal care effects on the hippocampal transcriptome and anxiety-mediated behaviors in the offspring that are reversible in adulthood. *Proc. Natl. Acad. Sci. USA* **103**, 3480–3485
49. Cao, X.-J., Huang, S.-H., Wang, M., Chen, J.-T., and Ruan, D.-Y. (2008) S-adenosyl-L-methionine improves impaired hippocampal long-term potentiation and water maze performance induced by developmental lead exposure in rats. *Eur. J. Pharmacol.* **595**, 30–34
50. Fan, G., Feng, C., Wu, F., Ye, W., Lin, F., Wang, C., Yan, J., Zhu, G., Xiao, Y., and Bi, Y. (2010) Methionine choline reverses lead-induced cognitive and N-methyl-D-aspartate receptor subunit 1 deficits. *Toxicology* **272**, 23–31
51. Tian, W., Zhao, M., Li, M., Song, T., Zhang, M., Quan, L., Li, S., and Sun, Z. S. (2012) Reversal of cocaine-conditioned place preference through methyl supplementation in mice: altering global DNA methylation in the prefrontal cortex. *PLoS One* **7**, e33435
52. Parrish, R. R., Buckingham, S. C., Mascia, K. L., Johnson, J. J., Matyjasik, M. M., Lockhart, R. M., and Lubin, F. D. (2015) Methionine increases BDNF DNA methylation and improves memory in epilepsy. *Ann. Clin. Transl. Neurol.* **2**, 401–416
53. Wright, K. N., Hollis, F., Duclot, F., Dossat, A. M., Strong, C. E., Francis, T. C., Mercer, R., Feng, J., Dietz, D. M., Lobo, M. K., Nestler, E. J., and Kabbaj, M. (2015) Methyl supplementation attenuates cocaine-seeking behaviors and cocaine-induced c-Fos activation in a DNA methylation-dependent manner. *J. Neurosci.* **35**, 8948–8958
54. Strüber, N., Strüber, D., and Roth, G. (2014) Impact of early adversity on glucocorticoid regulation and later mental disorders. *Neurosci. Biobehav. Rev.* **38**, 17–37
55. Herpfer, I., Hezel, H., Reichardt, W., Clark, K., Geiger, J., Gross, C. M., Heyer, A., Neagu, V., Bhatia, H., Atas, H. C., Fiebich, B. L., Bischofberger, J., Haas, C. A., Lieb, K., and Normann, C. (2012) Early life stress differentially modulates distinct forms of brain plasticity in young and adult mice. *PLoS One* **7**, e46004
56. Teicher, M. H., Anderson, C. M., and Polcari, A. (2012) Childhood maltreatment is associated with reduced volume in the hippocampal subfields CA3, dentate gyrus, and subiculum. *Proc. Natl. Acad. Sci. USA* **109**, E563–E572
57. Korosi, A., Naninck, E. F. G., Oomen, C. A., Schouten, M., Krugers, H., Fitzsimons, C., and Lucassen, P. J. (2012) Early-life stress mediated modulation of adult neurogenesis and behavior. *Behav. Brain Res.* **227**, 400–409
58. Suri, D., Veenit, V., Sarkar, A., Thiagarajan, D., Kumar, A., Nestler, E. J., Galande, S., and Vaidya, V. A. (2013) Early stress evokes age-dependent biphasic changes in hippocampal neurogenesis, BDNF expression, and cognition. *Biol. Psychiatry* **73**, 658–666
59. McClelland, S., Korosi, A., Cope, J., Ivy, A., and Baram, T. Z. (2011) Emerging roles of epigenetic mechanisms in the enduring effects of early-life stress and experience on learning and memory. *Neurobiol. Learn. Mem.* **96**, 79–88
60. Peña, C. J., Bagot, R. C., Labonté, B., and Nestler, E. J. (2014) Epigenetic signaling in psychiatric disorders. *J. Mol. Biol.* **426**, 3389–3412
61. Mirescu, C., Peters, J. D., and Gould, E. (2004) Early life experience alters response of adult neurogenesis to stress. *Nat. Neurosci.* **7**, 841–846
62. Zhang, T. Y., Labonté, B., Wen, X. L., Turecki, G., and Meaney, M. J. (2013) Epigenetic mechanisms for the early environmental regulation of hippocampal glucocorticoid receptor gene expression in rodents and humans. *Neuropsychopharmacology* **38**, 111–123
63. Hu, P., Oomen, C., van Dam, A. M., Wester, J., Zhou, J.-N., Joëls, M., and Lucassen, P. J. (2012) A single-day treatment with mifepristone is sufficient to normalize chronic glucocorticoid induced suppression of hippocampal cell proliferation. *PLoS One* **7**, e46224
64. Sarabdjitsingh, R. A., Zhou, M., Yau, J. L. W., Webster, S. P., Walker, B. R., Seckl, J. R., Joëls, M., and Krugers, H. J. (2014) Inhibiting 11 $\beta$ -hydroxysteroid dehydrogenase type 1 prevents stress effects on

- hippocampal synaptic plasticity and impairs contextual fear conditioning. *Neuropharmacology* **81**, 231–236
65. Bockmühl, Y., Patchev, A. V., Madejska, A., Hoffmann, A., Sousa, J. C., Sousa, N., Holsboer, F., Almeida, O. F. X., and Spengler, D. (2015) Methylation at the CpG island shore region upregulates Nr3c1 promoter activity after early-life stress. *Epigenetics* **10**, 247–257
  66. Francis, D., Diorio, J., Liu, D., and Meaney, M. J. (1999) Nongenomic transmission across generations of maternal behavior and stress responses in the rat. *Science* **286**, 1155–1158
  67. Uchida, S., Hara, K., Kobayashi, A., Funato, H., Hobara, T., Otsuki, K., Yamagata, H., McEwen, B. S., and Watanabe, Y. (2010) Early life stress enhances behavioral vulnerability to stress through the activation of REST4-mediated gene transcription in the medial prefrontal cortex of rodents. *J. Neurosci.* **30**, 15007–15018
  68. Morris, M. J., and Monteggia, L. M. (2014) Role of DNA methylation and the DNA methyltransferases in learning and memory. *Dialogues Clin. Neurosci.* **16**, 359–371
  69. Nijhout, H. F., Reed, M. C., Anderson, D. F., Mattingly, J. C., James, S. J., and Ulrich, C. M. (2006) Long-range allosteric interactions between the folate and methionine cycles stabilize DNA methylation reaction rate. *Epigenetics* **1**, 81–87
  70. Boku, S., Toda, H., Nakagawa, S., Kato, A., Inoue, T., Koyama, T., Hiroi, N., and Kusumi, I. (2015) Neonatal maternal separation alters the capacity of adult neural precursor cells to differentiate into neurons via methylation of retinoic acid receptor gene promoter. *Biol. Psychiatry* **77**, 335–344
  71. Mulligan, C. J., D'Errico, N. C., Stees, J., and Hughes, D. A. (2012) Methylation changes at NR3C1 in newborns associate with maternal prenatal stress exposure and newborn birth weight. *Epigenetics* **7**, 853–857
  72. Romens, S. E., McDonald, J., Svaren, J., and Pollak, S. D. (2015) Associations between early life stress and gene methylation in children. *Child Dev.* **86**, 303–309
  73. Gunanti, I. R., Marks, G. C., Al-Mamun, A., and Long, K. Z. (2014) Low serum vitamin B-12 and folate concentrations and low thiamin and riboflavin intakes are inversely associated with greater adiposity in Mexican American children. *J. Nutr.* **144**, 2027–2033
  74. van Oers, H. J., de Kloet, E. R., Whelan, T., and Levine, S. (1998) Maternal deprivation effect on the infant's neural stress markers is reversed by tactile stimulation and feeding but not by suppressing corticosterone. *J. Neurosci.* **18**, 10171–10179
  75. Craciunescu, C. N., Brown, E. C., Mar, M.-H., Albright, C. D., Nadeau, M. R., and Zeisel, S. H. (2004) Folic acid deficiency during late gestation decreases progenitor cell proliferation and increases apoptosis in fetal mouse brain. *J. Nutr.* **134**, 162–166
  76. Niculescu, M. D., Craciunescu, C. N., and Zeisel, S. H. (2006) Dietary choline deficiency alters global and gene-specific DNA methylation in the developing hippocampus of mouse fetal brains. *FASEB J.* **20**, 43–49
  77. Craciunescu, C. N., Albright, C. D., Mar, M.-H., Song, J., and Zeisel, S. H. (2003) Choline availability during embryonic development alters progenitor cell mitosis in developing mouse hippocampus. *J. Nutr.* **133**, 3614–3618
  78. Amarger, V., Lecouillard, A., Ancellet, L., Grit, I., Castellano, B., Hulin, P., and Parnet, P. (2014) Protein content and methyl donors in maternal diet interact to influence the proliferation rate and cell fate of neural stem cells in rat hippocampus. *Nutrients* **6**, 4200–4217

Received for publication July 19, 2016.  
Accepted for publication October 4, 2016.

## Early micronutrient supplementation protects against early stress-induced cognitive impairments

Eva F. G. Naninck, J. Efraim Oosterink, Kit-Yi Yam, et al.

*FASEB J* 2017 31: 505-518 originally published online October 21, 2016

Access the most recent version at doi:[10.1096/fj.201600834R](https://doi.org/10.1096/fj.201600834R)

---

**References** This article cites 77 articles, 20 of which can be accessed free at:  
<http://www.fasebj.org/content/31/2/505.full.html#ref-list-1>

**Subscriptions** Information about subscribing to *The FASEB Journal* is online at  
<http://www.faseb.org/The-FASEB-Journal/Librarian-s-Resources.aspx>

**Permissions** Submit copyright permission requests at:  
<http://www.fasebj.org/site/misc/copyright.xhtml>

**Email Alerts** Receive free email alerts when new an article cites this article - sign up at  
<http://www.fasebj.org/cgi/alerts>

---