

DOI: 10.5586/asbp.3558

**Publication history**

Received: 2016-11-22

Accepted: 2017-08-08

Published: 2017-09-14

**Handling editor**

Grażyna Kłobus, Faculty of Biological Sciences, University of Wrocław, Poland

**Authors' contributions**

AK: research design, conducting experiments, data collection and analysis, writing the manuscript; AA: supervision of the project, contribution to the idea and design of the experiments, critical revision of the manuscript

**Funding**

This research was financed by the Ministry of Agriculture and Rural Development of the Republic of Poland (grant No. HORhn-801-22/12).

**Competing interests**

No competing interests have been declared.

**Copyright notice**© The Author(s) 2017. This is an Open Access article distributed under the terms of the [Creative Commons Attribution License](#), which permits redistribution, commercial and non-commercial, provided that the article is properly cited.**Citation**Kielkowska A, Adamus A. Early studies on the effect of peptide growth factor phytosulfokine- $\alpha$  on *Brassica oleracea* var. *capitata* L. protoplasts. Acta Soc Bot Pol. 2017;86(3):3558. <https://doi.org/10.5586/asbp.3558>**Digital signature**

This PDF has been certified using digital signature with a trusted timestamp to assure its origin and integrity. A verification trust dialog appears on the PDF document when it is opened in a compatible PDF reader. Certificate properties provide further details such as certification time and a signing reason in case any alterations made to the final content. If the certificate is missing or invalid it is recommended to verify the article on the journal website.

## ORIGINAL RESEARCH PAPER

# Early studies on the effect of peptide growth factor phytosulfokine- $\alpha$ on *Brassica oleracea* var. *capitata* L. protoplasts

Agnieszka Kielkowska\*, Adela Adamus

Institute of Plant Biology and Biotechnology, Faculty of Biotechnology and Horticulture, University of Agriculture in Krakow, al. 29 listopada 54, 31-425 Krakow, Poland

\* Corresponding author. Email: [kielkowska@ogr.ur.krakow.pl](mailto:kielkowska@ogr.ur.krakow.pl)**Abstract**

Phytosulfokines (PSK) are peptidyl growth factors with the potential of inducing cell proliferation. We examined the effect of supplementation of liquid culture medium with 0.1  $\mu\text{M}$  phytosulfokine- $\alpha$  (PSK- $\alpha$ ) on protoplast viability and division frequencies in seven accessions of *Brassica oleracea* var. *capitata* L., including cultivars and breeding lines. Protoplasts were isolated from leaves and hypocotyls of in vitro grown plants and immobilized in calcium-alginate layers. Cabbage protoplast-derived cells cultured in medium supplemented with 0.1  $\mu\text{M}$  of PSK- $\alpha$  had higher viability and division frequencies compared to cells cultured in PSK- $\alpha$ -free control medium. The effect of PSK- $\alpha$  was more pronounced in low-responding accessions ('Sława z Gołębiewa', 'Ramkila F1', LM, and LM98); however, in two cultivars with very low response ('Badger Shipper' and 'Oregon 123'), although the division frequencies in the media supplemented with PSK- $\alpha$  were increased over the control, the differences were not significant. Obtained callus colonies were subjected to regeneration. PSK- $\alpha$  supplemented into the liquid culture medium had an indirect effect on shoot regeneration by inducing sustained cell divisions leading to an increase in shoot regeneration in Sława z Gołębiewa and both breeding lines.

**Keywords**

cabbage; peptide; phytosulfokine; PSK; protoplast culture

**Introduction**

*Brassica oleracea* var. *capitata* L. belongs to the family Brassicaceae, which includes many economically important vegetables such as broccoli, cauliflower, or brussels sprouts. Protoplast cultures have been intensively studied in the genus *Brassica*, with special emphasis on *Brassica napus* [1–3]; however, in *B. oleracea* var. *capitata*, positive results have been achieved using selected accessions with high division frequencies and regeneration capacity [4–6]. The effect of genotype has overridden many efforts to use *B. oleracea* var. *capitata* protoplast cultures in biotechnology [5,7,8]; therefore, it is crucial to develop improvements in protocols leading to an increase in mitotic activity and regeneration process.

Plant hormones, such as auxins and cytokinins, have been considered to be the main growth regulators involved in cell divisions [9]. In several studies, *B. oleracea* protoplasts were cultured in liquid media containing various proportions of auxins and cytokinins; however, in some genotypes, despite the presence of plant growth regulators, divisions of protoplast-derived cells were very low or the cells failed to divide [6,10–12]. Mitotic activity of protoplast-derived cells can also be improved by the application of a nurse culture, feeder layers, or conditioned media with addition of filtrate prepared from a suspension culture of rapidly dividing cells [13]. Chen and others [14] described a procedure for regenerating shoots from hypocotyl-derived protoplasts of *B. oleracea* var. *capitata* f. *rubra* in the presence of viable protoplasts of *B. juncea*. Another example

is the use of feeder layers from *B. campestris* suspension cultures for protoplast culture of *B. oleracea* ssp. *botrytis* and *B. oleracea* ssp. *italica* [15]. Such an approach might be beneficial; however, it is laborious and time consuming and requires the establishment of actively dividing suspension cultures [16]. Favorable effect of nurse cultures, feeder layers, or conditioned media suggested the presence of division-inducing factors that are secreted into the medium with cultured cells [17,18]. Many efforts have been made to identify and characterize such factors and initial amino acids, purine derivatives, highly hydrophilic and heat stable agents, or protease-resistant small compounds were considered as putative mitogenic inducers; however, the available assay methods were not sufficiently sensitive to define any specific compound [19–21]. The elaboration of highly sensitive bioassay systems for the characterization of mitotic activity in low-density cultures of asparagus in the presence of conditioned medium led to the discovery of phytosulfokines (PSK), biologically active components with mitogenic activity [22]. Phytosulfokines PSK- $\alpha$  and PSK- $\beta$  are small (five and four amino acids, respectively) peptides containing two post-translationally sulfated tyrosine residues. Matsubayashi and others [23] reported that PSK- $\beta$  is an enzymatically degraded product of PSK- $\alpha$ . They also observed that the unsulfated PSK- $\alpha$  analogue was less active than the native form of PSK- $\alpha$ . Moreover, the studies of Bahyrycz and others [24] showed that PSK analogues with different modifications in the position of the sulfur atoms had weak or no biological effects. These results point out that sulfur plays an important role in the creation of biological properties of phytosulfokine. PSK- $\alpha$  is a primary signal molecule responsible for cell to cell communication [25]. Studies in *Arabidopsis thaliana* on PSK signaling revealed two genes (*PSKR1* and *PSKR2*) encoding for PSK receptors. Analysis of *pskr1-3* and *pskr2-1* T-DNA insertion lines showed that root elongation was predominately controlled through *PSKR1*. *PSKR1* signaling regulated root growth mainly by increasing cell size [26]. Studies on *A. thaliana* hypocotyl-derived protoplasts showed that PSK signaling controls osmotically-driven cell expansion suggesting that PSK- $\alpha$  acts as an osmoregulator and involves  $K^+$  uptake [27].

It has been reported that PSK- $\alpha$  plays an important role in plant growth and development due to its contribution to the formation of somatic embryos [28,29], its role in adventitious buds and roots formation [25,30], and its contribution to cellular longevity and the cell division cycle [31,32]. The objective of this study was to analyze the effect of PSK- $\alpha$  on viability and division frequencies of *Brassica oleracea* protoplast-derived cells embedded in calcium alginate layers.

## Material and methods

### Plant material

As protoplasts source, we used seven accessions of *B. oleracea* var. *capitata*, including five cultivars: ‘Sława z Gołębiewa’ (Plantico, Poland), ‘Reball F1’ and ‘Ramkila F1’ (Syngenta Seeds, Poland), ‘Badger Shipper’ and ‘Oregon 123’ (Warwick Genetic Resources Unit, United Kingdom), and two breeding lines: LM and LM98 (KHiNO Polan, Poland). Seeds of all accessions were sterilized in 70% (v/v) ethanol for 2 min, 10% (w/v) chloramine T (Biochemie GmbH, Poland) for 20 min, and washed three times with sterile distilled water for 5 min each. Seeds were germinated on MS [33] medium supplemented with 0.8% (w/v) agar (Biocorp, Poland). Prior to isolation of protoplasts from leaves, seeds were placed in sterile 500-mL plastic boxes (Pakler Lerka, Poland) containing 80 mL of MS medium and kept at  $26 \pm 2^\circ\text{C}$  under a 16-h photoperiod with light intensity  $55 \mu\text{mol m}^{-2} \text{s}^{-1}$ . Etiolated hypocotyls were obtained from seeds placed in a 90-mm Petri dish with 25 mL of MS medium and kept in the dark at  $26 \pm 2^\circ\text{C}$ .

### Protoplast isolation and culture

Young leaves of 4-week-old plants and hypocotyls from 2-week-old seedlings were used as donor tissue for protoplast isolation. Protoplasts were isolated and cultured according to the protocol described by Kielkowska and Adamus [10]. The enzyme solution for

tissue digestion consisted of 0.5% (w/v) cellulase Onozuka R-10 (Duchefa Biochemie, the Netherlands), 0.1% (w/v) pectolyase Y-23 (Duchefa Biochemie), 3 mM  $\text{CaCl}_2$ , 2 mM of 2-(*N*-morpholino) ethanesulfonic acid (MES; Sigma-Aldrich, Poland), and 0.4 M mannitol, and was filter-sterilized (0.22  $\mu\text{m}$ ; Millipore, United Kingdom). Isolated and purified protoplasts were immobilized in filter-sterilized alginate according to the protocol of Kielkowska and Adamus [34]. Final density of protoplasts was  $4 \times 10^5$  per mL of culture medium. Alginate discs with embedded protoplasts were transferred to 60-mm Petri dishes containing 4 mL of culture medium. Basal culture medium was CPP medium according to Dirks et al. [35], supplemented with 0.1 mg L<sup>-1</sup> 2,4-dichlorophenoxyacetic acid (2,4-D) and 0.2 mg L<sup>-1</sup> zeatin, pH 5.6, filter sterilized (0.22  $\mu\text{m}$ ; Millipore). Basal medium was supplemented with 0.1  $\mu\text{M}$  of PSK- $\alpha$  (Peptide Institute, Inc., Japan). Alginate discs with embedded protoplasts cultured in basal medium were used as a control. Culture medium was renewed after 10 days and cultures were incubated in the dark at  $26 \pm 2^\circ\text{C}$ .

Protoplast viability was estimated on the first and fifth day after isolation by staining with fluorescein diacetate (FDA) according to Anthony and others [36]. Observations of yellow-green fluorescence of viable cells were performed using an Axiovert S 100 (Carl Zeiss, Germany) inverted microscope.

### Shoot regeneration

Protoplast-derived callus colonies were freed from alginate layers according to the protocol of Damm and Willmitzer [37] and transferred for further development into 90-mm Petri dishes with 20 mL of regeneration medium consisting of B5 [38], micro-, macroelements and vitamins with 750 mg L<sup>-1</sup>  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ , 1 mg L<sup>-1</sup> benzylaminopurine (BA), and 20 g L<sup>-1</sup> sucrose. The medium was adjusted to a pH of 5.7–5.8 and 0.28% (w/v) Gelrite was added (Sigma-Aldrich) prior to autoclaving (20 min at  $121^\circ\text{C}$ ; 0.1 MPa). Developing shoots were transferred to fresh medium every 3 to 4 weeks. Rooting was performed on MS [33] medium with 1 mg L<sup>-1</sup> of indoleacetic acid (IAA). Cultures were maintained at  $26 \pm 2^\circ\text{C}$  with a 16-h photoperiod at a light intensity of 55  $\mu\text{mol m}^{-2} \text{s}^{-1}$ .


### Statistical analysis

The single experiment consisted of four independent protoplast isolations with a single treatment represented by four Petri dishes. Experiment was repeated three times. Viability of protoplasts was calculated as the number of protoplasts with yellow-green fluorescence divided by the total number of observed cells ( $\times 100$ ). Division frequency was expressed as the number of dividing protoplast-derived cells per total number of observed cells ( $\times 100$ ). Regeneration frequency was calculated as the number of shoots regenerated from callus per total number of calli cultured on the regeneration medium ( $\times 100$ ). Analyses were performed using Statistica ver. 10.0 (Statsoft, Inc., Poland) with the ANOVA module. Mean separation was performed with Tukey's honestly significant difference (HSD).


## Results

### Protoplast viability

On the day following isolation, protoplasts of majority of tested accessions ('Oregon 123', 'Reball F1', 'Badger Shipper', and breeding line LM) had relatively high (88–98%) viability, irrespectively from the presence of PSK- $\alpha$  in the medium (Fig. 1). In 'Sława z Gołębiewa' and 'Ramkila F1', higher viability (respectively 91% and 86%) was observed in protoplasts cultured in PSK-free control medium. The lowest protoplast viability was observed in breeding line LM98; however, protoplasts of this accession cultured in the presence of 0.1  $\mu\text{M}$  of PSK- $\alpha$  were more viable (82%) in comparison with protoplasts



**Fig. 1** Effect of phytosulfokine- $\alpha$  and accession on the viability of *B. oleracea* protoplast-derived cells in the first and fifth day of culture. Bars represent means for leaf- and hypocotyl-derived cells jointly  $\pm$ SE. Black and grey bars should be considered separately. Values followed by the same letter are not significantly different ( $p \leq 0.05$ , HSD).



**Fig. 2** Effect of phytosulfokine- $\alpha$  and protoplast source on the viability of *B. oleracea* protoplast-derived cells in first and fifth day of culture. Bars represent means  $\pm$ SE. Black and grey bars should be considered separately. Values followed by the same letter are not significantly different ( $p \leq 0.05$ , HSD).


cultured in control medium (79%). On the fifth day of culture, the number of viable cells decreased in all tested accessions. At this period of time, the highest percent (82) of viable cells were observed in 'Reball F1' on the medium with PSK- $\alpha$ ; moreover, in this accession, the decrease of viability in relation to the first day of culture was lowest (7%). The highest decrease was observed in 'Oregon 123' on both tested media, where cell viability on the fifth day was approximately 55% lower than on the first day of culture. In 'Sława z Gołębiewa', 'Ramkila F1', and 'Badger Shipper', the highest decrease in cell viability (approximately 51–52%) was observed in the control medium. In these accessions, as well as in breeding line LM98, protoplasts cultured in the medium supplemented with PSK- $\alpha$  were more viable.

On the first day of culture, we observed no differences in viability between leaf- and hypocotyl-derived protoplasts nor the protoplast cultured in the presence of PSK- $\alpha$  and in control medium (Fig. 2). On the fifth day of culture, leaf-derived cells had similar viability on both tested media; however, hypocotyl-derived cells cultured in the medium supplemented with PSK- $\alpha$  (52%) were more viable compared to PSK-free control medium (43%).

## Division frequency

Mitotic divisions were observed in all tested accessions, albeit with various frequencies, depending on the PSK- $\alpha$  presence in the culture medium (Fig. 3). On the fifth day of culture, the average division frequencies in all accessions did not exceed 15% (Fig. 3a). The highest division frequency was observed in breeding line LM98 (15%) and 'Sława z Gołębiewa' (14%) on the medium supplemented with 0.1  $\mu$ M of PSK- $\alpha$ , while lowest one was found in 'Ramkila F1' (0.3%) on the control medium. At this period of time, protoplast-derived cells of 'Badger Shipper' cultured on the PSK-free control medium, did not divide; however, on the medium with PSK- $\alpha$  0.5% of cultured cells underwent mitosis.


On the fifteenth day of culture, the division frequencies in all tested accessions increased (Fig. 3b). Supplementation of the culture medium with PSK- $\alpha$  showed a beneficial effect on mitotic activity of protoplast-derived cells in the majority of the tested accessions in comparison with the control. On the fifteenth day of culture, the highest division frequency of protoplast-derived cells was observed for 'Reball F1' in the medium supplemented with PSK- $\alpha$  (56%), which was approximately 10% higher than that in the PSK-free control. The protoplast-derived cells of breeding line LM98 cultured in control medium divided with a frequency of 5%, while in the medium supplemented with PSK- $\alpha$  47% of protoplast-derived cells underwent divisions. The stimulating effect of PSK- $\alpha$  was also recorded for 'Ramkila F1', 'Sława z Gołębiewa', and breeding line LM, where the mitotic activity of protoplast-derived cells cultured in the



**Fig. 3** Effect of phytosulfokine- $\alpha$  and accession on division frequency of *B. oleracea* protoplast-derived cells in the fifth (a) and fifteenth (b) day of culture. Bars represent means for leaf- and hypocotyl-derived cells jointly  $\pm$ SE. Values followed by the same letter are not significantly different ( $p \leq 0.05$ , HSD).

control medium was 5%, 14%, and 23%, respectively, while activity of cells cultured in the medium supplemented with PSK- $\alpha$  was 15%, 38%, and 33%, respectively. For two cultivars, 'Oregon 123' and 'Badger Shipper', although the division frequencies of protoplast-derived cells cultured in the media supplemented with PSK- $\alpha$  were slightly increased compared to the control, the differences were not significant.

On the fifteenth day of culture, higher division frequency was observed among hypocotyl-derived cells cultured in the media supplemented with PSK- $\alpha$  (33%) compared to the PSK-free control (10%) (Fig. 4). Similar tendency was observed for leaf-derived cells, although frequencies of divisions were lower.



**Fig. 4** Effect of phytosulfokine- $\alpha$  and protoplast source on the division frequency of *B. oleracea* protoplast-derived cells in the fifth and fifteenth day of culture. Bars represent means  $\pm$ SE. Black and grey bars should be considered separately. Values followed by the same letter are not significantly different ( $p \leq 0.05$ , HSD).

**Tab. 1** Shoot regeneration from the protoplast-derived callus colonies of *B. oleracea*.

Factor	Shoot regeneration (% $\pm$ SE)
Accession	
'Sława z Gołębiewa'	6.7 $\pm$ 2.9 <sup>b</sup>
'Ramkila F1'	0.0 $\pm$ 0.0 <sup>c</sup>
'Reball F1'	20.9 $\pm$ 2.9 <sup>a</sup>
'Oregon 123'	0.0 $\pm$ 0.0 <sup>c</sup>
'Badger Shipper'	0.0 $\pm$ 0.0 <sup>c</sup>
LM	7.1 $\pm$ 3.1 <sup>b</sup>
LM98	5.7 $\pm$ 1.5 <sup>b</sup>
Source of Protoplasts*	
Leaves	8.3 $\pm$ 1.8 <sup>a</sup>
Hypocotyls	12.1 $\pm$ 3.1 <sup>a</sup>

Values of the factor Accession represent means for leaf- and hypocotyl-derived cells jointly. Values followed by the same letter are not significantly different ( $p \leq 0.05$ , HSD).


\* Calculation based only on accessions with regeneration.

## Shoot regeneration

Mitotic divisions of cultured protoplast-derived cells resulted in formation of multiple cell colonies, visible with the naked eye after approximately 4 weeks of culture. Colonies of calli were freed from alginate layers and placed on a solid regeneration medium. Although callus colonies were noted in all seven tested accessions, only in four of them the divisions were sustained, leading to shoot development (Tab. 1). The callus colonies of 'Ramkila F1', 'Oregon 123', and 'Badger Shipper' failed to regenerate. The highest regeneration (21%) was observed in 'Reball F1'. There were no differences in shoot regeneration between 'Sława z Gołębiewa' (7%) and breeding lines LM (7%) and LM98 (6%). We also observed no differences in shoot regeneration between protoplast sources.

Detailed analysis of shoot regeneration capacity in dependency from the presence or absence of PSK- $\alpha$  in the earlier cultivation stages (in the liquid medium for culturing the protoplasts) showed that for the high-responding 'Reball F1', the effect of PSK- $\alpha$  was insignificant (Fig. 5). In all remaining responsive accessions, regeneration capacity was higher among callus colonies derived from protoplasts cultured in medium supplemented with PSK- $\alpha$ . In 'Sława z Gołębiewa' and breeding line LM, the increased shoot regeneration was twofold higher compared to the control. In breeding line LM98, regeneration frequency of callus colonies developed from protoplasts cultured in the control media was 0.3%; however, from PSK- $\alpha$  supplemented cultures, regeneration frequency increased up to 11%.





**Fig. 5** Effect of phytosulfokine- $\alpha$  and accession on shoot regeneration from *B. oleracea* protoplasts. Bars represent means for leaf- and hypocotyl-derived cells jointly  $\pm$ SE. Values followed by the same letter are not significantly different ( $p \leq 0.05$ , HSD).

## Discussion

Recently, peptides have been recognized as signaling molecules playing an important role in plant cell growth and development [31,39]. Results showed that in planta, PSK signaling participates in the control of shoot and root growth. Hypocotyl elongation and root growth are promoted by PSK, mainly through the signaling of cell elongation rather than cell division [26,27]. However, in suspension cultures PSK promotes the proliferation of cells of both monocot and dicot plants [23,28,31,32] and due to this feature, phytosulfokine- $\alpha$  has been proposed as a candidate to supplementing plant in vitro cultures together with other well-known plant growth regulators [40,41]. These results were the assumption to perform our experiments. In our previous work, we have developed a simple and efficient protocol for *Brassica oleracea* protoplast isolation and culture with embedding protoplasts in calcium alginate layers [10,34]; however, using this method, some accessions were recalcitrant to applied culture conditions and no or very low division frequency was noted. 'Oregon 123' and 'Badger Shipper' were classified as recalcitrant (unpublished data), becoming suitable experimental material to test the potential of PSK- $\alpha$  in vitro. Grzebelus et al. [32] evaluated the effect of supplementation of the culture medium with 0–0.01–0.1 and 1  $\mu$ M of PSK- $\alpha$  in sugar beet protoplast cultures. The highest plating efficiency (20%) was observed at a concentration of 0.1  $\mu$ M of PSK- $\alpha$ . Moreover, the same concentration of PSK- $\alpha$  promoted microspore embryogenesis in triticale and wheat [40]; therefore, in our study we applied this concentration of PSK- $\alpha$  to the culture medium.

Our results showed that the viability of protoplasts was independent from protoplast source tissue. Protoplast viability of all tested accessions on Day 1 after isolation was high (approximately 80–90%), and in that time the effect of PSK- $\alpha$  on protoplast viability was less obvious; however, the differences were visible later, on fifth day of culture, when viability decreased in all tested accessions. Decrease in protoplast viability over the first days of culture was reported previously [32,34] and might be associated with the damage in plasma membranes as a result of enzyme digestion and purification, osmotic stress, or oxidative stress during isolation and culture establishment [13,42]. Here, on Day 5, protoplast-derived cells of majority of tested accessions were more viable when cultured in the medium with PSK- $\alpha$  in comparison with the control.

In this study, mitotic activity of protoplasts was genotype-dependent, but also PSK- $\alpha$  supplemented to the culture medium increased mitotic activity of cultured protoplast-derived cells over twofold compared to cells cultured in the PSK-free control medium. Detailed results of the interaction of PSK- $\alpha$  supplementation in the culture medium and the genotype showed interesting relations – in the well-responding cultivar 'Reball F1', the effect of PSK- $\alpha$  was not as strong as in the low-responding breeding line LM98, where mitotic activity of protoplast-derived cells cultured in the presence of PSK was near 10-fold higher, compare to PSK-free the control. A beneficial effect of PSK- $\alpha$  was also observed for 'Sława z Gołębiewa', 'Ramkila F1', and breeding line LM. In recalcitrant

cultivars used in this study ('Oregon 123' and 'Badger Shipper'), although statistically insignificant, an increase in division frequencies of protoplast-derived cells cultured on PSK- $\alpha$  supplemented medium compared to the PSK-free control was observed.

Matsubayashi et al. [31] demonstrated the effect of phytosulfokines supplemented to the culture media on rice protoplasts. Protoplasts embedded in agarose blocks in low density cultures ( $5 \times 10^4$  of cells per mL of medium) and cultured in the presence of PSK- $\alpha$  and PSK- $\beta$  divided frequently (58–66%) and gave rise to microcallus colonies after 4 weeks of cultivation. Results of experiments with mesophyll protoplast cultures of sugar beet [32] showed that plating efficiency of control cultures was 1%, while in the medium with PSK- $\alpha$  it increased up to 28%. The authors demonstrated that application of exogenous PSK reversed the recalcitrance of in vitro cultured mesophyll sugar beet cells. Recently, Maćkowska et al. [41] cultured protoplasts of different *Daucus* accessions in the presence of PSK. Their results showed that in five out of seven tested accessions, PSK- $\alpha$  at a concentration of 100 nM (0.1  $\mu$ M) led to an increase in plating efficiency (55–80%) in comparison with the control (40–65%). However, for two wild *Daucus* species, stimulation of cell divisions was observed only in early cultures and callus colonies were not formed. The above-mentioned results and observed in our study differences in culture efficiency among seven tested cabbage accessions suggest that supplementation of the liquid culture media with PSK- $\alpha$  might increase mitotic activity of protoplast-derived cells; however, the genotype plays the main role in the responsiveness of isolated plant cells to in vitro cultures [5,7,37,43].

In protoplast cultures, cell proliferation highly depends on culture density, and it was reported that cell divisions cease in low density cultures [28,44]. Yamakawa et al. [30] reported that mitotic activity in low-density suspension cultures is not stimulated by known plant hormones or defined nutrients; however, can be induced by the addition of conditioned medium prepared from cultures of rapidly proliferating cells. The successful stimulation of cell divisions in low density cultures was induced in *B. napus* [45], but not in *B. oleracea* var. *capitata* [2,46]. In this study, we cultured protoplasts in optimal density [10] and in the presence of PSK- $\alpha$ ; however, in two genotypes, we did not observed satisfactory stimulation of cell divisions. We used PSK- $\alpha$  in concentration of 0.1  $\mu$ M, and it might be likely that this concentration was not efficient for these two accessions for the promotion of cell divisions. In contrast, in high-responsive genotypes PSK- $\alpha$  had only a minor effect.

We also observed the beneficial effect of supplementation of the protoplast culture medium in PSK on the regeneration process. Phytosulfokines have the potential to induce cell divisions, but our results showed it was also important for its maintenance. In controls without PSK- $\alpha$ , even if cell divisions were induced, they ceased at some point, and as a result shoot regeneration was not observed. It seems that the mitotic divisions stimulated by PSK- $\alpha$  were sustained even after transferring protoplast-derived callus tissue on the solid regeneration medium, what resulted in a higher percentage of regenerated shoots obtained from callus colonies developed in the media supplemented with PSK- $\alpha$ .

To the best of our knowledge, this is the first report showing the effect of PSK- $\alpha$  on *Brassica oleracea* protoplast cultures. We demonstrate that mitotic activity of cabbage protoplasts can be increased by supplementation of culture medium with PSK- $\alpha$ . The effect of PSK- $\alpha$  was more pronounced in some low-responding genotypes, and we observed 10–47% increase in division frequency of protoplast-derived cells cultured in medium supplemented with PSK- $\alpha$  compared to PSK-free control medium. The supplementation of the culture medium in PSK- $\alpha$  had a minor effect on mitotic activity of two out of seven tested accessions, showing that the genotype plays a major role in the response of cabbage to protoplast culture. PSK- $\alpha$  supplemented into the liquid culture medium had an indirect effect on shoot regeneration by inducing sustained cell divisions leading to an increase in shoot regeneration. Further research will be focused on the evaluation of the effect of broader spectra of PSK- $\alpha$  concentrations in the liquid medium for culturing cabbage protoplasts as well as on the examination of the effect of direct supplementation of solid medium with phytosulfokines on the regeneration process.



## Acknowledgments

The authors would like to thank Marta Pałowska for help with performing protoplast cultures.

## References

1. Hansen LN, Earle ED. Novel flowering and fatty acid characters in rapid cycling *Brassica napus* L. resynthesized by protoplast fusion. *Plant Cell Rep.* 1994;14:151–156. <https://doi.org/10.1007/BF00233780>
2. Sundberg E, Glimelius K. A method for production of interspecific hybrids within Brassicaceae via somatic hybridization, using resynthesis of *Brassica napus* as a model. *Plant Sci.* 1986;43:155–162. <https://doi.org/10.1007/BF00221131>
3. Yamagishi H, Landgren M, Forsberg J, Glimelius K. Production of asymmetric hybrids between *Arabidopsis thaliana* and *Brassica napus* utilizing an efficient protoplast culture system. *Theor Appl Genet.* 2002;104:959–964. <https://doi.org/10.1007/s00122-002-0881-9>
4. Glimelius K. High growth rate and regeneration capacity of hypocotyl protoplasts in some Brassicaceae. *Physiol Plant.* 1984;61:38–44. <https://doi.org/10.1111/j.1399-3054.1984.tb06097.x>
5. Holme IB, Torp AM, Hansen LN, Andersen SB. Quantitative trait loci affecting plant regeneration from protoplasts of *Brassica oleracea*. *Theor Appl Genet.* 2004;108:1513–1520. <https://doi.org/10.1007/s00122-003-1570-z>
6. Zhao KN, Bittisnich DJ, Halloran GM, Whitecross MI. Studies of cotyledon protoplast cultures from *B. napus*, *B. campestris* and *B. oleracea*. II: callus formation and plant regeneration. *Plant Cell Tissue Organ Cult.* 1995;40:73–84. <https://doi.org/10.1007/BF00041121>
7. Hansen LN, Ortiz R, Andersen SB. Genetic analysis of protoplast regeneration ability in *Brassica oleracea*. *Plant Cell Tissue Organ Cult.* 1999;58:127–132. <https://doi.org/10.1023/A:1006359804328>
8. Jourdan PS, Earle ED, Mutschler MA. Improved protoplast culture and stability of cytoplasmic traits in plants regenerated from leaf protoplasts of cauliflower (*Brassica oleracea* ssp. *botrytis*). *Plant Cell Tissue Organ Cult.* 1990;21:227–236. <https://doi.org/10.1007/BF00047615>
9. Davies PJ. The plant hormones: their nature, occurrence and functions. In: Davies PJ, editor. *Plant hormones biosynthesis, signal transduction, action.* Dordrecht: Springer; 2010. p. 1–15. [https://doi.org/10.1007/978-94-009-3585-3\\_1](https://doi.org/10.1007/978-94-009-3585-3_1)
10. Kielkowska A, Adamus A. An alginate-layer technique for culture of *Brassica oleracea* L. protoplasts. *In Vitro Cell Dev Biol.* 2012;48:265–273. <https://doi.org/10.1007/s11627-012-9431-6>
11. Robertson D, Earle ED, Mutschler MA. Increased totipotency of protoplasts from *Brassica oleracea* plants previously regenerated in tissue culture. *Plant Cell Tissue Organ Cult.* 1988;14:15–24. <https://doi.org/10.1007/BF00029571>
12. Sheng X, Zhao Z, Yu H, Wang J, Xiaohui Z, Gu H. Protoplast isolation and plant regeneration of different double haploid lines of cauliflower (*Brassica oleracea* var. *botrytis*) *Plant Cell Tissue Organ Cult.* 2011;107:513–520. <https://doi.org/10.1007/s11240-011-0002-z>
13. Davey MR, Anthony P, Power JB, Lowe KC. Plant protoplasts: status and biotechnological perspectives. *Biotechnol Adv.* 2005;23:131–171. <https://doi.org/10.1016/j.biotechadv.2004.09.008>
14. Chen LP, Zhang MF, Xiao QB, Wu JG, Hirata Y. Plant regeneration from hypocotyl protoplasts of red cabbage (*Brassica oleracea*) by using nurse cultures. *Plant Cell Tissue Organ Cult.* 2004;77:133–138. <https://doi.org/10.1023/B:TICU.0000016811.29125.18>
15. Walters TW, Earle ED. A simple, versatile feeder layer system for *Brassica oleracea* protoplast culture. *Plant Cell Rep.* 1990;9:316–319. <https://doi.org/10.1007/BF00232859>
16. Mustafa NR, de Winter W, van Iren F, Verpoorte R. Initiation, growth and cryopreservation of plant cell suspension cultures. *Nat Protoc.* 2011;6:715–742. <https://doi.org/10.1038/nprot.2010.144>
17. Folling M, Madsen S, Olesen A. Effect of nurse culture and conditioned medium on colony formation and plant regeneration from *Lolium perenne* protoplasts. *Plant Sci.* 1995;108:229–239. [https://doi.org/10.1016/0168-9452\(95\)04146-L](https://doi.org/10.1016/0168-9452(95)04146-L)
18. Somers DA, Birnberg PR, Petersen WL, Brenner ML. The effect of conditioned medium on colony formation from black Mexican sweet corn protoplast. *Plant Sci.* 1987;53:249–

256. [https://doi.org/10.1016/0168-9452\(87\)90162-2](https://doi.org/10.1016/0168-9452(87)90162-2)
19. Bellincampi D, Morpurgo G. Evidence for the presence of a second conditioning factor in plant cell culture. *Plant Sci.* 1989;65:125–130. [https://doi.org/10.1016/0168-9452\(89\)90215-X](https://doi.org/10.1016/0168-9452(89)90215-X)
  20. Sargent PA, King J. Investigations of growth-promoting factors in conditioned soybean root cells and in the liquid medium in which they grow: ammonium, glutamine, and amino acids. *Can J Bot.* 1974;52:1747–1755. <https://doi.org/10.1139/b74-226>
  21. Stuart R, Street HE. Studies on the growth in culture of plant cells. *J Exp Bot.* 1971;22:96–106. <https://doi.org/10.1093/jxb/22.1.96>
  22. Matsubayashi Y, Sakagami Y. Phytosulfokine, sulfated peptides that induce the proliferation of single mesophyll cells of *Asparagus officinalis* L. *Proc Natl Acad Sci USA.* 1996;93:7623–7627. <https://doi.org/10.1073/pnas.93.15.7623>
  23. Matsubayashi Y, Hanai H, Hara O, Sakagami Y. Active fragments and analogs of the plant growth factor, phytosulfokine: structure–activity relationship. *Biochem Biophys Res Commun.* 1996;225:209–214. <https://doi.org/10.1006/bbrc.1996.1155>
  24. Bahyrycz A, Matsubayashi Y, Ogawa M, Sakagami Y, Konopinska D. Plant peptide hormone phytosulfokine (PSK- $\alpha$ ): synthesis of new analogues and their biological evaluation. *J Pept Sci.* 2004;10:462–469. <https://doi.org/10.1002/psc.492>
  25. Yang G, Shen S, Kobayashi T, Matsubayashi Y, Sakagami Y, Kamada H. Stimulatory effects of a novel peptidyl plant growth factor, phytosulfokine- $\alpha$  on the adventitious bud formation from callus of *Antirrhinum majus*. *Plant Biotechnol.* 1999;16:231–234. <http://doi.org/10.5511/plantbiotechnology.16.231>
  26. Kutschmar A, Rzewuski G, Stührwohldt N, Beemster GTS, Inzé D, Sauter M. PSK- $\alpha$  promotes root growth in *Arabidopsis*. *New Phytol.* 2009;181:820–831. <https://doi.org/10.1111/j.1469-8137.2008.02710.x>
  27. Stührwohldt N, Dahlke RI, Steffens B, Johnson A, Sauter M. Phytosulfokine- $\alpha$  controls hypocotyl length and cell expansion in *Arabidopsis thaliana* through phytosulfokine receptor 1. *PLoS One.* 2011;6(6):e21054. <https://doi.org/10.1371/journal.pone.0021054>
  28. Hanai H, Matsuno T, Yamamoto M, Matsubayashi Y, Kobayashi T, Kamada H, et al. A secreted peptide growth factor, phytosulfokine, acting as a stimulatory factor of carrot somatic embryo formation. *Plant Cell Physiol.* 2000;41:27–32. <https://doi.org/10.1093/pcp/41.1.27>
  29. Igasaki T, Akashi N, Ujino-Hara T, Matsubayashi Y, Sakagami Y, Shinohara K. Phytosulfokine stimulates somatic embryogenesis in *Cryptomeria japonica*. *Plant Cell Physiol.* 2003;44:1412–1416. <https://doi.org/10.1093/pcp/pcg161>
  30. Yamakawa S, Sakuta C, Matsubayashi Y, Sakagami Y, Kamada H, Satoh S. The promotive effects of a peptidyl plant growth factor, phytosulfokine- $\alpha$ , on the formation of adventitious roots and expression of a gene for a root-specific cystatin in cucumber hypocotyls. *J Plant Res.* 1998;111:453–458. <https://doi.org/10.1007/BF02507810>
  31. Matsubayashi Y, Takagi L, Sakagami Y. Phytosulfokine- $\alpha$ , a sulfated pentapeptide, stimulates the proliferation of rice cells by means of specific high- and low-affinity binding sites. *Proc Natl Acad Sci USA.* 1997;94:13357–13362. <https://doi.org/10.1073/pnas.94.24.13357>
  32. Grzebelus E, Szklarczyk M, Greń J, Śniegowska K, Jopek M, Kacińska I, et al. Phytosulfokine stimulates cell divisions in sugar beet (*Beta vulgaris* L.) mesophyll protoplast cultures. *Plant Growth Regul.* 2012;67:93–100. <https://doi.org/10.1007/s10725-011-9654-2>
  33. Murashige T, Skoog F. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plant.* 1962;15:473–497. <https://doi.org/10.1111/j.1399-3054.1962.tb08052.x>
  34. Kielkowska A, Adamus A. Embedding in filter sterilized alginate enhances *Brassica oleracea* L. protoplast culture. *Acta Biol Crac Ser Bot.* 2014;56(2):20–26. <https://doi.org/10.2478/abcsb-2014-0018>
  35. Dirks R, Sidorov V, Tulmans C. A new protoplast culture in *Daucus carota* L. and its applications for mutant selection and transformation. *Theor Appl Genet.* 1996;93:809–815. <https://doi.org/10.1007/BF00224080>
  36. Anthony P, Otoni W, Power JB, Lowe KC, Davey MR. Protoplast isolation culture, and plant regeneration from *Passiflora*. In: Hall RD, editor. *Plant cell culture protocols*. Totowa, NJ: Humana Press; 1999. p. 169–181. (Methods in Molecular Biology; vol 111). <https://doi.org/10.1385/1-59259-583-9:169>

37. Damm B, Willmitzer L. Regeneration of fertile plants from protoplasts of different *Arabidopsis thaliana* genotypes. *Mol Gen Genet*. 1988;213:15–20. <https://doi.org/10.1007/BF00333392>
38. Gamborg OL, Miller RA, Ojima K. Nutrient requirements of suspension cultures of soybean root cells. *Exp Cell Res*. 1968;50:151–158. [https://doi.org/10.1016/0014-4827\(68\)90403-5](https://doi.org/10.1016/0014-4827(68)90403-5)
39. Ryan CA, Pearce G. Polypeptide hormones. *Plant Physiol*. 2001;125:65–68. <https://doi.org/10.1104/pp.125.1.65>
40. Asif M, Eudes F, Randhawa H, Amundsen E, Spaner D. Phytosulfokine  $\alpha$  enhances microspore embryogenesis in both triticale and wheat. *Plant Cell Tissue Organ Cult*. 2014;116:125–130. <https://doi.org/10.1007/s11240-013-0379-y>
41. Maćkowska K, Jarosz A, Grzebelus E. Plant regeneration from leaf-derived protoplasts within the *Daucus* genus: effect of different conditions in alginate embedding and phytosulfokine application. *Plant Cell Tissue Organ Cult*. 2014;117:241–252. <https://doi.org/10.1007/s11240-014-0436-1>
42. Tiburcio AF, Kaur-Sawhney R, Galston AW. Polyamine metabolism and osmotic stress – relation to protoplast viability. *Plant Physiol*. 1986;82:369–374. <https://doi.org/10.1104/pp.82.2.369>
43. Kirti PB, Bhat SR, Kumar VD, Parkash S, Chopra VL. A simple protocol for regenerating mesophyll protoplasts of vegetable brassicas. *J Plant Biochem Biotechnol*. 2001;10:49–51. <https://doi.org/10.1007/BF03263106>
44. Eun CH, Ko SM, Matsubayashi Y, Sakagami Y, Kamada H. Phytosulfokine- $\alpha$  requires auxin to stimulate carrot non embryogenic cell proliferation. *Plant Physiol Biochem*. 2003;41:447–452. [https://doi.org/10.1016/S0981-9428\(03\)00052-4](https://doi.org/10.1016/S0981-9428(03)00052-4)
45. Simmonds DH, Long NE, Keller WA. High plating efficiency and plant regeneration frequency in low density protoplast cultures derived from an embryogenic *Brassica napus* cell suspension. *Plant Cell Tissue Organ Cult*. 1991;27:231–241. <https://doi.org/10.1007/BF00157586>
46. Jie EY, Kim SW, Jang HR, In DS, Liu JR. Myo-inositol increases the plating efficiency of protoplast derived from cotyledon of cabbage (*Brassica oleracea* var. *capitata*). *J Plant Biotechnol*. 2011;38:69–76. <https://doi.org/10.5010/JPB.2011.38.1.069>