

Ecosystem recovery after climatic extremes enhanced by genotypic diversity

Thorsten B. H. Reusch^{*†}, Anneli Ehlers[‡], August Hämmerli^{*}, and Boris Worm^{*§}

^{*}Max-Planck-Institut für Limnologie, August-Thienemann-Strasse 2, 24306 Plön, Germany; [†]Leibniz Institut für Meereswissenschaften, Düsternbrooker Weg 20, 24105 Kiel, Germany; and [§]Biology Department, Dalhousie University, Halifax, NS, Canada B3H 4J1

Communicated by Jane Lubchenco, Oregon State University, Corvallis, OR, January 6, 2005 (received for review April 22, 2004)

Contemporary climate change is characterized both by increasing mean temperature and increasing climate variability such as heat waves, storms, and floods. How populations and communities cope with such climatic extremes is a question central to contemporary ecology and biodiversity conservation. Previous work has shown that species diversity can affect ecosystem functioning and resilience. Here, we show that genotypic diversity can replace the role of species diversity in a species-poor coastal ecosystem, and it may buffer against extreme climatic events. In a manipulative field experiment, increasing the genotypic diversity of the cosmopolitan seagrass *Zostera marina* enhanced biomass production, plant density, and faunal abundance, despite near-lethal water temperatures due to extreme warming across Europe. Net biodiversity effects were explained by genotypic complementarity rather than by selection of particularly robust genotypes. Positive effects on invertebrate fauna suggest that genetic diversity has second-order effects reaching higher trophic levels. Our results highlight the importance of maintaining genetic as well as species diversity to enhance ecosystem resilience in a world of increasing uncertainty.

global change | ecosystem functioning | ecological resilience | seagrass

Concerns about the accelerating loss of biodiversity have motivated an influential research program on the consequences of biodiversity loss at the species and functional group level (summarized in refs. 1–4). Although there are some important exceptions, most studies have found that in primary producers in particular, local species richness is positively correlated with a number of important ecosystem properties, such as productivity and resilience (5–8). Here we ask whether these results can be generalized to the genetic level, which represents the most fundamental aspect of biodiversity.

We hypothesized that genetic diversity may functionally replace the role of species diversity in species-poor aquatic macrophyte stands. The term species poor as it is used here refers to the low number of structuring primary producers. Seagrass meadows are a prominent example because they are often composed of only one or two species (9). Despite low species richness, seagrasses form the structural foundation of productive marine communities worldwide (10, 11), providing essential services such as nutrient cycling, habitat for fish and invertebrate species, and coastal erosion control (10, 12). In contrast to their species richness, their genotypic diversity can be very high and differs dramatically across all spatial scales (13, 14). This fact prompts the question of the ecological significance of such variation in diversity.

In a manipulative field experiment, we studied the role of genotypic diversity for ecosystem functioning in a coastal ecosystem dominated by eelgrass *Zostera marina*, the most abundant seagrass species of the northern hemisphere (15). Our experiment coincided with a period of an unprecedented heat wave that hit Europe in 2003 (16, 17). According to historical climate records, the return time for this extreme warming event was predicted to be >10,000 years (17). Nevertheless, it can be viewed as precursor for increasing climatic variability predicted for the coming decades (17, 18), and as such it provided an

unparalleled opportunity to assess the short-term ecological response of a coastal community to rapid global warming. Because anthropogenic alterations of the world's climate are already resulting in range shifts and species extinctions (19–21), an understanding of factors that enhance resistance and resilience (22) of populations and communities in the face of climatic extremes is fundamental for biodiversity conservation and environmental management.

Methods

Study Species. Eelgrass (*Z. marina* L.), a monoecious marine flowering plant or seagrass, is the dominant macrophyte species of shallow sedimentary shorelines in the northern hemisphere (15). Like all seagrasses (and many land plants), eelgrass reproduces predominantly vegetatively (23). In the study area, the southwestern Baltic Sea, 2-year observations of genetic diversity in permanent plots of 1 m × 1 m ($n = 24$) indicated that only 7% of the yearly shoot turnover was due to sexual reproduction (seedlings) and 93% was due to vegetative (clonal) growth (T.B.H.R., unpublished data). The genetic diversity of eelgrass has two clearly defined components: the genomic diversity in a sample of genotypes (24) and the number of genotypes (clones) per area (25). In contrast to previous experiments (26), both components can now be precisely separated by using high-resolution molecular markers (27). Here, we use DNA microsatellites that are assumed to be selectively neutral to identify clones (genotypes) for a replicated manipulation of genotypic (clonal) diversity.

Study Site and Experimental Design. We tested the hypothesis that higher genotypic diversity increases ecosystem functioning in a field experiment that manipulated the number of eelgrass genotypes (clones) in plots situated at a shallow protected estuary of the southwestern Baltic Sea (Maasholm, Germany, 54°41'N, 10°12'E). The experiment was conducted by using SCUBA in a water depth of 1.6–1.8 m. At our high-latitude study site, growth of macrophytes is highly seasonal and confined to May–September. Thus, our experimental period from May 16 through Sep. 26, 2003, covered almost the complete growing season. To identify donor genotypes, we set up a marked grid of 10 m × 9.3 m in the vicinity (<20 m distance) of the experimental site in mid-April 2003. A leaf tip (3–5 cm) of candidate donor plants was sampled at 33.3-cm intervals (i.e., $31 \times 29 = 899$ grid points) and genotyped by using microsatellite markers (see below). This method allowed us to construct a detailed clonal map (see Fig. 4, which is published as supporting information on the PNAS web site). After selection of the appropriate subareas within the grid (Fig. 4), naturally occurring clusters of eelgrass leaf shoots (ramets) were harvested without breaking the rhizome connections and were reassembled into desired mixtures without leaving the ambient water. Shoots of all experimental units were planted within 24 h and secured in the sediment with wire staples.

[†]To whom correspondence should be addressed. E-mail: reusch@mpil-ploen.mpg.de.

multivariate analysis of variance with the six most abundant species to test for an overall effect on faunal richness. Univariate tests were then performed for total abundance, and for three functional groups (grazers, filter feeders, and detritivores). All data were transformed as appropriate (usually square root or \log_{10}) to meet assumptions of the analysis (homogeneity of variances/normality).

Results

At the study site, water temperatures frequently attained levels above which *Z. marina* ceases growing (ca. 20°C; refs. 26 and 32), or even starts to die off (ca. 25°C; ref. 33; Fig. 1A). Accordingly, heat-related mortality reduced shoot numbers in the experiment by $\approx 50\%$ from May through August (Fig. 1B and C). This result contrasts markedly with several independent experiments that used a similar planting technique during normal climatic conditions; these experiments consistently showed low mortality and a 3-fold increase in shoot density from May through August (30, 34), making it unlikely that transplant shock was responsible for this decline (but see ref. 35). During the first major warming event at the beginning of June, unusual signs of heat-stress-related mortality were already present (discolored meristems, root necrosis, and leaf loss). In a subsample of nonmanipulated plants, half (10/20) had apparently died from heat stress during the warmest period (beginning of August). In nearby meadows, these losses amounted to a 48–52% decline in leaf shoot density and a 56% reduction in shoot height compared with data from 2001, 2002, and 2004 ($n = 10\text{--}12$, t test, all $P < 0.0001$), years for which average summer temperatures were recorded. The frequency of plants that showed discolored meristems was highly correlated with the number of leaf shoots in experimental plots during peak perturbation, further supporting a role for heat stress for plant mortality (response shoot density on Aug. 28, 2004; $r^2 = 0.52$, $P = 0.0013$). In the experiment, shoot densities increased markedly only when water temperatures dropped below 20°C in mid-August. By then, the diverse experimental plots had attained natural shoot densities observed in nearby untouched meadows (Fig. 1B; t test, $P = 0.58$). After mid-August, three- and six-genotype treatments diverged markedly from the monocultures and showed faster recovery of shoot density (Fig.

1B). Higher recovery rates were statistically supported when we used the slopes of plotwise temporal regressions of shoot density (ln transformed) vs. time as response variable. Slopes were significantly higher with increasing genotypic diversity, suggesting faster recovery and possibly, higher ecological resilience [linear model, including blocking factor, $F_{(1,23)} = 8.2$, $P = 0.009$].

Our design allowed a comparative analysis of all clones because they were planted in replicated one-genotype treatments. The genotypes A–F showed very different responses to the climate event [one-way ANOVA; response variable final shoot density, $F_{(5,18)} = 4.04$, $P = 0.01$]. Although genotype A increased almost 3-fold over the experimental period, genotype F never recovered from a 50% loss in shoot density during the heat wave [Fig. 1C; linear contrast, $F_{(1,18)} = 14.1$, $P = 0.001$].

When the experiment was sampled at the end of the growth period, there was a positive correlation between genotypic diversity and both shoot number and dry biomass (Fig. 2A and B). The yield of experimental plots slightly exceeded the highest yields observed in monocultures whereas the variances were progressively reduced [F test, variance one-genotype vs. six-genotype treatment; biomass $F_{(24,12)} = 3.78$, $P = 0.01$; shoot number $F_{(24,12)} = 2.74$, $P = 0.035$]. Such variance reduction may be due either to a lack of combinatorial replication at the highest diversity level (six genotypes), or to niche complementarity, i.e., a true diversity effect (36). Positive diversity effects were supported in an analysis of net biodiversity effects (ΔY , or overyield) that compared the final shoot number or biomass expected from the one-genotype plots with the observed yield (31). Significant ΔY values were present for both variables, shoot density and biomass (t tests against null-expectation $\Delta Y = 0$, $P = 0.045$ and 0.007 for biomass and shoot density, respectively). There was, however, no additional increase in ΔY between mixtures of three and six genotypes, i.e., the slopes of the regressions were not significant. On average, six-genotype treatments had 14.9 more shoots (+34%) and 2.59 g of more dry mass (+26%) than that predicted from monocultures.

A separation of genotypic diversity effects into complementarity and selection based on the comparison of diversity-treatments with replicated one-genotype treatments (31) revealed significant complementarity for biomass and shoot




Fig. 2. Effects of increasing genotypic diversity on eelgrass (*Z. marina*) and associated fauna. Shoot number, square-root transformed (A), and dry biomass, \log_{10} transformed (B), in eelgrass (*Z. marina*) plots as functions of their genotypic diversity. (C–F) Comparisons of the abundance of associated invertebrates (abundance \log_{10} transformed). Regression models take spatial heterogeneity (blocking factor) into account. Shaded circles in A and B indicate treatment means. DW, dry weight.

term responses of local communities and their structuring species to increasing climate stress. This applies particularly to sessile species that cannot rapidly shift their distributional range, such as all plants.

Biodiversity experiments should ideally be replicated at multiple levels, including diversity, composition, and multiple sites (48). Our experiment was performed in an area of particularly high genetic and clonal diversity (14), with alternating low and high clonal diversities occurring side by side within the same meadow (Fig. 4). It will be interesting to conduct similar experiments at sites of lower clonal diversities to assess the robustness of our results. In our experiment, we opted for proper replication of a relatively small selection of genotypes to disentangle the contribution of complementarity and selection to overall biodiversity effects. As a consequence, the number of unique genotype combinations was relatively low, increasing the risk that some effects observed were due to specific interactions among genotypes (the composition effect; refs. 48 and 49). Compositional effects were identified in terrestrial grasslands particularly among members of different functional guilds, for example nitrogen fixing and nonfixing plants (49). *A priori*, such effects seem difficult to envisage among a sample of genotypes, given that individuals of the same species are much more alike. Notwithstanding, our data support a net biodiversity effect for the variable shoot density when all putative compositional effects are removed, i.e., when replicates of genotype combinations are averaged in an overyield analysis (paired *t* test, expected vs. observed yield, $n = 7$, $P = 0.019$).

Strong effects of genotypic diversity on fauna associated with eelgrass were surprising because the spacing of the experimental units was relatively close, permitting migration of crustaceans and gastropods between experimental units. In this respect, reported effects of eelgrass genotypic diversity on faunal abundance should be regarded as conservative estimates. Herbivorous gastropods and juvenile bivalves constitute important components of the food web in seagrass ecosystems as they crop epiphytic and planktonic algae, respectively, and may enhance the persistence of seagrass beds even in the face of eutrophication (50, 51). Larger, mobile consumers such as fish and crabs also use seagrass habitat as feeding and breeding habitat (12). Hence, the mediation of herbivore abundance by genotypic

diversity of a primary producer may constitute a positive feedback loop (52) that links consumer dynamics with resilience and persistence of ecologically important seagrass beds (51). The epifaunal species enhanced in the experiment are consumed by a number of fish species several of which are commercially important (see Table 2, which is published as supporting information on the PNAS web site). We speculate that genotypic diversity of seagrass could have far-reaching effects across several trophic levels, including grazers, their predators, and human fisheries.

In conclusion, our study shows that climatic extremes can have immediate effects on coastal communities, and that genetic diversity may enhance recovery after such perturbations. Although we focus here on short-term processes within one generation of a long-lived clonal plant, evolutionary responses of seagrass populations are likely given the striking differences in clonal performance. Ongoing losses of genetic diversity *sensu lato* through habitat fragmentation (53) or genetic erosion (54) may not only endanger the evolutionary potential of populations (55). In addition, they may have immediate consequences for associated communities, ecosystem functioning, and ecosystem services provided by primary producers. These results have implications for biodiversity conservation and environmental management, which so far has focused on the maintenance of species diversity in particularly species-rich ecosystems (56). Our results suggest that conserving genetic and genotypic diversity in species-poor ecosystems that have no redundancy at the species level may be just as important for strengthening the resilience of dependent communities in the face of global change and increasing climatic extremes.

We thank W. Lampert for continual encouragement and support; A. Bockelmann, M. V. Ruggiero, S. Carstensen, S. Liedtke, D. Albrecht, T. Sonntag, R. Neuhaus, M. Dahl, G. Corno, I. Dankert, and A. Hasselmeyer for technical or field assistance; F. Nevoigt and N. Langhanki for meteorological data; H. Hillebrand for statistical advice; H. Lotze, A. Bockelmann, and J. L. Olsen for comments; and the Institute of Marine Geology (GEOMAR) for logistical support. T.B.H.R. and B.W. were supported by individual grants from the Deutsche Forschungsgemeinschaft (Grants Re 1108/3 and -4 and Wo 818/1-2).

- Wardle, D. A. (2002) *Communities and Ecosystems: Linking the Aboveground and Belowground Components* (Princeton Univ. Press, Princeton).
- Tilman, D. (1999) *Ecology* **80**, 1455–1474.
- Kinzig, A. P., Pacala, S. W. & Tilman, D. (2002) *The Functional Consequences of Biodiversity* (Princeton Univ. Press, Princeton).
- Loreau, M., Naeem, S. & Inchausti, P. (2002) *Biodiversity and Ecosystem Functioning: Synthesis and Perspective* (Oxford Univ. Press, Oxford).
- Hector, A., Schmid, B., Beierkuhnlein, C., Caldeira, M. C., Diemer, M., Dimitrakopoulos, P. G., Finn, J. A., Freitas, H., Giller, P. S., Good, J., *et al.* (1999) *Science* **286**, 1123–1127.
- Engelhardt, K. A. M. & Ritchie, M. E. (2001) *Nature* **411**, 687–689.
- Tilman, D., Wedin, D. & Knops, J. (1996) *Nature* **379**, 718–720.
- Loreau, M., Naeem, S., Inchausti, P., Bengtsson, J., Grime, J. P., Hector, A., Hooper, D. U., Huston, M. A., Raffaelli, D., Schmid, B., *et al.* (2001) *Science* **294**, 804–808.
- Hemminga, M. A. & Duarte, C. M. (2000) *Seagrass Ecology* (Cambridge Univ. Press, Cambridge, U.K.).
- Duarte, C. M. (2002) *Environ. Conserv.* **29**, 192–206.
- Williams, S. L. & Heck, K., Jr. (2001) in *Marine Community Ecology*, eds. Bertness, M. D., Gaines, S. D. & Hay, M. E. (Sinaur, Sunderland, MA), pp. 317–338.
- Micheli, F. & Peterson, C. H. (1999) *Conserv. Biol.* **13**, 869–881.
- Reusch, T. B. H., Stam, W. T. & Olsen, J. L. (2000) *Mol. Ecol.* **9**, 127–140.
- Olsen, J. L., Stam, W. T., Coyer, J. A., Reusch, T. B. H., Billingham, M., Boström, C., Calvert, E., Christie, H., Granger, S., La Lumiere, R., *et al.* (2004) *Mol. Ecol.* **13**, 1923–1941.
- den Hartog, C. (1970) *Verh. K. Ned. Akad. Wet. Afd. Natuurkd. II* **59**, 1–275.
- Luterbacher, J., Dietrich, D., Xoplaki, E., Grosjean, M. & Wanner, H. (2004) *Science* **303**, 1499–1503.
- Schär, C., Vidale, P. L., Lüthi, D., Frei, C., Häberli, C., Liniger, M. A. & Appenzeller, C. (2004) *Nature* **427**, 332–335.
- Palmer, T. N. & Räsänen, J. (2002) *Nature* **415**, 512–514.
- Hughes, L. (2000) *Trends Ecol. Evol.* **15**, 56–61.
- Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J., Collingham, Y. C., Erasmus, B. F. N., de Siqueira, M. F., Graininger, A., Hannah, L., *et al.* (2004) *Nature* **427**, 145–148.
- Parmesan, C., Ryrholm, N., Stefanescu, C., Hill, J. K., Thomas, C. D., Descimon, H., Huntely, B., Kaila, L., Kullberg, J., Tammaru, T., *et al.* (1999) *Nature* **399**, 579–583.
- Pimm, S. L. (1984) *Nature* **307**, 321–326.
- Tomlinson, P. B. (1974) *Aquaculture* **4**, 107–130.
- Hämmerli, A. & Reusch, T. B. H. (2003) *Mol. Ecol.* **12**, 619–629.
- Widen, B., Cronberg, N. & Widen, M. (1994) *Folia Geobot. Phytotaxon.* **29**, 245–263.
- Williams, S. L. (2001) *Ecol. Appl.* **11**, 1472–1488.
- Reusch, T. B. H. (2000) *Mol. Ecol.* **9**, 371–373.
- Reusch, T. B. H. (2002) *Limnol. Oceanogr.* **47**, 78–85.
- Parks, J. C. & Werth, C. R. (1993) *Am. J. Bot.* **80**, 537–544.
- Worm, B. & Reusch, T. B. H. (2000) *Mar. Ecol. Prog. Ser.* **200**, 159–166.
- Loreau, M. & Hector, A. (2001) *Nature* **412**, 72–76.
- Rasmussen, E. (1973) *Ophelia* **11**, 1–507.
- Greve, T. M., Borum, J. & Pedersen, O. (2003) *Limnol. Oceanogr.* **48**, 210–216.
- Reusch, T. B. H. & Williams, S. L. (1998) *Oecologia* **113**, 428–441.
- Hughes, A. R. & Stachowicz, J. J. (2004) *Proc. Natl. Acad. Sci. USA* **101**, 8998–9002.
- Huston, M. A. & McBride, A. C. (2002) in *Biodiversity and Ecosystem Functioning*, eds. Loreau, M., Naeem, S. & Inchausti, P. (Oxford Univ. Press, New York), pp. 47–60.

37. Mulder, C. P. H., Uliassi, D. D. & Doak, D. F. (2001) *Proc. Natl. Acad. Sci. USA* **98**, 6704–6708.
38. Cardinale, B. J., Palmer, M. A. & Collins, S. L. (2002) *Nature* **415**, 426–429.
39. Huston, M. A. (1997) *Oecologia* **110**, 449–460.
40. Chambers, R. M., Meyerson, L. A. & Saltonstall, K. (1999) *Aquat. Bot.* **64**, 261–273.
41. Paine, R. T. (2002) *Science* **296**, 736–739.
42. Giller, P. S., Hillebrand, H., Berninger, U. G., Gessner, M. O., Hawkins, S., Inchausti, P., Inglis, C., Leslie, H., Malmqvist, B., Monaghan, M. T., *et al.* (2004) *Oikos* **104**, 423–436.
43. Kennedy, T. A., Naeem, S., Howe, K. M., Knops, J. M. H., Tilman, D. & Reich, P. (2002) *Nature* **417**, 636–638.
44. Gaines, S. D. & Denny, M. W. (1993) *Ecology* **74**, 1677–1692.
45. Holmgren, M., Scheffer, M., Ezcurra, E., Gutierrez, J. R. & Mohren, G. M. J. (2001) *Trends Ecol. Evol.* **16**, 89–94.
46. Barry, J. P., Baxter, C. H., Sagarin, R. D. & Gilman, S. E. (1995) *Science* **267**, 672–675.
47. Etterson, J. R. & Shaw, R. G. (2001) *Science* **294**, 151–154.
48. Schmid, B., Hector, A., Huston, M. A., Inchausti, P., Nijs, I., Leadley, P. W. & Tilman, D. (2002) in *Biodiversity and Ecosystem Functioning: Synthesis and Perspective*, eds. Loreau, M., Naeem, S. & Inchausti, P. (Oxford Univ. Press, New York), pp. 61–78.
49. Tilman, D., Knops, J., Wedin, D., Reich, P., Ritchie, M. & Siemann, E. (1997) *Science* **277**, 1300–1302.
50. Neckles, H. A., Wetzel, R. L. & Orth, R. J. (1993) *Oecologia* **93**, 285–295.
51. Duffy, J. E., Richardson, J. P. & Canuel, E. A. (2003) *Ecol. Lett.* **6**, 637–645.
52. Worm, B. & Duffy, J. E. (2003) *Trends Ecol. Evol.* **18**, 628–632.
53. Olesen, J. M. & Jain, S. K. (1994) in *Conservation Genetics*, eds. Loeschke, V., Tomiuk, J. & Jain, S. K. (Birkhäuser, Basel), pp. 417–426.
54. Avise, J. C. & Hamrick, J. L. (1996) *Conservation Genetics: Case Histories from Nature* (Chapman & Hall, New York).
55. Myers, N. & Knoll, A. H. (2001) *Proc. Natl. Acad. Sci. USA* **98**, 5389–5392.
56. Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B. & Kent, J. (2001) *Nature* **403**, 853–858.