

EDITED SYNOPTIC CLOUD REPORTS  
FROM SHIPS AND LAND STATIONS OVER THE GLOBE, 1982-1991

Carole J. Hahn  
Department of Atmospheric Sciences  
University of Arizona  
Tucson, AZ 85721

Stephen G. Warren  
Department of Atmospheric Sciences  
University of Washington  
Seattle, WA 98195

Julius London  
Department of Astrophysical, Planetary and  
Atmospheric Sciences  
University of Colorado  
Boulder, CO 80309

Environmental Sciences Division  
Publication No. 4367

Date Published: February 1996

Printed for the  
Global Change Research Program  
Environmental Sciences Division  
Office of Health and Environmental Research  
U.S. Department of Energy  
(KP 05 02 00 0)

Printed by the  
Carbon Dioxide Information Analysis Center  
World Data Center-A for Atmospheric Trace Gases  
OAK RIDGE NATIONAL LABORATORY  
Oak Ridge, Tennessee 37831-6335  
managed by  
LOCKHEED MARTIN ENERGY RESEARCH CORP.  
for the  
U.S. DEPARTMENT OF ENERGY  
under contract DE-AC05-96OR22464

**MASTER**

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

10121

## TABLE OF CONTENTS

LIST OF TABLES .....	5
LIST OF FIGURES .....	6
ABSTRACT .....	7
1. INTRODUCTION .....	9
2. DATA SOURCES .....	10
3. PROCESSING OF WEATHER REPORTS .....	10
A. Cloud Information in Synoptic Reports and the "Extended" Cloud Code .....	10
B. Processing Through the Total Cloud Stage .....	11
1) <i>Determination of Cloudiness at Night.</i> .....	12
C. Consistency Checks for Cloud Types and the Change Code .....	13
D. The Amounts of Upper Level Clouds .....	14
4. THE EDITED CLOUD REPORT AND THE DATA ARCHIVE .....	15
A. Contents and Format of the ECR .....	15
B. Organization of the Archive .....	16
5. COUNT SUMMARIES .....	16
A. Distribution of Reports over the 8 Synoptic Hours .....	16
B. Distribution of Code Values .....	16
C. Cases of Sky-obscured and Nimbostratus Cloud .....	17
D. Distribution of Reports over the Globe .....	18

6. COMMENTS ON USE OF THE DATA .....	18
A. Biases .....	18
1) <i>The Night-detection Bias and the Day-night Sampling Bias.</i> .....	18
2) <i>The Clear-sky Bias.</i> .....	18
3) <i>The Sky-obscured Bias</i> .....	19
B. Computing the Average Cloud Amounts and Frequencies .....	20
1) <i>Total Cloud Cover</i> .....	20
2) <i>Low Cloud Types</i> .....	20
3) <i>Upper Level Clouds</i> .....	21
7. HOW TO OBTAIN THE DATA .....	22
ACKNOWLEDGEMENTS .....	22
REFERENCES .....	22
TABLES .....	24
FIGURES .....	33
APPENDIX A. Percent Frequency of Occurrence of Extended Code Values for Cloud Variables in Edited Cloud Reports (Light). .....	44
APPENDIX B. Glossary of Terms and Abbreviations Used .....	45

## LIST OF TABLES

<u>Table</u>		<u>Page</u>
1.	Cloud Information Contained in Synoptic Weather Reports .....	24
2.	Cloud and Weather Type Definitions Used .....	25
3.	FORTTRAN Code for Checking Cloud Type Consistencies .....	26
4.	Change Codes for Edited Cloud Reports .....	27
5.	Number of Reports with Cloud Type Information .....	27
6.	FORTTRAN Code for Determining Upper Level Cloud Amounts .....	28
7.	Random Overlap Computation Table .....	29
8.	Number of Reports in which Upper Level Cloud Amounts were Computable .....	29
9.	Contents and Format of the 56-character Edited Cloud Report .....	30
10.	Sample Edited Cloud Reports in 56-character Format .....	31
11.	Distribution of Reports over the Synoptic Reporting Times .....	32
12.	Contribution of the Various Paths to Total Nimbostratus Frequency .....	32

## LIST OF FIGURES


<u>Figure</u>	<u>Page</u>
1.	Flow chart of report processing through the total cloud stage ..... 33
2a.	Frequency distribution of extended code values for indicated cloud variables in edited light reports from land stations over the globe for 1982-1991 ..... 34
2b.	Frequency distribution of extended code values for indicated cloud variables in edited light reports from ships over the globe for 1982-1991 ..... 35
3a.	Global distribution of log (number) of light reports for cloud types for 1982-1991 land data ..... 36
3b.	Global distribution of log (number) of light reports for cloud types for 1982-1991 ship data ... 37
4a.	Global distribution of occurrence of CL=/ or Nh=/ with N>0 (fb) in light reports for 1982-1991 land data (%) ..... 38
4b.	Global distribution of occurrence of CL=/ or Nh=/ with N>0 (fb) in light reports for 1982-1991 ship data (%) ..... 39
5a.	Global distribution of the clear-sky adjustment factor (AF0) in light reports for 1982-1991 land data ..... 40
5b.	Global distribution of the clear-sky adjustment factor (AF0) in light reports for 1982-1991 ship data ..... 41
6a.	Global distribution of occurrence of N=9 (f9, %) in light type reports for 1982-1991 land data ..... 42
6b.	Global distribution of occurrence of N=9 (f9, %) in light type reports for 1982-1991 ship data ..... 43

## ABSTRACT

Hahn, C.J., S.G. Warren, and J. London, 1996: *Edited Synoptic Cloud Reports from Ships and Land Stations Over the Globe, 1982-1991*. NDP026B, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, TN. (Also available from Data Support Section, National Center for Atmospheric Research, Boulder, CO.)

Surface synoptic weather reports for the entire globe for the 10-year period from December 1981 through November 1991 have been processed, edited, and rewritten to provide a data set designed for use in cloud analyses. The information in these reports relating to clouds, including the present weather information, was extracted and put through a series of quality control checks. Reports not meeting certain quality control standards were rejected, as were reports from buoys and automatic weather stations. Correctable inconsistencies within reports were edited for consistency, so that the "edited cloud report" can be used for cloud analysis without further quality checking. Cases of "sky obscured" were interpreted by reference to the present weather code as to whether they indicated fog, rain or snow and were given appropriate cloud type designations. Nimbostratus clouds, which are not specifically coded for in the standard synoptic code, were also given a special designation. Changes made to an original report are indicated in the edited report so that the original report can be reconstructed if desired. While low cloud amount is normally given directly in the synoptic report, the edited cloud report also includes the amounts, either directly reported or inferred, of middle and high clouds, both the non-overlapped amounts and the "actual" amounts (which may be overlapped). Since illumination from the moon is important for the adequate detection of clouds at night, both the relative lunar illuminance and the solar altitude are given, as well as a parameter that indicates whether our recommended illuminance criterion was satisfied.

This data set contains 124 million reports from land stations and 15 million reports from ships. Each report is 56 characters in length. The archive consists of 240 files, one file for each month of data for land and ocean separately. With this data set a user can develop a climatology for any particular cloud type or group of types, for any geographical region and any spatial and temporal resolution desired.





## 1. INTRODUCTION

Surface synoptic weather reports are readily available in data sets such as those prepared by the National Meteorological Center (NMC) and the U. S. Navy Fleet Numerical Oceanography Center (FNOC). For marine reports there is also the Comprehensive Ocean-Atmosphere Data Set (COADS). Those data sets are archived at the National Climatic Data Center (NCDC) in Asheville, North Carolina, and at the National Center for Atmospheric Research (NCAR) in Boulder, Colorado. The data set described here includes only the information from the synoptic weather report that directly relates to clouds, as well as some derived quantities that aid in cloud analysis. It was developed as an intermediate stage in our own global analysis of total cloud cover (Hahn et al., 1994a) and cloud types, but should be useful to other researchers who wish to compare individual surface cloud reports to concurrent satellite-derived cloud data for example, or who wish to obtain averages over time or space scales not already provided in existing archives.

The cloud report provided in this data set, referred to as the "edited cloud report" (ECR, or ECRA when referring to the archive of all reports), has several features that make it desirable and easy to use in cloud analyses:

1) Synoptic weather reports contain information in addition to clouds, such as air temperature, pressure, winds, humidity, visibility, past weather and, for ships, sea surface temperature and ocean wave parameters. The ECR excludes these data, thus reducing the data volume.

2) Data sets of synoptic weather reports include reports that do not contain cloud information, such as those from automated weather stations on land and buoys in the oceans. These are not included in the ECRA.

3) The cloud portion of the synoptic report occasionally contains obvious errors or inconsistencies which must be checked for to avoid inclusion of detectably erroneous data in an analysis. Quality control procedures developed over years of analyzing surface cloud reports have been applied so that erroneous or inconsistent reports have either been excluded or, if possible, corrected before inclusion in the archive.

4) While the amount of low cloud is directly coded in the synoptic report, the amounts of middle and high clouds are not, but may often be inferred. Where possible for upper level clouds, the ECR includes both the "actual" cloud amount (sometimes requiring use of the random-overlap assumption) and the non-overlapped amount, which is simply that portion of the cloud that is visible from below and requires no assumptions.

5) Although all reports that meet the above criteria are included in the ECRA, it has been shown that many of the nighttime reports are made under conditions of insufficient (moon) light for adequate detection of clouds. Use of such reports results in an underestimate of nighttime cloudiness by about 4% globally and has a profound influence on computed diurnal cycles in cloudiness (Hahn et al., 1994b). Reports made under conditions that satisfy the criterion for adequate illumination specified by Hahn et al. (1994b) are flagged in the ECR, and both the relative lunar illuminance and the solar altitude are given for each report.

The edited cloud report archive described here covers the ten-year period December 1981 to November 1991. These beginning and ending months were chosen to coincide with the boundaries of the traditionally-defined seasons December-January-February (DJF), March-April-May (MAM), June-July-August (JJA) and September-October-November (SON). Since December 1981 is considered to be part of DJF 1982, the 10-year period is referred to as 1982-1991. This particular 10-year period was selected to coincide with the International Satellite Cloud Climatology Project (ISCCP; Rossow and Schiffer, 1991) and to extend our previous climatologies (Hahn et al., 1988; Warren et al., 1986 {hereafter referred to as W86}; Warren et al., 1988 {hereafter referred to as W88}), which terminated in (November) 1981.

Non-standard terms used in the following discussions are defined in the glossary of terms and abbreviations in Appendix B.

## 2. DATA SOURCES

The NMC data set (obtained from NCAR) was the source of the synoptic weather reports used for land stations. About 144 million reports (22 gigabytes of data) were processed. Only those stations assigned official station numbers by the World Meteorological Organization (WMO) were used. Synoptic reporting hours are 00, 03, 06, 09, 12, 15, 18, 21 Greenwich Mean Time (GMT). In the NCAR archive the 6-hourly reports (00,06,12,18 GMT) are stored in separate files from the intermediate 3-hourly reports (03,09,15,21 GMT) and we processed them in tandem for each month. Within each of these two groups the reports are sorted by time.

The COADS Interim Product CMR5 Reports (Woodruff et al., 1987) were the source of the ship data used (also obtained from NCAR). About 22 million reports (540 megabytes of data in packed binary format) were processed. Many of these reports are from buoys, however, and do not contain cloud information. For an individual month these data are sorted first by 2-degree latitude x longitude boxes and then by day and hour. COADS merges a dozen or so subsets of ship data and, as such, is the most extensive record of climatic data for the sea surface. However, the Interim Products available at the time of our analysis lack much of the digitized ship logbook data from foreign countries which may be delayed 2 - 5 years (Woodruff et al., 1992). Thus future releases of COADS will contain more data than were utilized here.

## 3. PROCESSING OF WEATHER REPORTS

### A. Cloud Information in Synoptic Reports and the "Extended" Cloud Code

Synoptic weather reports are coded according to the system given by the World Meteorological Organization (WMO, 1988). The information in these reports that relates to cloud analysis is summarized in Table 1. All other parameters were ignored in our processing. A more detailed breakdown of the definitions of the cloud and weather types, as used here, is given in Table 2. The table shows the synoptic codes that correspond to various precipitation types (ww codes) as well as the codes that correspond to the various cloud types defined within each of the three reporting levels: low, middle and high ( $C_L$ ,  $C_M$ ,  $C_H$ ).

We give special consideration to the cloud type nimbostratus (Ns), which is not specifically defined in the synoptic code. Codes  $C_M=2$  or 7 may signify Ns but may also signify As or Ac, respectively. We consider these codes to signify Ns when there is concurrent precipitation in the form of drizzle, rain, or snow as indicated in the present weather code ww (symbolized in Table 2 as D, R and S, respectively). To distinguish Ns from As/Ac we "extend" the synoptic code for  $C_M$  to include the values 12 and 11 to represent these cases of  $C_M=2$  and 7, respectively. The extended code values (shown in Table 2) are entered in the edited cloud report (Section 4) without loss of the information content in the original report.

Nimbostratus is also considered to be present when  $C_M=/$  and specified combinations of precipitation and low cloud types are present (Table 2). These cases are given the extended code  $C_M=10$ . This definition for nimbostratus has been simplified from that used in our previous work (W86, W88). Cases of  $C_M=/$  with low stratus and drizzle are no longer defined to be Ns. This will cause a slight reduction in computed Ns amounts (Section 5C).

Special consideration is also given to the case of N=9 (sky obscured). If ww indicates that the sky was obscured due to F, Ts, or DRS (symbols defined in Table 2), the cloud type is considered to be Fg, Cb, or Ns, respectively, and is given the extended code  $C_L=11$ ,  $C_L=10$ , or  $C_M=10$ , respectively, and the value of N is set to 8 oktas.

All the changes described here are coded in a parameter called "the change code" (Section 3C below) which is also included in the edited cloud report (Section 4), so that unchanged reports could be reconstructed if desired.

## B. Processing Through the Total Cloud Stage

A cloud report may be suitable for total cloud analysis even if cloud type information is incomplete. Certain inconsistencies within the cloud type portion of the report may, however, make the whole report suspect and cause it to be rejected even for total cloud analysis. The processing and quality control checks performed on each weather report read from the original archives (NMC or COADS) and designed to ensure suitability for total cloud analysis are shown in the flow chart in Figure 1. The percentage of reports discarded at each stage of the processing, which is discussed in the following paragraphs, is indicated.

In the early stages of processing, land and ship reports required slightly different checks (upper portion of Figure 1). A land station that did not have a WMO station number was discarded (most of these were from the United States), thus ensuring more uniformity in reporting procedures. A ship report known to be from a buoy (by the "deck" number in the COADS data) was discarded. Any report that had no cloud information (N=/) was discarded.

In 1982 WMO introduced several changes in the coding procedure (WMO, 1988). One of these changes now instructs observers to set ww=/ if present weather was either "not available" or "observed phenomena were not of significance" (ww codes 00-03 are considered to represent phenomena without significance). The present weather indicator,  $I_x$ , is used to distinguish these cases. Land station reports with  $I_x$  values of 4, 5 or 6 signify automatic weather stations and were discarded. Reports with  $I_x=3$  (data not available) were also discarded because without ww it is not possible to interpret cases of N=9 (see W86) or to evaluate the occurrence of precipitation.  $I_x=2$  indicates that observed phenomena were not of significance, while  $I_x$  is coded as "1" when ww is given. Occasionally  $I_x=1$  when ww=/; these inconsistent reports were also discarded.

Examination of the NMC data set showed that while land stations adopted this new coding procedure almost immediately,  $I_x$  was not consistently coded in ship reports until 1985, as ship observers tended to continue reporting ww in accordance with the old rules. The COADS data set does not even contain  $I_x$ . Thus ship reports could not be screened on the basis of  $I_x$ .

At the upper horizontal dashed line in Figure 1, 125 million land reports and 15.8 million ship reports remained. The discard fractions below the line are fractions of these numbers. If the reported latitude and longitude of a land station put the station on water (rare) or if reported latitude and longitude of a ship put the ship on land (0.3%), the report was discarded. A land station was considered to be on water (or a ship on land) if the 5x5 degree latitude x longitude box in which it was reported to reside was 100% ocean (or land) as defined in W86. Exceptions to this are that a number of boxes with small islands were allowed to contribute to the land data and reports from the Great Lakes and the Caspian Sea were allowed to contribute to the ship data.

If the sky was obscured (N=9) by fog (ww=F; 1.1% land, 2.7% ship), thunderstorms-showers (ww=Ts, abbreviated as T in the figure; 0.05% land, 0.1% ship), or drizzle-rain-snow (ww=DRS, abbreviated here

as R; 0.4% land, 0.9% ship), the sky was considered to be overcast ( $N=8$ ). This source of "cloudiness" contributed about 1% to the total cloud cover globally, and much more in some locations and seasons (Hahn et al., 1992). Clouds could not be inferred if the sky was obscured for other reasons, such as blowing dust or snow, and such reports were discarded. The change code,  $IC=1$  (discussed in Section 3C below), signifies that a report came through the  $N=9$  branch of the processing. Thus 1.5% of the land reports and 3.7% of the ship reports had  $N=9$  with the ww codes D, R, S, F or Ts.

Other data consistency checks indicated in Figure 1 ensure that the low cloud amount is not greater than the total cloud cover, that precipitation (as defined in Table 2) is not reported with a clear sky, and that if cloud is present (and types are reported), some cloud type must be indicated in at least one of the three possible levels (this test actually discards a report if  $N>0$  and  $C_L=0$  and  $C_M\leq 0$  and  $C_H\leq 0$ ). The recoding indicated in the lower left box in the figure is necessary because of a 1982 code change (WMO, 1988) that instructs observers to set  $C_L=C_M=C_H=/$  when  $N=0$  (this requires special attention in cloud type analysis and will be discussed in Section 4).

The number of reports that survive these tests and are suitable for total cloud analysis (referred to as "total reports") is 124.2 million for land and 14.7 million for ships. Of these, 90.3 million and 11.1 million, respectively, were made under sufficient solar or lunar illuminance (referred to as "light reports") to meet the established illuminance criterion for adequate cloud visibility (Hahn et al., 1994b).

#### *1) Determination of Cloudiness at Night.*

The ability of surface observers to adequately detect clouds at night has been questioned for many years (e.g. Riehl, 1947; Schneider et al., 1989). In an attempt to find a practical solution to this "night-detection-bias", Hahn et al. (1994b) analyzed ten years of nighttime data for the zone 0-50° N and plotted reported cloud cover as a function of the illumination due to moonlight, which depends on the phase and altitude (angle above or below the horizon) of the moon and on the distance of the moon from the earth. The amount of total cloud reported at night increased as the lunar illuminance increased up to a certain threshold, after which the reported cloud cover leveled off. This threshold is referred to as "the illuminance criterion" and corresponds to the twilight produced by the sun at an altitude of about 9 degrees below the horizon. Thus the illuminance criterion is met when either the sun is at an altitude greater than -9° or the position of the moon is such that its illuminance exceeds the threshold. These conditions were determined for each report with the use of an ephemeris and the latitude, longitude, date and time of the report.

This illuminance criterion was applied in analyses of total cloud cover and clear-sky frequency (Hahn et al., 1994a). (Fog and precipitation frequencies were also analyzed in that study but their detection does not depend on illumination). Application of the illuminance criterion increased the computed global average total cloud cover at night by about 4% and thus increased the daily average computed cloud cover by about 2%. Diurnal cycles of total cloud cover over land, which typically show daytime maxima, were reduced in amplitude when compared to previous studies which did not use the illuminance criterion (W86). Over the oceans, the increased computed nighttime cloud cover was often sufficient to result in nighttime maxima, in contrast to the daytime maxima previously reported (W88). Preliminary surveys conducted in conjunction with the present work suggest that we should expect similarly dramatic effects on analyses of middle and high clouds but little effect on low clouds.

Because of the importance of moonlight in the detection of clouds at night, parameters relating to the illuminance criterion are included in the edited cloud report (Section 4). Reports for which the illuminance criterion is met are referred to as "light reports", as opposed to "dark reports" (for which the criterion is not met) or "all reports" (which includes both light and dark).

### C. Consistency Checks for Cloud Types and the Change Code

The reports that failed the cloud type consistency checks shown in Figure 1 were discarded. Other inconsistencies are possible which may be correctable or may provide cause simply to reject the report for cloud type analysis. As the synoptic reports were processed, any inconsistency encountered required a change to be made in the existing code before the report was entered into the edited cloud report archive. Any changes thus made are noted by assigning a "change code" (IC) to that report. This change code (with values 0 to 9) is preserved in the ECR (Section 4) so that modifications made to the original report can be identified.

Details of the cloud type processing following the total cloud stage shown in Figure 1 are presented in the form of the FORTRAN code in Table 3. Each segment of the table (delineated by a change code heading) describes the processing of a particular type of inconsistency or change. The changes associated with particular coded cases are described briefly in Table 4 along with the frequency of occurrence of each change type. (The changes referred to by IC=1 were discussed in section 3B.) Most of the inconsistencies under consideration have been discussed previously (W86, W88) but are summarized here.

For a report to provide useful cloud type information, both  $N_h$  and  $C_L$  must be given. If either is missing (and not correctable) then the report cannot be used for cloud type analysis and all cloud type variables are set to -1 for consistency (see segment IC\_5 in Table 3). Thus if any inconsistency is discovered that cannot be corrected, simply setting  $N_h=-1$  will result in exclusion of the report from cloud type analyses. For example, the IC\_2,4 segment in Table 3 examines the case in which there is middle cloud but no low cloud. In such a case  $N_h$  should give the middle cloud amount. In some reports  $N_h$  was improperly set to 0 (W86). If there is also no high cloud then it must be that  $N_h=N$  and the report can be corrected (IC=2). But if high cloud is present, then the value of  $N_h$  is indeterminate, and set to -1 (IC=4), and the report cannot be used for cloud types (in such cases all cloud type variables are set to -1 in segment IC\_5).

The situation is similar in segment IC\_3,4. If only high cloud is present,  $N_h$  should properly be 0 but is sometimes given the value of  $N$  by an observer. This is readily corrected (IC=3). But if  $N_h < N$  in such a case, the report is irreconcilably inconsistent and must be rejected for cloud type analysis (set  $N_h=-1$  and IC=4).

Segments IC\_6 and IC\_9 simply correct cases of sloppy reporting of middle and high clouds, such as reporting  $C_M=0$  rather than  $C_M=/$  when the sky is overcast with low cloud.

Segments IC\_7 and IC\_8 merely set  $C_M$  to represent our definition of  $N_s$  as a convenience for future cloud type analyses. As is usually the case, the original report can be reconstructed if desired.

The order in which some of these consistency checks is performed is also important to the outcome. For example, if segment IC\_8 were performed before segment IC\_6, then some cases of  $N_s$  would go undetected. Also, segment IC\_9 must be performed after IC\_8 for the same reason. However since it is not desirable to have the change code IC=8 overwritten by the relatively trivial change IC=9, this latter code is only entered in IC (although the change is made) if no previous change has been made (second part of segment IC\_9 in Table 3). This co-occurrence of change codes should be rare since  $N$  is usually large when  $N_s$  is present.

From Table 4 we see that some change is made in 12% of land reports and 16% of ship reports, but that roughly half of the changes just represent classifying an observation as  $N_s$ , Cb or fog. Thus less than 5% of land reports and less than 9% of ship reports have been changed due to inconsistencies and most of these are due to the relatively trivial cases with IC=6 (and IC=9 for ships).

After passing the cloud type consistency checks, the number of light reports available for cloud type analysis for 1982-91 is 88 million for land and 9.4 million for the ocean (Table 5). Reports suitable for cloud type analysis ( $C_L$  and  $N_h > -1$ ) are referred to as "type reports".

#### D. The Amounts of Upper Level Clouds

The synoptic code contains two cloud amount variables,  $N$  and  $N_h$ . The amount of low cloud, if present, is directly specified by  $N_h$ . While the amounts of upper level clouds are not directly specified, they may often be inferred. Thus when  $C_L=0$ , the amount of middle cloud is given by  $N_h$ , and when  $C_L=C_M=0$ , the amount of high cloud is given by  $N$ . If all three levels are present there are too few known variables to determine the upper cloud amounts. If two levels are present, the amount in the upper level may be estimated if the extent of overlap is assumed.

In the ECR we provide amounts that utilize the random overlap assumption, where necessary, in order to best represent the actual cloud amounts (the fraction of the sky covered by a cloud type, whether visible or not). We also provide the non-overlapped amounts which require no assumptions but which indicate only the amount of the upper level cloud visible from below. (Satellite-derived cloud amounts are typically given as the non-overlapped amounts visible from above.) Tian and Curry (1989) tested the minimum, maximum and random overlap assumptions and found the maximum overlap assumption to be best for adjacent cloud layers, while the random overlap assumption was best for vertically separated layers.

Table 6 gives our method, in the form of FORTRAN code, for determining the actual and non-overlapped amounts of middle and high clouds from a synoptic weather report. A few points should be noted. The random overlap equation (lines 17 and 38) is invoked only when  $N_h < 7$ . Table 7, which gives the outcomes of the possible combinations of  $N$  and  $N_h$  in the equation, shows that only 2 outcomes are possible for the upper cloud amount when  $N_h=7$ , namely 0 and 8 oktas, making this a highly inaccurate determination (W86). In such cases the upper cloud amount is left undetermined. If the upper cloud in question is  $N_s$ , however, the maximum overlap assumption is employed and the  $N_s$  amount is assigned the value of  $N$  (lines 13-14, Table 6). In this case the nimbostratus cloud layer is likely to be adjacent to or continuous with the low cloud, so the maximum overlap assumption is more appropriate (Tian and Curry, 1989). Also, certain arbitrary decisions are sometimes required, such as our choice, in line 7 of Table 6, to allow middle cloud to be computed when  $C_H \neq /$ . This choice is justifiable since such a case tends to occur with large  $N$  so that any error induced by this situation would be small.

The number of times reports were processed through the possible paths in Table 6 are listed in Table 8. Light reports (for which the illuminance criterion was met) and dark reports (for which the illuminance criterion was not met) are both shown, where possible, for comparison. Land and ocean data were processed separately. Non-overlapped (NOL) amounts were computable in more than 90% of the cloud type reports since one can know that a cloud cannot be seen even if one does not know whether it is present. Thus the non-overlapped amount of an upper level cloud is frequently zero.

Percentages are not explicitly shown in the table but it can be seen that upper level clouds are reported more frequently in the set of light reports than in the set of dark reports (38% and 30%, respectively, for land middle clouds, and 44% and 29%, respectively, for ocean middle clouds, for example). When upper clouds are present, they are more frequently computable within the set of dark reports and random overlap (ROL) is less often required. Comparing land and ocean, upper level clouds, when observed to be present, are less likely to be computable from ocean data and are more likely to require ROL because of the predominance of low level clouds over the oceans. (The percentages given here merely represent the fractions of reports within the data set and are not area-weighted global averages.)

## 4. THE EDITED CLOUD REPORT AND THE DATA ARCHIVE

### A. Contents and Format of the ECR

Table 9 shows the variables included in the ECR, the number of characters allocated for each, and the maximum and minimum values allowed. Each item in the table is discussed briefly below. Sample reports selected from ship and land data for 1981 December and 1982 January are provided in Table 10. These reports are in the order in which they appear in their respective files (see next section) though these selections are not consecutive within the file. The reports are numbered in the table for convenience.

Item 1: The first item in the report gives the year, month, day and GMT hour of the report, with 2 characters allotted for each. There are no spaces ("3", for example, is given as "03") so that the entire item can be read as a single integer. Only the last 2 digits of the year (1900's) are given. Months are coded as 01 through 12, representing January through December.

Item 2: The IB variable indicates whether the illuminance criterion of Hahn et al. (1994b) was satisfied (IB=1) at the time and place of the report or not (IB=0). This variable can be checked in lieu of SA and RI (items 19 and 20 below) if one accepts the criterion specified in Hahn et al. (1994b).

Item 3: The latitude (in degrees north and south) is given to 2 decimal places and written as a 5-digit integer, and thus must be divided by 100 to obtain actual latitude. Actual values range from +90 to -90 for 90N to 90S, respectively.

Item 4: The longitude (in degrees east) is given to 2 decimal places and written as a 5-digit integer, and thus must be divided by 100 to obtain actual longitude. Actual values range from 0 to 360E.

Item 5: For land stations, ID is the WMO station number (WMO, 1977). For ships, ID is the card deck assignment (Slutz et al., 1985).

Item 6: This parameter indicates whether a report is from a land station (LO=1) or a ship (LO=2).

Items 7-13: These weather and cloud variables are coded as specified by WMO (1988) except that items 11 and 12 have been "extended" as described in Section 3A (Table 2). Also, cases of N=9 (item 8) that were not discarded have been converted to N=8. Any such conversion is coded in the "change code" (item 18 below). The value "-1" indicates missing data. Item 8 (N) does not obtain a value of -1 in this data set since all such reports were discarded during processing. Item 10 (h) may have a value of 9 only when a cloud is present since h was set to -1 in cases of N=0 (Figure 1).

Items 14-15: These variables give the "actual" cloud amounts of middle and high clouds and utilize the random overlap equation where necessary (Section 3D). Values are given in oktas to 2 decimal places and written as 3-digit integers, so they must be divided by 100 to obtain the actual values. An actual value of 9 (coded value 900) indicates missing data.

Items 16-17: These variables give the non-overlapped cloud amounts of middle and high clouds and represent the amount of cloud visible from below (Section 3D). Values are given in oktas. A value of 9 indicates missing data.

Item 18: The change code indicates whether a change was made to the original report during processing. Code values are defined briefly in Table 4 and in detail in Figure 1 and Table 3 (Section 3C). A change code of 0 means that no change was made other than the trivial change of converting /s to 0's in the case of N=0. Examples of reports with each change code are provided in Table 10.

Items 19-20: These variables give the solar and lunar parameters needed to determine the illuminance provided by the sun or moon for the date, time and location of the report (Section 3B1). SA is the altitude of the sun above the horizon, given to a tenth of a degree (divide the coded value by 10 to obtain the actual value). RI is the lunar relative illuminance defined by Hahn et al., (1994b).  $RI = \Phi \sin(A) (R^2/r^2)$ , where A is the lunar altitude, r is the earth-moon distance, R is the mean earth-moon distance, and  $\Phi$  is the lunar phase function which varies from 0 to 1 in a concave shape such that a full moon is 10 times as bright as a half moon (Hapke, 1971). The illuminance criterion of Hahn et al. (1994b) is met (IB=1, item 2) when  $SA \geq -9^\circ$  or  $RI > 0.11$ . (A negative value of RI means the moon was below the horizon.)

## **B. Organization of the Archive**

The data are divided into 240 files, one for each month for ten years for land and ocean separately. These files range in size from 5.0 to 8.4 MB for the ocean data and from 49.8 to 63.5 MB for the land data.

In the NMC data set archived at NCAR, the 6-hourly reports and 3-hourly reports (Section 2) are stored on separate files. Each of these subsets is sorted first by time and then by latitude and longitude. Each land data file of the ECRA was formed by merging the 6-hourly and the 3-hourly data so that for each month the reports appear in time order.

The sort used in the COADS Interim Product (Section 2) was first by 2-degree box, then by time, and finally by latitude and longitude for each month. For the ECRA files, the reports were re-sorted by time and then by latitude and longitude.

## **5. COUNT SUMMARIES**

### **A. Distribution of Reports over the 8 Synoptic Hours**

Figure 1 and Tables 4, 5 and 8 showed the number of reports processed, deleted and changed, as well as the number of light reports, the number suitable for cloud type analysis, and the number of times upper level cloud amounts were computable. Table 11 shows how the reports are distributed over the synoptic reporting times. Land stations usually report 8 times per day but some do not (notably in the United States and Australia), so that 59% of all reports are made during the 6-hourly times. Ships, however, tend to report at the 6-hourly times and only about 10% of the ship reports are for the intermediate 3-hourly times. Having only 4 reports per day, rather than 8, limits the resolution possible in computations of the phase and amplitude of the diurnal cycle. It was also noted (W88) that regional averages formed from 6-hourly ship data may be uniformly different from averages formed from 3-hourly data, consistent with a tendency for some ships to give a 3-hourly report only in unusually stormy weather. A bias is also possible when averaging over a land grid box that has more than one station if stations within one climatic region report 8 times per day while stations within a different climatic region report 4 times per day.

### **B. Distribution of Code Values**

The histograms in Figures 2a for land and 2b for ocean show the frequency of occurrence of the extended code values for the six cloud variables for light reports in the archive of edited cloud reports. [In these figures N=9 is shown separate from N=8 although N=9 appears as N=8 (with IC=1) in the ECR.] The shaded areas show the occurrence of precipitation (DRSTs, Table 2). Numerical values for the data shown in these figures are provided in Appendix A. Several interesting features are evident in these figures. The distribution of codes for total cloud cover N is nearly U-shaped for land but strongly skewed towards the higher amounts for oceans. 96.5% of all precipitation occurs with  $N \geq 7$  over land and with  $N \geq 6$  over oceans. About 75% of precipitation occurs with  $N_h \geq 6$ . The reports with  $N_h = -1$  are not usable.



for cloud type analysis. The most commonly occurring cloud base height code is  $h=5$  (600-1000m) over land and  $h=4$  (300-600m) over oceans. The high frequency of  $h=9$  over land is a consequence of the high occurrence of  $C_L=0$  so that  $h=9$  often refers to the middle cloud level.

The lower panels in Figures 2a and 2b show the occurrences of various cloud types within the three reporting levels. The occurrence of  $C_L=-1$  is the same as  $Nh=-1$  due to the processing procedure and is the fraction of reports that do not contain information relating to cloud types (2.6% for land data, 15.3% for ocean data). A larger fraction of the reports have  $C_M=-1$  (17.7% for land, 36.0% for ocean) and  $C_H=-1$  (33.2% for land, 50.9% for ocean) because of lower overcast. Thus 97.4% of the land reports have information about cloud types but only 82% of those have information about the middle cloud level and 66% about high clouds. For the oceans, 84.7% of the reports have low cloud information but only 57% of those have middle cloud information and 40% have high cloud information.

The low cloud type most commonly reported over land is stratocumulus ( $C_L=5$ ). While this type is also relatively common over the oceans, it is exceeded by the cumulus types  $C_L=1$  and 2. About 25% of all precipitation occurs with the stratus cloud  $C_L=7$ . When  $C_L=7$  is reported over land, precipitation is present in 67% of the reports. Precipitation occurs in 34% of the ship reports of  $C_L=7$ .

While 58% (land) and 46% (ocean) of all precipitation occurs with the middle clouds defined to be nimbostratus ( $C_M=10,11,12$ ), 25% and 38%, respectively, of precipitation occurs when the middle cloud level is not given in the report (typically because of low overcast). Because of our definition of  $Ns$  shown in Table 2, most of these latter cases must have  $ww=D$  or  $Ts$ . In the high cloud level, 90% of all precipitation occurs in reports with  $C_H=-1$  (high cloud level not reported, usually because of lower overcast).

### C. Cases of Sky-obscured and Nimbostratus Cloud

The occurrence of reports of sky-obscured ( $N=9$ ) due to fog or precipitation is 1.5% for land and 3.5% for ocean, with fog ( $C_L=11$ ) accounting for more than two thirds of these values for both land and ocean (Figure 2 and Appendix A). These cases of sky-obscured due to fog make up 14% (land) and 48% (ocean) of all reported cases (light) of fog. Cases of thunderstorms or showers ( $C_L=10$ ) account for only about 3.5% of the reports of sky obscured, and sky-obscured due to thunderstorms or showers make up only 1.6% (land) and 5.8% (ocean) of the light reports of thunderstorms and showers. The remaining contribution to the reports of sky obscured (about one quarter) is due to drizzle, rain or snow. Sky-obscured due to drizzle, rain or snow make up 5.1% (land) and 12.9% (ocean) of the light reports of drizzle, rain and snow.

A report of sky-obscured due to drizzle, rain or snow is considered to indicate nimbostratus cloud and is given the extended code value  $C_M=10$  with  $IC=1$  (Tables 2 and 4). Two other sets of circumstances are considered to indicate  $Ns$  as well. Table 12 shows the contributions of the three major paths to the frequency of  $Ns$  within the light type reports. (Frequencies based on light type reports are slightly higher than the frequencies quoted in the last paragraph which were based on the total set of light reports.) The largest contributor to  $Ns$  in the land data is the path through  $C_M=2,7$  (with  $ww=DRS$ ). For both land and ocean  $C_M=2$  is far more important than  $C_M=7$  (see  $C_M=11,12$  in Appendix A). The largest contributor to  $Ns$  in the ocean data comes through the  $C_M=/$  path, which has several contributors itself, the largest being the case of  $C_L=7$  with  $DRS$ .

Excluded from the definition of  $Ns$  are the cases of  $C_M=/$  with  $C_L=4,5,6,8$  and  $ww=D$  (more than half of these cases are due to  $C_L=6$  alone). These cases were considered to indicate  $Ns$  in our previous climatologies (W86, W88) but after subsequent consideration and discussions with colleagues we concluded that since drizzle could occur from these low cloud types, the additional inference of  $Ns$  above

them was inappropriate. Thus the frequencies of occurrence of Ns computed under the current definition will be reduced to about 97% (land) and 90% (ocean) of the frequencies given in W86 and W88.

Another change associated with the simplification of our previous definition of Ns involves cases of  $C_L=6,7$  with DRS and  $C_M \neq$  other than 2,7,/. The  $C_L$  6,7 in these cases were previously reassigned as Ns, but are left unchanged in the present, simplified definition. This results in a further reduction in computed Ns frequency by factors comparable to those in the last paragraph. Thus the Ns frequencies computed under the present definition may be about 94% (land) and 81% (ocean) of those computed under the previous definition. Note that these percentages refer to the number of reports in the data set which contains a disproportionate contribution of reports from the densely populated northern mid-latitudes and thus do not represent the area-weighted global averages. Note also that the user of this data set is not restricted to the definitions assigned here since all the information necessary for any other interpretation is contained in the edited cloud reports. The definitions discussed above apply only if the reports are used exactly as written.

The cases of  $C_L=1,2,3,9$  with  $C_M \neq$  and  $ww=DRS$  are not considered to indicate nimbostratus cloud.

#### **D. Distribution of Reports over the Globe**

To show the global distribution of the reports, numbers (shown as  $\log_{10}$ ) of light type reports are displayed on a 5c grid (see Glossary in Appendix B) in Figures 3a (land) and 3b (ocean). Numbers from 1-9 appear as 0, numbers from 10-99 appear as 1, etc. Grid boxes with no light type reports are blank.

### **6. COMMENTS ON USE OF THE DATA**

#### **A. Biases**

A number of biases which can affect analyses of clouds from surface observations are summarized in W88. Four biases which we can measure and possibly correct for are described in more detail here.

##### *1) The Night-detection Bias and the Day-night Sampling Bias*

The night-detection bias is largely eliminated by using only data for which the illuminance criterion is met (Section 3B1). This, however, enhances the day-night sampling bias unless precautions are taken since less than half as many nighttime observations will be available compared to the number of daytime observations. Thus Hahn et al. (1994a) prepared nighttime and daytime averages separately and averaged the two together to obtain the average cloud cover.

##### *2) The Clear-sky Bias*

Another potential bias for cloud type analyses was introduced by the synoptic code change in 1982 which allowed observers to record a "/" for cloud types when  $N=0$ . Previously a report with  $N=0$  and  $C_L$  or  $Nh \neq$  would indicate a station that never reports cloud types and the report would be omitted from the cloud type analysis. Now every occurrence of  $N=0$  must be treated as  $C_L=0$  and  $Nh=0$ . Thus a report from a station or ship which never reports cloud types would contribute to a cloud type analysis only when the sky was clear, producing a clear-sky bias: the frequency of occurrence of clear sky computed from the cloud type reports would be too high and the frequencies of occurrence of the various cloud types would be too low. The magnitude of this bias can be estimated by counting the number of times observers within a grid box omit cloud types ( $Nh \neq$  or  $C_L \neq$ ) when  $N>0$  and assuming that the same fraction of the cases of

N=0 would be from stations or ships that do not report cloud types, and thus should be excluded from the denominator when determining the frequency of occurrence of a cloud type.

Figures 4a (land) and 4b (ocean) show the global distribution of the percent occurrence of  $C_L \neq /$  (or  $N_h \neq /$ ) in light reports with  $N > 0$  (this bias fraction is referred to as  $f_b$ ). (Note that when computing these fractions using the ECRA, reports of  $N_h = -1$  with  $IC = 4$  must not be counted in  $f_b$  since these values were set during data processing and not by the observer.) In the land data, extremely large values occur only in northern Alaska (87%, underlined in Figure 4a) and in northeast Greenland (72%). Ship data show large values in the Great Lakes region of North America where values exceed 90% and in some European waters where values are as great as 82%. Values of  $f_b$  average about 3% for land data and about 10% for ship data.

To determine how much effect this bias would have on computed cloud type frequencies, the "clear-sky adjustment factor" (AF0) was defined such that  $F_a = AF0 \cdot F_r$ , where  $F_a$  is the adjusted frequency of occurrence of some cloud type and  $F_r$  is the frequency that would be computed using the potentially biased cloud type reports. Since  $F_r = N_t / N_r$ , where  $N_t$  is the number of occurrences of a particular cloud type and  $N_r$  is the number of reports contributing to the cloud type analysis, and  $F_a = N_t / (N_r - f_b \cdot N_0)$ , where  $N_0$  is the number of occurrences of  $N=0$  and  $f_b$  is the fraction of  $N_0$  that should be discounted, then, using  $f_0 = N_0 / N_r$ ,  $F_a$  can be represented as  $F_a = N_t / (N_r - f_b \cdot f_0 \cdot N_r) = N_t / (N_r \cdot (1 - f_b \cdot f_0)) = F_r / (1 - f_b \cdot f_0)$ . Thus  $AF0 = 1 / (1 - f_b \cdot f_0)$  and is equal to one if either  $f_b$  or  $f_0$  is zero.

Figures 5a and 5b show the global distribution of AF0 {displayed as  $100 \times (AF0 - 1)$ } over land and ocean, respectively. This analysis shows that, on average, cloud type frequencies, if uncorrected, would be reduced only to about 99.5% of their correct values by this bias (average AF0 = 1.003 for land and 1.007 for ocean) and that most regions of the globe are essentially unaffected. However a few regions are greatly affected, namely northern Alaska, northeastern Greenland, the Great Lakes, and some seas around Europe. These are the regions with high values of  $f_b$  (Figures 4a and 4b). The two most extreme regions, the Great Lakes and northern Alaska, have biased values 41% and 27% of their correct values. Other than the AF0 values of 3.68 in Alaska and 1.31 in Greenland, no land box has a value greater than 1.08. In the ocean data, moderate values of AF0 (around 1.10) occur in the Middle East where clear-sky frequencies are high (W88), and several higher values are seen in the seas of Europe and in the Great Lakes of North America. While it may be desirable to apply this adjustment factor to regions of moderately large AF0 values, practical application of this correction will be complicated by the fact that AF0 values vary from year to year, season to season, and day to night. Fortunately correction is unnecessary over most regions of the globe and the regions noted in Figures 5a and 5b for large adjustment factors can simply be eliminated from a cloud type analysis.

Note that while cloud type frequencies are subject to the clear-sky bias because of the coding instructions and practices for cloud type reports, the frequency of occurrence of clear sky itself (and also the frequency of fog) can be computed without this bias by using the total cloud data set to which this bias does not pertain (as was done in Hahn et al., 1994a).

### 3) The Sky-obscured Bias

Our recognition of certain cases of  $N=9$  as overcast cloud (Section 3B) is important in obtaining accurate estimates of the amounts of fog and nimbostratus cloud, but may introduce the sky-obscured bias, which is similar in principle to the clear-sky bias discussed above. Since  $C_L \neq /$  whenever  $N=9$ , it is not possible to distinguish stations or ships that normally report cloud types from those that do not. Thus the latter stations will contribute to the cloud type analysis only when the sky is obscured (aside from the case of clear sky which was discussed above). This will tend to cause the computed frequencies of fog and nimbostratus cloud to be too large and the frequencies of other cloud types to be reduced. The fraction  $f_b$

shown in Figures 4a and 4b is again a measure of the potential of this bias. A "sky-obscured adjustment factor" (AF9) is defined in a manner similar to that for AF0 defined above such that  $F_a = AF9 \cdot F_r$  and  $AF9 = 1 / (1 - f_b \cdot f_9)$ , where  $f_9$  is the frequency of occurrence of N=9 in cloud type reports.

The global distribution of  $f_9$  is given in Figures 6a and 6b. The box in northern Alaska (underlined), which was shown previously to have  $f_b=0.87$  (Figure 4a), also has the relatively large value of 0.16 for  $f_9$  which gives  $AF9=1.16$ . Inspection of Figures 4a and 6a together shows that this is the largest AF9 value for land data and that in most regions the value of AF9 is near 1.00. Ship data have larger  $f_b$  values (Figure 4b) and larger  $f_9$  values (Figure 6b) than land data. Also the large values are distributed over larger regions. The largest  $f_9$  values occur, again, in the Great Lakes region where they combine with  $f_b$  to produce AF9 values that approach 2. In the North Pacific, where large amounts of fog occur during the summer season (W88), moderately large  $f_9$  values (~0.20) occur with moderate  $f_b$  values (~0.11), giving  $AF9=1.02$  which is a relatively small bias. The global average values for AF9 are 1.0003 for land and 1.003 for the ocean. Thus, aside from the few regions specially noted to be removed from cloud type analysis, the sky-obscured bias is generally small. Any bias in the frequency of fog itself can be eliminated by computing it from the total cloud reports, as mentioned above. Any bias towards increasing the nimbostratus frequency will be small since N=9 contributes only a small portion of the total nimbostratus (Table 12) and will be compensated somewhat by the tendency towards reducing the frequency of Ns contributed by the  $C_M=2,7$  and  $C_M=7$  paths.

## B. Computing the Average Cloud Amounts and Frequencies

The determination of the average cloud amounts and frequencies of occurrence from surface observations requires some special considerations to avoid various potential biases and to obtain representative values. Upper level clouds present special problems because they are sometimes partially or completely hidden from the view of the observer by lower clouds. These issues are discussed in detail in W86 and W88 but will be highlighted here.

### 1) Total Cloud Cover

Total cloud cover is basically the sum of the values of N in the synoptic code (converted to percent if desired) divided by the number of contributing reports. However, to avoid the day-night bias discussed above, some method of equalizing the contribution of reports between day and night is necessary. In W86, averages were obtained by first forming averages for the 8 synoptic hours and then averaging these 8 numbers. For oceans, where data are less plentiful, this method will result in significant loss of data because the 3-hourly times often do not have a sufficient number of reports to obtain a statistically reliable average. Therefore, Hahn et al. (1994a) divided the day into two 12-hour periods, 0600-1800 local time ("day") and 1800-0600 local time ("night"), and averaged these two numbers. Note that when using only the light reports (to avoid the night-detection bias) to form monthly averages, only about two weeks of data will contribute to the nighttime average in any single month. Due to this "monthly-sampling error" there will be more scatter in monthly averages from year to year although multi-year averages should become more statistically representative of climatological means as the number of contributing years is increased. Similarly, seasonal averages should be more representative of an individual season than monthly averages are of an individual month.

These considerations of the day-night bias, night-detection bias, and monthly-sampling error apply equally well to cloud type analyses discussed below. However, for quantities such as fog and precipitation, whose detection does not depend on illumination, all observations may be used, minimizing all three of these biases.

## 2) Low Cloud Types

Of the 90.3 million light reports suitable for total cloud analysis for land (Figure 1), 88.0 million have cloud type information (Table 5). For the ocean these numbers are 11.1 million and 9.4 million. In the type reports, the amount of a low cloud type (if present) is always given in the  $N_h$  variable of the report. The average amount for a particular low cloud type can be obtained, in a manner similar to that for total cloud amount, by summing the  $N_h$  values when the type is present and dividing by the number of contributing reports (using the precautions against the day-night bias discussed above and adjusting for the small clear-sky and sky-obscured biases if desired). The contributing reports consist of those with  $C_L \geq 0$  and  $N_h \geq 0$  and include reports of  $N=0$ . An alternative, but equivalent, method for obtaining the average is to compute the frequency of occurrence ( $f$ ) of the type (the number of occurrences of the type divided by the number of contributing reports) and the amount-when-present (awp; sum of  $N_h$ 's divided by the number of occurrences of the type) separately. Then the average amount is  $avg = f \times awp$ . This latter method is described because it is often of interest to know the frequency of occurrence of a type in addition to its amount, because awp tends to be characteristic of a cloud type, and because this is the method used in computing upper level cloud type amounts.

## 3) Upper Level Clouds

Cloud type reports do not always contain information about upper level clouds because they may be hidden by an overcast or near-overcast layer of lower clouds. Thus, of the 88.0 million light type reports for land (Table 8), only 74.4 million have information about the middle cloud level ( $C_M \geq 0$ ) and 60.3 million have information about the high level ( $C_H \geq 0$ ). Of the 9.4 million light type reports for the oceans, 7.1 million have  $C_M \geq 0$  and 5.5 million have  $C_H \geq 0$ .

The average amounts of upper level cloud types are obtained as described in the last section:  $avg = f \times awp$ . Since we want the actual frequency of occurrence of a cloud type, and not just the frequency with which it is visible,  $f$  is computed as the number of times the type was observed divided by the number of reports of  $C_U \geq 0$  (where  $C_U$  represents either  $C_M$  or  $C_H$ ). For land, middle and high cloud frequencies are determined from 84.5% and 68.5% of the light type reports, respectively. For the oceans these values are 75.5% and 58.5%, respectively. The question of the degree to which these portions of the data set represent the whole data set for types is discussed in W86 and W88. Based on a study of the frequency of occurrence of  $As/Ac$  [ $f(As,Ac)$ ] versus low cloud amount, Warren et al. (W88) applied an adjustment to  $f(As,Ac)$  which assigned to the cases of  $C_M = /$  (15.5% of the type reports for land and 24.5% for ocean) a value that is the average of  $f(As,Ac)$  of the reports that have low cloud amounts of 3 to 7 oktas. For high clouds,  $f$  was computed only from reports with  $N_h < 7$  in order to reduce the partial-undercast bias (W88).

The amount-when-present of an upper cloud type can be determined, when it is reported present ( $C_U > 0$ ), only if there are at most 2 cloud levels present. In addition, amounts are not computed for an upper cloud if it is undercast by a layer which covers 7 oktas or more of the sky (Section 3D). Therefore awp is computed from an even smaller pool of data than that used for frequency computations. Table 8 shows that, for land, 78% of the observed (light) occurrences of middle clouds and 75% of the observed occurrences of high clouds are computable. For the ocean data these values are 61% and 43%, respectively. Nevertheless, awp computed from these data is probably fairly representative of the actual awp (W86 and W88) and any error in awp results in a smaller error in avg by the factor  $f$ . Any systematic error inherent in the random-overlap assumption would produce a smaller error in computed amounts since this assumption is used for only a fraction of the computable observations. Table 8 shows that the random-overlap assumption is used in 39% of the computable observations (light) of middle cloud and in 55% for high cloud over land. These fractions are larger for ocean data with random overlap used for 60% of the computable middle clouds and 70% of the computable high clouds.

Special note about Ns: Because Ns is defined on the basis of the occurrence of precipitation (Table 2) which does not depend on the visibility of the middle cloud level for its detection, its presence or absence is known for every type report. Thus the number of contributing reports for f(Ns) is the same as that for low cloud types ( $C_L \geq 0$  and  $N_h \geq 0$ ). However, when present, its amount is not always known and a separate tally (which will be different from that for the As/Ac clouds) must be kept for determining its awp.

## 7. HOW TO OBTAIN THE DATA

This documentation and the data described herein are available from:

Carbon Dioxide Information Analysis Center  
Oak Ridge National Laboratory  
Post Office Box 2008  
Oak Ridge, TN 37831-6335, U.S.A.  
Telephone (423) 574-3645 or (423) 574-0390

or

Data Support Section  
National Center for Atmospheric Research  
Boulder, CO 80307, U.S.A.  
Telephone (303) 497-1215.

The following citation should be used for referencing this archive and/or this documentation report:

Hahn, C.J., S.G. Warren, and J. London, 1996: *Edited Synoptic Cloud Reports from Ships and Land Stations Over the Globe, 1982-1991*. NDP026B, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, TN. (Also available from Data Support Section, National Center for Atmospheric Research, Boulder, CO.)

The archive of our earlier climatology (Hahn et al., 1988) and the accompanying atlases (Warren et al., 1986, 1988) are also available from the same sources listed above.

## ACKNOWLEDGEMENTS

This work was supported by NASA Grant NAG-1-998 and by Battelle Pacific Northwest Laboratories, Atmospheric Radiation Measurement Program (Contract 144806-A-Q1). We also acknowledge the long term cooperation and provision of computing facilities by the Scientific Computing Division of the National Center for Atmospheric Research.

## REFERENCES

Hahn, C.J., S.G. Warren, J. London, and R.L. Jenne, 1988: *Climatological Data for Clouds Over the Globe from Surface Observations*. NDP-026, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, TN. (Also available from Data Support Section, National Center for Atmospheric Research, Boulder, CO.)

- Hahn, C.J., S.G. Warren, and J. London, 1992: The use of COADS ship observations in cloud climatologies. *Proceedings of the International COADS Workshop*, H.F. Diaz, K. Wolter, and S.D. Woodruff, Eds., NOAA/ERL, Boulder, CO, 271-280.
- Hahn, C.J., S.G. Warren, and J. London, 1994a: *Climatological Data for Clouds Over the Globe from Surface Observations, 1982-1991: The Total Cloud Edition*. NDP026A, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, TN. (Also available from Data Support Section, National Center for Atmospheric Research, Boulder, CO.)
- Hahn, C.J., S.G. Warren and J. London, 1994b: The effect of moonlight on observation of cloud cover at night, and application to cloud climatology. *J. Climate*, **8**, 1429-1446.
- Hapke, B., 1971: Optical properties of the lunar surface. *Physics and Astronomy of the Moon*, 2nd ed., Z Kopal, Ed., Academic Press, New York, 303 pp.
- Riehl, H., 1947: Diurnal variation of cloudiness over the subtropical Atlantic Ocean. *Bull. Amer. Meteor. Soc.*, **28**, 37-40.
- Rossow, W.B. and R.A. Schiffer, 1991: ISCCP cloud data products. *Bull. Amer. Meteor. Soc.*, **72**, 2-20.
- Schneider, G., P. Paluzzi and J.P. Oliver, 1989: Systematic error in the synoptic sky cover record of the South Pole. *J. Climate*, **2**, 295-302.
- Slutz, R.J., S.J. Lubker, J.D. Hiscox, S.D. Woodruff, R.L. Jenne, D.H. Joseph, P.M. Steurer and J.D. Elms, 1985: *Comprehensive Ocean-Atmosphere Data Set; Release 1*. NOAA Environmental Research Laboratories, Boulder, Colo., 268 pp. (NTIS PB86-105723).
- Tian, L. and J.A. Curry, 1989: Cloud overlap statistics. *J. Geophys. Res.*, **94**, 9925-9935.
- Warren, S.G., C.J. Hahn, J. London, R.M. Chervin and R.L. Jenne, 1986: *Global distribution of total cloud cover and cloud type amounts over land*. NCAR Technical Note TN-273+STR, Boulder, CO, 29 pp. + 200 maps (also DOE/ER/60085-H1).
- Warren, S.G., C.J. Hahn, J. London, R.M. Chervin and R.L. Jenne, 1988: *Global distribution of total cloud cover and cloud type amounts over the ocean*. NCAR Technical Note TN-317+STR, Boulder, CO, 42 pp. + 170 maps (also DOE/ER-0406).
- Woodruff, S.D., S.J. Lubker, and M.Y. Liu, 1992: Updating COADS<sup>-</sup> problems and opportunities. *Proceedings of the International COADS Workshop*, H.F. Diaz, K. Wolter, and S.D. Woodruff, Eds., NOAA/ERL, Boulder, CO, 19-36.
- Woodruff, S.D., R.J. Slutz, R.L. Jenne and P.M. Steurer, 1987: A comprehensive ocean-atmosphere data set. *Bull. Amer. Meteor. Soc.*, **68**, 1239-1250.
- World Meteorological Organization, 1977: *Weather Reporting/Messages Meteorologiques, Volume A*. (WMO Publ. No. 9), WMO, Geneva.
- World Meteorological Organization, 1988: *Manual on Codes, Volume I*. (WMO Publ. No. 306), WMO, Geneva.

Table 1. Cloud Information Contained in Synoptic Weather Reports

Symbol	Meaning	Codes *
N	total cloud cover	0-8 oktas 9= sky obscured
N <sub>h</sub>	lower cloud amount	0-8 oktas
h	lower cloud base height	0-9
C <sub>L</sub>	low cloud type	0-9
C <sub>M</sub>	middle cloud type	0-9
C <sub>H</sub>	high cloud type	0-9
ww	present weather	00-99
I <sub>x</sub>	present weather indicator	1-6

\* Any category for which information is lacking to the observer is coded as "/".



Table 2. Cloud and Weather Type Definitions Used

Level	Shorthand notation	Meaning	Synoptic codes	Extended codes*
	Tcc	total cloud cover	N = 0-9	0-8
	Clr	completely clear sky	N = 0	
	Ppt	precipitation	ww= 50-75,77,79,80-99	
	D	drizzle	50-59	
	R	rain	60-69	
	S	snow	70-75,77,79	
	Ts	thunderstorm, shower	80-99	
Low			$C_L =$	
	Fg	sky obscured by fog	/ with N=9 and ww=F*	11
	St	stratus	6,7	
	Sc	stratocumulus	4,5,8	
	Cu	cumulus	1,2	
	Cb	cumulonimbus	3,9, or N=9 with ww=Ts	10
Mid			$C_M =$	
	Ns	nimbostratus	2,7, or N=9, with ww=DRS / with ww=DRS and $C_L=0,7$ / with ww=RS and $C_L=4-8$	12,11,10 10 10
	As	altostratus	1; 2 if not DRS	
	Ac	altocumulus	3,4,5,6,8,9; 7	
High			$C_H =$	
	Cs	cirrostratus	5,6,7,8	
	Cic	cirrus, cirrocumulus	1,4,9	
	Cid	dense cirrus	2,3	

\* Extended codes shown where they differ from synoptic codes. In the extended code the value "-1", rather than "/", is used to signify missing information.

\* F represents the fog codes ww=10-12,40-49.

Table 3. FORTRAN Code for Checking Cloud Type Consistencies (All variables are integers. Cloud variables are defined in Table 1. Here DRS (Table 2) is "1" if true and "0" if false. IC is the Change Code (Table 4).)

```

c.IC_2,4----- CORRECT MISCODED Nc with CM -----
IF (CM.GE.0 .AND. Nc.EQ.0 .AND. CL.EQ.0) THEN
  IF (CM.LE.0) THEN
    Nc = 0
    IC = 2
  ELSE
    IC=4
    Nc= -1
  END IF
END IF
IF (CM.GT.0 .AND. Nc.LT.N .AND. CL.EQ.0.AND.CH.EQ.0) THEN
  IF (Nc.GE.0) IC=4
  Nc= -1
END IF

c.IC_3,4----- CORRECT MISCODED Nc with CH -----
IF (CH.GT.0 .AND. CM.EQ.0 .AND. CL.EQ.0) THEN
  IF (Nc.NE.0) THEN
    IF (Nc.EQ.N) THEN
      Nc=0
      IC = 3
    ELSE
      IF (Nc.GE.0) IC=4
      Nc=-1
    END IF
  END IF
END IF

c.IC_4----- EXCLUDE INCONSISTENCIES for TYPES -----
IF (CL.GT.0 .AND. CM.EQ.0.AND.CH.EQ.0) THEN
  IF (Nc.LT.N .AND. N.NE.0) THEN
    IF (Nc.GE.0) IC=4
    Nc=-1
  END IF
END IF

c.IC_5-----
IF (CL.LT.0 .OR. Nc.LT.0) THEN
  IF ((CM.GE.0 .OR. CH.GE.0) .AND. CL.LT.0) IC=5
  Nc=-1
  CL=-1
  CM=-1
  CH=-1
  h =-1
END IF

c.IC_6----- CORRECT MISCODED CM,CH 0->/ -----
IF (CH.EQ.0 .AND. CM.EQ.0 .AND.
1 (N.EQ.0 .OR. (N.EQ.7.AND.N.EQ.Nc))) THEN
  CM=-1
  IC=6
END IF
IF (CH.EQ.0 .AND.
1 (N.EQ.0 .OR. (N.EQ.7.AND.N.EQ.Nc))) THEN
  CH=-1
  IC=6
END IF

c.IC_7----- RESET Nc FROM CM 2,7 -----
IF (DRS.EQ.1 .AND. (CM.EQ.2 .OR. CM.EQ.7)) THEN
  IF (CM.EQ.2) CM=12
  IF (CM.EQ.7) CM=11
  IC = 7
END IF

c.IC_8----- SET Nc FROM CM/ -----
IF (DRS.EQ.1 .AND. CM.LT.0 .AND. CL.GE.0) THEN
  IF (.NOT.(CL.EQ.1 .OR.CL.EQ.2 .OR.CL.EQ.3 .OR.CL.EQ.9)) THEN
    IF (Nc.GE.0 .OR. CL.EQ.7 .OR. CL.EQ.0) THEN
      CM =10
      IC = 8
    END IF
  END IF
END IF

c.IC_9----- CORRECT MISCODED CM,CH /->0 -----
IF (CM.LT.0 .AND. CH.GE.0 .AND. CL.GE.0) THEN
  CM=0
  IF (IC.EQ.0) IC=9
END IF
IF (Nc.LE.4 .AND. N.EQ.Nc .AND. CL.GE.0) THEN
  IF ((CM.LT.0 .OR. CH.LT.0).AND.IC.EQ.0) IC=9
  IF (CM.LT.0) CM=0
  IF (CH.LT.0) CH=0
END IF

```

Table 4. Change Codes for Edited Cloud Reports

IC	* Case (brief description **)	Changes made #	Occurrence (%)			
			Land		Ocean	
			all	light	all	light
0		none##	88.2	88.2	84.4	84.2
1	N=9 with ppt or fog 3.7 3.5	N=8; CL=10,11 or CM=10			1.5	1.5
2	Nh=0 with CM>0 and CL=0 0.1 0.1	Nh=N		0.2	0.2	
3	Nh=N with CH>0 and CL=CM=0	Nh=0	0.1	0.1	0.3	0.4
4	Nh<N where should be Nh=N	Nh=/	0.4	0.4	0.9	0.9
5	CL =/ with CM or CH	CM,CH =/	0.1	0.1	0.7	0.7
6	CM or CH miscoded as 0	CM or CH =/	3.2	3.5	4.0	4.4
7	CM=7,2 for Ns	CM=11,12	3.7	3.5	1.3	1.4
8	CM=/ for Ns	CM=10	2.4	2.2	1.9	2.0
9	CM or CH miscoded as /	CM or CH =0	0.3	0.3	2.7	2.4

\* Also order in which changes made. IC=9 is recorded only if no previous change (possibly 7 or 8) occurred.

\*\* See Table 3 for details.

# The value "-1" is used to signify "/".

## Cases of N=0 for which Nh=CL=CM=CH=/ were set to 0 were not considered to be changes.

Table 5. Number of Reports with Cloud Type Information (Nh > -1, CL > -1)

	Land	Ocean
all reports	121 million	12.1 million
light reports	88 million	9.4 million

Table 6. FORTRAN Code for Determining Upper Level Cloud Amounts

```

*jum,juh are middle and high cloud non-overlapped amounts in octa (9=missing).
*AM,AH are "actual" amounts with possible use of random overlap.
*JAM,JAH are integer values of AM,AH to 2 decimal places for ECR (900=missing).
* Other variables are integers. Cloud variables are defined in Table 1.
    jum= 9
    juh= 9
    JAM=900
    JAH=900
    if (CL.ge.0 .and. Nh.ge.0) then
c_MID-----
    IF (CM.GT.0) THEN
c. . . . . present
        IF (CL.EQ.0 .OR. CH.LE.0) THEN
            IF (CL.EQ.0) THEN
                jum= Nh
c. . . . . computable
                JAM= Nh*100
            ELSE
                jum=(N-Nh)
                IF (CM.LE.12 .AND. CM.GE.10) THEN
c. . . . . Ns computable
                    JAM= N*100
                ELSE
                    IF (Nh.LT.7) THEN
c. . . . . computable,ROL
                        AM= 8.*(N-Nh) / (8.-Nh)
                        JAM= AM*100. +.5
                    END IF
                END IF
            ELSE
                IF (Nh.EQ.N) jum=0
            END IF
        ELSE IF (CM.EQ.0) THEN
            JAM=0
            jum=0
        ELSE
            IF (Nh.EQ.N .AND. CL.GT.0) jum=0
        END IF
c_HI-----
    IF (CH.GT.0) THEN
c. . . . . present
        IF (CL.EQ.0 .AND. CM.EQ.0) THEN
            juh= N
c. . . . . computable
            JAH= N*100
        ELSE IF (CL.EQ.0 .OR. CM.EQ.0) THEN
            juh= (N-Nh)
            IF (Nh.LT.7) THEN
c. . . . . computable,ROL
                AH= 8.*(N-Nh) / (8.-Nh)
                JAH= AH*100. +.5
            END IF
        ELSE
            IF (Nh.EQ.N) juh=0
        END IF
    ELSE IF (CH.EQ.0) THEN
        JAH=0
        juh=0
    ELSE
        IF (Nh.EQ.N .OR. (CL.GT.0 .AND. CM.GT.0)) juh=0
    END IF
end if

```

Table 7. RANDOM OVERLAP COMPUTATION TABLE  
 (For upper level cloud amount when 2 and only 2 levels present)  
 Amount Upper Cloud [octa] = 8. \* (N - Nh) / (8. - Nh)

N\Nh	7	6	5	4	3	2	1	0
8	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00
7	0	4.00	5.33	6.00	6.40	6.67	6.86	7.00
6	x	0	2.67	4.00	4.80	5.33	5.71	6.00
5	x	x	0	2.00	3.20	4.00	5.57	5.00
4	x	x	x	0	1.60	2.67	3.43	4.00
3	x	x	x	x	0	1.33	2.29	3.00
2	x	x	x	x	x	0	1.14	2.00
1	x	x	x	x	x	x	0	1.00

Table 8. Number of Reports in which Upper Level Cloud Amounts were Computable  
 (millions of reports)

Number of:	LAND				SHIPS			
	Middle Cloud		High Cloud		Middle Cloud		High Cloud	
	Light	Dark	Light	Dark	Light	Dark	Light	Dark
Type reports	88.0	33.0	88.0	33.0	9.4	2.7	9.4	2.7
NOL computed	81.2	31.7	81.1	31.7	8.4	2.5	8.4	2.5
Level reported	74.7	#	60.3	#	7.1	#	5.5	#
Cloud observed	33.0	10.0	27.6	6.5	4.1	0.8	2.3	0.8
Computable	25.7	8.7	20.7	5.3	2.5	0.6	1.0	0.15
ROL used	10.1	2.4	11.3	2.3	1.5	0.3	0.7	0.10

\* NOL signifies non-overlapped amounts and ROL signifies the random overlap assumption.

# Data not available.

Table 9. Contents and Format of the 56-character EDITED CLOUD REPORT

Item	Description	Abbreviation	Number of characters	Minimum value	Maximum value
1	year, month, day, hour	yr, mn, dy, hr	8	81120100	91113023
2	sky brightness indicator	IB	1	0	1
3	latitude x100	LAT	5	-9000	9000
4	longitude x100	LOn	5	0	36000
5	station number (land) or source deck (ships)	ID	5	01000 110	98999 999
6	land/ocean indicator	LO	1	1	2
7	present weather	ww	2	-1	99
8	total cloud cover	N	1	0	8 <sup>#</sup>
9	lower cloud amount	Nh	2	-1	8
10	lower cloud base height	h	2	-1	9
11	low cloud type	CL	2	-1	11 <sup>#</sup>
12	middle cloud type	CM	2	-1	12 <sup>#</sup>
13	high cloud type	CH	2	-1	9
14	middle cloud amount <sup>*</sup> x100	AM	3	0	900
15	high cloud amount <sup>*</sup> x100	AH	3	0	900
	non-overlapped amounts: <sup>*</sup>				
16	middle cloud amount	UM	1	0	9
17	high cloud amount	UH	1	0	9
18	change code	IC	2	0	9
19	solar altitude (deg x10)	SA	4	-900	900
20	relative lunar illuminance x100 RI		4	-110	117

# Cases of sky-obscured (N=9) due to fog (ww 10-12, 40-49), thunderstorms (ww 80-99) or drizzle/rain/snow (ww 50-75, 77, 79) have been converted to N=8 and CL=11 (fog) or CL=10 (thunderstorms) or CM=10 (Ns). Certain cases of CM=/ with drizzle/rain/snow have been converted to CM=10 also. Cases of CM=2, 7 with drizzle/rain/snow have been converted to CM=12, 11, respectively, to indicate Ns. Changes are coded in the IC parameter.

\* Upper level cloud type amounts can be computed only when 2 or less levels are present. AM and AH come from Nh, N, or the random-overlap equation and represent "actual" cloud amount. UM and UH are the amounts visible (the non-overlapped amounts) and come from Nh, N, or N-Nh. [Amounts are given in octa, AM and AH to two decimal places.]

Table 10. Sample Edited Cloud Reports in 56-character Format

	I			L							uu I		
	YrMnDyHrB	Lat	Lon	IdOwvNNh	hClCmCh	Am	AbMH	C	SA	RI			
Ship:	(1)	811205060	674030580	8902758	8	0	010-180090080	1-391	-4				
	(2)	811224180	5810 140	8882618	-1	-1	-1-190090099	0-204	0				
	(3)	811230181	586021640	9272	37	6	3 6 6-140090010	0 -31	-1				
	(4)	811209181	5510 600	8902857	-1	-1	-1-190090099	0-230	31				
	(5)	811203001	541019970	9262	36	6	2 7 1-1 090000	0 114	0				
	(6)	811218120	514017160	9272683	3	3	710 0300 000	8-613	-1				
	(7)	811220121	500035790	8902	28	8	3 5 2-190090000	0 165	0				
	(8)	811204000	498029810	9262458	8	0	11-1-190090000	1-363	2				
	(9)	811207181	468031380	8902468	8	9	0 1-180090080	6 91	3				
	(10)	811224121	428031810	8902804	2	0	0 8 520026722	0 134	0				
	(11)	811210001	398014250	8902	27	7	5 5 2-190090000	6 190	-38				
	(12)	811225060	397023360	8902463	-1	-1	-1-190090099	5-552	0				
	(13)	811224001	369014160	8882	14	0	9 0 0 6 040004	3 197	0				
	(14)	811219151	360034550	9262	25	4	3 3 2-120090010	0 236	-1				
	(15)	811216120	355014110	8902	00	0	-1 0 0 0 0 000	0-551	-3				
	(16)	811224061	349016210	9262	28	7	4 5 2-190090010	0 1	0				
	(17)	811209001	332016430	8902	38	8	4 7 8 190090000	0 324	-33				
	(18)	811221050	332033690	9272	12	-1	-1-1-190090099	4-433	0				
	(19)	811224120	245013560	9262508	8	3	4-1-190090000	0-496	0				
	(20)	811205001	250023650	9262	27	7	4 8 9 390090000	0 158	6				
	(21)	811221181	244028980	9262	56	4	5 8 3-140090020	0 382	0				
	(22)	811211181	218027640	9272	22	2	3 1 0 0 0 000	9 444	-72				
	(23)	811210121	143011600	8902	28	1	3 1 0 5 080007	0-308	58				
	(24)	811222181	121026940	9272	23	-1	-1-1-190090099	4 544	0				
	(25)	811230120	-47911170	8902807	3	4	3 7-164090040	0-172	0				
	(26)	811205180	-78911830	9262	28	3	3 6 1-180090050	0-473	-3				
	(27)	811221121	-123932380	8902508	8	4	9-1-190090000	0 543	1				
	(28)	811201181	-163934780	8902	26	6	7 0 5 0600 060	2 145	1				
	(29)	811218061	-2069 1280	8902	27	7	4 5-1-190090000	6 200	7				
	(30)	811209001	-2239 780	8902168	8	3	9 7-190090000	0-438	29				
	(31)	811226121	-238931790	9262	38	6	6 4 7-180090020	0 514	0				
	(32)	811213181	-3519 1940	8882	38	8	5 6 5-190090000	6 -23	-21				
	(33)	811217001	-351927230	8902	27	6	6 5-1-190090099	0 106	-13				
	(34)	811201181	-395930100	8902	56	-1	-1-1-190090099	4 563	1				
	(35)	820104180	448014800	8902638	8	5	712-180090000	7-391	-6				
Land:	(36)	820102180	411012615543771498	8	0	11-1-190090000	1-554	-4					
	(37)	820103121	6112 907 13711-11-1-1-1-190090099	0	58	0							
	(38)	820110181	-2087 5552619801-14	1	5	8 3 290090099	0-343	34					
	(39)	820109031	184329033784851-17	4	5	2 3 0600 030	0-649	89					
	(40)	820122091	6028 522 13111618	7	4	712-180090010	7 28	0					

Table 11. Distribution of Reports over the Synoptic Reporting Times

	Total Number	Percent of reports at reporting times (GMT)									
		00	03	06	09	12	15	18	21	6-hr	3-hr
<b>LAND</b>											
all reports	124,164,607	14.4	9.4	14.6	11.4	15.7	10.3	14.4	9.8	59.1	40.9
light reports	90,348,885	12.6	9.6	17.6	14.7	17.3	10.3	11.3	6.6	58.8	41.2
cloud type rpts (light)	87,982,297	12.6	9.7	17.6	14.6	17.3	10.3	11.3	6.6	58.8	41.2
<b>SHIP</b>											
all reports	14,721,941	22.7	2.6	22.4	2.8	22.6	2.9	21.7	2.3	89.4	10.6
light reports	11,093,710	22.0	2.2	21.1	3.0	22.9	3.4	23.2	2.2	89.3	10.7
cloud type rpts (light)	9,400,201	22.5	2.1	21.8	2.8	22.8	2.9	23.0	2.1	90.1	9.9

Table 12. Contribution of the Various Paths to Total Nimbostratus Frequency

Path to Ns	ww	LAND 88 million light type reports		OCEAN 9.4 million light type reports	
		frequency Ns, %	percent of Ns	frequency Ns, %	percent of Ns
Total		6.24		4.90	
CM=2,7	D,R,S	3.59	58	1.60	33
N=9	D,R,S	0.39	6	0.96	19
CM=/:		2.26	36 *	2.34	48
CL=7	D,R,S		(17)		(26)
CL=0	D,R,S		(.1)		(.02)
CL=6	R,S		(5)		(12)
CL=4,5,8	R,S		(14)		(10)
Exclusions *:					
CL=4,5,6,8	D	~0.2		~0.6	
CL=1,2,3,9	D,R,S	~0.08		~0.3	

\* Approximate values based on January data: for exclusions and values in parentheses.






Figure 1. Flow chart of report processing through the total cloud stage. (Discard percentages given in order: land, ships. T and R here abbreviate Ts and DRS of Table 2. Other symbols are defined in Tables 1 and 2 except IC is defined in Table 4.)




Figure 2a. Frequency distribution of extended code values for indicated cloud variables in edited light reports from land stations over the globe for 1982-1991. Shaded areas indicate occurrence of precipitation. (N=9 is relabeled as N=8 in the ECR.)









Figure 2b. Frequency distribution of extended code values for indicated cloud variables in edited light reports from ships over the globe for 1982-1991. Shaded areas indicate occurrence of precipitation. (N=9 is relabeled as N=8 in the ECR.)




Figure 3a. Global distribution of log(number) of light reports for cloud types for 1982-1991 land data. 87,982,297 reports in 860 5x5° grid boxes (see Glossary: "5c grid"). Blank indicates no reports.




Figure 3b. Global distribution of log(number) of light reports for cloud types for 1982-1991 ship data. 9,400,201 reports in 1502 5x5° grid boxes (see Glossary: "5c grid"). Blank indicates no reports.




Figure 4a. Global distribution of occurrence of CL= or Nh= with N>0 (fb) in light reports for 1982-1991 land data (%). Global average for 843 5c-grid boxes with 50 or more reports is 2.7%.




Figure 4b. Global distribution of occurrence of CL= or Nh= with N>0 (fb) in light reports for 1982-1991 ship data (%). Global average for 1487 5c-grid boxes with 50 or more reports is 9.7%.




Figure 5a. Global distribution of the clear-sky adjustment factor (AF0) in light reports for 1982-1991 land data. Global average for 843 5c-grid boxes with 50 or more reports is 1.007. Values mapped are  $100 \times (AF0 - 1)$  where  $AF0 = 1 / (1 - fb \cdot f_0)$ .






Figure 5b. Global distribution of the clear-sky adjustment factor (AF0) in light reports for 1982-1991 ship data. Global average for 1487 5c-grid boxes with 50 or more reports is 1.003. Values mapped are  $100x(AF0-1)$  where  $AF0=1/(1-fb*fo)$ .




Figure 6a. Global distribution of occurrence of N=9 (f9,%) in light type reports for 1982-1991 land data. Average of 843 5c- grid boxes with 25 or more reports is 1.1%.




Figure 6b. Global distribution of occurrence of N=9 (19.%) in light type reports for 1982-1991 ship data. Average of 1501 5c-grid boxes with 25 or more reports is 3.3%.

Appendix A. Percent Frequency of Occurrence of Extended Code Values for Cloud Variables in Edited Cloud Reports (Light). Also: percent that reported code occurs with precipitation and percent of total precipitation that occurs with each code value. (rpts=reports)

TOTAL CLOUD AMOUNT													
N CODE	-1	0	1	2	3	4	5	6	7	8	9		
LAND													
% of rpts	0.0	15.3	6.9	7.6	5.9	4.3	6.0	10.1	18.3	24.1	1.5		
% with ppt	0.0	0.0	0.7	0.2	0.3	0.5	1.0	2.4	8.0	34.3	28.8		
% of ppt	0.0	0.0	0.1	0.1	0.2	0.2	0.6	2.4	14.0	78.4	4.1		
SHIP													
% of rpts	0.0	5.8	5.0	7.6	9.0	8.0	7.4	11.8	14.0	28.0	3.5		
% with ppt	0.0	0.0	0.3	0.2	0.5	1.1	2.0	3.7	7.0	22.6	26.7		
% of ppt	0.0	0.0	0.2	0.2	0.5	1.0	1.7	4.9	10.9	70.3	10.4		
LOWER CLOUD AMOUNT													
Nh CODE	-1	0	1	2	3	4	5	6	7	8	9		
LAND													
% of rpts	2.6	25.6	11.2	10.9	7.9	5.7	5.8	6.7	8.9	14.6	0.0		
% with ppt	11.7	0.5	1.1	2.9	5.3	8.7	12.3	16.0	16.1	37.8	0.0		
% of ppt	2.9	1.2	1.2	3.0	4.0	4.7	6.8	10.3	13.7	52.3	0.0		
SHIP													
% of rpts	15.3	8.5	6.4	9.8	9.4	7.9	6.4	7.8	7.8	20.7	0.0		
% with ppt	9.0	0.1	0.6	1.1	2.1	3.9	6.0	8.8	10.0	24.6	0.0		
% of ppt	15.3	0.1	0.4	1.2	2.2	3.5	4.3	7.6	8.6	56.8	0.0		
LOWER CLOUD HEIGHT													
h CODE	-1	0	1	2	3	4	5	6	7	8	9		
LAND													
% of rpts	19.8	1.7	0.5	1.9	3.1	13.3	23.0	10.6	3.4	1.6	21.1		
% with ppt	3.4	28.6	38.0	42.3	39.3	23.1	9.6	5.8	5.8	8.2	4.3		
% of ppt	6.4	4.7	2.0	7.8	11.5	29.2	20.9	5.8	1.9	1.2	8.7		
SHIP													
% of rpts	22.5	4.1	1.1	4.0	12.2	24.3	18.6	5.8	2.2	2.0	3.1		
% with ppt	6.7	25.6	29.6	27.6	16.8	7.7	4.0	2.8	2.3	1.8	1.1		
% of ppt	16.9	11.6	3.8	12.5	22.9	21.0	8.3	1.8	0.6	0.4	0.4		
LOW CLOUD TYPE													
CL CODE	-1	0	Cu	Cu	Cb	Sc	Sc	St	St	Sc	Cb	Cb	Fg
LAND													
% of rpts	2.6	38.3	6.3	10.2	2.4	2.2	15.5	4.0	3.8	5.9	6.6	0.1	1.1
% with ppt	11.7	4.7	0.8	4.0	18.3	5.2	11.7	14.4	67.2	8.7	28.6	100	0.0
% of ppt	2.9	17.1	0.5	3.9	4.2	1.1	17.2	5.5	24.3	4.9	17.8	0.5	0.0
SHIP													
% of rpts	15.3	13.3	11.4	13.5	5.4	6.0	10.8	5.9	6.6	6.5	2.6	0.1	2.6
% with ppt	9.0	6.8	0.7	2.4	9.2	3.3	7.7	22.3	34.5	7.2	21.9	100	0.0
% of ppt	15.3	10.1	0.9	3.7	5.6	2.2	9.3	14.6	25.3	5.2	6.4	1.4	0.0
MIDDLE CLOUD TYPE													
CM CODE	-1	0	As	As	Ac	Ac	Ac	Ac	Ac	Ac	Ns	Ns	Ns
LAND													
% of rpts	17.6	45.9	1.4	2.6	10.8	4.4	1.6	1.5	7.1	1.0	0.1	2.6	0.7
% with ppt	15.0	1.0	17.7	7.4	2.6	2.0	3.8	7.3	4.6	2.2	9.2	100	100
% of ppt	25.0	4.6	2.4	1.8	2.6	0.8	0.6	1.0	3.1	0.2	0.1	24.5	6.9
SHIP													
% of rpts	36.1	26.8	3.5	4.6	5.1	3.8	2.9	4.9	4.8	2.0	1.4	2.8	0.2
% with ppt	9.4	0.9	3.9	3.5	2.2	2.4	2.9	6.4	3.0	3.1	7.6	100	100
% of ppt	37.5	2.7	1.5	1.8	1.2	1.0	1.0	3.5	1.6	0.7	1.2	31.2	2.4
HIGH CLOUD TYPE													
CH CODE	-1	0	Cic	Cid	Cid	Cic	Cs	Cs	Cs	Cs	Cic		
LAND													
% of rpts	33.2	36.3	11.5	10.5	0.9	0.8	0.8	1.8	1.9	1.9	0.4		
% with ppt	28.4	1.0	1.3	1.7	6.2	0.8	1.5	2.4	11.3	2.4	1.9		
% of ppt	89.6	3.6	1.4	1.7	0.5	0.1	0.1	0.4	2.1	0.4	0.1		
SHIP													
% of rpts	50.9	28.8	5.7	3.2	1.2	2.1	1.5	1.5	1.1	2.4	1.5		
% with ppt	15.9	1.4	2.4	2.5	4.4	1.7	1.4	2.4	3.8	2.5	2.5		
% of ppt	90.1	4.4	1.5	0.9	0.6	0.4	0.2	0.4	0.5	0.7	0.4		


## Appendix B. Glossary of Terms and Abbreviations Used\*

---

Term	Meaning and description
5c grid	5x5° (latitude x longitude) boxes between latitudes 50N and 50S, 5x10 for latitudes 50-70, 5x20 for latitudes 70-80, 5x40 for latitudes 80-85, and 5x360 for 85-90.
actual amount	Fraction of the sky covered by a cloud, visible or not.
all reports	All reports regardless of whether they are light or dark.
awp	Amount-when-present. The average fraction of the sky covered by a cloud type when it is present, whether it is visible or not.
dark reports	Reports for which illuminance criterion is not met (IB=0).
ECR	Edited Cloud Report. Synoptic cloud reports screened and edited for cloud code consistencies and written as the 56-character report described in Table 9.
ECRA	Archive made up of edited cloud reports.
extended code	The synoptic code extended beyond the usually allowed values of 0-9 to allow C <sub>L</sub> =10 to represent Cb, C <sub>L</sub> =11 to represent fog and C <sub>M</sub> =10,11,12 to represent Ns cloud.
f	Frequency of occurrence. For a cloud type it is the fraction of weather observations in which the cloud type is present, whether it can be seen or not.
light reports	Reports for which illuminance criterion is met (IB=1).
NOL	Non-overlapped; refers to method for determining upper level cloud amounts.
ROL	Random overlap; refers to method for determining upper level cloud amounts.
total reports	Reports suitable for total cloud analyses (and clear sky, fog, and precipitation); either all reports (the entire ECR data set) or light reports only.
type reports	Reports in which cloud type information is given (Nh≥0 and CL≥0). These may be light, dark, or all reports.

---

\* Terms not shown here may be defined in Table 1, 2, 4 or 9.



INTERNAL DISTRIBUTION

1. B. A. Berven
2. T. A. Boden
3. H. L. Boston
4. A. Brenkert
5. M.-D. Cheng
6. R. B. Cook
7. J. H. Cushman
8. R. M. Cushman
9. V. H. Dale
10. R. C. Daniels
11. M. P. Farrell
12. D. E. Fowler
13. K.N. Gibson
14. S. G. Hildebrand
15. G. K. Jacobs
16. S. B. Jones
17. P. Kanciruk
18. D. P. Kaiser
19. A. Kozyr
20. J. M. Loar
21. R. A. McCord
22. T. R. Nelson
23. D. E. Reichle
24. F. E. Sharples
25. D. E. Shepherd
26. P. Singley
27. D. S. Shriner
28. S. H. Stow
29. L. D. Voorhees
- 30-231. CDIAC
232. Central Research Library
- 233-247. ESD Library
- 248-249. Laboratory Records Department
250. Laboratory Records, ORNL-RC
251. ORNL Patent Section
252. Y-12 Technical Library

EXTERNAL DISTRIBUTION

253. Sereno Barr-Kumarakulasinghe, Marine Sciences Research Center, State University of New York, Stony Brook, New York 11794-5000

254. M. Broido, Acting Director, Environmental Sciences Division, ER-74, U.S. Department of Energy, 19901 Germantown Road, Germantown, MD 20874
255. Kaye L. Brubaker, Hydrology Laboratory USDA-ARS, BARC-W Building 007, Room 104, Beltsville, Maryland 20705
256. Bill Collins, Center for Clouds, Chemistry, and Climate, Scripps Institution of Oceanography, La Jolla, California 92093-0221
257. Roger C. Dahlman, Global Change Research Program, Environmental Sciences Division, Office of Health and Environmental Research, ER-74, U.S. Department of Energy, 19901 Germantown Road, Germantown, MD 20874
258. F.A. Donath, Director, Institute for Environmental Education, Geological Society of America, 1006 Las Posas, San Clemente, CA 92673
259. Marie Doutriaux, LMD/CNRS, Ecole Normale Supérieure, 24 rue Lhomond, 75231 Paris Cedex 05, France
260. Tom Farr, MS 300-233, Jet Propulsion Laboratory, Pasadena, California 91109
261. Diana W. Freckman, Director, College of Natural Resources, 101 Natural Resources Building, Colorado State University, Fort Collins, CO 80523
- 262-266. Carole J. Hahn, Department of Atmospheric Sciences, PAS Building, #81, University of Arizona, Tucson, AZ 85721
267. Anand Inamdar, Center for Clouds, Chemistry, and Climate, Scripps Institution of Oceanography, La Jolla, California 92093-0221
268. R. L. Jenne, NCAR, Scientific Computing Division, PO Box 3000, 1850 Table Mesa Drive, Boulder, Colorado 80307-3000
269. Dennis Joseph, NCAR, Scientific Computing Division, PO Box 3000, 1850 Table Mesa Drive, Boulder, Colorado 80307-3000
270. Thomas R. Karl, National Climatic Data Center, 151 Patton Avenue, Suite 120, Asheville, NC 28801
271. Jeffrey R. Key, Department of Geography, Boston University, 675 Commonwealth Avenue, Boston, Massachusetts 02215
272. Aulis Lind, Associate Professor, Vermont State Climatologist, Box 54170, University of Vermont, Burlington, Vermont 05405-4170
- 273-277. Julius London, Department of Astrophysical, Planetary and Atmospheric Sciences, University of Colorado, Boulder, Colorado 80309
278. Peter Lunn, US Department of Energy, Environmental Sciences Division, Office of Energy Research, ER-74, Washington, DC 20585



279. Farn Parnungo, ARL/NOAA/ERL, Boulder, CO 80303
280. Bobbi Parra, Environmental Sciences Division, Office of Health and Environmental Research, U.S. Department of Energy, G-165, Germantown, MD 20874
281. Ari Patrinos, Associate Director, Office of Health and Environmental Research, ER-70, U.S. Department of Energy, 19901 Germantown Road, Germantown, MD 20874
282. Stephen C. Piper, Scripps Institution of Oceanography, Mail Stop A-020, La Jolla, California 92093-0220
283. S. Ichtiaque Rasool, IGBP Data and Information System Office, Universite Paris, Tour 26, 4 Etage, Aile 26-16, 4 Place Jussieu, 75230 Paris, Cedex 06, France
284. Michael R. Riches, Office of Health and Environmental Research, ER-74, U.S. Department of Energy, 19901 Germantown Road, Germantown, MD 20874
285. Edward S. Sarachik, Department of Atmospheric Sciences, University of Washington, Box 351640, Seattle, Washington 98195-1640
286. Isaac Savdie, CMCC, 224 Glenmanor Way, Thornhill, Ontario L4J3E5, Canada
287. G. S. Saylor, Professor, 10515 Research Drive, Suite 100, The University of Tennessee, Knoxville, TN 37932-2567
288. Jerry Stokes, Battelle Pacific Northwest Lab, Battelle, Blvd, MS K1-74, PO Box 999, Richland, Washington 99352
289. Yichun Shan, Standing Deputy Director, National Marine Data and Information Service, 93 Liu Wei Road, He Dong District, Tianjin 300171, China
290. University Corporation for Atmospheric Research, Library Acquisitions, P.O. Box 3000, Boulder, Colorado 80307-3000
- 291-295. Stephen G. Warren, Department of Atmospheric Sciences, University of Washington, Seattle, Washington 98195
296. Li Wenhai, Deputy Director, National Marine Data and Information Service, 93 Liu Wei Road, He Dong District, Tianjin 300171, China
297. Office of Assistant Manager for Energy Research and Development, U.S. Department of Energy Oak Ridge Operations, P. O. Box 2001, Oak Ridge, TN 37831-8600
- 298-302. Office of Scientific and Technical Information, P. O. Box 62, Oak Ridge, TN 37831

[The page contains extremely faint and illegible text, likely bleed-through from the reverse side of the document. No specific content can be transcribed.]