

Effect of Acute Static Stretch on Maximal Muscle Performance: A Systematic Review

ANTHONY D. KAY^{1,2} and ANTHONY J. BLAZEVICH²

¹*Sport Exercise & Life Sciences, The University of Northampton, Northampton, UNITED KINGDOM;* and ²*School of Exercise, Biomedical & Health Sciences, Edith Cowan University, Joondalup, Western Australia, AUSTRALIA*

ABSTRACT

KAY, A. D., and A. J. BLAZEVICH. Effect of Acute Static Stretch on Maximal Muscle Performance: A Systematic Review. *Med. Sci. Sports Exerc.*, Vol. 44, No. 1, pp. 154–164, 2012. **Introduction:** The benefits of preexercise muscle stretching have been recently questioned after reports of significant poststretch reductions in force and power production. However, methodological issues and equivocal findings have prevented a clear consensus being reached. As no detailed systematic review exists, the literature describing responses to acute static muscle stretch was comprehensively examined. **Methods:** MEDLINE, ScienceDirect, SPORTDiscus, and Zetoc were searched with recursive reference checking. Selection criteria included randomized or quasi-randomized controlled trials and intervention-based trials published in peer-reviewed scientific journals examining the effect of an acute static stretch intervention on maximal muscular performance. **Results:** Searches revealed 4559 possible articles; 106 met the inclusion criteria. Study design was often poor because 30% of studies failed to provide appropriate reliability statistics. Clear evidence exists indicating that short-duration acute static stretch (<30 s) has no detrimental effect (pooled estimate = -1.1%), with overwhelming evidence that stretch durations of 30–45 s also imparted no significant effect (pooled estimate = -1.9%). A sigmoidal dose–response effect was evident between stretch duration and both the likelihood and magnitude of significant decrements, with a significant reduction likely to occur with stretches ≥ 60 s. This strong evidence for a dose–response effect was independent of performance task, contraction mode, or muscle group. Studies have only examined changes in eccentric strength when the stretch durations were >60 s, with limited evidence for an effect on eccentric strength. **Conclusions:** The detrimental effects of static stretch are mainly limited to longer durations (≥ 60 s), which may not be typically used during preexercise routines in clinical, healthy, or athletic populations. Shorter durations of stretch (<60 s) can be performed in a preexercise routine without compromising maximal muscle performance. **Key Words:** MUSCLE STRENGTH, WARM-UP, FORCE REDUCTION, PREPERFORMANCE STRETCH

It is well documented that both physical performance and injury risk can be altered by the performance of a complete preexercise routine (a warm-up) before intense physical work (2,113). Static stretching increases range of motion (ROM) and can also decrease musculotendinous stiffness, even during short-duration (5–30 s) stretches (6,52). Furthermore, a recent review (70) has suggested that there is evidence that preperformance stretching can reduce the risk of acute muscle strain injuries. However, given that multi-intervention preexercise routines commonly include

cardiovascular work, progressively intense muscular contractions and muscle stretching, the specific element or combination of elements responsible for improving performance and reducing injury risk is impossible to ascertain. This issue has been raised in several reviews of the literature, which report equivocal findings regarding the benefits of muscle stretching as a preventative tool for injury risk (70,99,112). Furthermore, numerous publications have reported that acute passive static muscle stretch can induce significant reductions in low-speed (strength), moderate-speed (power), and higher-speed (speed) force production (12,15,21,25,27,40,52,58,59,65,69,77,78,82,96,105,107,119). Accordingly, the inclusion of static stretching in a preexercise routine before the performance of maximal strength-, power-, and/or speed-dependent activities is thought to negatively affect our ability to maximally perform simple and complex movements (movement performance).

A growing body of research has highlighted a detrimental effect of muscle stretching on maximal muscular performance, with some authors specifically examining stretch-induced force deficits in an attempt to identify the possible mechanical, physiological, and neurological mechanisms underpinning these changes in force (40,53,54). This has resulted in the publication of a position statement by the European College

Address for correspondence: Anthony D. Kay, Ph.D., Sport, Exercise & Life Sciences, The University of Northampton, Boughton Green Road, NN2 7AL, United Kingdom; E-mail: tony.kay@northampton.ac.uk.

Submitted for publication April 2011.

Accepted for publication May 2011.

Supplemental digital content is available for this article. Direct URL citation appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site (www.acsm-msse.org).

0195-9131/12/4401-0154/0

MEDICINE & SCIENCE IN SPORTS & EXERCISE®

Copyright © 2012 by the American College of Sports Medicine

DOI: 10.1249/MSS.0b013e318225cb27

of Sport Sciences (63), which concluded that there was firm evidence that an acute bout of stretching could diminish performance in tests requiring maximal muscle efforts. This finding is in agreement with an earlier systematic review (94) examining acute and chronic responses of various stretch modalities on muscular performance. However, a subsequent review by Rubini et al. (89) revealed equivocal effects of static, ballistic, and proprioceptive neuromuscular facilitation stretching on maximal force production. The authors concluded that, although the majority of studies documented a deleterious effect on strength, the broad remit of their review (focusing on both acute and chronic effects, different stretch modalities, and various durations) resulted in equivocal findings. Simultaneously, Young (116) specifically addressed the use of acute static stretching in preexercise routines and concluded that there were equivocal results regarding the effects of acute stretch, possibly resulting from major issues in research design (including a lack of control or reliability analysis) and the long, practically irrelevant durations of the imposed stretches. A more recent review (70) examining the effects of various stretch modes on injury prevention and performance suggested that, although stretching may reduce the acute incidence of muscle strain injuries, there was an abundance of literature demonstrating a negative effect of stretch on performance. Although collectively these four articles report equivocal effects of stretch on maximal force and power production, there is a predominant theme that acute muscle stretch can significantly impair muscle performance and that it should be used with caution in a preexercise routine. A consequence of the detrimental reports in the literature was a recent change in the American College of Sports Medicine's guidelines (3) to suggest the removal of static stretching as part of a warm-up routine and to only include cardiovascular work when strength or power was important to performance.

Closer examination of these reviews revealed that relatively few studies, which specifically address the effects of acute static stretch ($n = 17$ [63], $n = 32$ [70], $n = 36$ [89], $n = 21$ [94], $n = 21$ [116]), were cited. To date, although other generic reviews examining the effects of various muscle stretching modes on performance and injury risk exist, no systematic review has focused specifically on the acute effects of static stretching on maximal muscle efforts. Given that static muscle stretching is the most common form of preexercise stretching to be used in clinical, normal, and athletic populations, there are a considerable number of methodological issues reported in the literature (116). Also, given that numerous articles have been published since Rubini et al. (89) and Young (116) published their findings in 2007 ($n = 64$), the aim of the present review was to provide a detailed systematic examination of the acute effects of passive static stretch on performance in strength-, power-, and speed-dependent tasks. Furthermore, given the equivocal findings reported previously in the literature, the specific effects of static stretch duration, test contraction mode, and the muscle group tested were examined.

METHODS

Search strategy. The latest PRISMA guidelines for conducting a systematic review (73) were followed, including the four-step systematic approach of identification, screening, eligibility, and inclusion. We used a federated search tool (MetaLib) to search four databases concurrently (MEDLINE (1966–2011), ScienceDirect (1823–2011), SPORTDiscus (1985–2011), and Zetoc (1993–2011)) for articles using an acute static stretch-based intervention examining a maximal muscular performance outcome measure; we completed our last search on February 16, 2011. Search terms within the article title were “static stretch*,” “acute stretch*,” “stretch* & effects,” “stretch* & force,” “stretch* & power,” and “stretch* & speed.” Additional searches were conducted on eligible articles using the first author's surname and the search term “stretch*” in the title, with recursive reference screening of eligible articles performed to identify other possibly relevant articles (*enables other “stretch” word derivatives, e.g., stretching, stretches, to be included).

Study selection and inclusion criteria. The review included original research articles examining the effects of an acute static stretch intervention on a maximal voluntary muscular performance outcome measure in strength-, power-, and speed-dependent tasks. Randomized and quasi-randomized control trials were included, which met the PEDro inclusion criteria: 1) the comparison of at least two interventions, 2) that interventions were currently part of physiotherapy practice, 3) that interventions were applied to human subjects, 4) there was randomization of interventions, and 5) the article was a full article published in a peer-reviewed journal. Intervention-based studies examining prestretch and poststretch data that did not meet the first criterion (comparison of at least two interventions) were also included. One reviewer excluded obviously irrelevant articles by screening the titles and abstracts, with a 5% sample of the excluded articles verified by a second reviewer. Abstracts of the remaining articles were assessed by one reviewer, with articles selected for exclusion being verified by a second reviewer. Full texts of the remaining articles were then obtained and independently assessed by two reviewers, with articles selected for exclusion agreed by both reviewers. Discrepancies were resolved by discussion.

Assessment of study validity. Included studies were assessed for methodological quality using the PEDro scale, which comprises 11 criteria, of which the first determines external validity (eligibility criteria) and the remaining 10 measure internal validity (randomization, allocation concealment, homogeneity, subject, therapist and assessor blinded, <15% attrition of subjects, intention to treat, statistical comparison, and measures of variability; for a detailed description of the PEDro scale and criteria, see Maher et al. [64]). The methodological quality of each study was established by awarding one point for each criterion satisfied with a total score out of 10. Two reviewers independently assessed the quality of studies, with disagreements resolved by discussion.

Data extraction. One reviewer extracted data from studies that met the inclusion criteria, while a second reviewer verified the validity of these data. Data that summarized the following factors were extracted: stretch duration, muscle group stretched, maximal muscular performance outcome measures, whether significance was or was not reached in each variable measured (within a realistic post-stretch time frame, ≤ 20 min), mean reduction in a performance variable, and whether appropriate control or reliability analyses were reported. Where multiple variables were reported within studies, each relevant finding was included in the analysis to remove any possible bias on our part and to ensure that reporting bias was not introduced to the review. Multiple analyses within studies were grouped according to stretch duration, performance variable, contraction mode, and muscle group. Where several significant or nonsignificant findings were reported within a specific grouping (e.g., concentric force at several velocities), only one of the significant or nonsignificant findings was tabulated for our synopsis, with the mean of the significant findings used for analysis. This was done to ensure we did not inflate the importance of such studies in relation to others and thus skew the analysis.

Data analysis. Two analyses are reported: 1) where all studies were included, to provide a holistic overview of the published literature; and 2) where studies without appropriate control or provision of reliability statistics were removed. This allowed us to determine whether the removal of studies based on experimental design influenced the findings of the review. Given the heterogeneity of intervention types (specifically differences in stretch duration and muscle group stretched), the diverse methods used to measure muscular performance (specifically isometric, concentric, eccentric or isokinetic muscle actions, drop-, countermovement-, or squat-jump techniques, sprint running over various distances, and free-weight or machine-based strength and power assessment) and that many studies failed to report specific statistical details of both their significant and nonsignificant findings, meta-analysis was deemed to be neither feasible nor appropriate (49). A systematic review of the literature was thus performed with studies pooled according to stretch duration by examining the total time the muscle was placed under stretch (<30 vs $30\text{--}45$ s vs $1\text{--}2$ vs >2 min) and examined for effects on performance in strength-, power-, or speed-dependent tasks. Further analyses were performed again examining duration-dependent effects by muscle contraction mode (e.g., isometric vs concentric vs eccentric) and by muscle group stretched (lower limb only; e.g., plantar flexors vs knee extensors vs knee flexors). The percentage of significant and nonsignificant findings and the magnitude of the changes in the performance variables were collated.

RESULTS

Search results. Our searches identified 4559 potentially relevant articles. By reviewing titles and abstracts, we identified 112 articles examining the effects of acute static

stretch on a maximal muscular performance variable; reference screening of these articles revealed a further 11 articles, giving a total of 123 articles. After examining the full text, 17 articles were removed because they failed to meet our methodological inclusion criteria, which resulted in 106 articles being included for review (see Table, Supplemental Digital Content 1, <http://links.lww.com/MSS/A106>, which presents the major findings of the stretch-based studies included for review).

Methodological quality of included studies. Not all of the PEDro criteria could be satisfied, as the experimental crossover design implemented by the majority of studies resulted in subject and therapist blinding not being possible. Given that therapist and assessor roles were normally performed by the same individuals, assessor blinding was also highly limited. Despite this limitation, the methodological quality of studies was found to be moderate, ranging from 3 to 7 (mean = 5.4 ± 0.9). The present review examined the study designs implemented from 106 randomized and quasi-randomized control trials and intervention-based studies. Careful examination of the study design revealed that 11 studies failed to include a control group or any reliability analyses and a further 21 inappropriately used a control condition (see Table, Supplemental Digital Content 1, <http://links.lww.com/MSS/A105>) that failed to determine reliability, which is a serious concern for the quality of their study design and validity of their data.

Overview—effects on maximal muscular performance. Analysis of the 106 articles revealed that 55% had reported a significant reduction in performances in strength-, power-, or speed-dependent tasks after acute static stretch, whereas 69% had reported no significant reduction in task performances. This apparent conflict in percentages can be explained by numerous studies reporting the effects of acute static stretch on several variables within the same study, including different muscle groups (11), muscle lengths (77), contraction modes (68), contraction velocities (78), durations of stretch (52,58,82,96,119), and performance tasks (91). In addition to equivocal data existing across studies, equivocal data also existed within 25 studies where significant and nonsignificant results were reported concurrently. By examining the findings within the studies rather than collating which studies report significant findings, we were able to remove the possibility of introducing reporting bias on our part. This approach yielded 149 findings from the 106 articles with only 44% of the findings indicating significant reductions in maximal strength-, power-, or speed-dependent performance (pooled estimate of reductions = $-3.7\% \pm 4.9\%$). When the studies without sufficient control or reliability were removed from the analysis, 74 studies reporting 104 findings remained. The percentage reporting significant reductions increased only slightly, to 50%, as a similar proportion of the studies removed reported significant and nonsignificant findings (pooled estimate of reductions = $-4.5\% \pm 5.2\%$). Thus, their removal did not markedly influence the results.

TABLE 1. Duration-dependent effects of acute static stretch on performance in strength-, power-, and speed-dependent tasks, across contraction modes and muscle groups (lower limb).

Comparison	Measure	Stretch Duration	No. of Findings	Percentage Reporting Significant Reduction (%)	Mean ± SD Reduction (%)
Duration	All measures	<30 s	7	14	-1.1 ± 1.8
		30–45 s	23	22	-1.9 ± 3.4
		60–120 s	36	61	-4.2 ± 5.0
		>120 s	38	63	-7.0 ± 5.7
		All durations	104	50	-4.5 ± 5.2
Task type	Speed or power	<30 s	4	25	-0.2 ± 1.1
		30–45 s	15	7	-0.6 ± 3.1
		60–120 s	23	48	-2.7 ± 4.7
		>120 s	2	50	-4.5 ± 6.4
		All durations	44	32	-1.8 ± 4.1
	Strength	<30 s	3	0	-2.3 ± 2.0
		30–45 s	8	50	-4.2 ± 2.7
		60–120 s	13	77	-7.0 ± 4.5
		>120 s	36	64	-7.1 ± 5.7
		All durations	60	62	-6.5 ± 5.1
Contraction mode	Concentric	≤45 s	5	40	-2.7 ± 3.1
		≥60 s	24	67	-5.2 ± 3.6
		All durations	29	62	-4.8 ± 3.6
	Isometric	≤45 s	8	38	-4.4 ± 2.7
		≥60 s	25	76	-8.9 ± 6.0
		All durations	33	67	-7.8 ± 5.7
	Eccentric	≤45 s	0	NA	NA
		≥60 s	6	50	-6.3 ± 5.8
		All durations	6	50	-6.3 ± 5.8
		All durations	4	25	-2.0 ± 3.1
Muscle group	Knee extensors	≤45 s	4	25	-2.0 ± 3.1
		≥60 s	25	64	-6.7 ± 4.5
		All durations	29	59	-6.0 ± 4.6
	Knee flexors	≤45 s	2	50	-4.2 ± 3.0
		≥60 s	11	82	-7.4 ± 3.8
		All durations	13	77	-6.9 ± 3.8
	Plantar flexors	≤45 s	1	0	-3.7
		≥60 s	13	62	-7.5 ± 7.8
		All durations	14	57	-7.2 ± 7.6

For seven studies where nonsignificant results were found but no data were provided, a nominal value of "0" was given for the mean reduction.

Dose-response relationship. To determine whether a dose-response effect of stretch was evident across the studies, we separated the research into groups where the total stretch duration imposed was either <30 or 30–45 s or either 1–2 or >2 min (Table 1). Surprisingly, only 10 studies reporting 11 findings, which examined the effects of stretch where duration was <30 s, were found. Nine studies did not reveal any significant reduction: five reported no change in power- or speed-dependent tasks including 20-m sprint time (8), vertical jump (19,50,76), and medicine ball throw (71); and two studies reported significant increases in five-step jump distance (2.5% [71]) and peak cycling power (5% [81]), although the last study failed to demonstrate appropriate control. Furthermore, three studies reported no significant reductions in maximal strength, including isometric plantar flexor maximum voluntary contraction (MVC) (52), handgrip strength (58), or isometric and concentric knee extensor MVC (95). Only one study reported a significant but small reduction in 20-m sprint velocity (-1.2% (38)), which is in conflict with Beckett et al. (8). Collectively, data from these studies demonstrate that short durations of stretch (<30 s) do not result in a meaningful reduction in muscular performance (pooled estimate = -1.1% ± 1.8%; Fig. 1).

When examining studies that used a longer total duration of stretch (30–45 s), 25 studies were found reporting 31 findings. Fifteen studies examined power- or speed-dependent performance, with only two studies reporting a significant reduction in vertical jump height (-4.2% [39], -4.3% [51]), although the latter study failed to

demonstrate appropriate control. In direct conflict with these findings, nine studies reported no significant reduction in vertical jump performance (18,31,32,42,57,62,86,103,120), with one study reporting a significant increase in jump performance (2.3% [75]). Furthermore, no significant effect was detected for 10- (62), 20- (97), or 30-m (18) sprint time, with a significant improvement in 20-m rolling sprint time reported (1.7% [62]), which reinforces the previous suggestion that short-duration stretch does not clearly influence maximal running performance. Also, no significant reductions were reported for throwing velocity (44), bench press and overhead throws (101), or leg extension power (114). Collectively, these data demonstrate no clear detrimental effect on performance in speed- and power-dependent tasks where stretch duration is 30–45 s (pooled estimate = -0.6% ± 3.1%; Fig. 1). This finding is especially important because the duration of stretch is reflective of normal preexercise routine practices (2,98) and the performance tasks examined are highly applicable to both clinical and athletic subjects.

Eleven studies examined the effects of 30–45 s of stretch on maximal strength, with equivocal findings being reported. Significant reductions were reported in handgrip strength (-7.8% [58], -6.7% [102]), concentric knee flexor MVC (-6.3% [109]), and isometric and concentric knee extensor MVC (-6.6% [95]). In contrast, three studies reported no significant effect on concentric knee extensor strength (9,121,122) following similar durations of stretch. Furthermore, no significant reductions were found in concentric plantar flexor MVC (1), chest press strength (9,74),




FIGURE 1—Mean percentage change ($*P < 0.05$, significant) in strength-, power-, and speed-dependent task performances after stretches of <30 s (top) or 30–45 s (bottom) in duration. Most studies found no significant reduction in muscle performance after shorter stretch durations, with small mean reductions across studies indicating no meaningful change in performance.

or isometric knee flexor MVC (82). Thus, while some studies have reported significant performance decrements in lower limb muscle groups, this is not a common finding. Overall, the majority of the findings suggest that no detrimental effect on strength is likely when stretch duration is 30–45 s (pooled estimate = $-4.2\% \pm 2.7\%$; Fig. 1).

When stretch durations were greater, the percentage of significant losses reported increased sharply after 60 s of stretch (61%) and then reached a plateau when stretch duration increased above 2 min, indicating a sigmoidal relationship (Fig. 2). This finding is congruent with previous dose–response studies (52,58,82,95,119). Clearly, the duration of stretch at which significant reductions are likely is approximately 60 s; however, longer durations (>2 min) did not further increase the likelihood of significant reductions. A linear relationship was evident in the average magnitude of reductions as the average reductions continued to increase with longer durations of stretch (Table 1).

Effect of contraction mode. Although most findings from studies using shorter static stretch durations indicated no significant effect, equivocal findings were reported in studies using longer durations (≥ 60 s). Accordingly, we examined whether stretch duration influenced results when studies were organized by muscle contraction mode (Table 1). Given that this reduced the sample size substantially, the four dose–response groups were merged into two (≤ 45 and ≥ 60 s). A similar proportion of studies reported significant reductions after ≥ 60 s stretch in concentric and isometric strength (67% and 76%, respectively); however, the size of the reductions was greater for isometric than for concentric (-8.9% and -5.2% , respectively; Table 1). The most interesting finding from this analysis was that only 6 of the 68 findings reported in studies examining the effect of contraction mode assessed changes in maximal eccentric strength (15,26,28,69,93,111) and all of these used stretch durations >60 s. Two studies reported significant force losses (-4.3% [15] and -9.7% [93]), whereas no change was reported in the remaining four studies that all used much longer stretch durations (3–9 min).

Muscle group–specific effects. A final analysis was conducted to determine whether the equivocal reports could be explained further by separating the studies by muscle




FIGURE 2—The sigmoidal relationship between (A) stretch duration and likelihood of a significant reduction and (B) the curvilinear relationship between stretch duration and the mean reduction in the performance of strength-, power-, and speed-dependent tasks. The likelihood of significant reductions was minimal after stretch durations of <30 s (14%) and 30–45 s (22%); this rose sharply after 1–2 min (61%) and then reached a plateau after >2 min (63%) of stretch. The average magnitude of losses also remained small for shorter-duration stretches (pooled estimate <30 s = $-1.1\% \pm 1.8\%$; 30–45 s = $-1.9\% \pm 3.4\%$) and then continued to increase with longer durations of stretch (pooled estimate 1–2 min = $-4.2\% \pm 5.0\%$; >2 min = $-7.0\% \pm 5.7\%$).

group. Most studies focused on lower limb strength, with few studies examining upper body strength; accordingly, studies measuring knee flexor, knee extensor, and plantar flexor strength were examined, and again, the dose–response groups were merged into two groups (≤ 45 and ≥ 60 s). Although similar findings were revealed across muscle groups for magnitude of loss (Table 1), the knee flexors (82%) seemed to be more regularly influenced by stretch compared with the knee extensors (64%) and plantar flexors (62%). This finding, in conjunction with the finding that the muscle contraction mode of the test exercise influenced the results, may partly explain the equivocal findings reported across the literature for longer-duration (≥ 60 s) stretches. However, although there is some evidence for a contraction mode– and muscle-specific effect, the lack of data does not allow firm conclusions to be drawn, and we cannot fully explain the equivocal findings reported for longer-duration stretches.

DISCUSSION

When all relevant studies are examined *in toto*, the results of the present review seem to largely agree with previous suggestions that acute static stretching can reduce maximal muscle performance (63,70,89,116). Forty-four percent of all variables included in our analyses (144 findings) from 106 studies showed significant reductions in maximal strength-, power-, or speed-dependent performance. However, a more detailed examination reveals clear evidence that no performance decrements in strength-, power-, or speed-dependent tasks occur when total stretch durations are < 45 s. Furthermore, there is only a moderate effect of stretch for durations > 60 s. We found there to be only minor differences in the effect across muscle contraction modes or muscle groups and no substantial effect of movement velocity.

Potential bias. We used a systematic review methodology to remove potential sources of bias as far as possible, although this procedure does not guarantee the absence of bias. Analyses such as those performed in the present review may be influenced by publication bias (100) because studies, reporting nonsignificant effects of stretch, may have been less likely to be accepted for publication. However, the potential inclusion of these studies would not have changed the main conclusion that shorter-duration (≤ 45 s) stretching has no effect on force production. Examination of the methodological quality of the literature revealed that experimental study design was often poor, where 30% of the studies reported no control group or reliability analyses. This supports the contention of Young (116), who previously highlighted this problem. Many studies did not include, or did not clearly report, a test reliability analysis, which is a major concern because it reduces the validity of the findings. Data presented in many of the included studies were collected during both control (rest) and experimental (stretch) conditions, and statistical analyses were then performed on the data sets to determine the level of significance between conditions. One problem, however, is that statistics for reli-

ability were rarely presented, so the potential exists for the magnitude of between-condition differences to have been within the limits of data variability, resulting from learning, motivation variability, fatigue, or some other external influence, and were not solely influenced by the stretch intervention. Nonetheless, several statistical methods to eliminate this problem, including comparison of mean tests (e.g., *t*-tests, ANOVA), intraclass correlation coefficients, and coefficients of variation (CV) to establish reliability from repeated testing during control conditions, were appropriately used by several researchers (39,106,121) and should provide an exemplar for future research. Regardless, and importantly, our analysis revealed that the removal of studies with the poorer design did not markedly affect the conclusions drawn from the review because a similar proportion of these studies reported significant versus nonsignificant results.

Acute effects of short-duration static stretch. The present systematic review revealed clear evidence that the widely reported negative effects of stretch on maximal strength performance are not apparent after stretch durations (≤ 30 s) (52,58,95) that are commonly performed in a pre-exercise routine (2,98), although there are a limited number of studies imposing this stretch duration. Nonetheless, equivocal results were found when durations increased to 30–45 s in knee extensor (9,95,121,122) and knee flexor MVC tests (82,109). Significant reductions were found in handgrip strength (58,102), but no change was found in plantar flexor MVC (1) or chest press one-repetition maximum (9,74). Examination of the literature revealed that, while some studies have reported significant losses in lower limb muscle groups, others did not. Overall, 50% of the findings indicated that no detrimental effect on strength was likely when stretch duration was 30–45 s, with the pooled estimate of the changes ($-4.2\% \pm 2.7\%$) well within the normal variability for maximum voluntary performance.

There was also clear evidence that stretch did not affect higher-speed force production when stretch durations were ≤ 45 s. Only two studies reported significant decreases in vertical jump height (39,51), with the latter failing to use an appropriate control. In direct conflict were 13 studies using similar durations of stretch that reported no significant reduction in jump performance (18,19,31,32,42,50,57,62,75,76,86,103,120). Similar patterns were evident in sprint performance, where again only one study reported a significant reduction (38), whereas four studies reported no significant reduction (8,18,62,97), and Little and Williams (62) reported an increase in sprint performance. Interestingly, Fletcher and Jones (38) did not use a control condition but determined reliability with intraclass correlation coefficient and CV calculations. The CV was calculated at 1.7%, which was greater than the significant difference reported; SEM was also similar in size to the reduction reported, and the effect size calculated from the reduction was small. Although the study design and the implementation of statistics were correct, the interpretation of their data and the practical

importance of their findings are debatable. Only 2 studies that demonstrated appropriate control or reliability reported a significant reduction in performance, as opposed to 15 studies that reported no difference in the same tasks and a further 5 studies reporting no difference in performance in other speed or power tests (44,71,81,101,114). Collectively, these data overwhelmingly indicate that there is no detrimental effect of short-duration static muscle stretch on speed- or power-dependent performance, with the pooled estimate of the change calculated at $-0.5\% \pm 2.8\%$.

Dose–response effects of stretch. The lack of consensus regarding the negative effects of static stretching is likely to be partly attributable to differences in the durations of stretch imposed across studies. Short-duration stretching tends not to result in significant impairments, whereas longer stretch duration more likely does, with the percentage of significant findings increasing concurrently with stretch duration (<30 s = 14%; 30–45 s = 22%; 1–2 min = 61%; >2 min = 63%). This is in agreement with several recent studies (52,58,82,90,95,119) that specifically examined the dose–response effect of static muscle stretch on active force production. For example, Ogura et al. (82) reported that 30 s of stretch did not reduce isometric knee flexor strength but that 60 s of stretch induced significant impairment, and Knudson and Noffal (58) found that repeated 10-s stretches did not reduce handgrip strength compared with control until 40 s of total stretch was accumulated. Similarly, 5, 15, and 20 s of static stretch did not significantly reduce isometric plantar flexor force, whereas 60 s of stretch did (52); the size of the force impairment was also significantly correlated with the stretch duration, clearly highlighting the importance of stretch duration in the magnitude of force loss. Those studies, and other evidence reported in the present review, indicate that a clear dose–response effect exists, with decrements becoming more likely for stretch durations ≥ 60 s but not continuing to increase beyond 2 min. Thus, the dose–response relationship seems to be sigmoidal, with turning points at approximately 60 s and 2 min (Fig. 2).

Interestingly, comparable dose–response trends were evident across tasks involving largely strength-, power-, or speed-dependent movements, which suggest that the effects of stretch duration are task independent. However, the number (percentage) of significant findings and the magnitude of the performance decrement were larger for strength-based than power- and speed-based tasks. Given that power- and speed-dependent tasks are more typically performed in activities of daily living or athletic pursuits than the laboratory-based slow-speed strength tests, these findings perhaps have more practical relevance. Regardless, the finding that short-duration stretches (≤ 45 s) did not seem to impair muscle force production is of even greater practical importance. This important finding suggests that static muscle stretching can be safely used in a preexercise routine without compromising physical performance, whereas longer durations (≥ 60 s) are more likely to be problematic.

Although most short-duration studies (≤ 45 s) revealed no significant change, significant improvements were reported in jumping (71,75), cycling (81), and sprinting (62) performances, which suggest that improvements are possible in some tasks. Furthermore, significant improvements in ROM and reduced musculotendinous stiffness after short-duration stretches (5–30 s) have also been reported (6,52), which may reduce muscle strain injury risk. Thus, the inclusion of short-duration preperformance stretching may be deemed useful by some practitioners, although more research is needed to clarify the effects of short-duration static stretching.

Although a similar influence was seen across muscle groups (lower limb) and contraction modes, no studies exist detailing the effects of moderate-duration stretches (≤ 45 s) on eccentric strength. This is important not only for its physical performance implications but also for its effect on injury risk. Muscle strength has been cited as a major influencing factor within the etiology of muscle strain injury (83), and with most muscle strain injuries suggested to occur within normal ROM during eccentric loading, the ability of the muscle to withstand eccentric loading may be crucial to injury risk. Given the equivocal data reported from much longer durations of stretch (e.g., >60 s) on eccentric strength and that there are presently no data describing the effects of shorter, more practically relevant, stretch durations (≤ 45 s), a clear research focus is needed to fully explore the influence of stretch on the muscle's ability to withstand eccentric loading.

CONCLUSIONS

Static muscle stretches totaling <45 s can be used in preexercise routines without risk of significant decreases in strength-, power-, or speed-dependent task performances. Longer stretch durations (e.g., ≥ 60 s) are more likely to cause a small or moderate reduction in performance. Interestingly, the effect of stretch on performances across a range of muscle contraction modes, muscle groups, and movement speeds was similar. Importantly, no studies exist detailing the effects of moderate-duration stretches (≤ 45 s) on eccentric strength, and there is little evidence for an effect after longer periods of stretch. This is important because of the purported influence of eccentric strength on both movement performance and injury risk. Several avenues of further research exist, including an examination of the effects of stretch on upper body musculature and on eccentric movement performance, and more data are required to determine the effect of short-duration stretches (≤ 30 s) to more clearly delineate the magnitude of effect. A comprehensive review of the existing literature examining the influence of other forms of muscle stretching (dynamic, proprioceptive neuromuscular facilitation, and ballistic) should also be performed because the effects of different stretching modalities are likely to be different. Finally, no attempt was made in the present review to determine whether the number of stretches

performed, in addition to the total duration of stretch, is a factor influencing the effects of stretch, so future reviews are required to clarify whether it is a factor influencing the stretch-induced loss of force.

A.D.K. performed the literature search, selected articles for exclusion and inclusion, assessed the risk of bias, extracted the data, and

performed the analysis. A.J.B. verified a percentage of articles selected for exclusion, verified all articles selected for inclusion, verified the extracted data, and assessed the risk of bias. Both authors were involved in the study design, contributed to the writing and revision of the manuscript, and are able to take responsibility for its accuracy.

No funding was received for this work.

The results of the present study do not constitute endorsement by the American College of Sports Medicine.

REFERENCES

- Alpkaya U, Koceja D. The effects of acute static stretching on reaction time and force. *J Sports Med Phys Fit.* 2007;47:147–50.
- Alter MJ. *Science of Flexibility.* 3rd ed. Champaign (IL): Human Kinetics; 2004. p. 9.
- American College of Sports Medicine. *ACSM's Resource Manual for Guidelines for Exercise Testing and Prescription.* 8th ed. Philadelphia (PA): Lippincott, Williams & Wilkins; 2010. p. 173.
- Babault N, Kouassi BYL, Debrosses K. Acute effects 15 min static stretch or contract-relax stretching modalities on plantar flexors neuromuscular properties. *J Sci Med Sport.* 2010;13:247–52.
- Bacurau RFP, Monteiro GA, Ugrinowitsch C, Tricoli V, Cabral LF, Aoki MS. Acute effect of a ballistic and a static stretching exercise bout on flexibility and maximal strength. *J Strength Cond Res.* 2009;23:304–8.
- Bandy WD, Irion JM, Briggler M. The effect of time and frequency of static stretching on flexibility of the hamstring muscles. *Phys Ther.* 1997;77:1090–6.
- Bazett-Jones DM, Winchester JB, McBride JM. Effect of potentiation and stretching on maximal force, rate of force development, and range of motion. *J Strength Cond Res.* 2005; 19:421–6.
- Beckett JRJ, Schneiker KT, Wallman KE, Dawson BT, Guelfi KJ. Effects of static stretching on repeated sprint and change of direction performance. *Med Sci Sports Exerc.* 2009;41(2):444–50.
- Beedle B, Rytter SJ, Healy RC, Ward TR. Pretesting static and dynamic stretching does not affect maximal strength. *J Strength Cond Res.* 2008;22:1838–43.
- Behm DG, Bambury A, Cahill F, Power K. Effect of acute static stretching on force, balance, reaction time, and movement time. *Med Sci Sports Exerc.* 2004;36(8):1397–402.
- Behm DG, Bradbury EE, Haynes AT, Hodder JN, Leonard AM, Paddock NR. Flexibility is not related to stretch-induced deficits in force or power. *J Sports Sci Med.* 2006;5:33–42.
- Behm DG, Button DC, Butt JC. Factors affecting force loss with prolonged stretching. *Can J Appl Physiol.* 2001;26:261–72.
- Behm DG, Kibele A. Effects of differing intensities of static stretching on jump performance. *Eur J Appl Physiol.* 2007; 101:587–94.
- Bradley PS, Olsen PD, Portas MD. The effect of static, ballistic, and proprioceptive neuromuscular facilitation stretching on vertical jump performance. *J Strength Cond Res.* 2007;21:223–6.
- Brandenburg JP. Duration of stretch does not influence the degree of force loss following static stretching. *J Sports Med Phys Fit.* 2006;46:526–34.
- Brandenburg JP, Pitney WA, Luebbbers PE, Veera A, Czajka A. Time course of changes in vertical-jumping ability after static stretching. *Int J Sports Physiol Perf.* 2007;2:170–81.
- Burkett LN, Phillips WT, Ziuraitis J. The best warm-up for the vertical jump in college-age athletic men. *J Strength Cond Res.* 2005;19:673–6.
- Chaouachi A, Castagna C, Chtara M, et al. Effect of warm-ups involving static or dynamic stretching on agility, sprinting, and jumping performance in trained individuals. *J Strength Cond Res.* 2010;24:2001–11.
- Church JB, Wiggins MS, Moode EM, Crist R. Effect of warm-up and flexibility treatments on vertical jump performance. *J Strength Cond Res.* 2001;15:332–6.
- Cornwell A, Nelson AG, Heise GD, Sidaway B. Acute effects of passive muscle stretching on vertical jump performance. *J Hum Mov Stud.* 2001;40:307–24.
- Cornwell A, Nelson AG, Sidaway B. Acute effects of stretching on the neuromechanical properties of the triceps surae muscle complex. *Eur J Appl Physiol.* 2002;86:428–34.
- Costa EC, dos Santos CM, Prestes J, da Silva J, Knackfuss MI. Acute effects of static stretching on the strength performance of jiu-jitsu athletes in horizontal bench press. *Fit Perf J.* 2009; 8:212–7.
- Costa PB, Ryan ED, Herda TJ, DeFreitas JM, Beck TW, Cramer JT. Effects of static stretching on the hamstrings-to-quadriceps ratio and electromyographic amplitude in men. *J Sports Med Phys Fit.* 2009;49:401–9.
- Costa PB, Ryan ED, Herda TJ, DeFreitas JM, Beck TW, Cramer JT. Effects of stretching on peak torque and the H:Q ratio. *Int J Sports Med.* 2009;30:60–5.
- Cramer JT, Beck TW, Housh TJ, et al. Acute effects of static stretching on characteristics of the isokinetic angle-torque relationship, surface electromyography, and mechanomyography. *J Sports Sci.* 2007;25:687–98.
- Cramer JT, Housh TJ, Coburn JW, Beck TW, Johnson GO. Acute effects of static stretching on maximal eccentric torque production in women. *J Strength Cond Res.* 2006;20:354–8.
- Cramer JT, Housh TJ, Johnson GO, Miller JM, Coburn JW, Beck TW. Acute effects of static stretching on peak torque in women. *J Strength Cond Res.* 2004;18:236–41.
- Cramer JT, Housh TJ, Johnson GO, Weir JP, Beck TW, Coburn JW. An acute bout of static stretching does not affect maximal eccentric isokinetic peak torque, the joint angle at peak torque, mean power, electromyography, or mechanomyography. *J Orthop Sports Phys Ther.* 2007;37:130–9.
- Cramer JT, Housh TJ, Weir JP, Johnson GO, Coburn JW, Beck TW. The acute effects of static stretching on peak torque, mean power output, electromyography, and mechanomyography. *Eur J Appl Physiol.* 2005;93:530–9.
- Cronin J, Nash M, Whatman C. The acute effects of hamstring stretching and vibration on dynamic knee joint range of motion and jump performance. *Phys Ther Sport.* 2008;9:89–96.
- Curry BS, Chengkalath D, Crouch GJ, Romance M, Manns PJ. Acute effects of dynamic stretching, static stretching, and light aerobic activity on muscular performance in women. *J Strength Cond Res.* 2009;23:1811–9.
- Dalrymple KJ, Davis SE, Dywer GB, Moir GL. Effect of static and dynamic stretching on vertical jump performance in collegiate women volleyball players. *J Strength Cond Res.* 2010; 24:149–55.
- Di Cagno A, Baldari C, Battaglia C, et al. Preexercise static stretching effect on leaping performance in elite rhythmic gymnasts. *J Strength Cond Res.* 2010;24:1995–2000.
- Egan AD, Cramer JT, Massey LL, Marek SM. Acute effects of static stretching on peak torque and mean power output in national collegiate athletic association division I women's basketball players. *J Strength Cond Res.* 2006;20:778–82.
- Evetovich TK, Cain RM, Hinnerichs KR, Engebretsen BJ, Conley DS. Interpreting normalized and nonnormalized data after

- acute static stretching in athletes and nonathletes. *J Strength Cond Res.* 2010;24:1988–94.
36. Evetovich TK, Nauman NJ, Conley DS, Todd JB. Effect of stretching of the biceps brachii on torque, electromyography, and mechanomyography during concentric isokinetic muscle actions. *J Strength Cond Res.* 2003;17:484–8.
 37. Favero J, Midgley AW, Bentley DJ. Effects of an acute bout of static stretching on 40 m sprint performance: influence of baseline flexibility. *Res Sports Med.* 2009;17:50–60.
 38. Fletcher IM, Jones B. The effect of different warm-up stretch protocols on 20 meter sprint performance in trained rugby union players. *J Strength Cond Res.* 2004;18:885–8.
 39. Fletcher IM, Monte-Colombo MM. An investigation into the possible physiological mechanisms associated with changes in performance related to acute responses to different pre-activity stretch modalities. *Appl Physiol Nutri Metab.* 2010;35:27–34.
 40. Fowles JR, Sale DG, MacDougall JD. Reduced strength after passive stretch of the human plantar flexors. *J Appl Physiol.* 2000;90:1179–88.
 41. Gavin J, Morse C. The acute effects of static and dynamic stretching on passive torque, maximal voluntary contraction and range of motion in the plantarflexors. *SportEX dyna.* 2009;21:18–23.
 42. González-Ravé JM, Machado L, Navarro-Valdivielso F, Vilas-Boas JP. Acute effects of heavy-load exercises, stretching exercises, and heavy-load plus stretching exercises on squat jump and countermovement jump performance. *J Strength Cond Res.* 2009;23:472–9.
 43. Gurjão ALD, Gonçalves R, De Moura RF, Gobbi S. Acute effects of static stretching on rate of force development and maximal voluntary contraction in older women. *J Strength Cond Res.* 2009;23:2149–54.
 44. Haag SJ, Wright GA, Gillette CM, Greany JF. Effects of acute static stretching of the throwing shoulder on pitching performance of national collegiate athletic association division III baseball players. *J Strength Cond Res.* 2010;24:452–7.
 45. Handrakis JP, Southard VN, Abreu JM, et al. Static stretching does not impair performance in active middle-aged adults. *J Strength Cond Res.* 2010;24:825–30.
 46. Herda TJ, Cramer JT, Ryan ED, McHugh MP, Stout JR. Acute effects of static versus dynamic stretching on isometric peak torque, electromyography, and mechanomyography of the biceps femoris muscle. *J Strength Cond Res.* 2008;22:809–17.
 47. Herda TJ, Ryan ED, Costa PB, et al. Acute effects of passive stretching and vibration on the electromechanical delay and musculotendinous stiffness of the plantar flexors. *Electromyogr Clin Neurophysiol.* 2010;50:277–88.
 48. Herda TJ, Ryan ED, Smith AE, et al. Acute effects of passive stretching vs vibration on the neuromuscular function of the plantar flexors. *Scand J Med Sci Sports.* 2009;19:703–13.
 49. Higgins J, Green S. *Reviews Including Non-Randomised Studies. Cochrane Handbook for Systematic Reviews of Interventions.* Hoboken (NJ): Wiley; 2006. p. 99.
 50. Holt BW, Lambourne K. The impact of different warm-up protocols on vertical jump performance in male collegiate athletes. *J Strength Cond Res.* 2008;22:226–9.
 51. Hough PA, Ross EZ, Howartson G. Effects of dynamic and static stretching on vertical jump performance and electromyographic activity. *J Strength Cond Res.* 2009;23:507–12.
 52. Kay AD, Blazevich AJ. Reductions in active plantar flexor moment are significantly correlated with static stretch duration. *Eur J Sport Sci.* 2008;8:41–6.
 53. Kay AD, Blazevich AJ. Moderate-duration static stretch reduces active and passive plantar flexor moment but not Achilles tendon stiffness or active muscle length. *J Appl Physiol.* 2009;106:1249–56.
 54. Kay AD, Blazevich AJ. Isometric contractions reduce plantar flexor moment, Achilles tendon stiffness and neuromuscular activity but remove the subsequent effects of stretch. *J Appl Physiol.* 2009;107:1181–9.
 55. Kay AD, Blazevich AJ. Concentric muscle contractions before static stretching minimize, but do not remove, stretch-induced force deficits. *J Appl Physiol.* 2010;108:637–45.
 56. Kistler BM, Walsh MS, Horn TS, Cox RH. The acute effects of static stretching on the sprint performance of collegiate men in the 60- and 100-m dash after a dynamic warm-up. *J Strength Cond Res.* 2010;24:2280–4.
 57. Knudson D, Bennett K, Corn R, Leick D, Smith C. Acute effects of stretching are not evident in the kinematics of the vertical jump. *J Strength Cond Res.* 2001;15:98–101.
 58. Knudson D, Noffal G. Time course of stretch-induced isometric strength deficits. *Eur J Appl Physiol.* 2005;94:348–51.
 59. Kokkonen J, Nelson AG, Cornwell A. Acute muscle stretching inhibits maximal strength performance. *Res Q Sport Exerc.* 1998;69:411–5.
 60. Kubo K, Kanehisa H, Kawakami Y, Fukunaga T. Influence of static stretching on viscoelastic properties of human tendon structures *in vivo*. *J Appl Physiol.* 2001;90:520–7.
 61. La Torre A, Castagna C, Gervasoni E, et al. Acute effects of static stretching on squat jump performance at different knee starting angles. *J Strength Cond Res.* 2010;24:687–94.
 62. Little T, Williams AG. Effects of different stretching protocols during warm-ups on high-speed motor capacities in professional soccer players. *J Strength Cond Res.* 2006;20:203–7.
 63. Magnusson P, Renström P. The European College of Sports Sciences Position statement: the role of stretching exercises in sports. *Eur J Sport Sci.* 2006;6:87–91.
 64. Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M. A description of the trials, reviews, and practice guidelines indexed in the PEDro database. *Phys Ther.* 2008;88:1068–77.
 65. Maisetti O, Sastre J, Lecompte J, Portero P. Differential effects of an acute bout of passive stretching on maximal voluntary torque and the rate of torque development of the calf muscle-tendon unit. *Isokin Exerc Sci.* 2007;15:11–7.
 66. Manoel ME, Harris-Love MO, Danoff JV, Miller TA. Acute effects of static, dynamic, and proprioceptive neuromuscular facilitation stretching on muscle power in women. *J Strength Cond Res.* 2008;22:1528–34.
 67. Marek SM, Cramer JT, Fincher AL, et al. Acute effects of static and proprioceptive neuromuscular facilitation muscle strength and power output. *J Athl Traini.* 2005;40:94–103.
 68. McBride JM, Deane R, Mimphuis S. Effect of stretching in agonist-antagonist muscle activity and muscle force output during single and multiple joint isometric contractions. *Scand J Med Sci Sports.* 2007;17:54–60.
 69. McHugh M, Nesse M. Effects of stretch on strength loss and pain after eccentric exercise. *Med Sci Sports Exerc.* 2008;40:566–73.
 70. McHugh MP, Cosgrave CH. To stretch or not to stretch: the role of stretching in injury prevention and performance. *Scand J Med Sci Sports.* 2010;20:169–81.
 71. McMillian DJ, Moore JH, Hatler BS, Taylor DC. Dynamic vs. static-stretching warm up: the effect on power and agility performance. *J Strength Cond Res.* 2006;20:492–9.
 72. McNeal JR, Sands WA. Acute static stretching reduces lower extremity power in trained children. *Pediatr Exerc Sci.* 2003;15:139–45.
 73. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ.* 2009;339:332–6.
 74. Molacek ZD, Conley DS, Evetovich TK, Hinnerichs KR. Effects of low- and high-volume stretching on bench press performance

- in collegiate football players. *J Strength Cond Res.* 2010;24:711–6.
75. Murphy JR, Di Santo MC, Alkanani T, Behm DG. Aerobic activity before and following short-duration static stretching improves range of motion and performance vs. a traditional warm-up. *Appl Physiol Nutr Metab.* 2010;35:679–90.
 76. Murphy JC, Nagle E, Robertson RJ, McCrory JL. Effect of single set dynamic and static stretching exercises on jump height in college age recreational athletes. *Int J Exerc Sci.* 2010;3:214–24.
 77. Nelson AG, Allen JD, Cornwell A, Kokkonen J. Inhibition of maximal voluntary isometric torque production by acute stretching is joint-angle specific. *Res Q Exerc Sport.* 2001;72:68–70.
 78. Nelson AG, Guillory IK, Cornwell A, Kokkonen J. Inhibition of maximal voluntary isometric torque production by acute stretching is velocity specific. *J Strength Cond Res.* 2001;15:241–6.
 79. Nelson AG, Driscoll NM, Landin DK, Young MA, Schexnayder IC. Acute effects of passive muscle stretching on sprint performance. *J Sports Sci.* 2005;23:449–54.
 80. Nelson AG, Kokkonen J, Eldredge C. Strength inhibition following an acute stretch is not limited to novice stretchers. *Res Q Exerc Sport.* 2005;76:500–6.
 81. O'Connor DM, Crowe MJ, Spinks WL. Effects of static stretching on leg power during cycling. *J Sports Med Phys Fit.* 2006;46:52–6.
 82. Ogura Y, Miyahara Y, Naito H, Katamoto S, Aoki J. Duration of static stretching influences muscle force production in hamstring muscles. *J Strength Cond Res.* 2007;21:788–92.
 83. Orchard J, Marsden J, Lord S, Garlick D. Preseason hamstring muscle weakness associated with hamstring muscle injury in Australian footballers. *Am J Sports Med.* 1997;25:81–5.
 84. Papadopoulos C, Kalapotharakos VI, Noussios G, Meliggas K, Gantiraga E. The effect of static stretching on maximal voluntary contraction and force–time curve characteristics. *J Sports Rehabil.* 2006;15:185–94.
 85. Papadopoulos G, Siatras TA, Kellis S. The effect of static and dynamic stretching exercises on the maximal isokinetic strength of the knee extensors and flexors. *Isokinet Exerc Sci.* 2005;13:285–91.
 86. Power K, Behm D, Cahill F, Carroll M, Young W. An acute bout of static stretching: effects on force and jumping performance. *Med Sci Sports Exerc.* 2004;36(8):1389–96.
 87. Robbins JW, Scheuermann BW. Varying amounts of acute static stretching and its effect on vertical jump performance. *J Strength Cond Res.* 2008;22:781–6.
 88. Rossi LP, Pereira R, Simão R, Brandalize M, Gomes ARS. Influence of static stretching duration on quadriceps force development and electromyographic activity. *Hum Mov.* 2010;11:137–43.
 89. Rubini EC, Costa AL, Gomes PS. The effects of stretching on strength performance. *Sports Med.* 2007;37:213–24.
 90. Ryan ED, Beck TW, Herda TJ, et al. Do practical durations of stretching alter muscle strength? A dose–response study. *Med Sci Sports Exerc.* 2008;40(8):1529–37.
 91. Samuel MN, Holcomn WR, Guadagnoli MA, Rubley MD, Wallmann H. Acute effects of static and ballistic stretching on measures of strength and power. *J Strength Cond Res.* 2008;22:1422–8.
 92. Sayers AL, Farley RS, Fuller DK, Jubenville CB. The effect of static stretching on phases of sprint performance in elite soccer players. *J Strength Cond Res.* 2008;22:1416–21.
 93. Sekir U, Arabaci R, Akova B, Kadagan SM. Acute effects of static and dynamic stretching on leg flexor and extensor isokinetic strength in elite women athletes. *Scand J Med Sci Sports.* 2010;20:268–81.
 94. Shrier I. Does stretching improve performance? A systematic and critical review of the literature. *Clin J Sport Med.* 2004;14:267–73.
 95. Siatras TA, Mittas VP, Mameletzi DN, Vamvakoudis EA. The duration of the inhibitory effects with static stretching on quadriceps peak torque production. *J Strength Cond Res.* 2008;22:40–6.
 96. Siatras TA, Papadopoulos G, Mameletzi D, Gerodimos V, Kellis S. Static and dynamic acute stretching effect on gymnasts' speed in vaulting. *Pediatr Exerc Sci.* 2003;15:383–91.
 97. Sim AY, Dawson BT, Guelfi KJ, Wallman KE, Young WB. Effects of static stretching in warm-up on repeated sprint performance. *J Strength Cond Res.* 2009;23:2155–62.
 98. Taylor DC, Dalton JD, Seabar AV, Garrett WE. Viscoelastic properties of muscle–tendon units. *Am J Sports Med.* 1990;18:300–9.
 99. Thacker SB, Gilchrist J, Stroup DF, Kimsey CD. The impact of stretching on sports injury risk: a systematic review of the literature. *Med Sci Sports Exerc.* 2004;36(3):371–8.
 100. Thornton A, Lee P. Publication bias in meta-analysis: its causes and consequences. *J Clin Epidemiol.* 2000;53:207–16.
 101. Torres EM, Kraemer WJ, Vingren JL, et al. Effects of stretching on upper-body muscular performance. *J Strength Cond Res.* 2008;22:1279–85.
 102. Torres JB, Conceição MCSC, Sampaio AO, Dantas EHM. Acute effects of static stretching on muscle strength. *Biomed Human Kin.* 2009;1:52–5.
 103. Unick J, Kieffer HS, Cheeseman W, Feeney A. The acute effects of static and ballistic stretching on vertical jump performance in trained women. *J Strength Cond Res.* 2005;19:206–12.
 104. Vetter RE. Effects of six warm-up protocols on sprint and jump performance. *J Strength Cond Res.* 2007;21:819–23.
 105. Viale F, Nana-Ibrahim S, Martin RJF. Effect of active recovery on acute strength deficits induced by passive stretching. *J Strength Cond Res.* 2007;21:1233–7.
 106. Wallmann HW, Mercer JA, Landers MR. Surface electromyographic assessment of the effect of dynamic activity and dynamic activity with static stretching of the gastrocnemius on vertical jump performance. *J Strength Cond Res.* 2008;22:787–93.
 107. Wallmann HW, Mercer JA, McWhorter JW. Surface electromyographic assessment of the effects of static stretching of the gastrocnemius on vertical jump performance. *J Strength Cond Res.* 2005;19:684–8.
 108. Weir DE, Tingley E, Elder J, Geoffrey CB. Acute passive stretching alters the mechanical properties of human plantar flexors and the optimal angle for maximal voluntary contraction. *Eur J Appl Physiol.* 2005;93:614–23.
 109. Winchester JB, Nelson AG, Kokkonen J. A single 30-s stretch is sufficient to inhibit maximal voluntary stretch. *Res Q Exerc Sport.* 2009;80:257–61.
 110. Winchester JB, Nelson AG, Landin D, Young MA, Schexnayder IC. Static stretching impairs sprint performance in collegiate track and field athletes. *J Strength Cond Res.* 2008;22:13–8.
 111. Winke MR, Jones NB, Berger CG, Yates JW. Moderate static stretching and torque production of the knee flexors. *J Strength Cond Res.* 2010;24:706–10.
 112. Witvrouw E, Mahieu N, Danneels L, McNair P. Stretching and injury prevention: an obscure relationship. *Sports Med.* 2004;34:443–9.
 113. Woods K, Bishop P, Jones E. Warm-up and stretching in the prevention of muscular injury. *Sports Med.* 2007;37:1089–99.
 114. Yamaguchi T, Ishii K. Effects of static stretching for 30 seconds and dynamic stretching on leg extension power. *J Strength Cond Res.* 2005;19:677–83.
 115. Yamaguchi T, Ishii K, Yamaga M, Yasuda K. Acute effect of static stretching on power output during concentric dynamic constant external resistance leg extension. *J Strength Cond Res.* 2006;20:804–10.
 116. Young WB. The use of static stretching in warm-up for training and competition. *Int J Sports Physiol Perf.* 2007;2:212–6.

117. Young WB, Behm DG. Effects of running, static stretching and practice jumps on explosive force production and jumping performance. *J Sports Med Phys Fit.* 2003;43:21–7.
118. Young WB, Clothier P, Otago L, Bruce L, Liddel D. Acute effects of static stretching on hip flexor and quadriceps flexibility, range of motion and foot speed in kicking a football. *J Sci Med Sport.* 2004;7:23–31.
119. Young WB, Elias G, Power J. Effects of static stretching volume and intensity on plantar flexor explosive force production and range of motion. *J Sports Med Phys Fit.* 2006;46:403–11.
120. Young WB, Elliott S. Acute effects of static stretching, proprioceptive neuromuscular facilitation stretching, and maximum voluntary contractions on explosive force production and jumping performance. *Res Q Exerc Sport.* 2001;72:273–9.
121. Zakas A, Doganis G, Galazoulas C, Vamvakoudis E. Effect of acute static stretching duration on isokinetic peak torque in pubescent soccer players. *Pediatr Exerc Sci.* 2006;18:252–61.
122. Zakas A, Doganis G, Papakonstandinou V, Sentelidis T, Vamvakoudis E. Acute effects of static stretching duration on isokinetic peak torque production of soccer players. *J Bodyw Mov Ther.* 2006;10:89–95.
123. Zakas A, Galazoulas C, Doganis G, Zakas N. Effect of two acute static stretching durations of the rectus femoris muscle on isokinetic peak torque in professional soccer players. *Isokinet Exerc Sci.* 2006;14:357–62.