

Effect of bioactive compounds from Sainfoin (*Onobrychis viciifolia* Scop.) on the *in vitro* larval migration of *Haemonchus contortus*: role of tannins and flavonol glycosides

E. BARRAU¹, N. FABRE¹, I. FOURASTE¹ and H. HOSTE^{2*}.

¹: Unité Mixte de Recherches 152, IRD /UPS "Pharmacochimie des substances naturelles et pharmacophores redox" Faculté des Sciences Pharmaceutiques, Université Toulouse III, 35 Chemin des Maraîchers, F31062 Toulouse Cedex 4. France.

²: Unité Mixte de Recherches 1225 INRA/DGER, "Interactions Hôte Agents Pathogènes". École Nationale Vétérinaire de Toulouse, 23 chemin des Capelles, F31076 Toulouse. France.

Running title : Effect of sainfoin tannins and flavonol glycosides on *Haemonchus contortus*

*Corresponding author: H. HOSTE

Unité Mixte de Recherches 1225 INRA/DGER, "Interactions Hôte Agents Pathogènes". École Nationale Vétérinaire de Toulouse, 23 Chemin des Capelles, F31076 Toulouse. France.

Email: h.hoste@envt.fr Tel: +(33) 05-61-19-38-75 Fax: +(33) 05-61-19-32-43.

SUMMARY

Anthelmintic bioactivity against gastrointestinal nematodes has been associated with leguminous forages supporting the hypothesis of a role of condensed tannins. However, the possibility that other compounds might also been involved has received less consideration. Using bio-guided fractionation, the current study aimed at characterising the biochemical nature of the active compounds present in sainfoin (*Onobrychis viciifolia*), previously identified as an anthelmintic leguminous forage. The effects of sainfoin extracts were evaluated on 3rd stage larvae (L3) of *Haemonchus contortus* by using a larval migration inhibition (LMI) assay. Comparison of extracts obtained with several solvent systems showed that the bioactivity was associated with the 70:30 acetone/water extract. Further fractionation of the later allowed the separation of phenolic compounds. By use of a dialysis method, compounds were separated with a molecular weight cut-off of 2000 Da. The *in vitro* anthelmintic effects of the fraction with condensed tannins was confirmed. In the fraction containing molecules of MW < 2000 Da, 3 flavonol glycosides were identified as rutin, nicotiflorin and narcissin. At 1200 µg/ml, each inhibited significantly migration of larvae. Addition of polyvinyl pyrrolidone (PVPP) to both fractions before incubation restore larval migration. These results confirmed the role of both tannins and flavonol glycosides in the anthelmintic properties of sainfoin.

Key words: Sainfoin (*Onobrychis viciifolia* Scop.), Bio-guided fractionation, *Haemonchus contortus*, Tannins, Flavonol glycosides.

INTRODUCTION

Parasitic nematodes are of major economic importance in livestock, and anthelmintics have to be employed to reduce the production losses caused by them. However, because of the widespread development of resistance to broad-spectrum anthelmintics, the control of gastrointestinal parasites based on chemical substances is now severely impaired (Roos, Kwa and Grant, 1995; Waller, 1997; Jackson and Coop, 2000). Therefore, there is an urgent need to seek alternative or complementary solutions to the control of parasitic nematodes of ruminants (Niezen *et al.*, 1996; Waller 1999). One of these novel approaches is represented by the use of nutraceutical forages (Waller and Thamsborg, 2004). Several legume forages (*Hedysarum coronarium* L., *Onobrychis viciifolia* Scop., *Lotus pedunculatus* Cav. or *Lotus corniculatus* L.) containing condensed tannins have been reported to have anthelmintic properties. This was shown either through *in vivo* or *in vitro* studies (see reviews by Kahn and Diaz-Hernandez, 2000; Athanasiadou, Kyriazakis and Jackson, 2003; Paolini and Hoste, 2003c; Min and Hart, 2003). The strategic use of condensed-tannin forages in grazing management may therefore, lead to a promising solution for the control of parasite infections both in conventional and in organic farming systems. However, some variability in results has been described which related either to the nematode species or to the sources of tannins. This emphasizes the need to understand the mechanisms of action of the plant secondary metabolites and to better characterise the bioactive compounds associated with the anthelmintic properties of these forages.

To date, the anthelmintic properties of tanniferous legume forages have mainly been related to their condensed tannins content. This conclusion was drawn from studies which demonstrated 1) the bioactivity on nematodes of condensed tannins purified from plants and 2) the inhibition of these antiparasitic effects by addition of specific inhibitors of tannins to plant extracts (Molan *et al.*, 2000b). In addition, Molan *et al.*, 2003 have recently shown that flavan-3-ols and flavan-3-ol gallates, the basic units of condensed tannin polymers, presented an inhibitory activity on egg hatching, larval development and the viability of *Trichostrongylus colubriformis* 3rd stage larvae (L3). Some authors have suggested that other plant compounds, in particular plant secondary metabolites, might also affect the biology of worms (Athanasiadou *et al.*, 2003; Molan *et al.*, 2003). However, no study has investigated in detail the role of the different fractions isolated from a bioactive forage to

determine whether and which compounds other than tannins might be associated with the effects.

Sainfoin is a tanniferous legume forages (Bate-Smith, 1973; Koupay-Abyazani *et al.*, 1993) which has shown positive effects against various parasitic nematode species in field studies (Paolini, Dorchies and Hoste, 2003b; Thamsborg *et al.*, 2003). In addition *in vitro* experiments have demonstrated that extracts of sainfoin showed an inhibitory activity against L3 of gastrointestinal nematodes, as measured by the larval migration inhibition assay (Molan *et al.*, 2000b; Paolini, Fouraste and Hoste, 2004). Sainfoin contains tannins but also other phenolic compounds, such as flavonoid glycosides (Lu *et al.*, 2000; Marais *et al.*, 2000).

The aim of the current study was to fractionate extracts of sainfoin in order to characterise the biochemical nature of the active compound(s) with anthelmintic effects and better to understand the mechanisms of action. *Haemonchus contortus* was used as the parasitic model, using a larval migration inhibition assay (LMI), to investigate the anthelmintic properties.

MATERIALS AND METHODS

Plant materials

Sainfoin, *Onobrychis viciifolia* Scop. cv. "Fakir" and ryegrass, *Lolium perenne* L. were harvested at the end of spring 2002 in the south east of France, and air-dried at room temperature. Rye grass hay was used as a negative control because of its low percentage of tannins. The water contents of the hays were 16 % for the sainfoin and 33% for the gramineous hay at the time of extraction.

Experimental design and biochemical procedure

In a first step, we compared results between different extracts of Sainfoin and ryegrass hays, with various solvents in order to determine the polarity of the active compounds. In a second step, we examined the best solvent to use to improve the extraction of those active components from sainfoin. In a third step, we used activity guided fractionation to isolate and characterize the biochemical substances presenting the anthelmintic properties.

Preparation of extracts : First, a "successive step extraction" was applied to determine the polarity of the active compounds. Hexane, methylene chloride, 30:70 alcohol/water (v/v)

and distilled water were used as solvents, known to present different polarity, in order to extract successively : lipids, sterols, isoflavons, polyphenols, vitamins, glycosides compounds, tannins, amino acids, proteins, sugars, mucilages. One hundred grams of ground hay was extracted with 2 x 600 ml of solvent at room temperature for 30 min. The extracts were concentrated under low pressure at 35 °C. Water was removed by freeze-drying.

Improvement of the extraction yield : After the polarity of the active compounds was determined, we focused on improving the yield of extraction by using another appropriate solvent. An acetone extract was then used for comparison with 30:70 alcohol/water. Five hundreds grams of dry, ground plant were extracted with 2 x 3 l of 70:30 acetone/water (v/v) containing ascorbic acid (1 g.l⁻¹). The acetone was removed under low pressure (at temperature < 35 °C), and the aqueous solution was washed four times with 500 ml methylene chloride to remove chlorophyll and lipids. The extract was freeze-dried. Compared with the 30:70 alcohol/water (v/v) (yield 15 %), the yield of extraction with 70:30 acetone/water was found to be approximately 30%.

Fractionation : In order to continue the identification of the active compounds, we have further fractionated the active extract using a polarity method. The defatted 70% aqueous acetone extract (300 g) was suspended in water and subjected to chromatography over a C-18 silica column using first 100% H₂O and then 100% MeOH. The results of fractionation was monitored by thin layer chromatography (TLC) on Al plates coated with Kieselgel 60F₂₅₄ (Merck), developed in CHCl₃/MeOH/H₂O, 65 : 45 : 10 v/v/v and detected using UV and vanillin sulphuric acid spray reagent followed by heating the plate at 110 °C for 5-10 min. Before being tested on the nematode larvae, using the LMI test, the two fractions, the aqueous one (Faq) and the methanolic one (Fmet), were concentrated under low pressure at 35 °C and the water was removed by freeze-drying. After the results of this first bioassay, a second fractionation on the Fmet fraction was performed using a dialysis method, in order to separate, the molecules with high or low molecular weight. A spectrapor 7 membrane of 2000 Da molecular weight cut off (MWCO), (Spectrum Industries, Laguna hills, CA) was washed with distilled water to remove the sodium azide. The Fmet fraction was suspended in water by sonication and was then pipetted into the dialysis bag and placed in a glass flask with 500 ml of distilled water. Dialysis was performed at room temperature for 48 h with two changes of the outer dialysate. The dialysate and dialyzed solutions were then freeze-dried before being tested on the nematode larvae, with the LMI test.

Identification of compounds

In order further to characterise biochemically the active compounds of the Fmet fraction, purifications were achieved by chromatography on a reverse-phase silica cartridge (Mega Bond Elut Varian[®] C18 cartridge), eluting with a water-MeOH gradient.

HPLC analysis was performed on Merck-Hitashi apparatus equipped with a LaChrom L-7100 quaternary gradient pump and a LaChrom L-7450 DAD detector using a Touzart and Matignon 100 Å (150 × 46 mm) column. Solvent A : 2% AcOH in H₂O, solvent B : CH₃CN; starting from 5% B up to 15% B in 20 min, to 20% B in 30 min, to 50% B in 50 min. The flow rate was set 1 ml/min and UV detection was at 280 nm.

ES-MS (positive and negative ion mode), 3.5 kV (water, MeOH) were measured on a Perkin-Elmer Siex API 365 mass spectrometer. ¹H-NMR (400 MHz) were recorded on a Bruker ARX – 400 spectrometer with TMS as internal standard and Dimethyl sulfoxide (DMSO) *d*₆ as solvent.

Larval Migration Inhibition (LMI) bioassay

In vitro experiments were undertaken to determine the effect of the different solutions, extracted or fractionated, on the mobility of ensheathed *Haemonchus contortus* L3 using the LMI bioassay developed by Wagland *et al.* (1992) and modified by Rabel, Mc Gregor and Douch (1994).

The third-stage larvae (L3) were obtained from a donor goat infected with a pure, susceptible strain of *H. contortus*. The larvae collected were stored at 4°C for 4 months before use in the assay. One thousand L3 were added to microtubes containing the negative control (distilled water, PBS) or anthelmintic control (levamisole at 1% concentration) or the solutions to be tested. All incubations were carried out for 3 h at 20°C. After this time, the L3 were washed and centrifuged (at 5000 rpm for 5 min) three times and then transferred to sieves (inserts equipped with a 20 µm mesh positioned in a conical tube). The 20 µm mesh size was selected in order to ensure that migration of larvae through the sieves was active. After 3 h at room temperature, the number of L3 that had migrated through the mesh, was counted at a 40x magnification. The percentage of migration was calculated as $\frac{M}{T} \times 100$ where T is the total number of L3 deposited on the sieve and M the number of L3 which had successfully migrated through the sieve.

LMI bioassay of extracts : The freeze-dried extracts were dissolved in distilled water alone or distilled water with 5% dimethyl sulfoxide (DMSO Sigma Ltd) for hexane and methylene

chloride ones. We first verified that the addition of 5% of DMSO had no significant effect on the migration of larvae (Rabel *et al.*, 1994). If needed, 5% DMSO was also added to the controls. The concentration of plant extracts solution used for incubation was 1200 µg/ml. Three replicates were run per assay.

LMI bioassay of fractions : The freeze dried samples of fractions were dissolved in phosphate buffer solution (PBS; 0.1M phosphate, 0.05M NaCl; pH 7.2) in order to buffer any acidity in the fractions. The concentration used for incubation was 1200 µg/ml. Five replicates were run, per assay.

LMI bioassay of flavonol glycosides : Three flavonol glycosides were identified in the Fmet fraction. For biological assay, Quercetin-3-rutinoside was obtained from Prolabo Co. (Nogent sur Marne, France), kaempferol-3-rutinoside and isorhamnetin-3-rutinoside were purchased from Extrasynthese Co. (Gemay, France). The three flavonol glycosides were dissolved in PBS with 5% of DMSO and 5% DMSO was added to the controls. Three concentrations were used for incubation, i.e. 300, 600, and 1200 µg/ml. Five replicates were run per flavonol glycoside and per concentration.

Binding with polyvinyl pyrrolidone (PVPP) : In order to confirm the role of phenolic compounds in anthelmintic activity, a series of incubations were undertaken using insoluble PVPP (Doner, Becard and Irwin, 1993; Makkar, Blummel and Becker 1995). Five hundred mg of PVPP was added to 10 ml of the incubation solution corresponding to the 2 fractions obtained after dialysis and also to the rutin solution before being used in the LMI test. The mixture was vortexed and then centrifuged (3000 xg for ten min). The supernatant was collected for use in the LMI assay. The LMI values obtained with the different fractions, with or without PVPP (Sigma Aldrich Co.), were compared.

Statistical analyses

Statistical analyses of data were performed using StatView 5.0. computer software (SAS Institute, Cary, NC). P values less than 0.05 were considered statistically significant. Significant differences between treatment means were assessed using the non-parametrical test of Kruskal-Wallis. In addition, comparisons were carried out by means of the Student's t test for the experiments when 5 replicates were available.

RESULTS

Effects of plant extracts

Fig. 1A No significant effects were found with the extracts of ryegrass irrespective of the solvent
Fig. 1B used (Fig. 1). For the sainfoin extracts, no difference was found in the percentage of
migration between the hexane (Fig. 1A) or methylene chloride (Fig. 1B) extracts and the
DMSO control values. In contrast, 30:70 alcohol/water (v/v) (Fig. 1C), distilled water (Fig.
Fig. 1C inhibition of L3 migration ($p < 0.05$). Those three extracts were then shown by TLC to have
Fig. 1D very similar compound profiles: flavonol glycosides, sugar and tannins which were visible
Fig. 1E under UV and developed specific colorations under vanillin sulphuric acid spray reagent
(data not shown). Since a better yield of extraction was achieved by the 70:30
acetone/water (v/v), this solvent was used in the rest of the study for further fractionation
steps.

Bio-guided fractionation

Fig. 2 The Fmet and the Faq fractions (Fig. 2) showed differences in their efficacy to inhibit
larvae migration. Compared with the PBS control values (90.45% of larval migration), the
migration was affected after incubation with 70:30 acetone/water (v/v) extracts ($p < 0.04$)
and with the Fmet fraction ($p < 0.01$). In contrast, no difference was found in the percentage
of migration between the Faq fraction and the PBS control values. By thin layer
chromatography, the Faq fraction was found to contain mainly a mixture of sugars, which
were detected based on a green coloration after treatment with vanillin sulphuric acid
spray reagent. In contrast, the Fmet fraction was shown to contain a complex mixture of
phenolic compounds (data not shown).

Fig. 3 The dialysate and the dialyzed parts of the Fmet fraction caused a significant reduction
($p < 0.001$) of the larval migration (Fig. 3). The most effective fraction (dialysate) produced a
50 % reduction in larval migration compared with the PBS control. As revealed by TLC, the
dialysate fraction contained a mixture of flavonol glycosides and tannins (< 2000 Da). The
dialyzed fraction, which developed a red coloration upon treatment with vanillin sulphuric
acid spray reagent, contained only tannins (data not shown).

Identification of compounds

In the Fmet fraction, the most abundant flavonoid was isolated and identified as quercetin-3-rutinoside (or rutin) from the ES-MS which yielded the $[M - H]^+$ peak at m/z 609. This was confirmed by NMR spectra comparison to an authentic sample (Lu *et al.*, 2000; Marais *et al.*, 2000). Kaempferol-3-rutinoside (or nicotiflorin) and the isorhamnetin-3-

rutinoside (or narcissin) were also identified through their ES-MS spectra. This was confirmed based on HPLC retention time and UV absorption of an authentic sample (Lu *et al.*, 2000).

Bioactive effects of flavonol glycosides

The inhibitory activities of the 3 flavonol glycosides against L3 larvae are presented in Fig. 4. No significant differences in activities were detected between these compounds and the PBS control values at low concentrations (300 and 600 µg/ml). In contrast, at 1200 µg/ml, the three compounds inhibited significantly ($p < 0.02$) the migration of *Haemonchus* L3. Compared with the PBS values, the migration was reduced respectively by 25% with rutin, 30% with nicotiflorin and 35% with narcissin.

Fig. 4

Effect of PVPP on the activity of the dialysis fractions and of rutin.

The dialysate and the dialyzed fractions caused a significant reduction in larval migration compared to the control. Addition of PVPP (Fig. 5) to these 2 fractions restored the percentage of migration to values that did not differ significantly from those recorded using PBS.

Fig. 5

Similarly, the addition of PVPP to the incubation solution of rutin at 1200 µg/ml (Fig. 6) eliminated its inhibitory effect against the L3 of *Haemonchus contortus*.

Fig. 6

DISCUSSION

Most of the previous studies examining the activity of sainfoin extracts against parasitic nematodes *in vitro* have used *Trichostrongylus colubriformis* (Molan *et al.*, 2000a; Athanasiadou *et al.*, 2001; Paolini *et al.*, 2004). In contrast, for *Haemonchus contortus*, there is a paucity of information despite its economic importance due to its high pathogenicity and to the severe production losses associated with its presence (Allonby and Urquhart, 1975; Perry *et al.*, 2002). Only Paolini *et al.* (2003a) assessed the impact of condensed tannins in goats infected with adult *H. contortus*.

The results of the current study confirmed the activity of sainfoin extracts on *Haemonchus* larvae. They also showed how the activity is attributable to different fractions of *Onobrychis viciifolia*. Different fractions of sainfoin significantly reduced the migration of L3 of *H. contortus* compared with control incubations, and the biochemical nature of these compounds was not restricted to condensed tannins.

The first aim of our study was to determine the best way to extract the active compounds from sainfoin. We showed that only extracts obtained with high polarity solvents retained activity. However, such extracts contained only a small proportion of the total proanthocyanidins present in tissues, noticeably in the case of sainfoin (Bate-Smith, 1973). This fact seems to be associated with the particular chemical composition of condensed tannins in this plant, and showed how the extraction system is a discriminating factor in evaluating *in vitro* activity.

By measuring the migration of L3, it was possible to screen the inhibitory effect of the different compounds of sainfoin during the whole fractionation procedure. The first fractionation obtained after chromatography on C-18 silica column allowed the separation of the extracts into two parts. The Faq (mainly composed of sugars) had no anthelmintic activity, but the bioactivity was associated with the Fmet fraction. The second fractionation of this active Fmet part, aimed to separate high and low MW compounds (> or < to 2000 Daltons). Min and Hart (2003) reported that the MW of condensed tannins of sainfoin ranged from 2040 to 3060. Using the separation, we confirmed that a part of the activity was due to high MW tannins of up to 2000 Da. However, other compounds whose MW was less than 2000 Da, such as flavonol, flavonol glycosides, proanthocyanidins, presented a higher inhibitory effect on the migration of *Haemonchus* L3.

Of the non-tannin compounds of the Fmet, the most abundant one was rutin. Moreover, several closely related glycosides were also identified (Lu *et al.*, 2000; Marais *et al.*, 2000), but were difficult to purify. After identification from the Fmet fraction, commercially purchased rutin, nicotiflorin and narcissin were used in order to measure their activity based on the LMI assay. All of these compounds had an activity against the *Haemonchus* L3 at the concentration of 1200 µg/ml. This concentration is low compared with previous studies. For example, Molan *et al.* (2000a) found that sulla extracts inhibited the migration of exsheathed *H. contortus* L3 by 37 % at the concentration of 100 mg condensed tannins/ml. In contrast, a 80 % inhibition on *H. contortus* ensheathed L3 was measured with sainfoin extracts at concentration of 1200 µg/ml (Paolini *et al.*, 2004), which is similar to the current results. This seems to correspond to some physiological values for small ruminants. In the abomasal digesta of a sheep fed with a diet containing condensed tannins, the concentrations of total condensed tannins were estimated to range between 1100 to 2800 µg/ml whilst the measurements of acetone extractable tannins was about

350 to 900 µg/ml (Terril *et al.*, 1994; Molan *et al.*, 2000b). Jones and Mangan (1977) have previously indicated the absence of any free tannin in the rumen fluid of sheep fed sainfoin, and due to this result, they questioned the possible *in vivo* activity of this forage. According to the current results, the presence of non tannin compounds active against an abomasal nematode could explain this apparent discrepancy.

The specific effect of condensed tannins was further assessed by using PVPP, which is known to bind and to inactivate the biological properties of tannins. As the capture of tannins with PVPP was achieved before incubation with *Haemonchus* L3 and since the tannins would have been then precipitated with PVPP, the supernatant might contain only simple phenolic compounds other than tannins. Makkar *et al.*, (1995), showed that the binding of tannins was maximum at pH 3. In LMI assay, PBS solvent buffered the fractions at pH 7.2, which could explain why the activity of dialyzed fraction did not disappear completely (Fig. 5). In contrast, addition of PVPP to the dialysate fraction (compounds < 2000 Da) also eliminated the effects, which is probably due to the capacity of PVPP to bind also with flavonoids (Doner *et al.*, 1993). This was also confirmed by the addition of PVPP to rutin (1200 µg/ml), which contributed to the restoration of control values of larval migration. The restoration of L3 migration to similar values of the controls was observed after PVPP addition, and supports the hypothesis that the bioactivity was associated with both high MW condensed tannins and flavonol glycosides.

From the current and previous studies (Molan *et al.*, 2000b, Paolini *et al.*, 2004), the evidence shows that the condensed tannins from sainfoin contribute to inhibit the mobility of nematode L3s. In some previous reports (Molan *et al.*, 2003), it has been underlined that, besides the concentration, the chemical structure of tannins, (i.e. the MW, the ratio between prodelphinidins and procyanidins) from different sources of condensed tannins may also be considered as being possible modulating factors for the anti-parasitic properties. Current results suggest that the presence of other compounds, such as flavonol glycosides, might also participate in the modulation of bioactivity. However, it is worth to underline the proximity of biochemical structure between flavonol glycosides and condensed tannins, which are polymers of flavan-3-ols. This similarity suggest a similar or close mechanism of action for both types of compounds. These data indicate that, to estimate the preventive value of any legume forage for the control gastrointestinal nematodes in livestock further analyses of the precise mechanisms involved and

measurements of phenolic compounds like flavonol glycosides in addition to condensed tannins are required.

ACKNOWLEDGMENTS

The authors would like to acknowledge the financial support from the European Union through the Contract WORMCOPS (number QLK5-CT 2001-01843), which is part of collaboration between Denmark, U.K., the Netherlands, Sweden, Spain and France. We also thank Dr. Virginie Paolini (Unité Mixte de Recherches 1225 INRA/DGER) for her help and fruitful discussion. The financial help of the Region Midi Pyrénées (contract 030121196) is also sincerely acknowledged.

REFERENCES

Allonby, E.W., and Urquhart, G.M. (1975) The epidemiology and pathogenic significance of haemonchosis in a merino flock in East Africa. *Veterinary Parasitology* **1**, 129-143.

Athanasiadou, S., Kyriazakis, I., Jackson, F. and Coop, R.L. (2001). Direct anthelmintic effects of condensed tannins towards different gastrointestinal nematodes of sheep *in vitro* and *in vivo* studies. *Veterinary Parasitology* **99**, 205-219. DOI: 10.1016/S0304-4017(01)00467-8

Athanasiadou, S., Kyriazakis, I. and Jackson, F. (2003). Can plant secondary metabolites have a role in controlling gastrointestinal nematode parasitism in small ruminants? *Proceedings of the satellite symposium : secondary compounds and browse utilization. The VI International Symposium on the Nutrition of Herbivores. Satellite Symposium 19th - 24th October 2003. Merida, Mexico.* 16-25.

Bate-Smith, E.C. (1973). Tannins of herbaceous leguminosae. *Phytochemistry* **12**, 1809-1812. DOI:10.1016/0031-9422(73)80409-1

Doner, L.W., Becard, G. and Irwin, P.L. (1993). Binding of flavonoids by polyvinylpolypyrrolidone *Journal of Agriculture and Food Chemistry* **41**, 753-757.

Jackson, F. and Coop, R.L. (2000). The development of anthelmintic resistance in sheep nematodes. *Parasitology* **120**, S95-S107.

Jones, W.T. and Mangan, J.L. (1977). Complexes of the condensed tannins of sainfoin (*Onobrychis viciifolia* Scop.) with fraction 1 leaf protein and with submaxillary mucoprotein,

and their reversal by polyethylene glycol and pH. *Journal of Science, Food and Agriculture* **28**, 126-136.

Kahn, L.P. and Diaz-Hernandez, A. (2000). Tannins with anthelmintic properties. *Tannins in livestock and human nutrition : Proceedings of an International workshop, Adelaide, Australia. ACIAR proceeding* **92** 140-149

Koupai-Abyazani, M.R., Muir, A.D., Bohm, B.A., Towers, G.H.N. and Gruber, M.Y. (1993). The proanthocyanidin polymers in some species of *Onobrychis*. *Phytochemistry* **34**, 113-117. DOI: 10.1016/S0031-9422(00)90791-X

Lu, Y., Sun, Y., Yeap Foo, L., McNabb, W.C. and Molan, A.L. (2000). Phenolic glycosides of forage legume *Onobrychis viciifolia*. *Phytochemistry* **55**, 67-75. DOI: 10.1016/S0031-9422(00)00143-6.

Makkar, H.P.S., Blummel, M. and Becker, K. (1995). Formation of complexes between polyvinyl pyrrolidones or polyethylene glycols and tannins, and their implication in gas production and the true digestibility in *in vitro* techniques. *British Journal of Nutrition* **73**, 897-913.

Marais, J.P.J., Mueller-Harvey, I., Brandt, E.V. and Ferreira, D. (2000). Polyphenols, condensed tannins, and other natural products in *Onobrychis viciifolia* (sainfoin). *Journal of Agriculture and Food Chemistry* **48**, 3440-3447. DOI : 10.1021/jf000388h S0021-8561(00)00388-5

Min, B.R. and Hart, S.P. (2003). Tannins for suppression of internal parasites. *Journal of Animal Science* **81** (E. Suppl.2), E102-E109.

Molan, A.L., Alexander, R.A., Brookes, I.M. and McNabb, W.C. (2000a). Effect of an extract from sulla (*Hedysarum coronarium*) containing condensed tannins on the migration of three sheep gastrointestinal nematodes *in vitro*. *Proceeding of New Zealand Society for Animal Production* **60**, 21-25.

Molan, A.L., Waghorn, G.C., Min, B.M. and McNabb, W.C. (2000b). The effect of condensed tannins from seven herbage on *Trichostrongylus colubriformis* larval migration *in vitro*. *Folia Parasitologica* **47**, 39-44.

Molan, A.L., Meagher, L.P., Spencer, P.A. and Sivakumaran, S. (2003). Effect of flavan-3-ol on *in vitro* egg hatching, larval development and viability of infective larvae of *Trichostrongylus colubriformis*. *International Journal for Parasitology* **33**, 1691-1698. DOI: 10.1016/S0020-7519(96)00207-8.

Niezen, J.H., Charleston, W.A.G., Hodgson, J., Mackay, A.D. and Leathwick, D.M. (1996). Controlling internal parasites in grazing ruminants without recourse to anthelmintics: approaches, experiences, prospects. *International Journal for Parasitology* **26**, 983-992. DOI: 10.1016/S0020-7519(96)00082-3.

Paolini, V., Bergaud, JP., Grisez, C., Prevot, F., Dorchies, Ph. and Hoste, H. (2003a). Effects of condensed tannins on goats experimentally infected with *Haemonchus contortus*. *Veterinary Parasitology* **113**, 253-261. DOI: 10.1016/S0304-4017(03)00064-5.

Paolini, V., Dorchies, Ph. and Hoste, H. (2003b). Effects of sainfoin hay on gastrointestinal nematode infections in goats. *Veterinary Record* **152**, 600-601.

Paolini, V. and Hoste, H. (2003c). Effects of condensed tannins on goats infected with gastrointestinal nematodes : a short review. *Proceedings of the satellite symposium : secondary compounds and browse utilization. The VI International Symposium on the Nutrition of Herbivores. Satellite Symposium 19th - 24th October 2003. Merida, Mexico.* 26-36.

Paolini, V., Fouraste I. and Hoste, H. (2004). *In vitro* effects of three woody plant and sainfoin on third-stage larvae and adult worms of three gastrointestinal nematodes. *Parasitology* **129**, 69-77. DOI: 10.1017/S0031-1820(04)00526-8

Perry, B.D., Randolph, T.F., McDermott, J.J., Sones, K.R. and Thornton, P.K. (2002) *Investing in animal health research to alleviate poverty*, ILRI, Nairobi Kenya.

Rabel, B., MC Gregor, R. and Douch, P.G.C. (1994). Improved bioassay for estimation of inhibitory effects of ovine gastrointestinal mucus and anthelmintics on nematode larval migration. *International Journal for Parasitology* **24**, 671-676. DOI: 10.1016/S0020-7519(93)E0011-W.

Roos, M.H., Kwa, M.S.G. and Grant, W.N. (1995). New genetic and practical implications of selection for anthelmintic resistance in parasitic nematodes. *Parasitology Today* **11**, 148-150. DOI: 10.1016/0169-4758(95)80136-7

Terrill, T.H., Waghorn, G.C., Woolley, D.J., McNabb, W.C. and Barry, T.N. (1994). Assay and digestion of ¹⁴C-labelled condensed tannins in the gastrointestinal tract of sheep. *British Journal of Nutrition* **72**, 467-477.

Thamsborg, S.M., Mejer, H., Bandier, M. and Larsen, M. (2003). Influence of different forages on gastrointestinal nematode infections in grazing lambs. *World Association for the Advancement of Veterinary Parasitology. The 19th International Conference, August 10-14, 2003. New Orleans*. 189.

Wagland, B.M., Jones, W.O., Hribar, L., Bendixsen, T. and Emery D.L. (1992). A new simplified assay for larval migration inhibition. *International Journal for Parasitology* **22**, 1183-1185. DOI:10.1016/0020-7519(92)90040-R

Waller, P.J. (1997). Anthelmintic resistance. *Veterinary Parasitology* **72**, 391-412. DOI: 10.1016/S0304-4017(97)00107-6.

Waller, P.J. (1999). International approaches to the concept of integrated control of nematode parasites of livestock. *International Journal for Parasitology* **29**, 155-164. DOI: 10.1016/S0020-7519(98)00178-7.

Waller, P.J. and Thamsborg, S.M. (2004). Nematode control in “green” ruminant production systems. *Trends in Parasitology* **20**, 493-497. DOI:10.1016/j.pt.2004.07.012



Figure 1 : The effect of  Sainfoin and  Ryegrass extracts on the larval migration inhibition (LMI) of infective 3rd-stage (L3) larvae of *Haemonchus contortus*. The incubations were performed with distilled water, levamisole 1%, and (A) hexane, (B) methyle chloride, (C) 30:70 alcohol/water, (D) Distilled water and (E) 70:30 acetone/water extracts. 5% of DMSO was added to controls, hexane and methylene chloride fractions. All incubations were made in triplicates. Results are presented as mean with standard error of the mean. The asterisks * indicate significant differences to control values ($p < 0.05$).

Figure 2 : The effect of Faq and Fmet fractions of acetone extract of sainfoin on the larval migration inhibition (LMI) of infective 3rd-stage (L3) larvae of *Haemonchus contortus*. The incubations were performed with PBS, levamisole 1%, and the 2 fractions of sainfoin. All incubations were made in five replicates. Results are presented as the mean \pm standard error of the mean. The asterisks indicate the significant differences to control values. (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$).

Figure 3 : The effect of dialysis fractions of Fmet fraction of sainfoin on the larval migration inhibition (LMI) of infective 3rd-stage (L3) larvae of *Haemonchus contortus*. The incubations were performed with PBS, levamisole 1%, and the dialysis fractions. All incubations were made in five replicates. Results are presented as the mean \pm standard error of the mean. The asterisks indicate the significant differences to control values (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$).




Figure 4 : The effect of  rutin  nicotiflorin and  narcissin on the larval migration inhibition (LMI) of infective 3rd-stage (L3) larvae of *Haemonchus contortus*. The incubations were performed with PBS, levamisole 1%, and 3 concentrations of each compound. Five % of DMSO was added. All incubations were made in five replicates. Results are presented as the mean \pm standard error of the mean. The asterisks indicate significant differences to control values (*p<0.05; **p<0.01;***p<0.001).



Figure 5 : The effect of dialysis fractions without  or with the addition of PVPP  on the larval migration inhibition (LMI) of infective 3rd-stage (L3) larvae of *Haemonchus contortus*. The incubations were performed with PBS, levamisole 1%, and the dialysis fractions. All incubations were made in five replicates. Results are presented as the mean \pm with standard error of the mean. The asterisks indicate significant differences to control values (*p<0.05; **p<0.01;***p<0.001).




Figure 6 : The effect of rutin compound without  or with the addition of PVPP  on the larval migration inhibition (LMI) of infective 3rd-stage (L3) larvae of *Haemonchus contortus*. The incubations were done with PBS + 5% DMSO, levamisole 1%, and the rutin. All incubations were made in five replicates. Results are presented as the mean \pm with standard error of the mean. The asterisks indicate the significant differences to control values (*p<0.05; **p<0.01;***p<0.001).


Figure 1



E. BARRAU, N. FABRE, I. FOURASTE AND H. HOSTE

Effect of sainfoin tannins and flavonol glycosides on *Haemonchus contortus*


Figure 2



E. BARRAU, N. FABRE, I. FOURASTE AND H. HOSTE

Effect of sainfoin tannins and flavonol glycosides on *Haemonchus contortus*


Figure 3



E. BARRAU, N. FABRE, I. FOURASTE AND H. HOSTE

Effect of sainfoin tannins and flavonol glycosides on *Haemonchus contortus*


Figure 4



E. BARRAU, N. FABRE, I. FOURASTE AND H. HOSTE

Effect of sainfoin tannins and flavonol glycosides on *Haemonchus contortus*


Figure 5



E. BARRAU, N. FABRE, I. FOURASTE AND H. HOSTE

Effect of sainfoin tannins and flavonol glycosides on *Haemonchus contortus*

Figure 6



E. BARRAU, N. FABRE, I. FOURASTE AND H. HOSTE

Effect of sainfoin tannins and flavonol glycosides on *Haemonchus contortus*