

Effect of Calcium Channel Blockers on Increased Plasma Testosterone Level Induced by Arecoline in Male Albino Rats

Ahmed M. Kabel^{1,2*}, Hany M. Borg³ and Mohamed S. Omar^{4,5}

¹Department of Clinical Pharmacy, College of Pharmacy, Taif University, Taif, Saudi Arabia

²Department of Pharmacology, Faculty of Medicine, Tanta University, Tanta, Egypt

³Department of Physiology, Faculty of Medicine, Kafrelsheikh University, Egypt

⁴Chemistry Department, Faculty of Science, Benha University, Benha, Egypt

⁵Division of Biochemistry, Pharmacology and Toxicology Department, College of Pharmacy, Taif University, Taif, Saudi Arabia

***Corresponding author:** Dr. Ahmed M. Kabel, Department of Clinical Pharmacy, College of Pharmacy, Taif University, Al-Haweiah, P.O. Box 888, Zip Code 21974, Saudi Arabia; Tel: +201009041488 or +966564833526; Email: drakabel@gmail.com

Article Type: Research, **Submission Date:** 10 October 2015, **Accepted Date:** 12 January 2016, **Published Date:** 11 April 2016.

Citation: Ahmed M. Kabel, Hany M. Borg and Mohamed S. Omar (2016) Effect of Calcium Channel Blockers on Increased Plasma Testosterone Level Induced by Arecoline in Male Albino Rats. *J. Drug* 1(1): 10-13. doi: <https://doi.org/10.24218/jod.2016.2>.

Copyright: © 2015 Ahmed M. Kabel and Hany M. Borg. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: Arecoline is one of the major components of betel nuts, which have been consumed as chewing gum in Southeast Asia. Arecoline might affect the endocrine system but the mechanism is still unclear. Some studies suggest that calcium channels may play a role in this effect.

Aim: The present study was carried out to investigate the effect of calcium channel blockers on increased plasma testosterone level induced by Arecoline of betel nuts in rats.

Methods: 80 male albino rats were divided into 8 equal groups as follows: control group; arecoline group; Human chorionic gonadotropin (HCG) group; arecoline + HCG group; nifedipine group; nifedipine + arecoline group; tetrandrine group and tetrandrine + arecoline group. After 2 hours of injection, rats were sacrificed and blood samples were collected and plasma was separated for determination of testosterone, androstendione and leutinizing hormone (LH).

Results: Arecoline with and without HCG resulted in significant increase of plasma testosterone and androsteindione, and only significant increase after HCG injection of LH levels. Injection of nifedipine or tetrandrine alone caused non-significant change in plasma testosterone and androsteindione compared with the control group. Injection of nifedipine or tetrandrine with arecoline resulted in significant reduction in plasma testosterone and androsteindione levels compared with arecoline treated group.

Conclusion: Arecoline caused significant increase in testosterone secretion in male albino rats that may be due to calcium channel

activation in Leydig cells.

Keywords: Arecoline, Calcium channel blockers, Testosterone, Rats.

Introduction

Arecoline is an alkaloid extracted from betel nuts (Areca catechol leaves). Four major alkaloids are found in betel nuts: Arecoline (7.5 mg/gm weight), arecaidine (1.5 mg/gm weight), guvacoline (2 mg/gm weight) and guvacine (2.9 mg/gm weight) [1]. Epidemiological studies showed a strong correlation between oral cancer and betel nut chewing habit [2]. Arecoline and its alkaloids have been shown to be carcinogenic [3], immunotoxic [4], genotoxic [5] and teratogenic in animal models [6]. The increased frequency of micronucleated cells, chromosomal aberrations and sister chromatid exchanges in exfoliated cells of buccal mucosa was observed in arecanut consumers [7,8]. In addition, it was shown that arecoline enhances the frequently of chromosomal aberrations and micronuclei in mouse bone marrow cells in vivo [9].

Arecoline mimics the actions of acetylcholine and exerts its effects at both muscarinic and nicotinic receptors [10]. Studies showed that arecoline readily penetrates the blood brain-barrier [11] and exerts its excitatory action by binding to M2 muscarinic receptors on the cell membrane of neurons of the locus coeruleus [12]. Also it was proved that cholinergic agonist arecoline stimulates the hypothalamo-pituitary-adrenal axis in rats [13], and this effect is mediated mainly by release of endogenous corticotropin-releasing hormone. It has been confirmed by the dose-dependent ability of arecoline to cause hypothalamic corticotropin releasing hormone secretion in vitro [12]. Arecoline significantly elevated

the number of micronucleated cells in a dose dependent manner [13]. Moreover, morphological abnormality and unscheduled DNA synthesis were observed in arecoline treated germ cells of the mouse [14]. Intravenous arecoline administration was followed by the increase in plasma epinephrine and ACTH [15]. It was proved that low dose of betel leaf extract was found to increase the tri-iodothyronine and decrease T_4 [16], but high dose of betel leaf extract increased T_4 and decreased T_3 [16]. Moreover, in Alzheimer patients, elevated plasma ACTH and cortisol were observed in high dose arecoline administration [17]. These studies indicated that arecoline might affect the endocrine system but the mechanism is still unclear. Some studies suggest that calcium channels may play a role in this effect [18].

Nifedipine and tetrandrine are calcium channel blockers used as antihypertensive and in treatment of angina. Calcium is vital in many biological processes including hormonal secretion, mitosis, reproduction, fertility, and regulation of gene expression [18]. Reports exist which suggest that calcium-channel blockers may have anti-reproductive effects in males on long-term treatment. Problems with sexual function have been a long-standing concern in the treatment of hypertension and may influence the choice of treatment regimens and decisions to discontinue drugs [19]. The aim of this work was to investigate the effect of calcium channel blockers on increased plasma testosterone level induced by arecoline of betel nuts in male albino rats.

Materials and Methods

The present study was carried out on 80 male albino rats weighing 200-250 grams. All the experiments were conducted according to the National Research Council's guidelines. Animal handling was followed according to Helsinki declaration of animal ethics. The rats were housed singly and were fed milk bread and had free access to water and food.

They were divided into 8 equal groups as follows:

Group (1): The control group, received 1 ml saline intraperitoneally.

Group (2): The arecoline treated group, received arecoline (Sigma) in a dose of 1 microgram/kg intraperitoneally [20].

Group (3): The human chorionic gonadotropin (HCG) treated group, received HCG (Sigma) in a dose of 5 IU/kg intraperitoneally [20].

Group (4): The arecoline and HCG treated group, received 1 microgram/kg arecoline and 5 IU/kg HCG intraperitoneally [20].

Group (5): Nifedipine treat group, received 25 mg/kg nifedipine (Sigma) intraperitoneally [21].

Group (6): Tetrandrine treated group, received 25 mg/kg tetrandrine (Sigma) intraperitoneally [21].

Group (7): Nifedipine and arecoline treated group, received 1 microgram/kg arecoline and nifedipine 25 mg/kg intraperitoneally [21].

Group (8): Tetrandrine and arecoline treated group, received 1 microgram/kg arecoline and 25 mg/kg tetrandrine intraperitoneally [21].

After 2 hours of injection, all rats were sacrificed and blood samples were collected and plasma was separated for determination of plasma testosterone hormone according to the method of Cox et al. [22], plasma leutinizing hormone (LH) and plasma androsteindione according to the method of Kullin and Santiner [23].

Statistical analysis

Data were presented as mean \pm standard error of mean (SEM). Student's t-test was used for evaluating the statistical significance of differences in means. P value of less than 0.05 was considered to be statistically significant [24].

Results

The results are shown in Figures 1-3. They showed that administration of 1 microgram/kg arecoline intraperitoneally resulted in significant increase in plasma testosterone and androstenedione levels ($P < 0.05$) and non-significant change in plasma LH level compared to control group. The results also showed significant increase in plasma levels of testosterone, androstenedione and LH after injection of HCG alone and injection of HCG and arecoline ($P < 0.05$) compared to the control group.

The results showed non-significant change in plasma testosterone, androstenedione and LH levels ($P > 0.05$) after injection of nifedipine or tetrandrine compared with the control group. There were significant reduction in plasma levels of testosterone and androstenedione after injection of both arecoline and nifedipine or arecoline and tetrandrine ($P < 0.05$) and non-significant change in plasma LH levels compared with the arecoline treated group.




Figure 1: Effect of arecoline, HCG and calcium channel blockers on plasma testosterone in male albino rats (Mean \pm SEM)

* Non significant compared to the control group ($P > 0.05$)

** Significant compared to the control group ($P < 0.05$)

Non significant compared to arecoline group ($P > 0.05$)

Significant compared to arecoline group ($P < 0.05$)




Figure 2: Effect of arecoline, HCG and calcium channel blockers on plasma androstenedione in male albino rats (Mean ± SEM)
 *Non significant compared to the control group (P>0.05)
 ** Significant compared to the control group (P<0.05)
 #Non significant compared to arecoline group (P>0.05)
 ## Significant compared to arecoline group (P<0.05)




Figure 3: Effect of arecoline, HCG and calcium channel blockers on plasma LH in male albino rats (Mean ± SEM)
 *Non significant compared to the control group (P>0.05)
 ** Significant compared to the control group (P<0.05)
 #Non significant compared to arecoline group (P>0.05)
 ## Significant compared to arecoline group (P<0.05)

Discussion

Betel is a masticatory substance and betel chewing is a popular oral habit in southern Taiwan, although the significant correlation of betel chewing and the incidence of oral cancer and oral submucosal fibrosis [3]. Arecoline is one of the major ingredients in betel nuts in addition to the other three ingredients; arecaidine, guvacoline and isoguvacine [1]. Each betel quid contains 7.5 mg/gm of Arecoline [25]. Although a number of studies showed that betel chewing is strongly related to oral cancer, betel has been used for the treatment of diarrhea, edema, throat inflammation and tapeworm infection [26].

The results of the present work showed that arecoline with or without HCG, significantly stimulated testosterone secretion. The increased secretion of testosterone may be explained by the ability of arecoline to stimulate steroidogenesis and also HCG-induced plasma testosterone and androstenedione was enhanced by arecoline injection. The effect of arecoline on testosterone was observed by Chiao et al. [27] who noted that significant increase in plasma testosterone in rats after 60 min from arecoline after the LH secretion from the anterior pituitary gland alone or after HCG treatment, in spite of the increase in plasma testosterone which suggested that the action of arecoline on testosterone is not LH dependent [28]. It was observed that, arecoline stimulated testosterone secretion from isolated Leydig cells which may be through nicotinic or muscarinic receptors in the tissues [29]. Testosterone is synthesized in Leydig cells by several metabolic steps known as steroidogenesis. The results of the present work showed significant increase in androstenedione by arecoline injection which can explain the significant increase in testosterone secretion [30], because androstenedione, testosterone precursor, which can be converted to testosterone by 17 beta hydroxylase enzyme [31].

The mechanism of testosterone hormone secretion can be explained by that, stimulation of LH on the steroidogenesis in the cells is through binding to specific receptors, activation of adenylate cyclase, formation of cAMP and protein kinase A activation, phospholipids turnover, inositol triphosphate formation and the increase of intracellular calcium [31]. The increase of cytoplasmic Ca²⁺, either released from endoplasmic reticulum or transported from calcium channels in the cell membrane, is involved in enhancement of testosterone production by rat Leydig cells [32]. The results of the present study showed that arecoline-induced testosterone secretion was reduced by administration of calcium channel blockers. These results can be explained by that, HCG induced a slow onset and sustained monophasic intracellular Ca²⁺ concentration that was dependent on extracellular Ca²⁺.

Rossato et al. [32] reported that, the transient but not sustained rise in Ca²⁺ was observed in Leydig cells treated with thapsigargin and cyclopiazonic acid, which are two sarcoplasmic reticulum calcium adenosine triphosphatase inhibitors, in Ca²⁺ free medium. This rise in Ca²⁺ indicated that thapsigargin and cyclopiazonic acid are able to empty the intracellular Ca²⁺ influx [32]. This could be explained by that, the influx of external Ca²⁺ induced by the emptying of internal stores in Leydig cells, that could occur through Na⁺ dependent depolarization of the plasma membrane [33], not through voltage activated Ca²⁺ channels [34], which significantly reduced by administration of either T and L calcium channel blockers, such as nifedipine, with arecoline.

Conclusion

Arecoline stimulates testosterone production in rats via activation of calcium channels in Leydig cells that can be blocked significantly by calcium channel blockers.

References

- Papke RL, Horenstein NA, Stokes C. Nicotinic Activity of Arecoline, the Psychoactive Element of "Betel Nuts", Suggests a Basis for Habitual Use and Anti-Inflammatory Activity. *PLoS ONE*. 2015; 10(10):e0140907. doi: 10.1371/journal.pone.0140907.
- Chiao YC, Lin H, Wang SW, Wang PS. Inhibition of testosterone production by propylthiouracil in rat Leydig cells. *Biol Reprod*. 2002; 67(2):416-422.
- Wang CK, Lee WH, Peng CH. Contents of phenolics and alkaloids in areca rate chulinn- during maturation. *J. Agric. Food Chem*. 1997; 45:1185-1188.
- Selvan RS, Selvokomaran M, Rao AR. Influence of arecoline on immune system. II. Suppression of thymus dependent immune responses and parameters on non-specific resistance after short term exposure. *Immunopharmacol Immunotoxicol*. 1991; 13(3):281-309.
- Pangraphi GB, Rao AR. Induction of *in vivo* sister chromatid exchanges by arecaidine, a betel nut alkaloid, in mouse bone marrow cells. *Cancer Lett*. 1984; 23(2):189-192.
- Sinha A, Rao AR. Embryotoxicity of betel nuts in mice. *Toxicol*. 1985; 37(3-4):315-326.
- Chang YC, Tsai CH, Taik W, Yung SH, Chou MY, Li CK. Elevated vimentin expression in buccal mucosal fibroblasts by arecoline *in vitro* as a possible pathogenesis for oral submucous fibrosis. *Oral Oncol*. 2002; 38(5):425-430.
- Dare BJ, Trivedi AH, Adharyu SG. Cytogenetic studies reveal increased genomic damage among "pan masala" consumers. *Mutagenesis*. 1991; 6(2):159-163.
- Deb S, Chatterjee A. Influence of glutathione and reduced glutathione on arecoline induced chromosomal damage and sister chromatid exchange in mouse bone marrow cells *in vivo*. *Mutagenesis*. 1998; 13(3):243-248.
- Brown JH, Taylor P. Muscarinic receptor agonists and antagonists. In: Hardman JG, Limbird LE editors. *Goodman and Gilman's the pharmacological basis of therapeutics*. 10th Ed. New York: McGraw Hill; 2003. 158-161 p.
- Rapoport SI, Ohno K, Pettigrew KD. Drug entry into the brain. *Brain Res*. 1979; 172(2):354-359.
- Young YR, Chang KC, Chen KC, Chen CL, Chiu TH. Arecoline excites rat locus coeruleus neurons by activating the M2-muscarinic receptor. *Clin J Physiol*. 2000; 43(1):23-28.
- Lee CH, Lin RH, Liu SH, Lin-Shiau SY. Mutual interactions among ingredients of betel quid in inducing genotoxicity on Chinese hamster ovary cells. *Mutat Res*. 1996; 367(2):99-104.
- Sinha A, Rao AR. Induction of shape abnormality and unscheduled DNA synthesis by arecoline in the germ cells of mice. *Mutat Res*. 1985; 158(3):189-192.
- Polinsky RJ, Brown RT, Curras MT, Baser SM, Baucom CE, Hooper DR, et al. Central and peripheral effects of arecoline in patients with autonomic failure. *J Neurol Neurosurg Psychiatry*. 1991; 54(9):807-812.
- Panda S, Kar A. Dual role of betel leaf extract on thyroid function in male mice. *Pharmacol Res*. 1998; 38(6):493-496.
- Asthana S, Raffaele KC, Greig NH, Berardi A, Morris PP, Schapiro MB, et al. Neuroendocrine responses to intravenous infusion of arecoline in patients with Alzheimer's disease. *Psychoneuroendocrinol*. 1995; 20(6):623-636.
- Lee JH, Kim H, Kim DH, Gye MC. Effects of calcium channel blockers on the spermatogenesis and gene expression in peripubertal mouse testis. *Arch Androl*. 2006; 52(4):311-318.
- Rabia L, Ghulam ML, Muhammad A. Effects of amlodipine on serum testosterone, testicular weight and gonado-somatic index in adult rats. *J. Ayub Med. Col. Abbottabad*. 2008; 20:8-10.
- Wang SW, Hwang GH, Chen TJ, Wang PS. Effects of arecoline testosterone release in rats. *Am J Physiol Endocrinol Metab*. 2008; 295(2):E497-E504. doi: 10.1152/ajpendo.00045.2008.
- Gatch N, Wallis C, Lal H. Effects of calcium channel blockers on pentylentetrazol drug discrimination in rats. *Alcohol*. 2001; 23(3):141-147.
- Cox JE, Williams JH, Rowe PH, Smith JA. Testosterone in normal, cryptorchid and castrated horses. *Equine Vet J*. 1973; 5(2):85-90.
- Kullin HE, Santiner FJ. The assessment of diminished testicular function in boys of pubertal age. *Clin Endocrinol (Oxf)*. 1996; 25(3):283-292.
- Stallard, Torrie JH. *Principals and procedures of statistics*. 2nd Ed. New York: McGraw-Hill; 1981.
- Wang CK, Wang LS. Study on the separation and hydrolysis of alkaloids and tumorigenicity of betel quid ingredients. *J Agric Food Chem*. 1996; 44:2014-2019.
- Thomas SJ, MacLennan R. Skald lime and betel nut cancer in Papua New Guinea. *Lancet*. 1992; 340(8819):577-8.
- Chiao YC, Lin H, Wang SW, Wang PS. Direct effects of propylthiouracil on testosterone secretion in rat testicular interstitial cells. *Br J Pharmacol*. 2000; 130(7):1477-1482.
- Yang NY, Kapide K, Wang PH, Jong DS, Wu LS. Effects of aqueous extracts of "betel quid" and its constituents on testosterone production by dispersed mouse interstitial cells. *Am J Chin Med*. 2004; 32(5):705-715.
- Clu NS. Effects of betel chewing on the central and autonomic nervous systems. *J Biomed Sci*. 2001; 8(3):229-236.
- Garg P, Chaturvedi PC, Gupta PC. A review of the systemic adverse effects of areca nut or betel nut. *Indian J Med Paediatr Oncol*. 2014; 35(1):3-9. doi:10.4103/0971-5851.133702.
- Aguirre PC, Jayes JD, Veldhuis. Luteinizing hormone (LH) drives diverse intracellular calcium second messenger signals in isolated porcine ovarian theca cells preferential recruitment of intracellular Ca²⁺ oscillatory cells by high concentrations of LH. *Endocrinology*. 2000; 4:2220-2228.
- Rossato M, Nogara A, Merico M, Ferlin A, Garolla A, Foresta C. Store operated calcium influx and stimulation of steroidogenesis in rat Leydig cells, role of Ca²⁺ activated K⁺ channels. *Endocrinology*. 2001; 142(9):3865-3872.
- Rossato M, Nogara A, Gottardello F, Bordon P, Foresta C. Pituitary adenylate cyclase activating polypeptide stimulates rat Leydig steroidogenesis through a novel transduction pathway. *Endocrinology*. 1997; 138(8):3228-3235.
- Tomic M, Dufau MI, Catt KI, Stojkovic SS. Calcium signaling in single rat Leydig Cells. *Endocrinol*. 1995; 136(8):3422-3429.