

Effect of Chemical Treatments and Incorporation of Organic Matter on the Pathogenicity and Survival of *Pseudomonas solanacearum* (Smith) in Soil

J. M. R. S. BANDARA

Department of Agricultural Biology, Faculty of Agriculture, University of Peradeniya, Peradeniya, Sri Lanka

(Date of receipt: 13 June 1983)

(Date of acceptance: 10 December 1984)

Abstract: All solanaceous crops cultivated in Sri Lanka except chilli cv. MI 1 were found to be susceptible to biotype 3 of *Pseudomonas solanacearum*. Bacterium was also proven pathogenic to some of the improved cultivars of AVRDC tomato selections. Organic manure (chicken manure, cowdung, and *Gliricidia* leaves) when incorporated into inoculated soil significant increase in number of propagules of *P. solanacearum* and severity of wilt disease in subsequent tomato cv. marglobe were observed in contrast to chemical treatments (lime, acetic acid, Benomyl and Captan). Significant reduction of bacterial population and disease severity was seen in Captan and Benomyl treated soil only. Organic manure treatments stimulated growth of both *P. solanacearum* and other saprophytic bacteria. The ratio of *P. solanacearum* to other bacteria showed a direct relationship to percentage wilt in a subsequent tomato crop.

1. Introduction

The bacterial wilt caused by *Pseudomonas solanacearum* E.F. Smith is one of the major constraints in the production of solanaceous crops. Bacterium survives in soils for varying periods of time even in the absence of the host.¹⁰ The ability of the primary inoculum to survive during the non-crop period is therefore one of the important aspects to be considered in disease control. Survival of the pathogen may depend on the genetic and environmental factors it is subjected to. An attempt has been made to simulate a control practice based on the survival of *P. solanacearum* in soil under control conditions.

2. Materials and Methods

Highly virulent isolate of *Pseudomonas solanacearum* E.F. Smith biotype 3 sensu Hayward was used throughout these experiments. Pathogen was isolated from potato and its pathogenicity to potato plants was confirmed by stem and soil inoculation.

The nutrient medium used for the isolation of the bacterium and its routine culture was glucose peptone agar (Glucose 5.0 g, Peptone 5.0 g, Yeast extract 2.0 g, KHPO₄ 0.2 g, MgSO₄·7H₂O 0.2 g, CaCO₃ 0.1 g, Agar 18.0 g in 1 litre of distilled

water). Virulence of the isolates were maintained by passing the cultures at least once in 6 weeks through tomato (*Lycopersicon esculentum* Mill) cv. marglobe. Cultures in agar or as suspensions in sterile distilled water were maintained at $21 \pm 1^\circ\text{C}$ in Precision low temperature incubators.

2.1 Pathogenicity tests

Pathogenicity of the bacterial isolates were confirmed by re-inoculating tomato plants. Five weeks old tomato cv. Marglobe plants raised in sterile soil were stem inoculated with the bacterial suspension containing 10^6 cells/ml. Bacterial suspension was prepared with 48 h cultures grown in glucose peptone agar. Three, $10\mu\text{l}$ drops of the suspension were placed on the axil of the third leaf from the top and the stem was then pricked with a syringe needle (gauge 25) through the inoculum drop.

Inoculated plants were placed in a green house (24°C night $28 - 33^\circ\text{C}$ day). Plants were observed for the wilt and yellowing after three days from inoculation.

2.2 Evaluation of control measures

Cultivars of popular solanaceous crops were screened for resistance to *P. solanacearum* infection. Plants were inoculated either directly by stem inoculation or by transplanting into soil inoculated with *P. solanacearum*. In the latter to each plastic pot (diameter 15.5 cm) filled with 1.3 kg of sand loam soil (about 550 ml), 25 ml of *P. solanacearum* suspension 10^8 cells/ml was added. Soils were mixed thoroughly and left in the green house for three days before transplanting the test plants. Host plants used were tomato (*Lycopersicon esculentum* Mill.) cv. Marglobe, Katugastota selection and cvs. AVRDC selection No. CL 141.0.10.3, CL 113.2.4, CL 1561.6.0.22.4, CL 1591.5.0.1.6, CL 1591.5.0.1.7, CL 1094, F₄-57, CL 11d.0.12.1, L1 and L 387. Egg plant (*Solanum melongena*) cv. SM 164, Chilli (*Capsicum annum*) cv. MI 1, (*Capsicum annum* var. *grossum*) cv. CA 8, Potato (*Solanum tuberosum*).

2.3 Chemical and cultural methods

A sand loam soil collected from virgin land under a thick cover of grass near Meewatura research station of the Faculty of Agriculture was used. The physical analysis of soil reveals the following mechanical composition. Sand 75.40%, silt 10.32%, clay 14.32% and C = 3.05%. The soil was passed through a 2 mm sieve (End Cott. Ltd., London) prior to inoculation with *P. solanacearum*. Soil was inoculated as described above.

Inoculated pots were left in the green house for five days before transplanting five tomato cv. Marglobe seedlings raised in moist heat sterilized soil. One month after transplanting dead plant material was incorporated back into soil to develop "sick" soil. Pots were later emptied and the soil was mixed thoroughly before refilling the pots with "sick soil" for experimentation.

The pot trial consists of eight treatments. Each treatment was replicated five times. To each plastic pot (diameter 15.5 cm) 1.3 kg of "sick" soil was added.

In organic manure treatments 26 g of either sun-dried cattle manure, poultry manure or leaves of *Gliricidia maculata* was added to each pot. Two fungicides Captan and Benlate were used in addition to lime and acetic acid as chemical treatments. To each pot either 0.5 g in 100 ml of Captan (a.i. N-trichloro methyl thio tetrahydrophthalimide), 0.3 g in 100 ml water of Benlate (Dupont, Methyl (1-butylcarbamoyl) benzimidazol -2-ylcarbamate or Benomyl), 15 g of CaCO_3 or 25 ml of 10% acetic acid was added. Inoculated and uninoculated controls were used. Five weeks old tomato cv. Marglobe was transplanted as a test plant. Each treatment was replicated five times and pots were kept in the green house in completely randomized design.

2.4 Quantitative assessment of microflora

Population levels of *Aspergillus* sp., *Penicillium*, sp., *Pythium*, *Trichoderma* and *P. solanacearum* were estimated.

Five soil samples were taken from each pot at two weeks interval for 2 months using a No. 5 cork borer. Samples from all the pots were bulked together to form a composite sample. These samples were collected in polypropylene wide neck bottles and mixed by agitating on a Vortex Genie mixer before taking sub samples for the preparation of soil suspensions. Five sub samples were taken from each composite sample using No. 1 cork borer. This soil was used to make soil dilution 10^4 (w/v) in sterile distilled water.

Population of *Aspergillus* spp., *Trichoderma* spp., and other fungi were estimated by the dilution plate technique using Martin's medium.¹¹ One ml aliquots from the soil dilution ($1:10^4$) were pipetted into sterile petri dishes followed by 10 ml autoclaved Martin's medium to each plate and the soil suspension was mixed thoroughly with melted agar by swirling the freshly poured plates. Plates were incubated at 25°C in Precision low temperature incubator for 5-7 days and number of colonies of *Aspergillus* spp., *Penicillium* spp., *Trichoderma* spp. and other fungi were counted. Different species were identified by the characters described by Barnett³ and Gilman.⁷

Population of actinomycetes were estimated by procedure adopted by William and Cross.¹⁸ Inoculated plates were incubated for 8 - 10 days at 25°C before counting the number of actinomycetal colonies per plate.

The survival of *P. solanacearum* in soils under different treatments was studied by monitoring the population levels at 3 weeks intervals. Selective medium proposed by Karganilla and Buddenhagen⁹ was used for the estimation of the bacterial

populations. Colony counts as an average of five replicates at 10^{-4} dilution were recorded. Number of tomato cv. Marglobe plants wilted as percentage was recorded as estimates of pathogenicity.

3. Results and Discussion

Results of the pathogenicity tests showed that all cultivated varieties of *Solanacea* except chilli (pepper) cv. MI 1 are susceptible to *P. solanacearum* infection. Although initial symptoms of wilt appeared on the 3rd day after axil inoculation in most susceptible cultivars of the crops tested 100% wilt was recorded at a very early stage in commonly cultivated potato variety Arka 1 (Figure 1).

All the improved cultivars of tomato supplied by AVRDC and almost all other solanaceous crops tested were found to be susceptible to biotype 3 of *P. solanacearum*. It was found that 100% of the plants were wilted on the 13th day after axil inoculation. Since biotype 3 has an island-wide distribution and especially in tomato growing areas² any attempt to introduce new cultivars without proper soil sterilization or control measures will not be desirable.

Bacterial wilt thus remains potentially the most serious bacterial disease in solanaceous crops.

Effect of incorporation of organic manures and chemicals on the final disease severity in tomato plants are summarized in Table 1. There was a significant reduction in disease severity in Poultry manure treatment in comparison to control, although it remained at a high % of 40. However all organic manure additions have increased the disease severity.

TABLE 1. Effect of various soil treatments on the population of *Pseudomonas solanacearum*, other bacteria and bacterial wilt of Tomato (Marglobe) plants

Treatment	pH	<i>P. solanacearum</i> ^a x 10 ⁸ /g soil	Other bacteria ^a	OB/PS ratio	%wilt
Poultry manure	7.5	11.3	99.6	10	40
Cowdung	7.9	20.0	98.0	5	85
Glyricidia	7.2	42.3	166.0	4	75
Lime	7.4	7.0	126.0	18	5
Benlate	7.3	3.6	60.3	15	10
Captan	6.5	3.0	35.0	12	0
Acetic acid	5.0	3.0	22.3	7	0
Control	6.6	9.8	74.6	8	65
LSD at P = 0.01		8.02	22.6		5.84
0.05		6.05	18.7		5.10

a. Mean of 5 replicates after 9 weeks from treatments.




FIGURE 1 — Rate of wilting of few selected solanaceous crops artificial inoculation with *Pseudomonas solanacearum* (Smith) biotype III.

- △ Tomato — Marglobe
- ▲ Brinjal — SM 164
- Capsicum — CA 8
- Potato — Arka




FIGURE 2—Effect of incorporation of organic matter (Glirescencia leaves ■, Cowdung ○ Poultry manure ●) and chemicals (Lime △, Benomyl ◇, Captan □, Acetic acid ◆) on the number of *Pseudomonas solanacearum* in soil. Control ▲

Benomyl, Captan and Acetic acid applications to soil have significantly ($P=0.05$) reduced the disease level in subsequent tomato crops. It is evident from Table 1 that pH had little effect in disease development. The literature concerning the importance of pH however is contradictory.¹⁰ In some respects it appears that the disease is more severe in soils with moderate to acid pH values (pH 5.0-5.5) but it also may be severe in alkaline soil with pH 7.0 - 7.5.¹⁴

Okabe¹³ reported that *P. solanacearum* reproduces in soils with a pH of 5.0 but that the population decreased with an increase in soil pH. Untreated, inoculated control soil with pH 6.6 produced 65% wilt compared to Benomyl treated or lime incorporated soils with pH 7.3 - 7.4 (Table 1). However pH changes due to organic matter incorporation had no direct relationship with percentage wilt. Okabe¹³ attributed low bacterial population at pH 7.1 - 7.7 to greater activity of microflora. The pH changes in organic matter incorporated soils are temporary changes and could be easily reversed to its original pH with time.

Since the solanaceous crops are short age irrigated and are commonly cultivated it is vital to know the behaviour of *P. solanacearum* in a short period of non crop seasons in the field. Further examinations of treated soil for the survival of *P. solanacearum* and the biology of other common microorganisms in soil revealed that addition of organic matter to soil was favourable for the multiplication and survival *P. solanacearum* (Figure 2). Table 1 shows that the change in population of other bacteria was similar to that of *P. solanacearum*. This indicates that various organic manure used in tomato-vegetable cultivation provide favourable soil environment for the multiplication of bacteria and their survival. However, it is interesting to note the ratio of *P. solanacearum* to other bacteria was directly related to the disease level. This could be due to the saprophytic competition offered by the non-pathogenic bacteria.

When the microflora was studied it was found that percentage wilt was not directly related to *P. solanacearum* in soil but its relative density in soil with respect to saprophytic bacteria (Table 1). Percentage wilt showed direct relation to other bacteria/*P. solanacearum* ratio (OB/PS) in all treatments except in acetic acid treated soil (Figure 3). This was probably due to extremely low pH 5.0 toxic effect of acid treatment that completely inhibited bacterial multiplication. Poultry manure in soil at pH 7.5 gave a OB/PS ratio 10. All soils with a OB/PS ratio higher than 10 were suppressive for wilt production.

Gliricidia leaves when incorporated into soil had a boosting effect on the *P. solanacearum* population. In almost all organic manure treatments it remains significantly higher than control even after 9 weeks. Although *P. solanacearum*




FIGURE 3 — Relationship between (No. of other bacteria/No. of *Pseudomonas solanacearum* OB/SP ratio and percentage wilt increased in tomato cv. Marglobe.




FIGURE 4 — Distribution of saprophytic fungi in soils treated with chemicals on the 3rd and 9th week after treatment.

- A — Aspergillus spp.
- Py — Pythium spp.
- U — Unidentified fungi

- P — Penicillium spp.
- T — Trichoderma spp.

numbers in chemical amended soils were reduced, a significant reduction (at $P=0.05$) was obtained only in Captan, Acetic acid and Benomyl treatments (Figure 2). Effect of acetic acid could probably be due to its low pH and toxicity and therefore not desirable as a control practice.

The behaviour of *P. solanacearum* population in Captan and Benomyl treatments needs further discussion. Initial increase in bacterial population in Captan treatment was very prominent (Figure 2). A Captan spray on apple leaves was reported to increase the relative abundance of bacteria.⁸ Subsequent reduction in *P. solanacearum* and severity (%) of wilt could be due to biological suppression/control initiated by the chemical.¹ They suggested that Captan may stimulate saprophytic *Penicillium Trichoderma*, *Fusarium*, *Actinomyces* spp and bacteria. Captan was reported to be toxic to fungi, bacteria, higher plants and even insects.¹⁷ Domsch⁵ observed that it has no effect on bacteria. Picci¹⁵ however noted its toxic properties on nitrifying and ammonifying bacteria. Rich¹⁶ and Diener and Carlton⁴ reported that Captan controls Stewart's wilt in Maize (*Xanthomonas stewartii*) and bacterial spot of peach respectively.

It is suggested that the reduction in *P. solanacearum* levels in soil could be due to Captan initiated biological control probably by actinomycetes or saprophytic fungi (Figure 4).

However, a gradual decline in *P. solanacearum* population was shown in Benomyl treated soil (Figure 2), which is probably due to a direct effect on bacteria. Fassin⁶ stated that Benomyl had direct effect on soil bacteria and might cause shift in bacterial flora of the soil. It is also known to reduce Rhizobia in soil.¹²

The numerous materials and the microbial habitats available in soil to serve as energy sources provide a range of niches for plant pathogenic bacteria. Most pathogenic bacteria however could be easily displaced by more efficient nutrient utilizing saprophytes. Therefore the control of *P. solanacearum* may be achieved by selective stimulation of saprophytic population by the use of Captan or Benomyl rather than incorporation of organic matter.

Acknowledgements

I wish to express my sincere thanks to Miss S. Rombage for her helpful assistance. The financial assistance provided by the Natural Resources, Energy and Science Authority is gratefully acknowledged.

References

1. BAKER, K. F. & COOK, R. J. (1974). *Biological control of Plant Pathogens*. 433 p. W. H. Freeman & Co., San Francisco.
2. BANDARA, J. M. R. S. (1982). Biotype distribution of Vascular wilt pathogen *Pseudomonas solanacearum* in Sri Lanka. *J. Natn. Sci. Coun. Sri Lanka*. 11 (1), 65-76.
3. BARNETT, H. L. (1960). *Illustrated genera of imperfect fungi*. Burges Publishing Co., U.S.A.
4. DIENER, U. L. & CARLTON, C. C. (1960). Dodine - Captan combination controls bacterial spot of peach. *Pl. Dis. Repr.* 44, 136-138.
5. DOMSCH, K. H. (1968). The effect of soil fungicide III. Quantitative changes in soil flora. *Z. Pflanzenkrankh Pflanzenschutz* 66, 451-456.
6. FASSEN, H. G. VAN. (1974). Effect of the fungicide Benomyl on some metabolic process and on number of bacteria in the soil. *Soil Biol. Biochem.* 6, 131-133.
7. GILMAN, J. C. (1957). *A manual of soil fungi*. Iowa State University Press. 450 pp.
8. HISLOP, E. C. & COX, T. W. (1969). Effects of Captan on the nonparasitic microflora of apple leaves. *Trans. Brit. Mycol. Soc.* 52: 223-235.
9. KARAGANILLA, A. D. & BUDDENHAGEN, I. W. (1972). Development of a selective medium for *Pseudomonas solanacearum*. *Phytopathology*, 62: 1373-1376.
10. KELMAN, A. (1953). The bacterial wilt caused by *Pseudomonas solanacearum* N.C. *Agric. Exp. Sta. Tech. Bull.* 99: 194.
11. MARTIN, J. P. (1950). Use of acid, rose bengal and streptomycin in the plate method for estimating soil fungi. *Soil Sci.*, 69, 215-233.
12. NARAYANA, Y. D., RADAKRISHNA, D. & RAI, P. V. (1981). *Current Res.*, 10 (3) 47-48.
13. OKABE, N. (1971). Population changes of *Pseudomonas solanacearum* and soil microorganism in artificially infected natural field soil. *Rev. Plant Protection Res.*, 4: 105-108.
14. PARK, M. & FERNANDO, M. (1938). The relative resistance of some tomato varieties to bacterial wilt (*Bacterium solanacearum* F. F. Smith). *Trop. Agric. (Ceylon)* 91: 333-337.
15. PICCI, G. (1956). Effect of Captan on soil microorganisms. *Agr. Ital. (Pisa)* 56, 376-382.
16. RICH, S. (1956). Seed treatments to protect corn seedling against stewarts wilt. *Pl. Dis. Repr.* 40, 417-420.
17. SOSNOVOSKY, G. (1958). The Chemistry of trichloro methane sulphenyl chloride. *Chem. Rev.* 58, 509-540.
18. WILLIAMS, S. T. & CROSS, T. (1974). Actinomycetes. p 298-334. In *methods in microbiology* Vol. 4. ed. by C. Booth. p. 795. Academic press, London.