



Docherty, K. et al. (2020) Effect of dapagliflozin in DAPA-HF according to background glucose-lowering therapy. *Diabetes Care*, (doi: [10.2337/dc20-1402](https://doi.org/10.2337/dc20-1402))

The material cannot be used for any other purpose without further permission of the publisher and is for private use only.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/222797/>

Deposited on 04 September 2020

Enlighten – Research publications by members of the University of  
Glasgow

<http://eprints.gla.ac.uk>

**Full title:** Effect of dapagliflozin in DAPA-HF according to background glucose-lowering therapy

**Short title:** Background glucose-lowering therapy in DAPA-HF

**Authors:** Kieran F. Docherty MB ChB<sup>1</sup>;  
Pardeep S. Jhund MBChB MSc PhD<sup>1</sup>;  
Olof Bengtsson Ph. Lic.<sup>2</sup>;  
David L. DeMets, PhD<sup>3</sup>;  
Silvio E. Inzucchi, MD<sup>4</sup>;  
Lars Køber, MD, DMSc<sup>5</sup>;  
Mikhail N. Kosiborod, MD<sup>6</sup>;  
Anna Maria Langkilde, MD, PhD<sup>2</sup>;  
Felipe A. Martinez, MD<sup>7</sup>;  
Marc S. Sabatine, MD, MPH<sup>8</sup>;  
Mikaela Sjöstrand, MD, PhD<sup>2</sup>;  
Scott D. Solomon, MD<sup>9</sup>  
John J.V. McMurray MD<sup>1</sup>

On behalf of the DAPA-HF Investigators and Committees

**Affiliations:** <sup>1</sup>BHF Cardiovascular Research Centre, University of Glasgow, Glasgow, UK;  
<sup>2</sup> AstraZeneca R&D, Gothenburg, Sweden;  
<sup>3</sup> Department of Biostatistics & Medical Informatics, University of Wisconsin, Madison, WI, USA;

<sup>4</sup> Section of Endocrinology, Yale University School of  
Medicine, New Haven, CT, USA;

<sup>5</sup> Rigshospitalet Copenhagen University Hospital, Copenhagen,  
Denmark;

<sup>6</sup> Saint Luke's Mid America Heart Institute and University of  
Missouri-Kansas City, Kansas City, Missouri, USA;

<sup>7</sup> National University of Cordoba, Cordoba, Argentina;

<sup>8</sup> TIMI Study Group, Cardiovascular Division, Brigham and  
Women's Hospital, and Harvard Medical School, Boston, MA,  
USA;

<sup>9</sup> Cardiovascular Division, Brigham and Women's Hospital,  
Boston, MA, USA;

**Correspondence:**

Professor John J.V. McMurray,  
British Heart Foundation Cardiovascular Research Centre,  
University of Glasgow,  
126 University Place,  
Glasgow, G12 8TA,  
United Kingdom.

Tel: +44 141 330 3479

Fax: +44 141 330 6955

Email: [john.mcmurray@glasgow.ac.uk](mailto:john.mcmurray@glasgow.ac.uk)

**Word Count:**

133 (abstract)  
1217 (manuscript text)  
1350 (total)

## Abstract

**Objective:** To determine whether the benefits of dapagliflozin in patients with heart failure and reduced ejection fraction (HFrEF) and type 2 diabetes in DAPA-HF varied by background glucose-lowering therapy (GLT).

**Research design and methods:** We examined the effect of study treatment by the use or not of GLT, and by GLT classes and combinations. The primary outcome was a composite of worsening HF (hospitalization or urgent visit requiring intravenous therapy) or cardiovascular death.

**Results:** In the 2139 type 2 diabetes patients, the effect of dapagliflozin on the primary outcome was consistent by GLT use/no use (hazard ratio 0.72 [95%CI 0.58-0.88] versus 0.86 [0.60-1.23]; P-interaction=0.39) and across GLT classes.

**Conclusions:** In DAPA-HF, dapagliflozin improved outcomes irrespective of use/no use of GLT or by GLT type used in patients with type 2 diabetes and HFrEF.

**Trial Registration** - ClinicalTrials.gov Identifier: NCT03036124

## INTRODUCTION

Although sodium glucose cotransporter 2 inhibitors (SGLT2i) have been shown to improve cardiovascular outcomes in patients with type 2 diabetes, they are usually prescribed as second-line glucose-lowering therapy (GLT), most often in addition to metformin.<sup>1-3</sup>

Uncertainty about the place of SGLT2i in the management of patients with type 2 diabetes is reflected in differing recommendations in recent guidelines.<sup>4-8</sup> The placebo-controlled Dapagliflozin And Prevention of Adverse-outcomes in Heart Failure trial (DAPA-HF), in which the SGLT2i dapagliflozin reduced the risk of worsening heart failure (HF) and cardiovascular mortality in patients with heart failure and reduced ejection fraction (HFrEF), provides a unique opportunity to examine the efficacy of SGLT2i alone and in combination with other GLT in patients with type 2 diabetes.<sup>9</sup>

## RESEARCH DESIGN AND METHODS

DAPA-HF was a prospective, randomized, double-blind, placebo-controlled trial in patients with HFrEF, which evaluated the efficacy and safety of dapagliflozin 10 mg once daily, compared with placebo, added to standard care.<sup>9,10</sup>

In this *post-hoc* analysis, we included randomized patients with either undiagnosed (defined as central laboratory HbA1c  $\geq 6.5\%$  (48mmol/mol) at both screening and randomisation visits) or a medical history of type 2 diabetes. We examined the effect of dapagliflozin, compared with placebo, in subgroups (limited to those with >200 individuals to minimize the likelihood of type 1 error) by the use, or not of background GLT and by individual GLT classes (biguanides [hereafter referred to as metformin], sulfonylureas, dipeptidyl peptidase 4 [DPP-4] inhibitors and insulin). We examined the primary outcome, a composite of an episode of worsening HF (either an unplanned hospitalization or an urgent visit resulting in intravenous therapy for HF) or cardiovascular death, along with the individual components of cardiovascular death and HF hospitalization, and the prespecified secondary endpoints of all-cause mortality and the composite of total (first and recurrent) HF hospitalizations and cardiovascular death.

The cumulative incidence of the primary endpoint by treatment group in subgroups of interest was plotted using the Kaplan-Meier method. The effect of dapagliflozin compared with placebo was examined using Cox proportional-hazards models adjusted for history of hospitalization for HF and treatment-group assignment. An interaction test using a subgroup-by-randomized treatment interaction term was performed to assess for treatment effect modification within each subgroup. Analyses were performed using Stata version 16 (College Station, TX, USA). A p-value <0.05 was considered statistically significant.

## RESULTS

Of the 4744 randomized patients in DAPA-HF, 1983 (41.8%) had a documented medical history of type 2 diabetes and 156 (3.3%) had undiagnosed type 2 diabetes. Therefore, 2139 (45.1%) patients with type 2 diabetes were included in the analysis. Of these, 1596 (74.6%) were treated with GLT: metformin (47.7%), insulin (25.2%), sulfonylurea (20.6%), DPP-4 inhibitor (14.5%) and glucagon-like peptide-1 (GLP-1) receptor agonist (1.0%) (each alone or in combination). The baseline characteristics of patients by use of GLT and type of GLT are summarised in Supplemental Table 1 and 2.

Supplemental Figure 1 shows the cumulative incidence of the primary composite endpoint by randomized treatment in the subgroups of interest. The effect of dapagliflozin on the primary endpoint was consistent in patients taking GLT (hazard ratio 0.72 95%CI 0.58-0.88), and in those who were drug-naïve (0.86, 0.60-1.23; interaction  $p=0.39$ ) (Figure 1). When considering individual GLT classes (Figure 1) or combinations (Supplemental Figure 2) there was no statistically significant interaction between background GLT and the effect of randomized therapy on the primary composite outcome.

In general, the effect of dapagliflozin on each of cardiovascular death and HF hospitalization, was similar for individual GLT (Supplemental Figure 3) and combinations of these (Supplement Figure 2). Furthermore, no modification of treatment effect by background GLT was observed for the composite endpoint of total (first and recurrent) HF hospitalizations and cardiovascular death (Supplement Figure 4) or all-cause mortality (Supplement Figure 5).

## DISCUSSION

In this *post-hoc* analysis of DAPA-HF we found that the benefit of dapagliflozin, compared with placebo, in patients with type 2 diabetes and HFrEF was not influenced by background GLT use. The benefit of dapagliflozin was consistent in drug-naïve patients and across all classes of commonly used GLT, including metformin.

Perhaps the most interesting group of participants were the approximately 25% of individuals with type 2 diabetes in DAPA-HF who were not prescribed any GLT at baseline i.e. those in which randomized dapagliflozin became “first-line” GLT and pharmacological “monotherapy”. Despite limited power for subgroup analysis due to a relatively small number of patients and a lower event rate, the benefit of dapagliflozin on the primary endpoint seemed to be consistent with the effect in type 2 diabetes patients overall.

Metformin was the most commonly used GLT in DAPA-HF, taken by approximately half of patients with type 2 diabetes and HFrEF, despite limited evidence for its cardiovascular safety in this patient group.<sup>11</sup> Nevertheless, international HFrEF management guidelines support the use of metformin as first-line GLT in patients with type 2 diabetes.<sup>12</sup> It has been suggested that the benefit of SGLT2i is contingent on metformin use, based upon a subgroup analysis of the CANVAS trials.<sup>13</sup> This is clearly not the case from the present analysis of DAPA-HF or a post-hoc analysis of the EMPA-REG OUTCOME trial.<sup>14</sup>

Examination of outcomes in patients receiving the other major classes of GLT was also of interest. After metformin, insulin was the most widely used GLT and dapagliflozin was as effective in these participants, as compared to patients not taking insulin. Given the substantially higher event rate experienced by patients receiving insulin, compared to those



receiving other GLT, the *relative* risk reduction in insulin-treated individuals translated into an even larger *absolute* risk reduction and an NNT of only 16 to prevent one patient having the primary outcome over the median 18.2 months of follow-up. Furthermore, the benefits of dapagliflozin were again consistent whether added to a sulfonylurea or a DDP-4 inhibitor.

We believe our findings are relevant to the discussion that followed recent updated guidance on management of diabetes issued by the European Society of Cardiology (ESC) and jointly by the American Diabetes Association (ADA) and European Association for the Study of Diabetes (EASD).<sup>4-7</sup> Both recommendations emphasised that the cardiovascular benefits of SGLT2i, and GLP-1 receptor agonists, are obtained independently of starting HbA1c, an approach supported by the strategy employed in DAPA-HF. More controversially, the ESC guidance supported the use of SGLT2i, and GLP-1 receptor agonists, as “first-line” GLT and not necessarily as an adjunct to metformin, which had previously been the recommended initial GLT in most patients with cardiovascular disease.<sup>7</sup> Our data also support this recommendation, at least in patients with HFrEF and provide further evidence, along with the evidence of benefit in HFrEF patients without diabetes, to the view that the mechanisms of action underlying the cardiovascular benefits of dapagliflozin are independent of any glucose-lowering effect.<sup>15</sup>

As with all studies of this nature there are inherent limitations. The analyses were not prespecified and some had limited power, despite only including subgroups with >200 individuals. The small number of patients taking a GLP-1 receptor agonist at baseline prohibited further examination of this subgroup.

## **Conclusion**

In patients with type 2 diabetes and HFrEF, the reductions in risk of worsening HF and cardiovascular death with dapagliflozin were consistent across a range of background of GLT, and in patients receiving no GLT. Our data provide support for the use of dapagliflozin as first-line monotherapy in type 2 diabetes, at least in patients with HFrEF.

## Acknowledgements

**Authors contributions:** All authors contributed to the study design. KFD, PSJ, OB and JJVM contributed to the data analysis. All authors were involved in data interpretation and the writing or editing of the report. All authors read and approved the submitted version of the report.

**Guarantor's statement:** KFD and JJVM are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

**Sources of funding:** The DAPA-HF trial was funded by AstraZeneca. Prof McMurray is supported by a British Heart Foundation Centre of Research Excellence Grant RE/18/6/34217.

**Declarations of interest:** Dr Docherty reports his employer (the University of Glasgow) is paid by AstraZeneca for his involvement in the DAPA-HF trial during the conduct of the study; grants from Novartis, personal fees from Eli Lilly, outside the submitted work. Dr Jhund reports his employer (University of Glasgow) is paid by AstraZeneca for involvement in the DAPA-HF trial during the conduct of the study; consulting, advisory board, and speaker's fees from Novartis, advisory board fees from Cytokinetics, and a grant from Boehringer Ingelheim, outside the submitted work. Dr Bengtsson is an employee of AstraZeneca. Dr. DeMets reports personal fees from Frontier Science, personal fees and other from D L DeMets Consulting, personal fees from Actelion, personal fees from Population Health Research Institute, personal fees from Duke Clinical Research Institute, personal fees from BMS, personal fees from Medtronic, personal fees from Boston Scientific, personal fees from GSK, personal fees from Merck, personal fees from NIH/NIAID, personal fees from NIH/NHLDI, personal fees from Astra Zeneca, personal fees from Intercept, personal fees from Mesoblast, personal fees from Liva Nova, personal fees from DalCor, personal fees from Sanofi, outside the submitted work; . Dr. Inzucchi reports personal fees and non-

financial support from AstraZeneca, during the conduct of the study; personal fees from AstraZeneca, personal fees and non-financial support from Boehringer Ingelheim, personal fees and non-financial support from Sanofi/Lexicon, personal fees and non-financial support from Merck, personal fees from Zafgen, personal fees and non-financial support from VTV Therapeutics, personal fees and non-financial support from Abbott/Alere, personal fees and non-financial support from Novo Nordisk, outside the submitted work. Dr. Kober reports he is an executive committee member for the DAPA-HF study, payment from which will be administered by Rigshospitalet University Hospital, from Astra-Zeneca, during the conduct of the study; personal fees from Novartis as speaker, personal fees from Bristol-Myers Squibb as speaker, outside the submitted work. Dr. Kosiborod reports personal fees from AstraZeneca, during the conduct of the study; grants, personal fees and other from AstraZeneca, grants and personal fees from Boehringer Ingelheim, personal fees from Sanofi, personal fees from Amgen, personal fees from NovoNordisk, personal fees from Merck (Diabetes), personal fees from Janssen, personal fees from Bayer, personal fees from Glytec, personal fees from Novartis, personal fees from Applied therapeutics, personal fees from Amarin, personal fees from Eli Lilly, personal fees from Vifor Pharma, outside the submitted work. Dr Langkilde is an employee and shareholders of AstraZeneca. Dr Martinez reports personal fees from AstraZeneca, during the conduct of the study. Dr. Sabatine reports grants and personal fees from AstraZeneca, during the conduct of the study; personal fees from Althera, grants and personal fees from Amgen, personal fees from Anthos Therapeutics, grants from Bayer, personal fees from Bristol-Myers Squibb, personal fees from CVS Caremark, grants from Daichii-Sankyo, personal fees from Dalcour, personal fees from Dr Reddy's Laboratories, personal fees from Dyrnamix, grants from Eisai, personal fees from Esperion, personal fees from IFM Therapeutics, grants and personal fees from Intarcia, grants and personal fees from Jansen Research and Development, grants and personal fees from Medicine Company, grants and personal fees from Medimmune, grants and personal fees from Merck, grants and personal fees from Novartis, grants from Pfizer, grants from Quark Pharmaceuticals, grants from Takeda, outside the submitted work; and is a member of the TIMI Study Group, which has also received institutional research grant support through Brigham and Women's Hospital from: Abbott, American Heart Association, Aralez, Roche, and Zora Biosciences. Dr Sjöstrand is an employee and shareholder of AstraZeneca. Dr. Solomon reports grants from AstraZeneca, during the conduct of the study; grants and personal fees from Alnylam, grants and personal fees from Amgen, grants and personal fees from AstraZeneca, grants from Bellerophon, grants and personal fees from BMS, grants from

Celladon, grants and personal fees from Gilead, grants and personal fees from GSK, grants from Ionis, grants from Lone Star Heart, grants from Mesoblast, grants and personal fees from MyoKardia, grants from NIH/NHLBI, grants and personal fees from Novartis, grants from Sanofi Pasteur, grants and personal fees from Theracos, personal fees from Akros, grants and personal fees from Bayer, personal fees from Corvia, personal fees from Ironwood, personal fees from Merck, personal fees from Roche, personal fees from Takeda, personal fees from Quantum Genomics, personal fees from AoBiome, personal fees from Janssen, personal fees from Cardiac Dimensions, grants from Eidos, grants and personal fees from Cytokinetics, personal fees from Tenaya, personal fees from Daichi-Sankyo, personal fees from Cardurion, personal fees from Eko.Ai, outside the submitted work. Dr McMurray reports his employer (University of Glasgow) being paid by AstraZeneca during the conduct of the study; and his employer (University of Glasgow) being paid by Bayer, Cardiorentis, Amgen, Oxford Univeristy/Bayer, Theracos, Abbvie, Dalcor, Pfizer, Merck, Novartis, GlaxoSmithKline, Bristol-Myers Squibb, Vifor-Fresenius and Kidney Research UK (KRUK) outside the submitted work.

## REFERENCES

1. NHS Digital. Prescribing for Diabetes, England 2008/09 to 2018/19. 2019. <https://digital.nhs.uk/data-and-information/publications/statistical/prescribing-for-diabetes/2008-09---2018-19> (Accessed 15 May 2020)
2. Zelniker TA, Wiviott SD, Raz I, Im K, Goodrich EL, Bonaca MP, Mosenzon O, Kato ET, Cahn A, Furtado RHM, Bhatt DL, Leiter LA, McGuire DK, Wilding JPH, Sabatine MS. SGLT2 inhibitors for primary and secondary prevention of cardiovascular and renal outcomes in type 2 diabetes: a systematic review and meta-analysis of cardiovascular outcome trials. *Lancet* 2019;**393**:31–39.
3. Perkovic V, Jardine MJ, Neal B, Bompoint S, Heerspink HJL, Charytan DM, Edwards R, Agarwal R, Bakris G, Bull S, Cannon CP, Capuano G, Chu PL, Zeeuw D De, Greene T, Levin A, Pollock C, Wheeler DC, Yavin Y, Zhang H, Zinman B, Meininger G, Brenner BM, Mahaffey KW. Canagliflozin and renal outcomes in type 2 diabetes and nephropathy. *N Engl J Med* 2019;**380**:2295–2306.
4. Davies MJ, D'Alessio DA, Fradkin J, Kernan WN, Mathieu C, Mingrone G, Rossing P, Tsapas A, Wexler DJ, Buse JB. Management of hyperglycaemia in type 2 diabetes, 2018. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetologia* 2018;**61**:2461–2498.
5. Garber AJ, Handelsman Y, Grunberger G, Einhorn D, Abrahamson MJ, Barzilay JI, Blonde L, Bush MA, DeFronzo RA, Garber JR, Timothy Garvey W, Hirsch IB, Jellinger PS, McGill JB, Mechanick JI, Perreault L, Rosenblit PD, Samson S, Umpierrez GE. Consensus statement by the American Association of clinical Endocrinologists and American College of Endocrinology on the comprehensive type 2 diabetes management algorithm - 2020 executive summary. *Endocr. Pract.*

- 2020;**26(1)**:107-139.
6. Buse JB, Wexler DJ, Tsapas A, Rossing P, Mingrone G, Mathieu C, D'Alessio DA, Davies MJ. 2019 update to: Management of hyperglycaemia in type 2 diabetes, 2018. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetologia* 2020;**63**:221–228.
  7. Cosentino F, Grant PJ, Aboyans V, Bailey CJ, Ceriello A, Delgado V, Federici M, Filippatos G, Grobbee DE, Hansen TB, Huikuri H V., Johansson I, Jüni P, Lettino M, Marx N, Mellbin LG, Östgren CJ, Rocca B, Roffi M, Sattar N, Seferović PM, Sousa-Uva M, Valensi P, Wheeler DC, ESC Scientific Document Group. 2019 ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD. *Eur Heart J* 2020;**41**:255–323.
  8. Sattar N, McMurray JJ, Cheng AY. Cardiorenal risk reduction guidance in diabetes: can we reach consensus? *Lancet Diabetes Endocrinol* 2020; 2020 May;**8(5)**:357-360.
  9. McMurray JJV, Solomon SD, Inzucchi SE, Køber L, Kosiborod MN, Martinez FA, Ponikowski P, Sabatine MS, Anand IS, Bělohávek J, Böhm M, Chiang C-E, Chopra VK, Boer RA de, Desai AS, Diez M, Drozd J, Dukát A, Ge J, Howlett JG, Katova T, Kitakaze M, Ljungman CEA, Merkely B, Nicolau JC, O'Meara E, Petrie MC, Vinh PN, Schou M, Tereshchenko S, Verma S, Held C, DeMets DL, Docherty KF, Jhund PS, Bengtsson O, Sjöstrand M, Langkilde A-M. Dapagliflozin in Patients with Heart Failure and Reduced Ejection Fraction. *N Engl J Med* 2019;**21**:1995–2008.
  10. McMurray JJV, DeMets DL, Inzucchi SE, Køber L, Kosiborod MN, Langkilde AM, Martinez FA, Bengtsson O, Ponikowski P, Sabatine MS, Sjöstrand M, Solomon SD, McMurray JJV, DeMets DL, Inzucchi SE, Køber L, Kosiborod MN, Langkilde AM, Martinez FA, Ponikowski P, Sabatine MS, Sjöstrand M, Solomon SD, Investigators D-HC and. A trial to evaluate the effect of the sodium–glucose co-transporter 2 inhibitor

- dapagliflozin on morbidity and mortality in patients with heart failure and reduced left ventricular ejection fraction (DAPA-HF). *Eur J Heart Fail* 2019;**38**:665–675.
11. MacDonald MR, Eurich DT, Majumdar SR, Lewsey JD, Bhagra S, Jhund PS, Petrie MC, McMurray JJV, Petrie JR, McAlister FA. Treatment of type 2 diabetes and outcomes in patients with heart failure: A nested case-control study from the U.K. general practice research database. *Diabetes Care* 2010;**33**:1213–1218.
  12. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JGF, Coats AJS, Falk V, González-Juanatey JR, Harjola VP, Jankowska EA, Jessup M, Linde C, Nihoyannopoulos P, Parissis JT, Pieske B, Riley JP, Rosano GMC, Ruilope LM, Ruschitzka F, Rutten FH, Meer P Van Der. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J* 2016;**37**:2129–2200.
  13. Neuen B, Heerspink HJL, Neal B, Matthews DR, Zeeuw D de, Mahaffey K, Davidson J, Jardine MJ, Zoungas S, Perkovic V. Cardiovascular and renal outcomes with canagliflozin in people with type 2 diabetes according to baseline use of metformin. *Endocr Pract* 2019;**25**:99–100.
  14. Inzucchi SE, Fitchett D, Jurišić-Eržen D, Woo V, Hantel S, Janista C, Kaspers S, George JT, Zinman B. Are the cardiovascular and kidney benefits of empagliflozin influenced by baseline glucose-lowering therapy? *Diabetes, Obes Metab* 2020 Apr;**22**(4):631-639.
  15. Petrie MC, Verma S, Docherty KF, Inzucchi SE, Anand I, Belohlávek J, Böhm M, Chiang C-E, Chopra VK, Boer RA de, Desai AS, Diez M, Drozd J, Dukát A, Ge J, Howlett J, Katova T, Kitakaze M, Ljungman CEA, Merkely B, Nicolau JC, O’Meara E, Vinh PN, Schou M, Tereshchenko S, Køber L, Kosiborod MN, Langkilde AM, Martinez FA, Ponikowski P, Sabatine MS, Sjöstrand M, Solomon SD, Johanson P, Greasley PJ, Boulton D, Bengtsson O, Jhund PS, McMurray JJ V. Effect of



Dapagliflozin on Worsening Heart Failure and Cardiovascular Death in Patients With Heart Failure With and Without Diabetes. *JAMA* 2020; 2020; **323(14)**, 1353-1368.

## FIGURE LEGENDS

### **Figure 1: Effect of dapagliflozin compared to placebo on the risk of the primary composite outcome by background glucose-lowering therapy in patients with diabetes**

\*Overall effect calculated in all randomized patients (n=4744)

The primary outcome was a composite of worsening heart failure (hospitalization or an urgent visit resulting in intravenous therapy for heart failure) or death from cardiovascular causes. Patients on multiple glucose-lowering medications are included in each individual medication subgroup.

DPP-4 = dipeptidyl peptidase-4; 12 patients were prescribed saxagliptin.  
CI = confidence interval.