

Effect of dark chocolate on plasma epicatechin levels, DNA resistance to oxidative stress and total antioxidant activity in healthy subjects

A. Spadafranca^{1*}, C. Martinez Conesa², S. Sirini¹ and G. Testolin¹

¹Department of Food Science and Microbiology (di.S.T.A.M), Division of Human Nutrition, University of Milan, Via Celoria, 2, 20133 Milan, Italy

²Murcian Institute of Investigation and Agricultural Development (IMIDA), La Alberca (Murcia), Spain

(Received 18 February 2009 – Revised 2 October 2009 – Accepted 6 October 2009 – First published online 5 November 2009)

Dark chocolate (DC) may be cardioprotective by antioxidant properties of flavonoids. We investigated the effect of DC (860 mg polyphenols, of which 58 mg epicatechin) compared with white chocolate (WC; 5 mg polyphenols, undetectable epicatechin) on plasma epicatechin levels, mononuclear blood cells (MNBC) DNA damage and plasma total antioxidant activity (TAA). Twenty healthy subjects followed a balanced diet (55 % of energy from carbohydrates, 30 % from fat and 1 g protein/kg body weight) for 4 weeks. Since the 14th day until the 27th day, they introduced daily 45 g of either WC (*n* 10) or DC (*n* 10). Whole experimental period was standardised in antioxidant intake. Blood samples were collected at T₀, after 2 weeks (T₁₄), 2 h and 22 h after the first chocolate intake (T_{14+2h} and T_{14+22h}), and at 27th day, before chocolate intake (T₂₇), 2 h and 22 h after (T_{27+2h} and T_{27+22h}). Samples, except for T_{14+2h} and T_{27+2h}, were fasting collected. Detectable epicatechin levels were observed exclusively 2 h after DC intake (T_{14+2h} = 0.362 (SE 0.052) μmol/l and T_{27+2h} = 0.369 (SE 0.041) μmol/l); at the same times corresponded lower MNBC DNA damages (T_{14+2h} = -19.4 (SE 3.4) % v. T₁₄, *P* < 0.05; T_{27+2h} = -24 (SE 7.4) % v. T₂₇, *P* < 0.05; T_{14+2h} v. T_{27+2h}, *P* = 0.7). Both effects were no longer evident after 22 h. No effect was observed on TAA. WC did not affect any variable. DC may transiently improve DNA resistance to oxidative stress, probably for flavonoid kinetics.

Dark chocolate: Epicatechin: DNA damage: Total antioxidant activity

Free radicals are implicated in degenerative human diseases such as cancer and cerebro-cardiovascular pathologies by multiple mechanisms⁽¹⁾. There is evidence that reactive oxygen species, and in particular oxidative DNA damage, play a role in aetiology of atherosclerosis^(2,3). Recently, a wide number of studies have shown that the consumption of chocolate may induce cardioprotective effects by favourable action on blood pressure⁽⁴⁾, platelet function^(5,6), HDL cholesterol⁽⁷⁾, endothelium function⁽⁸⁾ and oxidative status⁽⁹⁾. The majority of work that investigated the antioxidant properties of chocolate focused on the effect on LDL oxidation susceptibility⁽¹⁰⁾, while fewer and more conflicting are the studies about the effect on plasma total antioxidant capacity^(11,12) and, although it has been widely shown whether oxidative DNA damage contributes to atherogenesis, to our knowledge, there are no studies *in vivo* that have investigated the effect of chocolate intake on DNA protection from oxidative damage.

Beneficial effects attributed to chocolate consumption and in particular antioxidant properties depend on flavonoid content. Cocoa and chocolate products have a much higher flavonoid concentration and total antioxidant capacity per weight than other flavonoid food sources such as red wine, green tea and black tea⁽¹³⁾. Among milk and dark chocolate (DC), the latter contains higher amounts of cacao and higher

amounts of flavonoids (i.e. 951 mg polyphenols/40 g serving compared to 394 mg in milk chocolate)⁽¹³⁾. In particular, the predominant flavonoids are catechin and epicatechin, mostly present in the form of procyanidins; epicatechin is absorbed from human intestine better than catechin⁽¹⁴⁾.

In the present study, we evaluated, in apparently healthy human volunteers, the effects of regular consumption of a 45 g serving of DC on plasma epicatechin concentrations, mononuclear blood cells (MNBC) DNA resistance to oxidative stress and plasma total antioxidant activity (TAA), in comparison with a white chocolate (WC) containing negligible amounts of flavonoids. Since data in the literature suggest that epicatechin plasma levels increase markedly 2 h after DC ingestion⁽¹⁵⁾, we also investigated the effect on DNA resistance to oxidative stress and total plasma antioxidant activity 2 h after consumption of the first and last servings.

Materials and methods

Reagents

All the chemicals used were AnalaR grade. Acetonitrile, methanol, ethanol, *n*-hexane, water (HPLC grade) and metaphosphoric acid were purchased from Merck Sharp

Abbreviations: DC, dark chocolate; MNBC, mononuclear blood cells; SD, standardised diet; TAA, total antioxidant activity; Trolox, (±)-6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxylic acid; WC, white chocolate.

* **Corresponding author:** Dr Angela Spadafranca, fax +39 02 503 16 191, email angela.spadafranca@unimi.it

& Dohme Italia S.p.A. (Rome, Italy). 2,2-Azinibis-(3-ethylbenzothiazoline-6-sulphonic acid), (\pm)-6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxylic acid (Trolox), potassium persulphate ($K_2S_2O_8$), Folin-Ciocalteu, L-ascorbic acid, α -glucuronidase and sulphatase (G 7017), Trizma base, perchloric acid, Na_2HPO_4 and Na_2 EDTA were purchased from Sigma-Aldrich (St Louis, MO, USA). Authentic standards of gallic acid, α -tocopherol, all-*trans*-retinol and (-)-epicatechin were obtained from Sigma-Aldrich.

Subjects

Twenty healthy individuals (ten men and ten women, mean age 24.2 (SE 0.7) years, range 20–32 years) were recruited from an Italian University campus by a written questionnaire about food habits, health and lifestyle.

All subjects were non-smokers, normal weight ($18.5 \text{ kg/m}^2 < \text{BMI} < 24.9 \text{ kg/m}^2$) and in good general health. Exclusion criteria were: diseases or serious medical conditions; supplements or medications (oral contraceptives included); special diets (vegan, vegetarian or macrobiotic diet, gluten-free diet); non-habitual fruit and vegetable consumption; sport d'elite. The present study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving human subjects were approved by the ethics committee of University of Milan. Written informed consent was obtained from all subjects.

Materials

DC with 70% cocoa and WC (cocoa butter) was supplied by Ferrero (Alba, Italy) in single servings of 45 g. The nutrient content of the two kinds of chocolates is presented in Table 1.

Experimental design

Study design was described in Fig. 1. The dietary intervention lasted for 4 weeks; it was characterised by an isocaloric diet balanced (55–60% of energy from carbohydrates, 25–30% from fat and 0.8–1 g protein/kg body weight) according to Recommended daily Assumption Levels of Nutrients⁽¹⁶⁾.

During the whole experimental period, intake of food antioxidants was limited to only two servings of fruit and vegetables each day, and not more than the amounts specified in Table 2 per week. Amount per serving was similarly

restricted to 50 g salad, 200 g other kinds of vegetables, 200 g apple and pear, 150 g banana and 250 g citrus fruit.

The volunteers were asked to complete a daily diary to verify, within the limitation of the technique, compliance with our dietary intervention.

The standardised diet (SD) was followed for 2 weeks before the intervention period to match the volunteers' short-term dietary habits and intake of antioxidant sources. In the final 2 weeks, the volunteers were given their preferred choice of either DC (45 g/serving) or WC (45 g/serving) chocolate to include. Of the twenty volunteers, five men and five women selected DC and the remainder WC. Volunteers were advised to consume chocolate as a snack with 40 g white bread. To avoid excess energy consumption while including the chocolate in their diet, the SD was adjusted; energy and in particular fat intake was reduced by replacing full-fat cows' milk with skimmed cows' milk. Crackers were eliminated as a snack, and the amounts of permitted sugar and pasta were also reduced.

Blood samples were collected at the beginning of the experimental period (T_0), the end of the SD alone (T_{14}), 2 and 22 h after the initial consumption (T_{14+2h} and T_{14+22h}) and again before and after the final chocolate serving (T_{27} , T_{27+2h} and T_{27+22h}). With the exception of T_{14+2h} and T_{27+2h} , all samples were collected at 08:30 h after an overnight fast.

Epicatechin extraction

Chocolate. Chocolate samples (1 g) were extracted following the method described by Gu *et al.*⁽¹⁷⁾. The analysis was carried out in duplicate using four samples of DC and four samples of WC. Each sample was from a different batch. Total polyphenols content was measured at 760 nm using the Folin-Ciocalteu reagent diluted tenfold before use, with gallic acid as the standard (5–10–25–50–100 $\mu\text{g/ml}$), and measurement after 5 min at 50°C⁽¹⁸⁾.

Plasma. Plasma samples were extracted and analysed by the method described by Ritchelle *et al.*⁽¹⁵⁾ and Rein *et al.*⁽¹⁹⁾ with some modifications; samples (400 μl) were added to 40 μl vitamin C–EDTA solution (200 mg vitamin C and 1 mg EDTA in 1 ml water) and 40 μl β -glucuronidase and sulphatase. The mixture was incubated at 37°C for 45 min. Plasma was extracted with acetonitrile (1 ml) and centrifuged at 10000 g for 5 min at 4°C. The supernatant was transferred to a tube containing 100 mg alumina and washed twice with 2 ml 2-amino-2-(hydroxymethyl)propane-1,3-diol buffer followed by a further wash with 2 ml methanol. Between each wash, the mixture was centrifuged at 4000 g for 5 min at 4°C and the supernatant discarded. After evaporation of the methanol, 250 μl perchloric acid was added to the alumina, which was vortexed for 1 min and centrifuged at 4000 g also for 1 min at 4°C. The supernatant was filtered using a polyvinylidene fluoride syringe filter (Whatman, Clifton, NJ, USA) and 100 μl were injected on to the HPLC system.

Quantification of epicatechin

HPLC analysis. The HPLC system Waters (Waters Corporation, Milford MA, USA) consisted of a 996 W photodiode and array detector, 1525 W Binary pump, 717 W plus

Table 1. Macronutrient composition of study chocolates

Nutrient/100 g chocolate	WC	DC
Energy (kJ)	2297	2063
Protein (g)	8.4	9.8
Carbohydrates (g)	53.6	30.0
Saccharose	41.9	30.0
Lactose	11.7	–
Fat (g)	33.4	37.1
Saturated	22.1	23.7
Polyunsaturated	10.3	12.3
Monounsaturated	1.0	1.1

WC, white chocolate; DC, dark chocolate.




Fig. 1. Experimental protocol of dietary intervention. T₀ (fasting), start experiment protocol; T₁₄ (fasting), end standard diet and start chocolate diet; T_{14+2h}, 2 h after the first chocolate intake; T_{14+22h} (fasting), 22 h after first chocolate intake; T₂₇ (fasting), last day of chocolate intake; T_{27+2h}, 2 h after last chocolate intake.

auto-sampler, array detector controlled by a data acquisition Empower software (Waters).

The HPLC analysis was performed on a reverse phase ODS HYPERSIL SB-C18 column (4.6 × 250 mm², 5-µm pore size, Thermo, San Jose, CA, USA) using a guard column ODS HYPERSIL SB-C18 (4.6 × 125 mm², 5-µm pore size, Thermo) at room temperature. Spectroscopy data from all peaks were captured in the range of 210–400 nm, and chromatograms were recorded at 279 nm. Separations were carried out at a flow rate of 1.5 ml/min with an isocratic mobile phase of 85 % Na₂PO₄ 10 mol/l, pH 3 and 15 % acetonitrile. The identification of the epicatechin was made by comparison of retention times and spectra with those of commercially available standard compound ((-)-epicatechin, E-1753, Sigma). For the purpose of quantification, linear regression models were determined using standard dilution techniques.

Evaluation of mononuclear blood cell DNA resistance to oxidative damage

The resistance of MNBC DNA against oxidative stress (strand breaks after treatment with H₂O₂ – 500 µmol/l – for 5 min) was evaluated by means of the Comet assay as reported in detail by Riso *et al.*⁽²⁰⁾. One hundred cells for each slide were electronically captured (400 × magnification) using an epifluorescence microscope (BX 60 OLYMPUS) equipped with an excitation filter BP520–550, dichroic beam-splitter DM565 and BA580-IF barrier filter (OLYMPUS, Olympus Italia S.R.L., Milan, Italy). The light source was a 100 W Hg lamp (Olympus Italia S.R.L.). The microscope was attached to a high-sensitivity charged coupled device video camera and to a computer provided with an image analysis system (Comet Program exploited on Image Pro-Plus;

Table 2. List of allowed and not allowed foods to standardise dietary antioxidant intake during the whole dietary intervention (frequencies of consumption for each week)

Food not allowed	Beer, chocolate, cocoa powder, tea, wine, fruit juices
Food allowed	Apple/banana/pear (13), citrus fruit (1), beet/fennel cooked (1), zucchini/aubergines (2), asparagus/French beans (2), mixed salad (7), tomatoes (2), tomato sauce (2), pesto sauce (1)

Immagini e Computer, Bareggio, Milan, Italy) and set to calculate the levels of strand breaks as percentage DNA in tail.

For each subject, the average percentage DNA in tail of control cells (cells that were not treated with H₂O₂) was subtracted from the average percentage DNA in tail of treated cells.

Analysis of total antioxidant activity

TAA of plasma samples was measured using the method of Re *et al.*⁽²¹⁾. Results were expressed as µmol Trolox equivalent/ml.

Antioxidant vitamins in plasma

To evaluate the basal levels of the main vitamins with antioxidant power and the effect of the standardised intake of food antioxidants during the dietary intervention, we analysed ascorbic acid, α-tocopherol and retinol plasma levels at every experimental time.

Plasma ascorbic acid was determined by the method of Mannino & Cosio⁽²²⁾. Plasma α-tocopherol and retinol were detected following a previously published method by Vuilleumier *et al.*⁽²³⁾.

Statistical analysis

All samples were prepared and analysed in duplicate. Statistical analysis was performed using STATISTICA (Statsoft Inc., Tulsa, OK, USA). A repeated-measures ANOVA design was used to investigate the effect of treatments. Differences between means were further evaluated by the least significant differences test and P values < 0.05 considered to be significant. Results are reported as the means with their standard errors.

The analysis of simple regression was used to evaluate the correlation between variables.

Results

All subjects finished the dietary intervention. Supplementation with DC or WC was well tolerated, and no subject reported untoward side effects. The collection of daily dietary diaries indicated that the volunteers were compliant with antioxidant standardisation.

Ascorbic acid, α -tocopherol and retinol plasma levels were within normal range at all times, and no difference was observed between the two groups. In both, the dietary standardisation caused a significant reduction in plasma ascorbic acid and α -tocopherol (T_{14} v. T_0 , $P < 0.05$), which was maintained until T_{27+22h} (Fig. 2). Retinol levels detected at T_0 were maintained similar for the whole experimental period.

Polyphenols in chocolate samples

Total polyphenol intake from DC (45 g) was 860 (SE 19) mg (expressed as gallic acid equivalents), of which epicatechin intake was 58 (SE 3.6) mg.

Ingestion of WC brought 5.1 (SE 0.7) mg total polyphenols, and epicatechin was not detectable.

Plasma epicatechin concentrations

Fig. 3 shows epicatechin plasma levels detected at all time points.

Two hours after the first DC consumption (T_{14+2h}), plasma epicatechin rose to 105 (SE 15) ng/ml (0.362 (SE 0.052) μ mol/l);




Fig. 2. Plasma ascorbic acid, α -tocopherol and retinol levels during the dietary intervention. Results are expressed as means with their standard errors (n 10). * $P < 0.05$ T_0 v. all other experimental points. ●, dark chocolate (DC); ◇, white chocolate (WC). Normal range: 6–20 mg/l (for ascorbic acid), 5.5–17.0 (for α -tocopherol), 0.4–1.2 μ g/ml (for retinol).




Fig. 3. Plasma epicatechin levels during the whole dietary intervention. Results are expressed as means with their standard errors (n 10). * $P < 0.05$ T_0 v. all other experimental points. ND, not detectable. ●, dark chocolate (DC); ◇, white chocolate (WC).

maximum level detected was 155 ng/ml (0.534 μ mol/l) and the minimum was 41 ng/ml (0.141 μ mol/l). Similarly, 2 h after the last DC consumption (T_{27+2h}), plasma epicatechin rose to 107 (SE 12.1) ng/ml (0.369 (SE 0.041) μ mol/l); maximum level detected was 163 ng/ml (0.562 μ mol/l) and the minimum was 71 ng/ml (0.243 μ mol/l).

Fasting plasma, collected before the dietary intervention (T_0), after SD (T_{14}), 22 h after the first DC intake (T_{14+22h}), at T_{27} before and 22 h after the last DC intake (T_{27+22h}), contained no detectable epicatechin in any subject (limit detection was 6 ng/ml). In the plasma of subjects in the WC group, epicatechin was undetectable at all time points during the investigation.

Mononuclear blood cells DNA damage

Fig. 4 shows MNBC DNA damage (as % of DNA in tail) evaluated after oxidative treatment at T_0 , T_{14} , T_{14+2h} , T_{14+22h} , T_{27} , T_{27+2h} and T_{27+22h} . SD consumption did not affect DNA resistance to oxidative stress at T_0 in either group.

Two hours after the first DC consumption (T_{14+2h}), MNBC DNA damage decreased significantly (-19.4 (SE 3.4) % v. T_{14} , $P < 0.05$). However, this effect was no longer evident after 22 h ($T_{14+22h} = 62$ (SE 1.4) % v. $T_{14} = 65$ (SE 2.8) %, $P = 0.65$).

Two hours after the last DC consumption (T_{27+2h}), we observed a significant reduction in MNBC DNA damage (-24 (SE 7.4) % v. T_{27} , $P < 0.05$), similar to that detected after the first DC consumption (T_{14+2h} v. T_{27+2h} , $P = 0.7$).




Fig. 4. Mononuclear blood cells DNA damage (%) evaluated by comet assay in white chocolate (WC, □; n 10) and dark chocolate (DC, ■) groups (n 10) during the experimental protocol. The results are expressed as means with their standard errors. * $P < 0.05$ T_{14+2h} , T_{27+2h} v. T_{14} , T_{14+22h} , T_{27} , T_{27+22h} .

and no longer evident 22 h after ($T_{27+22h} = 67.9$ (SE 0.9) % *v.* $T_{27} = 64$ (SE 2.8) %, $P=0.42$).

An inverse correlation between epicatechin concentrations, detected at T_{14+2h} and T_{27+2h} , and DNA damage was observed ($R^2 0.27$, $P<0.05$).

WC intake did not affect DNA resistance to oxidative damage at any time point.

Regular daily consumption of DC also did not affect DNA resistance to oxidative damage (fasting blood samples; T_{27+22h} *v.* T_{14} , $P=0.38$).

Total antioxidant activity

DC did not affect TAA after 2 h or following 14 d of daily intake (Table 3) as well as WC.

No significant correlation was observed between epicatechin concentrations, detected 2 h after the first and the last DC intake, and TAA ($R^2 0.009$, $P=0.65$).

SD did not affect TAA in any group detected at T_0 (T_{14} *v.* T_0 , $P=0.17$).

Discussion

Recent studies suggest healthy benefits associated with DC^(24–26). In particular, great attention has been directed to the antioxidant potential of cocoa and chocolate flavonoids, epicatechin especially, and the potential protective effects against risk of CVD⁽¹⁰⁾. Epicatechin from flavonoid-rich chocolate has the capacity to quench OH· (rather than approximately 100 times more effective than mannitol, a typical OH· scavenger⁽²⁷⁾, and it is rapidly absorbed in human subjects⁽²⁸⁾).

Plasma epicatechin concentrations typically return to baseline values within 6–8 h after consumption of a flavonoid-rich chocolate meal⁽²⁸⁾. Richelle *et al.*⁽¹⁵⁾, 2 h after consumption of DC (40 g) containing 892 mg polyphenols (82 mg were epicatechin), found that epicatechin plasma concentrations were comprised between 0.3 and 0.7 $\mu\text{mol/l}$ with mean value of 0.383 $\mu\text{mol/l}$; Wang *et al.*⁽²⁹⁾ administered DC (53 g) containing about 69 mg epicatechin to thirteen healthy volunteers, after 2 h, detected mean epicatechin

plasma concentration of 0.258 (SE 0.104) $\mu\text{mol/l}$. The present study confirmed that epicatechin plasma half-life is relatively short (<24 h), and showed that regular epicatechin intake over 2 weeks is not enough to increase fasting epicatechin plasma levels. Engler *et al.*⁽⁸⁾ reported that plasma epicatechin concentrations were markedly increased at 2 weeks (0.204 (SE 0.06) $\mu\text{mol/l}$, $P<0.001$) in eleven healthy adult subjects that consumed daily 46 g DC containing 1 mg/g epicatechin. However, this result considered chocolate consumption not fasting levels. In contrast, in the present study, we considered fasting plasma levels. Moreover, similar epicatechin plasma levels at 2 h following consumption of DC on the first and last occasions are not associated with a long-term increase in epicatechin plasma concentrations, and suggest that flavonoid plasma levels are dependent upon intake from recent food sources.

The present study showed an increase of DNA resistance to oxidative stress in healthy subjects, although this effect was transient and limited. The significant reduction of DNA damage, induced by oxidative treatment, observed at 2 h from the DC consumption and no longer evident 22 h later, may be related to epicatechin and its plasma kinetics⁽²⁷⁾ or indeed other bioactive compounds in the chocolate.

Epicatechin concentrations achieved at 2 h from chocolate intake in the present study were in accordance with flavonoid concentrations resulted in protection against DNA oxidation *in vitro*⁽³⁰⁾. Min & Ebeler⁽³⁰⁾ showed that the treatment of calf thymus DNA with catechin at low concentration (0.1 $\mu\text{mol/l}$) inhibited 8-OH guanine formation by 92%. Nair & Salvi⁽³¹⁾ showed that epicatechin offered protection to DNA against ionising radiation-induced damages; in particular, they reported that in mice administered intra peritoneally with epicatechin (40 mg/kg body weight) 1 h before γ -irradiation, DNA strand breaks in peripheral blood leucocytes were 37% less damage than those in mice administered with distilled water prior irradiation. The present experimental evidence and the present results support the hypothesis that decreased DNA damage depends on, at least in part, epicatechin levels. The shape of the molecule and the presence of OH· may enable epicatechin to interact with DNA to prevent damage as well as being a free radical scavenger. Planarity of the molecule and the presence of –OH groups might let epicatechin DNA intercalating activity and free radical scavenging action⁽³⁰⁾.

We did not observe any significant effect of DC consumption on plasma TAA. Although there is a consensus about the effect of several cocoa products in increasing plasma flavonoids, evidence concerning the resultant changes in total antioxidant capacity is conflicting^(10,12,32). The present results are in agreement with previous studies showing no increase^(12,33) in antioxidant capacity after ingestion of a cocoa snack. Experimental studies have shown that when human plasma is subjected to an *in vitro* oxidation, epicatechin and related catechins, at concentrations between 0.01 and 1 mmol/l, can, in a dose-dependent manner, prevent depletion of endogenous lipid-soluble antioxidants⁽³⁴⁾. In the present *in vivo* study, epicatechin concentrations detected were at least tenfold lower than those tested in previous studies; this could be one reason we did not observe any positive correlation between epicatechin plasma concentration and TAA in plasma. Retinol and α -tocopherol levels unchanged after DC consumption.

Table 3. Total antioxidant activity (TAA) in dark chocolate (DC) and white chocolate (WC) groups at all experimental times

(Mean values with their standard errors)

	TAA (mmol Trolox equivalents/l)			
	DC group (n 10)		WC group (n 10)	
	Mean	SE	Mean	SE
T_0	0.89	0.06	0.84	0.04
T_{14}	0.82	0.02	0.85	0.02
T_{14+2h}	0.77	0.04	0.83	0.03
T_{14+22h}	0.78	0.04	0.82	0.04
T_{27}	0.83	0.02	0.85	0.04
T_{27+2h}	0.85	0.04	0.84	0.03
T_{27+22h}	0.84	0.03	0.82	0.06

Trolox, (\pm)-6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxylic acid. DC intake did not significantly affect TAA at any time; no significant difference was observed between DC and WC groups.

However, this was not as a result of depletion by epicatechin, but more probably continued consumption during the SD and homeostatic control. In fact, we detected similar levels of the two lipid-soluble vitamins at 2 h when epicatechin levels peaked rather than 22 h when it was undetectable. Moreover, no differences were observed between DC and WC groups either in terms of antioxidant vitamin plasma levels or TAA.

Rein *et al.*⁽¹⁹⁾ reported that plasma antioxidant capacity increased at the 6-h time point following 53 and 80 g DC, although epicatechin concentrations peaked at 2 h reaching concentrations similar to those we observed. In the present study, TAA was evaluated only at 2 h and again at 22 h after ingestion. Thus, we cannot determine the TAA response between 2 and 22 h.

Furthermore, antioxidant status evaluated as plasma antioxidant activity reflects modifications of the whole antioxidant system rather than of single compound⁽³⁵⁾. Thus, other studies are necessary in order to clarify the effect of DC consumption on TAA, considering epicatechin, other flavonoids and their metabolites as insertion into a physiological antioxidant network.

In conclusion, a DC snack within a balanced diet can improve DNA resistance to oxidative stress in healthy subjects. However, this effect is transient because of flavonoid kinetics. Regular DC intake was no more beneficial than a single occasional intake. These findings suggest that the DNA protection provided by DC against chronic degenerative disease is via an unknown mechanism. The oxidative stress in fact affects inflammatory process and atherogenesis⁽³⁶⁾, but has also mutagenic power to the human genome with important consequences in oncogene expression and cancer pathogenesis⁽³⁷⁾. The present results are clinically encouraging especially in the field of the diet therapy of obesity, pathology related to greater incidence of CVD and cancer⁽³⁸⁾. In fact, DC, habitually excluded by hypoenergetic diets for its high-fat and energy content, is a sweet food that should be reconsidered: if included in controlled amounts, in a weight loss programme it could have healthy effects, and could improve the compliance of patients to diet therapy. However, further studies about bioavailability, kinetics of flavonoids from DC and protective effects against DNA oxidative damage are warranted. In fact, data about the dose, number of servings, times of administrations, dietary contexts and frequency of consumption could be important in order to give clear information to consumers about the benefits and risks of chocolate consumption.

Acknowledgements

There was no specific funding for the present work. The authors thank Ferrero (Soremartec, Alba, Italy) for supplying chocolate products. The idea of the present work was generated by A. S. and G. T. A. S. was involved in the planning of the work and she wrote the manuscript. C. M. C. contributed for writing the manuscript and carried out epicatechin analysis in plasma and chocolate samples. S. S. was involved with A. S. in comet assay, TAA and antioxidant vitamin analysis in plasma. G. T. was involved in funding providing. There are no conflicts of interest among members of the group.

References

- Ames BN, Shigenaga MK & Hagen TM (1993) Oxidants, antioxidants, and the degenerative diseases of aging. *Proc Natl Acad Sci U S A* **90**, 7915–7922.
- Halliwell B, Zhao K & Whiteman M (1999) Nitric oxide and peroxynitrite. The ugly, the uglier and the not so good: a personal view of recent controversies. *Free Radic Res* **31**, 651–669.
- Olinska R, Gackowska D, Foksinska M, *et al.* (2002) Oxidative DNA damage: assessment of the role in carcinogenesis, atherosclerosis, and acquired immunodeficiency syndrome. *Free Radic Biol Med* **33**, 192–200.
- Grassi D, Lippi C, Necozione S, *et al.* (2005) Short-term administration of dark chocolate is followed by a significant increase in insulin sensitivity and a decrease in blood pressure in healthy persons. *Am J Clin Nutr* **81**, 611–614.
- Innes AJ, Kennedy G, McLaren M, *et al.* (2003) Dark chocolate inhibits platelet aggregation in healthy volunteers. *Platelets* **14**, 325–327.
- Murphy KJ, Chronopoulos AK, Singh I, *et al.* (2003) Dietary flavanols and procyanidins oligomers from cocoa (*Theobroma cacao*) inhibit platelet function. *Am J Clin Nutr* **77**, 1466–1473.
- Mursu J, Voutilainen S, Nurmi T, *et al.* (2004) Dark chocolate consumption increases HDL cholesterol concentration and chocolate fatty acids may inhibit lipid peroxidation in healthy humans. *Free Rad Biol Med* **37**, 1351–1359.
- Engler MB, Engler MM, Chen CY, *et al.* (2004) Flavonoid-rich dark chocolate improves endothelial function and increases plasma epicatechin concentrations in healthy adults. *J Am Coll Nutr* **23**, 197–204.
- Gu L, House SE, Wu X, *et al.* (2006) Procyanidin and catechin contents and antioxidant capacity of cocoa and chocolate products. *J Agric Food Chem* **54**, 4057–4061.
- Wan Y, Vinson JA, Etherton TD, *et al.* (2001) Effects of cocoa powder and dark chocolate on LDL oxidative susceptibility and prostaglandin concentrations in humans. *Am J Clin Nutr* **74**, 596–602.
- Vlachopoulos C, Aznaouridis K, Alexopoulos N, *et al.* (2005) Effect of dark chocolate on arterial function in healthy individuals. *Am J Hypertens* **18**, 785–791.
- Wang JF, Shramm DD, Holt RR, *et al.* (2000) A dose response effect from chocolate consumption on plasma epicatechin and oxidative damage. *J Nutr* **130**, 2115S–2119S.
- Lee KW, Kim YJ, Lee HJ, *et al.* (2003) Cocoa has more phenolic phytochemicals and a higher antioxidant capacity than teas and red wine. *J Agric Food Chem* **51**, 7292–7295.
- Keen CL (2001) Chocolate: food as medicine/medicine as food. *J Am Coll Nutr* **20**, 436S–439S.
- Ritchelle M, Tavazzi I, Enslin M, *et al.* (1999) Plasma kinetics in man of epicatechin from black chocolate. *Eur J Clin Nutr* **53**, 22–26.
- Società Italiana di Nutrizione Umana (Italian Society of Human Nutrition) (1996) *LARN – Livelli di Assunzione Raccomandati di Nutrienti per la Popolazione Italiana (LARN – Levels of Recommended Intakes for the Italian Population)*. Rome: LARN.
- Gu L, House SE, Wu X, *et al.* (2006) Procyanidin and catechin contents and antioxidant capacity of cocoa and chocolate products. *J Agric Food Chem* **54**, 4057–4061.
- Singleton VL & Rossi JA (1965) Colorimetry of total phenolics with phosphomolibdic–phosphotungstic reagents. *Am J Enol Vitic* **16**, 144–158.
- Rein D, Lotito S, Holt RR, *et al.* (2000) Epicatechin in human plasma: *in vivo* determination and effect of chocolate consumption on plasma oxidant status. *J Nutr* **130**, 2109S–2114S.
- Riso P, Pinder A, Santangelo A, *et al.* (1999) Does tomato consumption effectively increase the resistance of lymphocyte DNA to oxidative damage? *Am J Clin Nutr* **69**, 712–718.

21. Re N, Pellegrini N, Proteggente A, *et al.* (1999) Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Rad Biol Med* **26**, 1231–1237.
22. Mannino S & Cosio MS (1997) Determination of ascorbic acid in foodstuffs by microdialysis sampling and liquid chromatography with electrochemical detection. *Analyst* **122**, 1153–1154.
23. Vuilleumier JP, Keller HE, Gysel D, *et al.* (1983) Clinical chemical methods for the routine assessment of the vitamin status in human populations. Part I: the fat-soluble vitamins A and E, and beta-carotene. *Int J Vitam Nutr Res* **53**, 265–272.
24. Vlachopoulos C, Alexopoulos N, Stefanadis C, *et al.* (2006) Effect of dark chocolate on arterial function in healthy individuals: cocoa instead of ambrosia? Effect of dark chocolate on arterial function in healthy individuals. *Curr Hypertens Rep* **8**, 205–211.
25. Umemura T, Ueda K, Nishioka K, *et al.* (2006) Effects of acute administration of caffeine on vascular function. *Am J Cardiol* **98**, 1538–1541.
26. Kay CD, Kris-Etherton PM & West SG (2006) Effects of antioxidant-rich foods on vascular reactivity: review of the clinical evidence. *Curr Atheroscler Rep* **8**, 510–522.
27. Hanasaki Y, Ogawa S & Fukui S (1994) The correlation between active oxygens scavenging and antioxidative effects of flavonoids. *Free Radic Biol Med* **66**, 845–850.
28. Holt RR, Lazarus S, Sullards MC, *et al.* (2002) Dimeric procyanidins in human plasma after the consumption of a flavanol-rich cocoa. *Am J Clin Nutr* **76**, 798–804.
29. Wang JF, Shramm DD, Holt RR, *et al.* (2000) A dose response effect from chocolate consumption on plasma epicatechin and oxidative damage. *J Nutr Suppl* **8 S**, 130, 2115S–2119S.
30. Min K & Ebeler SE (2008) Flavonoid effects on DNA oxidation at low concentrations relevant to physiological levels. *Food Chem Toxicol* **46**, 96–104.
31. Nair CKK & Salvi VP (2008) Protection of DNA from gamma-radiation induced strand breaks by epicatechin. *Mut Res* **650**, 48–54.
32. Serafini M, Bugianesi R, Maiani G, *et al.* (2003) Plasma antioxidant from chocolate. *Nature* **424**, 1013.
33. Fraga CG, Actis-Goretta L, Ottaviani JI, *et al.* (2005) Regular consumption of a flavanol-rich chocolate can improve oxidant stress in young soccer players. *Clin Dev Immunol* **12**, 11–17.
34. Lotito SB & Fraga CG (1998) (+)-Catechin prevents human plasma oxidation. *Free Radic Biol Med* **24**, 435–441.
35. Simonetti P, Gardana C & Pietta P (2001) Plasma levels of caffeic acid and antioxidant status after red wine intake. *J Agric Food Chem* **49**, 5964–5968.
36. Van Gaal LF, Mertens IL & De Block CE (2006) Mechanism linking obesity with cardiovascular disease. *Nature* **444**, 875–880.
37. Matés JM & Sánchez-Jiménez FM (2000) Role of oxygen species in apoptosis: implications for cancer therapy. *IJBCB* **32**, 157–170.
38. Calle EE & Kaas R (2004) Overweight, obesity and cancer: epidemiological evidence and proposed mechanism. *Nat Rev Cancer* **4**, 579–591.