

Effect of High-Intensity Interval Versus Continuous Exercise Training on Functional Capacity and Quality of Life in Patients With Coronary Artery Disease

A RANDOMIZED CLINICAL TRIAL

[AQ01]

Koldobika Villedabeitia Jaureguizar, MD; Davinia Vicente-Campos, PhD; Lorena Ruiz Bautista, MD; Cesar Hernández de la Peña, MD; María José Arriaza Gómez, MD; María José Calero Rueda, MD; Mahillo Fernández Ignacio, PhD

■ **PURPOSE:** There is strong evidence that exercise training has beneficial health effects in patients with cardiovascular disease. Most studies have focused on moderate continuous training (MCT); however, a body of evidence has begun to emerge demonstrating that high-intensity interval training (HIIT) has significantly better results in terms of morbidity and mortality. The aim of this study was to compare the effects of MCT versus HIIT on functional capacity and quality of life and to assess safety.

■ **METHODS:** Seventy-two patients with ischemic heart disease were assigned to either HIIT or MCT for 8 weeks. We analyzed cardiopulmonary exercise stress test data, quality of life, and adverse events.

■ **RESULTS:** High-intensity interval training resulted in a significantly greater increase in $\dot{V}O_{2peak}$ (4.5 ± 4.7 mL·kg⁻¹·min⁻¹) compared with MCT (2.5 ± 3.6 mL·kg⁻¹·min⁻¹) ($P < .05$). The aerobic threshold (V_{T1}) increased by 21% in HIIT and 14% in MCT. Furthermore, there was a significant ($P < .05$) increase in the distance covered in the 6-minute walk distance test in the HIIT group (49.6 ± 6.3 m) when compared with the MCT group (29.6 ± 12.0 m). Both training protocols improved quality of life. No adverse events were reported in either of the groups.

■ **CONCLUSIONS:** On the basis of the results of this study, HIIT should be considered for use in cardiac rehabilitation as it resulted in a greater increase in functional capacity compared with MCT. We also observed greater improvement in quality of life without any increase in cardiovascular risk.

KEY WORDS

coronary artery disease

functional capacity

high-intensity interval training

quality of life

Author Affiliations: Physical Medicine and Rehabilitation (Drs Villedabeitia Jaureguizar, Hernández de la Peña, and Arriaza Gómez), Infanta Elena University Hospital, Madrid, Spain; Physiotherapy (Dr Vicente-Campos), Francisco de Vitoria University, Madrid, Spain; Cardiology (Drs Calero Rueda and Ruiz Bautista), Infanta Elena University Hospital, Madrid, Spain; and Epidemiology (Dr Fernández Ignacio), IIS Fundación Jimenez Díaz, Madrid, Spain.

All authors have approved the final manuscript.

The authors declare no conflicts of interest.

Correspondence: Koldobika Villedabeitia Jaureguizar, MD, Infanta Elena University Hospital, Avda. Reyes Católicos, 21 (28342), Valdemoro, Madrid, Spain (koldovilledabeitia@hotmail.com).

DOI: 10.1097/HCR.0000000000000156

In Europe, cardiovascular diseases cause 1.8 million deaths per year.¹ Exercise has a class I recommendation² for the management of these patients and results in a 15% to 31% reduction in mortality because of cardiac causes.³

Physical exercise increases functional capacity such as peak oxygen uptake ($\dot{V}O_{2peak}$), which is considered to be the best predictor of survival in cardiovascular disease.^{4,5} In their meta-analysis of 33 studies involving 84 323 individuals, Kodama et al⁶ concluded that a 3.5 mL·kg⁻¹·min⁻¹ increase in $\dot{V}O_{2peak}$ (1 metabolic equivalent or MET) is associated with a 13% reduction in the risk of all-cause mortality and a 15% reduction in cardiovascular events. Similarly, Gullati et al⁷ confirmed in their study with 5721 asymptomatic women that, for every increase in exercise capacity of 1 MET, the risk of death was reduced by 17%. In addition, patients with myocardial infarction who were followed up for 19 years, an increase of 1 MET was associated with an 8% to 14% reduction in mortality.⁸

The majority of studies published on the benefits of exercise in cardiovascular disease use moderate continuous training (MCT) at 60% to 80% of $\dot{V}O_{2peak}$. These studies have shown a significant improvement in functional capacity between 12% and 31% of $\dot{V}O_{2peak}$.⁹

Over the past decade, high-intensity interval training (HIIT) has raised great interest in the context of cardiac rehabilitation because of excellent results with regard to morbidity and mortality outcomes.^{10,11} High-intensity interval training consists of a repeating series of high-intensity (peak interval) exercises, alternating with periods of low-intensity exercise (recovery interval). These brief recovery intervals require the patient to perform aerobic exercise practically without requiring energy production by the lactate-producing glycolytic system for energy. This avoids prolonged acidosis and prevents the sympathoadrenergic system from increasing cardiovascular effort, as the heart is already overworked.¹² Therefore, patients can comfortably maintain these high-intensity workloads, for a prolonged period and without any significant risk of cardiovascular complications.^{12,13} Recent reviews and meta-analyses have shown that using HIIT better results are obtained for certain prognostic variables related to morbidity and mortality, such as aerobic capacity, left ventricular function, endothelial function, and quality of life, without any additional cardiovascular risk.^{10,11}

The principal objective of this study was to compare the effect of 2 exercise training protocols (MCT vs HIIT) on functional capacity and quality-of-life variables. Our hypothesis was that HIIT would increase $\dot{V}O_{2peak}$ more than MCT in stable patients with

coronary artery disease. The secondary outcome was to determine the impact of the type of exercise on quality of life and to verify the safety of following these 2 exercise programs.

METHODS

Participants

A prospective, randomized clinical trial (NCT02168712) was conducted with patients referred by the Cardiac Rehabilitation Department who were diagnosed with stable New York Heart Association functional class I or II coronary artery disease with angina pectoris or myocardial infarction and no heart failure. To be included in the study, patients had to achieve a respiratory exchange ratio ≥ 1.10 during the initial cardiopulmonary exercise test (CPET). This respiratory exchange ratio value is often used as a criterion for achieving a maximum exercise effort.¹⁴ Patients who had residual ischemia (by electrocardiogram [ECG] criteria or angina symptoms), severe ventricular arrhythmias, uncontrolled hypertension, permanent pacemakers, or implanted cardiac defibrillators were excluded.

After signing an informed consent form, patients were randomized on a one-to-one basis to either the MCT or the HIIT group. The mode of exercise training was a cycle ergometer with 40 minutes per sessions, 3 days per week (total of 24 sessions over 2 months).

Patients entered the study within 6 weeks from the revascularization procedure. Clinical variables including the 6-minute walk test (6MWT) distance, selected CPET variables, and variables related to quality of life were recorded before and after the exercise training. Cardiopulmonary exercise tests were administered by staff blinded as to which exercise training group the patients were assigned.

CPET

All patients underwent exercise testing with a cycle ergometer (Ergoline900S, Ergoline GmbH, Bitz, Germany) including analysis of exhaled gases (UltimaCardiO₂, Medical Graphics Corporation, St Paul, Minnesota). The exercise test protocol was tailored to each patient's physical condition, with gradual increments of 10, 15, or 20 W/min. The same protocol was applied before and after the exercise training program. The objective of the exercise tests was to achieve a sustained effort for 8 to 12 minutes, with the aim of proper oxygen uptake ($\dot{V}O_2$) kinetics and maintaining a linear relationship between $\dot{V}O_2$, exercise workload, and heart rate (HR).

A 12-lead ECG was continuously monitored, and blood pressure was measured every 3 minutes during the exercise tests. Exercise workloads in watts and metabolic, cardiac, ventilatory, and electrocardiographic parameters were analyzed. The ECG was continuously monitored during the first 5 minutes of recovery. Recorded HR was determined from the computerized test reports and was the average of the last 5 RR intervals.

The first (V_{T1}) and second (V_{T2}) ventilatory thresholds were considered to be indicators of the aerobic and anaerobic thresholds, respectively, and were determined after the ventilatory equivalent method described by Skinner et al.¹⁴ The $\dot{V}O_2$ in $\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ and HR in $\text{beats}\cdot\text{min}^{-1}$ at V_{T1} were the parameters used to determine the MCT exercise intensity.

6MWT

We performed 2 consecutive 6MWT using a standardized protocol¹⁵ with the better of the 2 results being recorded. This process was used before and after the exercise training programs.

Steep Ramp Test

To design the HIIT program, we used the steep ramp test (SRT) protocol, according to the methodology described by Meyer et al.¹⁶ This exercise test protocol is composed of 2 minutes of free pedaling at 25 W followed by progressive 25-W increments every 10 seconds, maintaining a constant pedal cadence of between 50 and 60 rpm. The test was stopped when the patient could not maintain continuous pedal cadence for >40 rpm after encouragement to increase to 50 rpm and/or experienced hemodynamic and/or electrical alterations. The maximum exercise load achieved, as measured in watts, was the exercise parameter that was used to design the HIIT program for each patient.

MCT and HIIT Program Designs

The metabolic parameters obtained during the pre-training CPET were used to design the MCT program. Patients were asked to keep their training HR below an HR corresponding to the HR at V_{T1} during the first month. During the second month, the intensity of the exercise was adjusted, increasing to a training HR that corresponded to V_{T1} plus 10%.

The HIIT protocol used the methodology initially described by Meyer et al.¹⁶ and which was recently published by our group.¹⁷ In this type of exercise, the intensity was established using workload (watts), without taking HR into consideration as a measure for regulating the intensity of the exercise. The training workloads depended on the maximum workload achieved during the SRT. The intervals were designed as follows. In the first month of training, 20-second repetitions at an intensity corresponding to 50% of the maximum load reached with the SRT (peak intervals) were followed by 40-second recovery periods at 10%. In the second month of training, the intensity of exercise was adjusted using the results of a new SRT (Figure 1).

The total duration of both types of training was 40 minutes per session throughout the exercise program (including warm-up and cool-down). Table 1 summarizes the exercise time and intensity progression for both MCT and HIIT. Patients rated the peak level of exertion during each training session using the Borg Rating of Perceived Exertion Scale.¹⁸

Both types of exercise were reviewed and approved by the local Research Ethics Committee. Patients enrolled in the study participated in other activities established in our cardiac rehabilitation program that were aimed at managing psychological stress and learning about cardiac health habits. They were also taught to devise a home walking program for the days on which they did not have to attend sessions in




Figure 1. Interval training: 20-second repetitions at an intensity corresponding to 50% of the maximum workload reached with the steep ramp test, followed by 40-second intervals at 10%.

Table 1 • Program Designs for MCT and HIIT Groups

Week	Warm-Up Time and Intensity (MCT and HIIT)	Exercise Time and Intensity (MCT/HIIT)	Cool-Down Time and Intensity (MCT and HIIT)
1	12 min (25 W)	MCT: 15 min at V_{T_1} HIIT: 15 repetitions ^a	13 min (25 W)
2	10 min (25 W)	MCT: 20 min at V_{T_1} HIIT: 20 repetitions ^a	10 min (25 W)
3	7 min (25 W)	MCT: 25 min at V_{T_1} HIIT: 25 repetitions ^a	8 min (25 W)
4	5 min (25 W)	MCT: 30 min at V_{T_1} HIIT: 30 repetitions ^a	5 min (25 W)
5-8	5 min (25 W)	MCT: 30 min at ($V_{T_1} + 10\%$) HIIT: 30 repetitions ^b	5 min (25 W)

Abbreviations: HIIT, high-intensity interval training; MCT, moderate continuous training; SRT, steep ramp test.
^aIntervals of 50% (20 seconds) to 10% (40 seconds) of the maximum load reached in the first SRT.
^bIntervals of 50% (20 seconds) to 10% (40 seconds) of the maximum load reached in the second SRT.

hospital. The recommended intensity of walking was a perceived exertion of 11 to 13 on the Borg scale.

Safety of the Exercise Training Programs

To verify the safety of using this kind of aerobic exercise training, we made a daily record of any incidents or adverse effects that could limit the planned exercise. An incident was considered low if there were no repercussions and it was possible to start and/or restart training (eg, muscle overload, fatigue, muscular pain, and dyspnea without oxygen desaturation). A moderate incident was defined as one that limited the planned training (dyspnea with desaturation <94%, muscle injury, vasovagal conditions), and an incident was defined as severe if it was potentially life-threatening (ischemia, ventricular arrhythmia, hypertensive emergencies).

Quality-of-Life Questionnaires

A general quality-of-life (QOL) questionnaire (36-Item Short Form Health Survey [SF-36]) was given to the patients before and after the exercise training program. A disease-specific health-related QOL questionnaire, MacNew Heart Disease Health-related Quality of Life, was also administered before and after the exercise program.¹⁹

Statistical Analysis

Quantitative variables were described using means and standard deviations, and the qualitative variables were reported using frequency distributions. To evaluate the effect of each exercise protocol on the quantitative variables, pre- and postprogram values

were compared using Student's dependent samples *t* test. The effect was measured in absolute terms via the difference between the postprogram values and those obtained before training. These changes were described with the mean and standard deviation. Comparisons between the 2 training programs were made using Student's *t* test in the case of quantitative variables and using the χ^2 test of association or Fisher exact test for qualitative variables. All the comparisons were made using 2-tailed tests, and the level of significance was set at $P < .05$. The statistical analyses were done using R-3.0.1 for Windows (Microsoft Corporation, Redmond, Washington).

RESULTS

A total of 72 patients were included and studied (36 patients per group). At the start of the study, there were no significant differences between the groups with regard to clinical characteristics and medication use (Table 2).

Training Data

The intensity of exercise in the MCT group in the first month was $64.2\% \pm 8.5\%$ of the $\dot{V}O_{2peak}$ reached during the initial CPET (corresponding to the V_{T_1}) and $69.5\% \pm 8.7\%$ in the second month (corresponding to $V_{T_1} + 10\%$). The exercise workload applied at the peak intervals in the HIIT group after the Meyer et al methodology¹⁶ was $104.5\% \pm 22.2\%$ (first month) and $134.5\% \pm 29.7\%$ (second month) of the maximum load reached in the initial CPET corresponding to 50%

Table 2 • Patient Characteristics and Medication Use^a

	MCT (n = 36)	HIIT (n = 36)	P Value
Age, y	58 ± 11	58 ± 11	.82
Men	92	78	.21
Body mass index, kg/m ²	29.5 ± 4.1	29.6 ± 4.4	1.00
Waist circumference, cm	104 ± 9	104 ± 11	1.00
Hip circumference, cm	105 ± 8	104 ± 9	.88
Waist-to-hip ratio	0.99 ± 0.05	1.00 ± 0.07	.90
Cardiovascular risk factors			
Family history	47	46	1.00
Hypertension	57	65	.67
Diabetes mellitus	31	27	.88
Dyslipidemia	66	43	.09
History of smoking	81	73	.74
Active smoker	10	14	.64
Medical history			
Angina pectoris	50	43	.36
Myocardial infarction	50	57	.45
LVEF, %	59 ± 14	62 ± 11	.29
PCI	73	59	.35
CABG	15	22	.53
Conservative medical management	12	19	.44
Time from procedure to start of exercise program, d	54 ± 6	50 ± 4	.35
Medications			
β-Blockers	89	86	1.00
Calcium channel blockers	14	27	.27
ACE inhibitors	71	54	.20
Angiotensin receptor antagonists	20	19	1.00
Nitrates	11	11	1.00
Antiplatelet agents	97	97	1.00
Statins	94	100	.23
Antidiabetics	22	22	1.00

Abbreviations: ACE, angiotensin-converting enzyme; CABG, coronary artery bypass graft; HIIT, high-intensity interval training; LVEF, left ventricular ejection fraction; MCT, moderate continuous training; PCI, percutaneous coronary intervention.

^aContinuous data were reported as mean ± standard deviation; categorical data were reported as percentage.

of the SRT in both months. The resulting HR during the first and second months in the HIIT group was between V_{T1} and V_{T2} .

No significant differences were observed in perception of exercise between the HIIT and MCT groups during the first (RPE 11-13) and second (RPE 14-16) months of training. Adherence to the treatment sessions (the number of sessions attended compared with the number of sessions scheduled) was 87.5% in

the MCT group and 92% in the HIIT group. There was no significant difference between groups for adherence.

CPET and 6MWT

Results of testing for both groups before and after exercise training are summarized in Table 3. After 8 weeks of exercise training, both exercise programs significantly increased $V\cdot O_{2peak}$, with a greater increase

[AQ02]

Table 3 • CPET Variables and 6MWT Distance in Both Study Groups

	MCT Group			HIIT Group		
	Pretraining	Posttraining	Change	Pretraining	Posttraining	Change
CPET results						
Total exercise time, min	9.00 ± 2.02	10.1 ± 2.59	1.1 ± 1.8 ^c	7.80 ± 1.75	9.55 ± 2.26	1.7 ± 1.9 ^c
$\dot{V}O_2$ peak, mL·kg ⁻¹ ·min ⁻¹	20.3 ± 5.0	22.8 ± 6.5	2.5 ± 3.6 ^c	19.4 ± 4.7	24.0 ± 4.8	4.5 ± 4.7 ^{c,d}
$\dot{V}O_2$, % predicted	80 ± 17	89 ± 20	9.0 ± 13.2 ^c	80 ± 21	96 ± 19	16.5 ± 17.8 ^{c,d}
Resting HR, beats·min ⁻¹	61 ± 9	59 ± 8	-1.6 ± 8.2	66 ± 14	64 ± 10	-2.2 ± 10.0
Resting SBP, mm Hg	122 ± 18	116 ± 15	-5.6 ± 18.3	126 ± 15	123 ± 15	-3.1 ± 17.7
Resting DBP, mm Hg	74 ± 10	73 ± 9	-0.4 ± 11.3	76 ± 10	74 ± 8	-1.8 ± 11.0
Resting DP	7 368 ± 1 746	6 783 ± 1 388	-585 ± 1 614 ^a	8 333 ± 1 945	7 794 ± 1 421	-539 ± 1 727
Maximum HR, beats·min ⁻¹	118 ± 20	119 ± 21	1.0 ± 11.6	115 ± 14	126 ± 14	10.6 ± 11.7 ^{c,e}
Maximum SBP, mm Hg	171 ± 23	165 ± 26	-6.1 ± 22	171 ± 21	178 ± 23	7.2 ± 23.9 ^d
Maximum DBP, mm Hg	92 ± 12	89 ± 8	-3.0 ± 12.4	90 ± 11	92 ± 11	1.6 ± 10.9
Maximum DP	20 205 ± 4 880	20 098 ± 5 044	-107 ± 4 111	19 973 ± 4 099	22 312 ± 4 572	2 339 ± 4 386 ^{b,d}
Maximum HR, %	73 ± 14	73 ± 11	-0.2 ± 10.0	72 ± 10	77 ± 8	5.5 ± 8.7 ^{c,d}
Recovery HR at 1 min, beat·min ⁻¹	20 ± 9	21 ± 8	1.3 ± 8.0	15 ± 7	21 ± 7	5.8 ± 8.1 ^{c,d}
Maximum RER	1.12 ± 0.09	1.15 ± 0.10	0.03 ± 0.1	1.15 ± 0.09	1.19 ± 0.12	0.04 ± 0.1 ^a
Peak workload, W	109 ± 37	123 ± 41	13.1 ± 19.3 ^c	103 ± 38	129 ± 46	26.3 ± 23.5 ^{c,d}
HR at VT ₁ , beats·min ⁻¹	90 ± 14	89 ± 14	-0.9 ± 7.0	90 ± 11	93 ± 11	3.5 ± 9.2 ^{a,d}
$\dot{V}O_2$ at VT ₁ , mL·kg ⁻¹ ·min ⁻¹	12.6 ± 2.9	14.4 ± 3.5	1.8 ± 2.1 ^c	12.0 ± 2.6	14.5 ± 2.7	2.5 ± 3.0 ^c
Power at VT ₁ , W	55 ± 24	62 ± 27	7.9 ± 17 ^b	49 ± 22	65 ± 22	15.3 ± 16.1 ^c
HR at VT ₂ , beats·min ⁻¹	106 ± 17	107 ± 18	0.1 ± 7.0	102 ± 14	109 ± 14	6.9 ± 11.8 ^{a,d}
$\dot{V}O_2$ at VT ₂ , mL·kg ⁻¹ ·min ⁻¹	17.2 ± 4	19.1 ± 4.1	1.9 ± 2.9 ^b	17.0 ± 3.8	19.9 ± 3.9	2.9 ± 3.7 ^b
Power at VT ₂ , W	92 ± 38	106 ± 33	14.0 ± 15.5 ^c	88 ± 30	106 ± 28	17.2 ± 19.0 ^b
6MWT						
Distance, m	528 ± 96	558 ± 109	29.6 ± 12.0 ^c	531 ± 74	580 ± 80	49.6 ± 6.3 ^{c,d}

Abbreviations: BP, blood pressure; CPET, cardiopulmonary exercise test; DBP, diastolic blood pressure; DP, double product; HIIT, high-intensity interval training; HR, heart rate; MCT, moderate continuous training; RER, respiratory exchange ratio; SBP, systolic blood pressure; $\dot{V}O_2$, oxygen uptake; VT₁, first ventilatory threshold; VT₂, second ventilatory threshold; 6MWT, 6-minute walk test.
 Within-group differences: ^a*P* < .05; ^b*P* < .01; ^c*P* < .001.
 Between-group differences: ^d*P* < .05; ^e*P* < .001.

noted in the HIIT group (4.5 ± 4.7 mL·kg⁻¹·min⁻¹ vs 2.5 ± 3.6 mL·kg⁻¹·min⁻¹, for patients in the HIIT and MCT groups, respectively; *P* < .05). Both groups also showed a significant increase in the peak exercise workload achieved (MCT 13.1 ± 19.3 W vs HIIT 26.3 ± 23.5 W), with a significantly higher increase in the HIIT group (*P* < .001). A significant increase was observed in maximal HR in the HIIT group only (10.6 ± 11.7 beats·min⁻¹; *P* < .001). The $\dot{V}O_2$ and exercise workload at VT₁ and VT₂ significantly increased in both groups, but HR only increased in the HIIT group.

Thirty-two HIIT and 34 MCT patients, respectively, achieved VT₁ on the first CPET, and 35 and 34 patients, respectively, achieved it on the second, posttraining CPET. Concerning the achievement of VT₂, 20 HIIT and 25 MCT patients, respectively, achieved it on the initial CPET. On the CPET after the training programs, 29 HIIT and 32 MCT patients, respectively, achieved VT₂.

With regard to the HR recovery in the first minute after the exercise test, the only significant change was observed in the HIIT group (5.8 ± 8.1 ; *P* < .001).

Both groups showed significant (*P* < .001) increases in 6MWT distance (Table 4) after training, with a

Table 4 • Quality-of-Life Results in MCT and HIIT Groups^a

	MCT Group			HIIT Group		
	Pretraining	Posttraining	Change	Pretraining	Posttraining	Change
SF-36 ^a						
Physical functioning	73 ± 24	77 ± 23	4.3 ± 13.6	78 ± 15	83 ± 16	4.6 ± 13.3
Role—physical	51 ± 43	59 ± 44	8.1 ± 51.4	49 ± 42	54 ± 45	4.4 ± 49.4
Body pain	67 ± 30	73 ± 25	5.8 ± 28.2	72 ± 23	74 ± 25	2.4 ± 26.2
General health	58 ± 19	62 ± 22	3.9 ± 14.8	58 ± 18	63 ± 19	5.0 ± 17.4
Vitality	62 ± 18	67 ± 18	4.1 ± 14.7	57 ± 19	63 ± 23	6.2 ± 22.4
Social functioning	83 ± 22	83 ± 22	0.6 ± 23.4	82 ± 19	89 ± 17	7.7 ± 24.0
Role—emotional	73 ± 38	75 ± 40	2.9 ± 45.2	48 ± 44	73 ± 36	25.5 ± 47.9 ^c
Mental health	70 ± 20	73 ± 22	3.3 ± 14.6	64 ± 17	73 ± 18	9.3 ± 19.0 ^c
Self-reported health status	3 ± 1	2 ± 1	−0.6 ± 0.9 ^c	3 ± 1	2 ± 1	−0.9 ± 1.2 ^d
Physical health index	43 ± 11	46 ± 12	2.6 ± 7.3 ^b	47 ± 8	47 ± 7	−0.5 ± 7.6
Mental health index	48 ± 12	50 ± 14	1.6 ± 11.8	41.0 ± 12.4	49 ± 11	7.8 ± 14.0 ^c
MacNew						
Emotional domain	5.5 ± 1.1	5.7 ± 1.1	0.3 ± 0.8	5.3 ± 0.9	5.8 ± 0.9	0.5 ± 0.9 ^c
Physical domain	5.6 ± 0.9	5.9 ± 0.9	0.3 ± 0.7 ^b	5.5 ± 1.0	5.9 ± 1.0	0.4 ± 0.7 ^c
Social domain	5.7 ± 0.9	6.0 ± 0.8	0.4 ± 0.8 ^b	5.6 ± 0.9	6.0 ± 0.9	0.4 ± 0.7 ^c
Global domain	5.5 ± 0.9	5.8 ± 0.9	0.3 ± 0.7 ^b	5.3 ± 0.9	5.8 ± 0.9	0.5 ± 0.7 ^c

Abbreviations: HIIT, high-intensity interval training; MacNew, MacNew Heart Disease Health-related Quality of Life questionnaire; MCT, moderate continuous training; SF-36, 36-Item Short Form Health Survey.
^aAll data reported as mean ± standard deviation.
 Within-group differences: ^b*P* < .05; ^c*P* < .01; ^d*P* < .001.

greater increase in distance for the HIIT group compared with the MCT group (*P* < .05).

Quality of Life

After the training program, scores in all domains of the MacNew questionnaire significantly improved in both groups except for the emotional domain in the MCT group. With regard to the SF-36 QOL questionnaire, significant increases in the role—emotional, mental health, and self-reported health status scales—and the mental health index were observed in the HIIT group (Table 4).

Safety of the Training Intervention

No incidents or adverse events were recorded that limited the ability of patients to perform the prescribed exercise in either of the training programs.

DISCUSSION

Recent studies have revealed that HIIT is more effective than MCT for improving functional capacity and

other variables predictive of cardiovascular risk.^{10,11,20-22}

In our study, despite the increase in $\dot{V}O_{2peak}$ in both groups after 8 weeks of aerobic training, the improvement was significantly greater in the HIIT group. The mean increase in $\dot{V}O_{2peak}$ was $4.5 \pm 4.7 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ in the HIIT group and $2.5 \pm 3.6 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ in the MCT group. These results are important not only for the improved $\dot{V}O_{2peak}$ but also for the benefit that increased functional capacity has on survival. Before $\dot{V}O_{2peak}$ is considered to be the best predictor of survival in patients with cardiovascular diseases, HIIT may potentially contribute to the reduction of morbidity and mortality because its effect on increasing $\dot{V}O_2$.

The 24% increase (HIIT group) and the 12% increase (MCT group) in $\dot{V}O_{2peak}$ are similar to those reported by other authors,²¹ but are far from the results obtained by Wisloff et al,²⁰ who reported improvements in $\dot{V}O_{2peak}$ of 46% with HIIT and 15% with MCT. These greater differences in favor of the HIIT group could possibly be due to the fact that Wisloff's study was conducted on patients with heart failure and severe left ventricular dysfunction (average left ventricular ejection fraction ≈29%), with a lower baseline

functional capacity ($\dot{V}O_{2\text{peak}} = 13 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) and the use of a different interval protocol compared with this study.

VT_1 expressed in $\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ increased in both groups after the exercise program, but the HR at VT_1 in the HIIT group was significantly higher than in the MCT group. This finding reflects important peripheral and central adaptations to the exercise training as it allows patients to exercise longer at submaximal levels before the onset of fatigue and represents an improved aerobic capacity without increasing cardiovascular work.^{23,24} This shift in the VT_1 toward increased $\dot{V}O_2$ could account for the increase ($P < .05$) in meters walked during the 6MWT after training in the HIIT group ($49.6 \pm 6.3 \text{ m}$) compared with the MCT group ($29.6 \pm 12 \text{ m}$).

On the other hand, an improvement in acidosis tolerance was observed in both training groups, with a shift of the VT_2 closer to the $\dot{V}O_{2\text{peak}}$. This trend in the HIIT group could be due to a higher metabolic rate during exercise and a greater capacity of for eliminating or buffering the lactate accumulated during incremental exercise.²⁵ This change in the anaerobic threshold provides a possibility of training at higher intensity without producing undesirable metabolic changes when exceeding the VT_2 .²⁶

It is interesting to note that the maximum HR reached in the CPET after completing the exercise programs was significantly greater than at baseline testing in the HIIT group with no change in the MCT group. These findings in the HIIT group could be due to the fact that higher workloads were obtained after completing the program in the HIIT group ($103 \pm 38 \text{ W}$ vs $129 \pm 46 \text{ W}$) and were due to a significant increase in the exercise time on the second CPET (7.80 ± 1.75 minutes vs 9.55 ± 2.26 minutes; $P < .001$) and to an improved cardiovascular adaptation to exercise.

HR recovery in the first minute after maximum effort is considered by many to be predictive for future morbidity and mortality. A decrease in HR of $\leq 12 \text{ beats}\cdot\text{min}^{-1}$ during the first minute of recovery is considered abnormal.²⁷⁻²⁹ Improved HR recovery was observed only in the HIIT group, and this could be due to an increase in the vagal tone in patients in the HIIT group, as has been previously demonstrated.²⁸ These results suggest a greater contribution of the autonomic nervous system because of an increase in vagal tone that occurs after HIIT.²⁸ However, both the short- and long-term effects of HIIT on the autonomic nervous system are still unclear because of diversity in the results obtained in different studies.

Many of our results are similar to those reported by other authors, but it is difficult to compare them because of the different designs used for interval train-

ing programs. Different methods for determining the exercise intensity used during the peak and the recovery intervals include percentage of $\dot{V}O_{2\text{peak}}$ ²⁰; percentage of $\dot{V}O_2$ reserve³⁰; percentage of HR_{max} ^{20,31}; percentage HR reserve³²; or, as used in this study, percentage of workload.^{16,17} There are also differences between studies in the duration of the intervals, the number of peak and recovery periods per session, and the number of sessions per week.^{10,11,22} All of the above constitute important limitations because these variables significantly alter the total volume of training.^{10,11,22}

Our workgroup established the intensity of interval training exercise using exercise workload in watts as a percentage of the maximum workload achieved on the SRT. Our results show that the workloads applied in the peak intervals in this study were of high intensity and ranged between 104% and 134% of the maximum workload achieved during the initial CPET. Meyers et al²⁵ reported that using this methodology for the HIIT design, the exercise times during training at an intensity greater than 85% of the $\dot{V}O_{2\text{peak}}$ were prolonged and safe. This methodology described by Meyers et al²⁵ has been suggested as an alternative method for establishing the intensity of the short intervals,^{25,33} but we are aware that using it could be a limitation in our study because the SRT is not fully validated and is not a widely used test in cardiac rehabilitation.

Physiological improvement with both types of exercise training was accompanied by an improvement in QOL as measured by the MacNew and SF-36 questionnaires. Although no significant differences were observed between the 2 groups, the most analyzed domains in both questionnaires showed greater improvements in the HIIT group. All 3 domains in the MacNew questionnaire (emotional, physical, and social) improved demonstrated greater improvement in the HIIT group. This is a very important fact because it is related to adherence to exercise programs³⁴⁻³⁷ and it reflects good tolerance for and acceptability of the HIIT protocol.

In general, HIIT programs have been shown to be safe, although the number of studies is limited. As described in other studies,^{10,11,22,38} we observed good acceptability and tolerability by all of the patients in the HIIT group and no incidents or complications were recorded during the training program.

Although the absence of incidents in our study suggests that HIIT is safe way to prescribe exercise and the data are reassuring, we cannot draw a clear conclusion that HIIT is safe. Further long-term studies are necessary in order for HIIT to be accepted as a standard methodology for exercise training in cardiac rehabilitation. We recommend that large-scale, randomized clinical trials be conducted to investigate the impact of

HIIT on morbidity and mortality in patients with cardiovascular diseases. Fortunately, 2 multicenter, randomized studies are already underway in Europe: the SAINTEX-CAD coronary artery disease study³⁹ and the SMARTEX-HF heart failure study.⁴⁰ The results from these studies will be an important advance in being able to make definitive recommendations regarding HIIT and the previously mentioned problems of patient safety and future morbidity and mortality risk.

CONCLUSIONS

The results of our study documented that HIIT in low-risk patients diagnosed with chronic ischemic heart disease resulted in a greater increase in functional capacity compared with MCT and favored the HIIT group in both maximal ($\dot{V}O_{2peak}$) and submaximal (V_{T1} and V_{T2}) values. The acceptance and tolerability by patients, absence of adverse incidents, and its positive impact on the quality of life justify the use of HIIT as an alternative to standard exercise training protocols used in cardiac rehabilitation programs.

References

- Nichols M, Townsend N, Luengo-Fernandez R, et al. *European Cardiovascular Disease Statistics 2012*. European Heart Network, Brussels, Belgium and European Society of Cardiology, Sophia Antipolis, France; 2012.
- Perk J, De Backer G, Gohlke H, et al. European Guidelines on cardiovascular disease prevention in clinical practice (version 2012). The Fifth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of nine societies and by invited experts). Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *Eur Heart J*. 2012;33:1635-1701.
- Taylor RS, Brown A, Ebrahim S, et al. Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials. *Am J Med*. 2004;116:682-692.
- Kavanagh T, Mertens DJ, Hamm LF, et al. Prediction of long-term prognosis in 12169 men referred for cardiac rehabilitation. *Circulation*. 2002;106:666-671.
- Myers J, Prakash M, Froelicher V, et al. Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med*. 2002;346:793-801.
- Kodama S, Saito K, Tanaka S, et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA*. 2009;301:2024-2035.
- Gulati M, Pandey DK, Arnsdorf MF, et al. Exercise capacity and the risk of death in women: the St. James women take heart project. *Circulation*. 2003;108:1554-1559.
- Dorn J, Naughton J, Imamura D, et al. Results of a multicentre randomized clinical trial of exercise and long-term survival in myocardial infarction patients: the National Exercise and Heart Disease Project (NEHDP). *Circulation*. 1999;100:1764-1769.
- VanTol BA, Huijsmans RJ, Kroon DW, Schothorst M, Kwakkel G. Effects of exercise training on cardiac performance, exercise capacity and quality of life in patients with heart failure: a meta-analysis. *Eur J Heart Fail*. 2006;8:841-850.
- Smart NA, Dieberg G, Giallauria F. Intermittent versus continuous exercise training in chronic heart failure: a meta-analysis. *Int J Cardiol*. 2013;166:352-358.
- Aimee KC, Broadbent S, Birinder C. Interval training for patients with coronary artery disease: a systematic review. *Eur J Appl Physiol*. 2010;111:579-589.
- Billat LV. Interval training for performance: a scientific and empirical practice. *Sports Med*. 2001;31:75-90.
- Guiraud T, Nigam A, Gremeaux V, et al. High-intensity interval training in cardiac rehabilitation. *Sports Med*. 2012;42:587-605.
- Skinner JS, McLickell T. The transition from aerobic to anaerobic metabolism. *Res Q Exer Sport*. 1980;51:234-248.
- Hanson LC, McBurney H, Taylor NF. The retest reliability of the six minute walk test in patients referred to a cardiac rehabilitation programme. *Physiother Res Int*. 2012;17:55-61.
- Meyer K, Samek L, Schwaibold M, et al. Interval training in patients with severe chronic heart failure: analysis and recommendations for exercise procedures. *Med Sci Sports Exerc*. 1997;29:306-312.
- Villelabetia Jaureguizar K, Díaz-Buschmann I, Vaquerizo I, Calero MJ, Mahillo I. Entrenamiento interválico en pacientes con cardiopatía isquémica: metodología y análisis de resultados ergoespirométricos. *Rehabilitación*. 2011;45:327-334.
- Borg G. *Borg's Perceived Exertion and Pain Scales*. Champaign, IL: Human Kinetics; 1998.
- Höfer S, Lim LL, Guyatt GH, Oldridge N. The MacNew Heart Disease Health-Related Quality of Life Instrument: a summary. *Health Qual Life Outcomes*. 2004;2:3.
- Wisloff U, Stoylen A, Loennechen JP, et al. Superior cardiovascular effect of aerobic interval training versus moderate continuous training in heart failure patients: a randomized study. *Circulation*. 2007;115:3086-3094.
- Rognmo O, Hetland E, Helgerud J, Hoff J, Slørdahl SA. High intensity aerobic interval exercise is superior to moderate intensity exercise for increasing aerobic capacity in patients with coronary artery disease. *Eur J Cardiovasc Prev Rehabil*. 2004;11:216-222.
- Arena R, Myers J, Forman DE, Lavie CJ, Guazzi M. Should high-intensity aerobic interval training become the clinical standard in heart failure? *Heart Fail Rev*. 2013;18:95-105.
- Myers J, Ashley E. Dangerous curves. A perspective on exercise, lactate, and the anaerobic threshold. *Chest*. 1997;111:787-795.
- Sullivan MJ, Higginbotham MB, Cobb FR. Exercise training in patients with chronic heart failure delays ventilatory anaerobic threshold and improves submaximal exercise performance. *Circulation*. 1989;79:324-329.
- Meyer P, Normandin E, Gayda M, et al. High-intensity interval exercise in chronic heart failure: protocol optimization. *J Cardiac Fail*. 2012;18:126-133.
- Díaz-Buschmann I, Villelabetia KJ, Calero MJ, Sanchez-Aquino R. Programming exercise intensity in patients on beta-blocker treatment: the importance of choosing an appropriate method. *Eur J Prev Cardiol*. 2014;21:1474-1480.
- Dimopoulos S, Anastasiou-Nana M, Sakellariou D, et al. Effects of exercise rehabilitation program on heart rate recovery in patients with chronic heart failure. *Eur J Cardiovasc Prev Rehabil*. 2006;13:67-73.
- Kiviniemi AM, Tulppo MP, Eskelinen JJ, et al. Cardiac autonomic function and high-intensity interval training in middle-age men. *Med Sci Sports Exerc*. 2014;46(10):1960-1967.
- Currie KD, Rosen LM, Millar PJ, McKelvie RS, Macdonald MJ. Heart rate recovery and heart rate variability are unchanged in patients with coronary artery disease following 12 weeks of

- high-intensity interval and moderate-intensity endurance exercise training. *Appl Physiol Nutr Metab.* 2013;38:644-650.
30. Warburton D, McKenzie DC, Haykowsky MJ, et al. Effectiveness of high-intensity interval training for the rehabilitation of patients with coronary artery disease. *Am J Cardiol.* 2005;95:1080-1084.
 31. Munk P, Staal E, Butt N, Isaksen K, Larsen A. High-intensity interval training may reduce in-stent restenosis following percutaneous coronary intervention with stent implantation: a randomized controlled trial evaluating the relationship to endothelial function and inflammation. *Am Heart J.* 2009;158:734-741.
 32. Iellamo F, Manzi V, Caminiti G, et al. Matched dose interval and continuous exercise training induce similar cardiorespiratory and metabolic adaptations in patients with heart failure. *Int J Cardiol.* 2013;167:2561-2565.
 33. Guiraud T, Juneau M, Nigram A, et al. Optimization of high intensity interval exercise in coronary heart disease. *Eur J Appl Physiol.* 2009;108:733-740.
 34. Kloczek M, Kubinyi A, Bacior B, Kawecka-Jaszcz K. Effect of physical training on quality of life and oxygen consumption in patients with congestive heart failure. *Int J Cardiol.* 2005;103:323-329.
 35. Gibala MJ. High intensity interval training: a time-efficient strategy for health promotion. *Curr Sports Med Rep.* 2007;6:211-13.
 36. Bartlett JD, Close GL, MacLaren DP, Gregson W, Drust B, Morton JP. High-intensity interval running is perceived to be more enjoyable than moderate-intensity continuous exercise: implications for exercise adherence. *J Sports Med.* 2011;29:547-553.
 37. Puhon MA, Busching G, Schunemann HJ, et al. Interval versus continuous high intensity exercise in chronic obstructive pulmonary disease: a randomized trial. *Ann Intern Med.* 2006;45:816-825.
 38. Guiraud T, Nigram A, Juneau M, et al. Acute responses to high-intensity intermittent exercise in CHD patients. *Med Sci Sports Exerc.* 2011;43:211-217.
 39. Conraads VM, Van Craenenbroeck EM, Pattyn N, et al. Rationale and design of a randomized trial on the effectiveness of aerobic interval training in patients with coronary artery disease: the SAINTEX-CAD study. *Int J Cardiol.* 2013;168:3532-3536.
 40. Stoylen A, Conraads V, Halle M, et al. Controlled study of myocardial recovery after interval training in heart failure: SMARTEX-HF—rationale and design. *Eur J Cardiovasc Prev Rehabil.* 2012;19:813-821.

AUTHOR QUERIES

TITLE: Effect of High-Intensity Interval Versus Continuous Exercise Training on Functional Capacity and Quality of Life in Patients With Coronary Artery Disease: A Randomized Clinical Trial

AUTHORS: Koldobika Villeda Jaureguizar, Davinia Vicente-Campos, Lorena Ruiz Bautista, Cesar Hernández de la Peña, María José Arriaza Gómez, María José Calero Rueda, and Mahillo Fernández Ignacio

[AQ01]: Please verify the author affiliations and correct spelling of all author names.

[AQ02]: Please expand "RPE," if possible.