

## **Effect of *in vitro* and *in vivo* induction of polyethelene glycol-mediated osmotic stress on hybrid taro (*Colocasia esculenta* (L.) Schott)**

**Manas Ranjan Sahoo, Madhumita DasGupta and Archana Mukherjee**

*Regional Centre of Central Tuber Crops Research Institute,  
Bhubaneswar - 751019, Orissa, India*

### **ABSTRACT**

Sahoo, M. R., M. DasGupta and A. Mukherjee. 2006. Effect of *in vitro* and *in vivo* induction of polyethelene glycol-mediated osmotic stress on hybrid taro (*Colocasia esculenta* (L.) Schott). *Ann. Trop. Res.* 28(2):1-11.

Taro hybrid TSL (Topi X Satasankha), along with its parental lines, was evaluated for stress tolerance under *in vitro* and *in vivo* polyethelene glycol (PEG) -mediated osmotic stress conditions. Aseptic cultures were raised *in vitro* under control and PEG-mediated stress conditions. Stress index for survival and growth parameters like rooting ability in cultures and number of leaves produced per plantlet revealed the better response of hybrid line TSL than its parental lines. The plantlets were hardened and transferred to pots for evaluation of morpho-physico-biochemical parameters under PEG-mediated osmotic stress conditions. Significant variations were observed in stress index for plant height, number of leaves, leaf area, % relative water content (RWC), chlorophyll stability index (CSI), % injury by dessication and yield. Decline in biochemical traits like protein content but increase in catalase and peroxidase activities was observed under the osmotic stress conditions. Less variation in morpho-physico-biochemical characters were recorded in TSL under stress conditions as compared to its parents. The hybrid line-TSL showed tolerance to osmotic stress with a minimum yield reduction. TSL can be used for the future breeding program for the development of drought tolerant lines and can become the source of favorable genes for drought tolerance in taro.

**Keywords:** hybrid taro, *in vitro*, *in vivo*, PEG-6000, osmotic stress, stress index

*Correspondence:* M. R. Sahoo *Present Address:* Regional Centre of Central Tuber Crops Research Institute, Dumduma Housing Board, Bhubaneswar - 751019, Orissa, India 6521-A Tel. No +91 674 2470528 Email: mrsahoo2004@indiatimes.com

**DOI:** 10.32945/atr2821.2006



## INTRODUCTION

Taro (*Colocasia esculenta* (L.) Schott), a crop of immense commercial value, ranks fourteenth among vegetable crops worldwide. India is said to be the secondary center of origin of taro enriched with tolerant traits to abiotic stresses. Taro grows in a wide range of agroecological conditions - swampy lands, uplands under irrigation and also as a rainfed crop (Wilson, 1984). Despite its wide adaptive nature, production of taro can be affected severely by stress.

Drought is a major constraint to successful agriculture in the arid and semi-arid tropics. It has been found that from 1939 to 1978, over 40% of the total insurance indemnities for crop losses involved drought (Boyer, 1982). Hence, efforts are necessary to develop crop plants that can tolerate hostile environments. Some of the local cultivars of taro are found to have certain degree of tolerance to water deficit stress (Ravi and Chowdhury, 1997). However, cultivation under rainfed conditions can lead to severe yield loss (90-100%) due to erratic monsoon. In order to augment the taro breeding program for genetic improvement, studies have already been initiated to screen 115 taro accessions for tolerance to drought (Sahoo *et al.*, 2002). Polyethylene glycol 6000 (the number signifying molecular mass) is most frequently used in plant water deficit studies (Ng and Ekanayake, 1997) to induce dehydration by decreasing the water potential of the nutrient solution (Jacomini *et al.*, 1988). Hence, the present investigation was undertaken to evaluate the effects of *in vitro* and *in vivo* induction of polyethelene glycol-mediated osmotic stress on growth of hybrid-TSL and its parental lines- Topi and Satasankha.

## MATERIALS AND METHODS

The experiment was conducted at the Regional Centre of Central Tuber Crops Research Institute, Bhubaneswar, India. The Centre is located at 20° 15' N, 85° 52' E at an altitude of 26 m above the mean sea level. It is about 60 km away from the Bay of Bengal. The climatic condition of the place is warm, moist with hot summer and comparatively mild winter. The taro accession Satasankha that shows moderate tolerance to drought was crossed with Topi that has good potential to develop the hybrid line TSL. Cormels



obtained from the hybrid and its parents were used as materials for the present investigation.

### *In vitro* induction and evaluation for PEG-mediated stress

*In vitro* cultures were raised by inoculating cormel tips of TSL, Topi and Satasankha in agar-solidified (8%) Murashige and Skoog (MS) basal medium (Murashige and Skoog, 1962) supplemented with growth regulators [0.25 mg/l naphthalene acetic acid (NAA), 0.5 mg/l benzyl adenine (BA) and 0.25 mg/l gibberellic acid ( $GA_3$ )]. The MS media with all growth regulators and 3% sucrose were used as control. In order to induce osmotic stress (-2 bars), the same media were supplemented with PEG-6000 (118g/l). The pH of the media was adjusted to 5.8 and then, the media were sterilized at 121°C for 20 minutes. Surface sterilization of taro was done following the procedure of Mukherjee *et al.* (1998). For each treatment, 20 replications were maintained at  $25 \pm 1^\circ\text{C}$  with 12 hours photoperiod and  $30\mu\text{Em}^{-2}\text{s}^{-1}$  irradiance.

Survival rate of plantlets exposed to osmotic stress were calculated as the percentage of cultures with living plantlets. Other growth parameters such as rooting ability and number of leaves were recorded after 6 weeks of inoculation. *In vitro* raised plantlets were acclimatized, hardened in cups containing autoclaved soil and sand mixture (1:1 v/v) and then, were subsequently established in pots under control and PEG-mediated stress conditions.

### *In vivo* induction and evaluation for PEG-mediated stress

Potted cultures were laid out with *in vitro* plantlets to evaluate the effect of *in vivo* induction of PEG-mediated stress on TSL and its potential lines. The treated pots were irrigated with PEG-treated water (118 g/l) while normal irrigation was given to control pots. Six replications were maintained for both conditions. The variations in morphological characters such as plant height, number of leaves per plant and leaf area were determined at 30, 60 and 90 days after planting (DAP). The leaf area was calculated according to the linear measurement method of Biradar *et al.* (1978). Various physiological parameters like percentage relative water content (%RWC), chlorophyll stability index (CSI) and percentage injury by desiccation (% injury) were



studied following the methodologies developed by Barrs and Weatherley (1962), Kolcyorcas (1958) and Sullivan (1971), respectively. Biochemical parameters like protein content of the PEG-treated and control plants were recorded using Bradford method as explained by Mahadevan and Sridhar (1996). The activity of the antioxidant enzymes, catalase and peroxidase, was assayed by following the methodologies of Aebi (1983) and Kar and Feirabend (1984) using a UV spectrophotometer at 240 nm and 470 nm, respectively. The yield was recorded at harvest. Stress index (S.I.) for each parameter was calculated following the methodology developed by Ng and Ekanayake (1997). Data were transformed by angular transformation and analyzed for significance using ANOVA.

## RESULTS AND DISCUSSION

### *In vitro* induction and evaluation for PEG-mediated osmotic stress

Significant variations in growth parameters were observed among the lines evaluated *in vitro* under PEG-mediated stress conditions. Application of PEG to differentiate crop responses to osmotic stress has been well documented (Ng and Ekanayake, 1997). The stress index (SI) for survival was found to be lower in TSL (22.16%) than Satasankha (30.29%) and Topi (37.12%) [Fig. 1]. SI for number of roots produced per plantlet was less in TSL (40.69%) as compared to Satasankha (45.20%) and Topi (47.26%), whereas SI for the number of leaves produced per plantlet was recorded as low as 32.3% in TSL followed by 37.51% in Satasankha and 40.75% (Fig. 1). Similar results in survival of plantlets and growth retardations were noticed in cassava under PEG-mediated stress conditions (Ng and Ekanayake, 1997). Responses of cormel tips of hybrid and parental lines cultured on MS medium with growth regulators along with PEG-6000 are presented in Fig. 2A-C. Under *in vitro* stress conditions, multiple shoots formation was observed in the cormel tip cultures of hybrid taro (Fig. 2A). Satasankha responded well under stress conditions (Fig. 2B), whereas necrosis due to osmotic stress was observed in Topi (Fig. 2C). The present study indicates that the developed hybrid line TL






Figure 1. Effect of PEG on stress index for survival and growth parameters under in vitro conditions

showed more tolerance to osmotic stress with low SI for survival, rooting and leaf formation *in vitro*. In vitro raised plantlets were acclimatized, hardened in cups (Fig. 2D) containing sterilized soil and sand in equal proportion (1:1) and transferred subsequently to the pots for evaluation for stress tolerance (Fig. 2E).

*In vivo induction and evaluation for PEG-mediated osmotic stress*

Significant differences in retardations in growth parameters were observed among the lines evaluated under *in vivo* induced stress conditions as compared to the control. Plant height stress index (PHSI) was found to be significantly lower in TSL (28.53%, 27.03% and 17.92%) than in Satasankha (41.94%, 31.56% and 29.71%) and Topi (44.51%, 37.71% and 34.80%) at 30, 60 and 90 days after planting (DAP), respectively (Table 1). SI for number of leaves produced per plant was recorded to be 13.37%, 24.82% and 17.50%






Figure 2. Multiple shoot produced from cormel tip in hybrid-TSL under PEG-mediated osmotic stress (A); cormel tip culture of Satasankha and Topi under stress (B and C); acclimatization of *in vitro* plantlets; and evaluation for osmotic stress tolerance (E)



in TSL at 30, 60, and 90 DAP, respectively. Corresponding values in Satasankha (22.50%, 32.51% and 25.12%) and in Topi (39.13%, 33.46% and 33.04%) were higher. The stress index for leaf area was non significant at 30 DAP was significantly less in TSL at 60 and 90 DAP (31.81% and 15.85%). The leaf area stress index was recorded to be 56.12% and 15.94% in Satasankha and 47.29% and 38.08% in Topi at 60 and 80 DAP, respectively. This suggests decreased leaf area is an early adaptive response to water deficit and can be considered as a first line of defense against drought.

Significant variations in stress index for physiological parameters like percentage relative water content (%RWC), chlorophyll stability index (CSI) and percentage injury by desiccation (% injury) were observed among the lines. The SI for %RWC was lowest (1.56%) in TSL followed by Satasankha (1.84%) and Topi (3.77%) (Table 2). Reduction in %RWC was reported earlier in taro accessions growing under osmotic stress (Ravi and Chowdhury, 1997). Similar trends among the three accessions were noted in stress indices for CSI and yield. Reduction in total chlorophyll content under water stress conditions was discussed earlier by Indira and Kabeerathumma (1998) in sweet potato and Ravi and Chowdhury (1997) in taro. A high CSI value reflects that the stress did not have much effect on chlorophyll content of plants which in turn helps the plants to withstand stress through better availability of chlorophyll.

Biochemical analysis revealed that the protein content (mg/mg of leaves) was decreased in PEG-treated plants (Fig. 3A). It has been found that production of reactive oxygen species (ROS) is common of water stress leading to degradation of lipids, proteins and DNA. However, results indicate increase in catalase and peroxidase activities (n kat/ $\mu$ g protein) under stress conditions (Fig. 3B and C). The rate of increase in catalase and peroxidase activities under PEG-mediated osmotic stress conditions was higher in TSL than its parental lines which reflects its tolerance towards osmotic stress conditions. Probably higher level of these antioxidant enzymes in TSL play a vital role in scavenging of ROS and thereby minimizing the effect of stress.

A significant reduction in yield was observed among the lines under stress. Stress index for yield was recorded to be 24.20% in TSL, 44.45% in Satasankha and 52.31% in Topi (Table 2). The reduction in corm yield under water deficit stress was also attributed to reduction in crop growth rate, leaf






Figure 3. Effect of PEG on biochemical parameters in taro (A-C).



Table 1. Effect of PEG on morphological parameters in taro

		Stress Index (%)								
		Plant Height			Number of leaves /plant			Leaf Area		
SI No.	Accessions	30 DAP	60 DAP	90 DAP	30 DAP	60 DAP	90 DAP	30 DAP	60 DAP	90 DAP
1	Topi	44.51	37.71	34.80	39.13	33.46	33.04	41.73	47.29	38.08
2	Satasankha	41.94	31.56	29.71	22.50	32.51	25.12	40.66	56.12	15.94
3	TSL	28.53	27.03	17.92	13.37	24.82	17.50	41.67	31.81	15.85
	Mean	38.3	32.1	27.5	25.0	30.3	25.2	41.4	45.0	23.3
	SEm (0.05)	4.19	2.51	3.28	9.03	5.52	7.72	NS	3.13	3.48
	C. D. at 5%	12.63	7.55	9.88	27.21	16.64	23.26	NS	9.44	10.50

Table 2. Effect of PEG on physiological parameters and yield in taro

		Stress Index (%)			
SI No.	Accessions	%RWC	CSI	% Injury	Yield
1	Topi	3.77	4.47	21.24	52.31
2	Satasankha	1.84	4.39	11.36	44.45
3	TSL	1.56	2.04	2.88	24.20
	Mean	2.4	3.6	11.8	40.3
	SEm(0.05)	0.59	0.67	2.08	2.34
	C. D. at 5%	1.77	2.01	6.28	7.05



area and net assimilation rate (Ravi and Chowdhury, 1991).

## CONCLUSION

From the results of the present investigation, it can be concluded that the hybrid line TSL responded well to *in vitro* induced osmotic stress conditions with high survivability and less growth retardations than its parental lines Topi and Satasankha. The hybrid line also showed tolerance with regard to morpho-physico-biochemical parameters under *in vivo* PEG mediated osmotic stress conditions. The hybrid TSL developed from moderately susceptible parent Satasankha and high yielding line of Topi can be included in the genetic improvement program to develop elite lines tolerant to osmotic stress.

## ACKNOWLEDGMENT

The authors gratefully acknowledge the financial support of ICAR, NATP-CGP to carry out the present investigation. The authors are also thankful to the Head, Regional Centre of CTCRI, and Director of CTCRI for providing the necessary facilities to carry out the present study.

## REFERENCES

- AEBI, H. E. 1983. In: *Methods of Enzymatic Analysis* (H. U. Bergmeyer, J. Bergmeyer and M. Bl. Gra, eds.) 3rd Ed. Vol. 3. VCH, Weinheim, New York. 273-286 pp.
- BARRS, H. C. and P. E. WEATHERLEY. 1962. A re-examination of the relative turgidity technique for estimating water deficits in leaves. *Australian Biological Sciences*. **15**:413-428.
- BIRADAR, R. S., T. VENKATESWARLU and N. HIRISH. 1978. Leaf area estimation in Colocasia. *Root Crops*. **4**:51-53.
- BOYER, J. S. 1982. Plant productivity and environment. *Science*. **218**:443-448.
- INDIRA, P. and S. KABEERATHUMMA. 1988. Physiological response of sweet potato under water stress: 1. Effect of water stress during different phases of tuberization. *Root Crops*. **14**:47-50.



- JACOMINI, E., A. BERTANI and S. MAPELLI. 1988. Accumulation of PEG-6000 and its effects on water content and carbohydrate level in water stressed tomato plants. *Canadian J. Bot.* **66**:970-973.
- KAR, M., and J. FEIERABEND. 1984. Changes in the activities of the enzymes involved in amino acid metabolism during the senescence of detached wheat leaves. *Physiol. Plant.* **62**:39-44.
- KOLEYOREAS, S. A. 1958. A new method for determining drought resistance. *Plant Physiol.* **3**:22.
- MAHADEVAN, A. and R. SRIDHAR. 1996. Enzymes of infected plants and of parasites. In: *Methods in Physiological Plant Pathology*. Sivakami Publications, Chennai. 32 pp.
- MUKHERJEE, A., B. K. DEBATA and S. K. NASKAR. 1998. Somatic embryogenesis and chromosomal ability of regenerants in sweet potato and taro. *J. Scientific and Industrial Research.* **57**:709-715.
- MURASHIGE, T. and F. SKOOG. 1962. A revised medium for rapid growth and bioassays with tobacco tissue culture. *Physiol. Plant.* **15**:473-497.
- NG, S. Y. C. and I. J. EKANAYAKE. 1997. Screening cassava (*Manihot esculenta* Crantz) for polyethylene glycol-mediated osmotic stress *in vitro*. *J. Root Crops.* **23**(1):15-18.
- RAVI, V. and S. R. CHOWDHURY. 1997. Physiological response of taro to different soil moisture regimes. *Root Crops.* **17**:129-133.
- SAHOO, M. R., S. SAHU, A. MUKHERJEE, S. K. NASKAR and R. MISRA. 2002. In vitro screening of taro genetic resources for tolerance to biotic and abiotic stresses. In: *Plant Resources Utilization* (S. Sahoo, D. B. Ramesh, P. K. Panda and V. N. Misra, eds.) Allied Publishers, New delhi. 217-223 pp.
- SULLIVAN, C. Y. Techniques for measuring plant drought stress. In: *Drought Injury and resistance in crops* (K. L. Larson and J. D. Eastin, eds.) Crop Science Society of America. Madison, Wisconsin.
- WILSON, J. E. 1984. Cocoyam. In: *The Physiology of tropical field crops* (Goldworthy P. R. and N. M. Fisher, eds.) John Wiley and Sons Ltd. 589-604 pp.