

Effect of intimate partner violence on birth outcomes.

Tariku Laelago¹, Tefera Belachew², Meseret Tamrat²

1. Department of Nursing, Hossana College of Health Sciences, Hossana, Ethiopia.
2. Department of Population and Family Health, Jimma University, Jimma, Ethiopia.

Abstract

Background: Violence by intimate partner during pregnancy has many adverse pregnancy outcomes. Thus, that's why we sought to determine association between intimate partner violence during pregnancy and adverse birth outcomes.

Methods: A facility based cross-sectional study was conducted among 183 recently delivered women from March 31- April 30, 2014 in public health facilities of Hossana Town. The data were collected through structured questionnaire and record review. Women who were not mentally and physically capable of being interviewed and those admitted for abortion were excluded. Ethical clearance was obtained from Jimma University. Logistic regression analysis was employed to determine the association between intimate partner violence and adverse birth outcomes.

Results: About 23 % of women experienced intimate partner violence during pregnancy. The result of this study indicated an association of intimate partner violence with low birth weight of the new born (AOR:14.3,95% CI: (5.03, 40.7). Intimate partner violence was not associated with still birth, pre-term birth and Apgar score less than 7 at 5 minutes.

Conclusion: The findings of this study showed that intimate partner violence during pregnancy was associated with a low birth weight of the new born. Health sectors should train health care providers on how to screen, counsel, treat and follow up abused women.

Keywords: Intimate partner violence, birth outcomes, Ethiopia.

DOI: <https://dx.doi.org/10.4314/ahs.v17i3.10>

Cite as: Laelago T, Belachew T, Tamrat M. Effect of intimate partner violence on birth outcomes. *Afri Health Sci.* 2017;17(3): 681-689. <https://dx.doi.org/10.4314/ahs.v17i3.10>

Background

Intimate partner violence consists the use of physical, sexual or emotional abusive acts and controlling behaviours. An especially concerning form of physical violence during pregnancy is when abusive partners target a woman's abdomen which can cause hurting the woman and potentially jeopardizing the pregnancy^{1,2}. Intimate partner violence against women is a global public health concern, with 15% to 71% of women experiencing such abuse globally^{3,4}.

The prevalence of intimate partner violence during pregnancy among ever-pregnant women ranged from ap-

proximately 2% in Australia, Denmark, Cambodia and Philippines to 13.5% among ever-pregnant women in Uganda. More than half of the surveys had a prevalence estimate between 3.8% and 8.8%. However; prevalence appeared to be higher in the African and Latin American countries relative to the European and Asian countries though estimates within regions and countries were highly variable⁵. A systemic review of African studies on intimate partner violence against pregnant women reported that the overall prevalence of intimate partner violence during pregnancy ranges from 2.3% to 57.1%. The studies also reported prevalence rates of 23% to 40% for physical, 3% to 27% for sexual and 25% to 49% for emotional intimate partner violence during pregnancy⁶.

Domestic violence is common in Ethiopia in both urban and rural families. About 68% women agree that wife beating is justified in at least one of the specified situations⁷. About 71% of women experienced life time physical and sexual violence by their intimate partner⁸ while 8 % of women experienced intimate partner violence during pregnancy¹.

Corresponding author:

Tariku Laelago,
Department of Nursing,
Hossana College of health sciences,
Hossana, Ethiopia
Email: tarikulalago@gmail.com

Intimate partner violence during pregnancy has been found to be associated with adverse birth outcomes. The risk of being with born low birth weight and pre-term are associated with intimate partner violence. The effect of being born with low birth weight and preterm can result in immediate and long term health and developmental problems. The effects are not limited to women who experience only physical abuse; even psychological abuse has been linked to poor birth outcomes. Physical, mental health problems and negative health behaviours are some of maternal factors that are associated with intimate partner violence during pregnancy^{9,10}. Studies revealed the long-term effect of intimate partner violence as extremely premature and low birth weight infants. Such children commonly have cognitive deficits, motor delays including cerebral palsy, academic difficulties, language delays, and increased rates of attention problems, behavioural difficulties and psychological problems^{11,12,13}.

A longitudinal study demonstrates that maternal antenatal stress predicts children's behavioural and emotional problems four years later, with the underlying mechanism including the effect of maternal mood on the fetal brain development, which affects the child's behavioural development¹⁴.

Women who were reporting physical, sexual and emotional violence were more likely to deliver by caesarean section and to have abnormal progress of labour, premature rupture of membranes, low birth weight baby, preterm birth and any hospitalization before delivery than women who did not¹⁵.

A meta-analysis of 30 studies on maternal exposure to abuse and birth outcomes found that maternal exposure to domestic violence was significantly associated with an increased risk of low birth weight, as well as an increased risk of preterm birth¹⁶. Contrary to the meta-analysis study, study conducted in Canada identified no statistically significant associations for preterm birth and experience of violence¹⁷. Similarly, a study conducted in Karachi found the proportion of infants with birth weight < 2500 grams, gestational age < 37 weeks, or 5-minute Apgar < 7 was not significantly greater in abused women than in non-abused women¹⁸.

Understanding the relationship between violence during pregnancy and adverse birth outcomes could have important clinical and public health implications. Early iden-

tification and intervention to prevent violence against pregnant women might reduce adverse birth outcomes. In Ethiopia, information on effects of intimate partner violence on birth outcomes is limited. Therefore, the aim of this research was to determine association between intimate partner violence during pregnancy and adverse birth outcomes among recently delivered women in public health facilities of Hossana Town.

Methods

Study area and period

The study was conducted in Hossana Town, which is situated in Hadiya zone of South nation's nationalities regional state of Ethiopia. The Town is located 232 Kilometres far from Addis Ababa. In Hossana Town, there are three governmental health centres, one governmental hospital, and 25 private health facilities. The study was conducted in public health facilities. According to the information of Hadiya zone health department, 2,176 women gave birth in the four public health facilities of Hossana Town in 2013. The proportion of birth attended by health professional in the town was 27% in 2013. The study period lasted from March 31-April 30, 2014.

Study participants

Women who were giving birth in public health facilities of Hossana Town were recruited as source population for this study. The study population was all women who recently gave birth irrespective of whether the pregnancy outcome was alive or still birth. Women who were not mentally and physically capable of being interviewed and those admitted for abortion or for other pregnancy complications and later discharged undelivered were excluded in this study.

Data collection and measurement

Facility based cross sectional study was conducted among 195 recently delivered women in public health facilities of Hossana Town. The required number of sample was allocated proportionally among the four public health facilities. Consecutive sampling technique was undertaken by taking every woman who was presented in selected health facilities until allocated sample size was reached.

Pre-tested structured questionnaire and record review checklist were used to collect data from the study subjects. The questions were adopted from WHO multi-country

study on domestic violence⁸. Record review checklist was developed by principal investigator to collect data on new born birth outcomes. The questionnaire was first prepared in English, translated into Amharic, and then re-translated back to English to check for its consistency. Pre-test was carried out among ten study subjects who were not included in the actual study. Based on the result of pre-test, appropriate modifications were made to have the final version.

The data collection instrument comprises of socio-demographic and economic factors, experience of intimate partner violence during pregnancy, and record review checklist. The record reviews were conducted on delivery summary book of each respondent. The data collected on new born outcomes included alive birth, still birth, birth weight, Apgar score at 5 minutes, term and preterm birth.

Four female nurses and two supervisors' were recruited as interviewers and as supervisors respectively. Data collectors and supervisions were trained for one day on interviewing techniques, purpose of the study and ethical aspects.

The principal investigator and supervisor made a day to day on site supervision during the whole period of data collection and checked each questionnaire daily for its completeness and consistency.

Intimate partner was considered as current husband, co-habited (live in the same house without formal marriage), or boyfriend. Prevalence of intimate partner violence during pregnancy was measured by any act of physical, sexual or psychological abuse during last pregnancy by intimate partner.

We defined intimate partner violence according to the WHO Multi-Country Study⁸, as physical violence includes any of one or more (slapped, pushed or shoved, hit with fist or something else that could hurt her, beaten abdomen, choked or burnt on purpose, used or threatened to use knife, gun or weapon).

Sexual violence includes any of one or more (forced into sexual intercourse when she did not want, had sexual intercourse when she did not want to because she was afraid of what partner might do, forced to do something sexual that she found degrading or humiliating).

Psychological violence includes any of one or more (insult, humiliation, intimidated on purpose, threatened to hurt her or someone she cared about). Overall preva-

lence of intimate partner violence was measured by any act of physical, sexual or psychological violence during last pregnancy.

Adverse birth outcomes were measured by low birth weight, preterm birth, still birth or Apgar score <7 at 5 minutes. Low birth weight was defined as a live birth weighing < 2500 grams. Preterm was defined as a neonate born before 37 completed weeks. The cut off points used for the birth outcomes were based on standards¹⁹. Planned pregnancy was assessed by asking women whether her last pregnancy was planned by her and her partner. Recently delivered women were defined as women who were within 7 days of post-delivery in public health facility of Hossana Town.

Data processing and analysis

Data were entered into Epi-data version 3.1 20, then, it was exported into SPSS version 16.0 statistical software for further analysis²¹. Frequency tables and descriptive summaries were used to describe the study variables. Bivariate logistic regression analysis was used to see significance of association between the outcome and independent variables. Variables with P-value < 0.05 on bivariate analysis were transferred to multivariate logistic regression. Odds ratios at 95% CI were computed to measure the strength of the association between the outcome and the independent variables.

Multivariate logistic regression was performed to assess the relationship between intimate partner violence during pregnancy and adverse birth outcomes with adjustments for potential confounders. Confounding was assessed by entering potential confounders into a logistic model at a time, and then compared the unadjusted and adjusted ORs. We considered the following variables as possible confounders in these analyses. These were age of women, planned pregnancy, educational status, occupation, average monthly income, number of children, number of pregnancies and marital status. Final logistic regression models included covariates that altered unadjusted ORs by at least 10%²². P-value < 0.05 was considered as a statistically significant

Ethical consideration

We followed World Health Organization ethical and safety recommendations²³. Ethical clearance was taken from ethical clearance committee of the Jimma University.

Informed oral consent was taken from the respondents. The oral (non-written) consent was obtained because written consent demanded a certain level of education to read and sign the consent. The consent was documented by asking if the respondents want to participate in the study or not. The respondents were assured that they have full right to participate or withdraw from the study. The consent procedure was assured by ethical committee of Jimma University.

Interviewers were trained to be aware of the effects that the questions may have on informants and how best to respond, based on a woman's level of distress. Privacy and confidentiality of the respondents were protected.

Results

Of the 195 recently delivered women, 183 (94 %) gave an informed consent and were included in the study. The mean age women were 25.4 (SD \pm 4.6). Regarding to educational status, 64(35%) attained primary education. Occupation status of respondent's showed that house wife accounted 98(53.6%). One hundred seventy four (95%) of women were married. Ninety two (50.3%) of respondent became pregnant three or more times. For majority of women, 137(75%) their last pregnancy was planned. More than half of respondents 114(62.3%) had one to three children. The mean children women had were 2.8(SD \pm 1.9). The median monthly income of the respondent's was 1200 Ethiopian birr (table 1).

Table 1: Characteristics of the study population, Hossana, April 2014.

Variables	Intimate partner Violence(n=43)		No intimate partner violence(n=140)	
	Frequency	Percentage	Frequency	Percentage
Age groups(n=183)				
18-24	14	32.6	61	43.6
25-29	14	32.6	53	37.9
30-34	10	23.3	18	12.8
35-38	5	11.6	8	5.7
Educational status(n=183)				
No formal education	12	27.9	21	15
Primary education(grade 1-8)	17	39.5	51	36.4
Secondary education(grade 9-12)	8	18.6	34	24.3
Tertiary education(Above 12 grade)	6	14	34	24.3
Occupational status(n=183)				
Housewife	27	62.8	71	50.6
Student	3	7	14	10
Merchant	5	11.6	25	17.9
Government employee	7	16.3	26	18.6
Non-government employee	0	0	3	2.1
Other*	1	2.3	1	0.7
Marital status(n=183)				
Married	40	93	134	95.7
Never married	1	2.3	4	2.9
Divorced	2	4.7	2	1.4
Number of pregnancy(n=183)				
One or two	15	34.9	76	54.3
Three or above	28	65.1	64	45.7
Last pregnancy planned(n=183)				
Yes	19	44.2	118	84.3
No	24	55.8	22	15.7
Number of children(n=180)				
No child	0	0	1	0.7
One to three child	20	47.6	100	71.9
Four and above	22	52.4	38	27.3
Average monthly income(n=180)				
<1200 Ethiopian birr	29	67.4	70	51.1
\geq 1200 Ethiopian Birr	14	32.6	67	48.9

*daily workers

Forty three (23%) of women experienced at least one form of intimate partner violence during pregnancy. Psychological violence was the most common form 36 (20%) followed by physical 27(15%) and sexual 22

(12%). From physical violence, slapping 27(14.8 %) was the commonest form of violence. Physically forced to have sexual intercourse 22(12%), and insulting 36(19.7) were commonest form of sexual and emotional violence respectively (Table 2).

Table 2: Forms of intimate partner violence during pregnancy

Forms of violence (n=183)	No	Percent
All forms	43	23
Physical	27	15
Sexual	22	12
Psychological	36	20

Birth outcomes

One hundred eighty three records were reviewed to obtain the birth outcomes of the newborn. Of the 183 births, 175(95.6%) were live births and 8(4.4%) were still births. Of the live births, 39 (22.3%) were low birth weight. Only 16(8.7%) of the new born were preterm. Nine (5%) of new born Apgar scores were less than 7 at 5 minutes. From the total 183 births, 47(25.6%) had at least one adverse birth outcomes and sixteen (34%) had more than one adverse birth outcomes (Table 2).

Association between intimate partner violence during pregnancy and adverse birth outcomes.

Bivariate association was done using enter method to identify association between intimate partner violence during pregnancy and adverse birth outcomes. On bivar-

iate analysis, intimate partner violence during pregnancy showed significant association with low birth weight (COR=16.3(6.6, 40). There was no association between IPV and still birth (COR=0.29(0.07, 1.12), Apgar score less than 7 at 5 minutes (COR=1(0.2, 5.0) and preterm birth (COR=0.36(0.12.1.014).

A multivariate analysis was done to see significant association between intimate partner violence during pregnancy and low birth weight. After adjustment for age of women, number of pregnancy, planned pregnancy, educational status of women and monthly income, intimate partner violence during pregnancy was associated with low birth weight. The new born whose mothers were abused during pregnancy were 14.3 times more likely to be having low birth weight as compared to new born whose mothers were not abused during pregnancy (AOR=14.3(5.1,40.7) (Table 3).

Table 3: Association between birth outcomes and intimate partner violence during pregnancy

Birth outcomes	Violence (n=43)	No violence (n=140)	COR (95% CI)	AOR (95% CI)**
Low birth weight				
Yes	22(56.4)	17(43.6%)	16.3(6.6,40)*	14.3(5.1,40.7)*
No	10(7.3%)	126(92.7%)	1	1
Preterm birth				
Yes	7(43.8)%	9(56.2 %)	0.36(0.12,1.014)	-
No	36(21.6 %)	131(78.4 %)	1	
Apgar score less than 7 at 5 minutes				
Yes	2(22.2%)	7(77.8)	1(0.2,5.0)	-
No	37(22.3%)	129(77.7%)	1	
Still birth				
Yes	4(50%)	4(50%)	0.29(0.07,1.12)	-
No	39(22.3%)	136(77.7%)	1	
* p-value<0.05 **Adjusted for the following variables: age of women, number of pregnancy, pregnancy planned or unplanned, educational status of women and monthly income.				

Discussion

The overall prevalence of intimate partner violence during pregnancy in current study was 23%. The psychological violence was the most common form (20%) followed by physical (15%) and sexual (12%). This finding is consistent with studies done in other African countries. A review of literature on IPV in African countries found that its prevalence ranges from 2% to 57 %. The overall prevalence of intimate violence during pregnancy in current study is lower than Kenyan study, which showed 37% of women were abused during pregnancy^{6,24}. The difference may be due to the difference in culture and norms between the countries.

The prevalence of intimate partner violence during pregnancy in current study is higher than the WHO study of Ethiopia, which revealed 8% of women were abused during pregnancy¹. The difference may be due to difference in study setting and populations' characteristics. The current study has publicized that there is significant association between intimate partner violence during pregnancy and low birth weight of newborn. Studies

done in United States of America on maternal exposure to abuse and birth outcomes showed that there was an association of intimate partner violence and low birth weight^{9,16,25}. Similarly, studies done in Iran found an association between violence and low birth weight¹⁸. Other studies have also exhibited consistent result as current study^{26,27}.

The pathway through which violence may lead to risk of adverse birth outcomes has not been fully understood. One possible pathway is the direct effect of blunt physical trauma and subsequent adverse pregnancy outcome²⁸. A second potential pathway is elevated maternal stress levels and poor nutrition. Thus, these two factors are associated with low birth weight or preterm delivery and are well known risk factors for perinatal and infant mortality²⁹⁻³¹.

On the other hand, our study is inconsistent with the study done in Pakistan¹⁸ and Unites States of America, which presented that domestic violence during pregnancy, was not associated with low birth weight³². The differences may be due difference in study design and difference in population characteristics.

In current study, Apgar score less than 7 at 5 minutes, still birth and preterm birth were not greater in abused women than non-abused women. This finding is in line with study in Pakistan, which showed that proportion of infant with Apgar score less than 7 at 5 minutes, still birth and preterm birth was not significantly greater in abused women than non-abused women¹⁸. Similarly, Canadian and USA study found no statistically significant associations of preterm birth and violence^{17,32,33}.

Contrary to current study, study done in Peru and meta-analysis of 30 studies found maternal exposure to domestic violence was significantly associated with an increased risk of preterm birth^{10,16,34}. Previous study also showed the increased odds of low Apgar scores in infants of abused women³⁵. A research finding suggests that perinatal death (i.e., deaths in the first week of life and fetal deaths/stillbirths) is more common among women who experience violence during pregnancy³⁴. The discrepancies may be related to the large sample size of previous study. It may also be related to the difference in study setting, study population, and difference in operational definition of intimate partner violence.

Limitation of the study

Since this study was a cross sectional in design, it is difficult to establish causes and effect relationships among outcome of interest and explanatory variable. The sensitive nature of the topic and its proneness to response bias may lead to under-reporting of the true extent of the abuse.

Since we collected data from a secondary source there may be quality problems. Factors that can influence birth outcomes such as antenatal follow up, alcoholism, smoking and substance abuse were not collected in the survey, therefore adjusting for these factors was not possible.

Since the study was facility based and women who delivered at home were excluded in this study. Because of this reason, applicability of findings to the general population is uncertain. Besides, the wide confidence intervals were observed in this study and it may affect precision. This may be due to small sample size. Hence, interpretation of the finding shall take in to account all the factors mentioned above.

Conclusion

The findings of this study revealed that intimate partner violence during pregnancy is associated with adverse

birth outcome (low birth weight of new born). Thus, health sectors should train health care providers on how to screen, counsel, treat, and follow up of abused women. The study further calls for longitudinal research regarding the pathway through which violence may lead to risk of adverse birth outcomes.

Competing interests

The authors declare that they have no competing interests

Authors' contributions

Tariku Laelago wrote the proposal, participated in data collection, analysed the data, and drafted the paper. Tefera Belachew and Meseret Tamrat approved the proposal, participated in data analysis and revised subsequent draft of the paper. All authors read and approved the final manuscript.

Acknowledgements

We would like to thank Jimma University, college of public health and medical science for providing us ethical clearance. We would also like to show our gratitude to clients for their cooperation during our data collection time. Lastly, we would also extend our gratitude to the staff of public health facilities of Hossana town for their good cooperation and our data collectors for their full commitment during data collection period.

References

1. WHO. Intimate partner violence during pregnancy, information sheet. Available: http://whqlibdoc.who.int/hq/2011/WHO_RHR_11.35_eng.pdf.
2. Michelle Rice, Nat'l Ctr. for Posttraumatic Stress Disorder, Domestic Violence Fact Sheet. 2007. Available: http://www.ncptsd.va.gov/ncmain/ncdocs/fact_shfts/fs_domestic_violence.html #Anchor -Harway-46919.
3. Garcia-Moreno C, Jansen HA, Ellsberg M, Heise L, Watts CH. WHO Multi- country Study on Women's Health and Domestic Violence against Women Study Team. Prevalence of intimate partner violence: findings from the WHO multi-country study on women's health and domestic violence. *Lancet*. 2006; 368:1260–9. PubMed
4. Ellsberg M, Jansen HA, Heise L, Watts CH, Garcia-Moreno C. Intimate partner violence and women's physical and mental health in the WHO multi- country study on women's health and domestic violence: an observational study. *Lancet*. 2008; 371:1165-1172 PubMed .
5. Karen M Devries, Sunita Kishor, Holly Johnson,

- Heidi Stöckl, Loraine J Bacchus, Claudia Garcia-Moreno, Charlotte Watts: Intimate partner violence during pregnancy: analysis of prevalence data from 19 countries. *Reproductive Health Matters*. 2010; 18(36):158–70.
6. Shamu S, Abrahams N, Temmerman M, Musekiwa A, Zarowsky C. A Systematic Review of African Studies on Intimate Partner Violence against pregnant Women: Prevalence and Risk Factors. *PLoS ONE*. 2011; 6(3): e17591. PubMed doi:10.1371/journal.pone.0017591
 7. Central Statistical Agency Ethiopia and ICF International. Ethiopia Demographic and Health Survey, Addis Ababa, Ethiopia and Calverton, Maryland, USA: Central Statistical Agency and ICF International.2012.
 8. Garcia-Moreno C, Jansen HA, Ellsberg M, Heise L, Watts C.WHO multi- country study on women's health and domestic violence against women: initial result on prevalence, health outcomes and women's responses. Geneva, World Health Organization, 2005.
 9. Bailey BA. Partner violence during pregnancy: prevalence, effects, screening, and management, *International Journal of Women's Health*. 2010; 2,188.
 10. Sixto E Sanchez, Andrea V Alva ,Guillermo Diez Chang, Chungfang Qiu, David Yanez, Bizu Gelaye, and Michelle A.Williams:Risk of Spontaneous Preterm Birth in Relation to Maternal Exposure to Intimate Partner Violence During Pregnancy in Peru; *Maternal Child Health J*. 2013 April ; 17(3): 485–492. Doi: 10.1007/s10995-012-1012-0.
 11. Kilbride HW, Thorstad K, Daily DK. Preschool outcome of less than 801gram preterm infants compared with full-term siblings. *Pediatrics*. 2004; 113(4):742–747. PubMed
 12. Marlow N. Neurocognitive outcome after very preterm birth. *Arch Dis Child Fetal Neonatal* Ed. 2004; 89: 224–228.
 13. Taylor HG, Minich NM, Klein N, Hack M. Longitudinal outcomes of very low birth weight: neuropsychological findings. *J Int Neuropsychology Soc*. 2004; 10: 149–163.
 14. O'Connor TG, Heron J, Golding J, Beveridge M, Glover V. Maternal antenatal anxiety and children's behavioural/emotional problems at 4 years. Report from the Avon Longitudinal Study of Parents and Children. *British Journal of Psychiatry*. 2002; 180(6):502-508.
 15. M Faramarzi, S Esmaelzadeh, S Mosavi. Prevalence, maternal complications and birth outcome of physical, sexual and emotional domestic violence during pregnancy. *Acta Medica Iranica*. 2005; 43(2): 115-122.
 16. Shah PS, & Shah J, on Behalf of the Knowledge Synthesis Group on Determinants of Preterm/LBW Births. Maternal exposure to domestic violence and pregnancy and birth outcomes: A systematic review and meta-analysis. *Journal of Women's Health*. 2010. Available: <http://www.ncbi.nlm.nih.gov/pubmed/20919921> (accessed March 22, 2014).
 17. Marcelo L Urquia, Patricia J O'Campo, Maureen I Heaman, Patricia A Janssen, Kellie R Thiessen: Experiences of violence before and during pregnancy and adverse pregnancy outcomes: An analysis of the Canadian Maternity Experiences Survey: *BMC Pregnancy and Childbirth*. 2011; 11:42.
 18. Nabila Zareen, Nasreen Majid, Sonia Naqvi, Sadia Saboochi , Hassan Fatima: Effect of Domestic Violence on Pregnancy Outcome: *Journal of the College of Physicians and Surgeons Pakistan*. 2009 ;19 (5): 291-296.
 19. F Gary Cunningham, John C Hauth, Kenneth J Leveno, Larry Gilstrap III, Steven L Bloom, Katharine D Wenstrom. Williams's obstetrics. 22nd Ed. McGraw-Hill Medical Publishing Division; 2005.
 20. EpiData 3.1, <Http://www.epidata.dk>
 21. SPSS 16:00 for windows, www.SPSS.com
 22. Rothman, KJ. Greenland, S. Modern epidemiology. Philadelphia, PA: Lippincott-Raven; 1998.
 23. WHO: Putting women first: ethical and safety recommendations for research on domestic violence against women. WHO/FCH/GWH/01.1. Geneva: Department of Gender and Women's Health, Family and Community Health, World Health Organization; 2001.
 24. Lyndah A Makayoto, Jared Omolo, Abel M Kamweya, Valarie S Harder, Joseph Muta. Prevalence and Associated Factors of Intimate Partner Violence among Pregnant Women Attending Kisumu District Hospital. *Matern Child Health J*. 2013; 17:441–447.
 25. Sarkar NN: The impact of intimate partner violence on women's reproductive health and pregnancy outcome. *J Obstetric Gynaecology*. 2008; 28: 266–271.
 26. Grimstad H, Schei B, Backe B, Jacobsen G. Physical abuse and low birth weight: a case-control study. *Br J Obstet Gynaecology*. 1997; 104(11):1281-1287.
 27. Cokkinides VE, Coker AL, Sanderson M, Addy C, Bethea L. Physical violence during pregnancy: maternal complications and birth outcomes. *Obstetrics & Gynaecology*. 1999; 93(5 Pt 1):661-666. PubMed
 28. Connolly AM Katz VL, Bash KL, McMahan MJ, Hansen WF. Trauma and pregnancy. *Am J Perinatal*. 1997; 14: 331–336. PubMed

29. Jasinski JL. Pregnancy and domestic violence: a review of the literature. *Trauma Violence Abuse*. 2004; 5:47–64.
30. Hedin LW, Janson PO. The invisible wounds: the occurrence of psychological abuse and anxiety compared with previous experience of physical abuse during the childbearing year. *J Psychosomatic Obstet Gynaecology*. 1999; 20:136–144. PubMed
31. Campbell J, Torres S, Ryan J, King C, Campbell DW, Stallings RY, Fuchs SC. Physical and nonphysical partner abuse and other risk factors for low birth weight among full term and preterm babies: a multi-ethnic case control study. *Am J Epidemiol*. 1999; 150:714-726 PubMed .
32. Margaret H Kearney, Lois A Haggerty, Barbara H Munro, Joellen W Hawkins: Birth Outcomes and Maternal Morbidity in Abused Pregnant Women With Public Versus Private Health Insurance; *Journal of nursing scholarship*. 2003; 35:4:345-349. PubMed
33. Silverman J, Gupta J, Decker M, Kapur N, Raj A. Intimate partner violence and unwanted pregnancy, miscarriage, induced abortion, and stillbirth among a national sample of Bangladeshi women. *BJOG*. 2007; 114:1246–1252. PubMed
34. Janssen PA, Holt VL, Sugg NK, Emanuel I, Crichlow CM, & Henderson AD. Intimate partner violence and adverse pregnancy outcomes: A population-based study. *American Journal of Obstetrics and Gynaecology*. 2003; 188(5): 1341-1347.
35. Covington DL, Hage M, Hall T, Mathis M. Preterm delivery and the severity of violence during pregnancy. *Journal of Reproductive Medicine*. 2001; 46:1031-1039.