

RESEARCH ARTICLE

Effect of JAK Inhibitors on Release of CXCL9, CXCL10 and CXCL11 from Human Airway Epithelial Cells

Peter S. Fenwick¹, Patricia Macedo¹, Iain C. Kilty², Peter J. Barnes¹, Louise E. Donnelly^{1*}

1 Airway Disease, National Heart and Lung Institute, Imperial College London, London, United Kingdom, **2** Pfizer Inc, Cambridge, Massachusetts, United States of America

* l.donnelly@imperial.ac.uk



Abstract

Background

CD8⁺ T-cells are located in the small airways of COPD patients and may contribute to pathophysiology. CD8⁺ cells express the chemokine receptor, CXCR3 that binds CXCL9, CXCL10 and CXCL11, which are elevated in the airways of COPD patients. These chemokines are released from airway epithelial cells via activation of receptor associated Janus kinases (JAK). This study compared the efficacy of two structurally dissimilar pan-JAK inhibitors, PF956980 and PF1367550, and the glucocorticosteroid dexamethasone, in BEAS-2B and human primary airway epithelial cells from COPD patients and control subjects.

Methods

Cells were stimulated with either IFN γ alone or with TNF α , and release of CXCL9, CXCL10 and CXCL11 measured by ELISA and expression of CXCL9, CXCL10 and CXCL11 by qPCR. Activation of JAK signalling was assessed by STAT1 phosphorylation and DNA binding.

Results

There were no differences in the levels of release of CXCL9, CXCL10 and CXCL11 from primary airway epithelial cells from any of the subjects or following stimulation with either IFN γ alone or with TNF α . Dexamethasone did not inhibit CXCR3 chemokine release from stimulated BEAS-2B or primary airway epithelial cells. However, both JAK inhibitors suppressed this response with PF1367550 being ~50-65-fold more potent than PF956980. The response of cells from COPD patients did not differ from controls with similar responses regardless of whether inhibitors were added prophylactically or concomitant with stimuli. These effects were mediated by JAK inhibition as both compounds suppressed STAT1 phosphorylation and DNA-binding of STAT1 and gene transcription.

OPEN ACCESS

Citation: Fenwick PS, Macedo P, Kilty IC, Barnes PJ, Donnelly LE (2015) Effect of JAK Inhibitors on Release of CXCL9, CXCL10 and CXCL11 from Human Airway Epithelial Cells. PLoS ONE 10(6): e0128757. doi:10.1371/journal.pone.0128757

Academic Editor: Paul Proost, University of Leuven, Rega Institute, BELGIUM

Received: January 13, 2015

Accepted: May 1, 2015

Published: June 19, 2015

Copyright: © 2015 Fenwick et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This study was supported by the NIHR Respiratory Disease Biomedical Research Unit at the Royal Brompton and Harefield NHS Foundation Trust and Imperial College London and Pfizer Inc. The funders provided the molecules under assessment. However, the funders had no role in study design, data collection and analysis, or decision to publish.

Competing Interests: ICK is an employee of Pfizer Inc; PJB has served on Scientific Advisory Boards of AstraZeneca, Boehringer-Ingelheim, Chiesi, Daiichi-

Sankyo, GlaxoSmithKline, Glenmark, Johnson & Johnson, Merck, Novartis, Takeda, Pfizer, Prosonix, RespiVert, Sun Pharmaceuticals, Teva and UCB and has received research funding from Aquinox Pharmaceuticals, AstraZeneca, Boehringer-Ingelheim, Chiesi, Daiichi-Sankyo, GSK, Novartis, Takeda, Pfizer, Sun Pharmaceuticals. He is also a cofounder of RespiVert (now part of Johnson & Johnson), which has discovered novel inhaled anti-inflammatory treatments for asthma and COPD; LED has received unrestricted research funding from Pfizer Inc, Takeda, Cempra Inc. This does not alter the authors' adherence to PLOS ONE policies on sharing data and materials.

Conclusions

These data suggest that the novel JAK inhibitor, PF1367550, is more potent than PF956980 and that JAK pathway inhibition in airway epithelium could provide an alternative anti-inflammatory approach for glucocorticosteroid-resistant diseases including COPD.

Introduction

Type-1 helper (Th1) lymphocytes and CD8⁺ T cells are elevated in a number of inflammatory diseases including chronic obstructive pulmonary disease (COPD) [1] where these cells are located at the sites of airways obstruction [2, 3] and may contribute to emphysema via the production of granzyme B and perforins [4]. Recently, these cells have been shown to exhibit reduced apoptosis in COPD patients [5] leading to the persistence of these inflammatory cells in the airways. COPD is currently the fifth leading cause of death globally [6] and is increasing in prevalence with estimates that it affects ~10% of the population over the age of 40 [7]. Although inflammation underpins the pathophysiology of COPD, current anti-inflammatory treatments, including glucocorticosteroids, are ineffective [8]. Therefore, alternative strategies are required, for example, reducing recruitment of CD8⁺ cells to the airways of patients with COPD might prove to be beneficial.

The chemokine receptor, CXCR3 is highly expressed by activated Th1 and CD8⁺ lymphocytes and is thought to be involved in recruitment of these cells to the sites of inflammation [9]. CXCR3 binds to three distinct ELR negative ligands, CXCL9 (monokine induced by interferon γ (IFN γ); MIG), CXCL10 (interferon inducible protein of 10 kDa; IP10) and CXCL11 (interferon inducible T-cell α chemoattractant; ITAC) [10], all of which are elevated in the airways of patients with COPD [11] with CXCL10 being elevated in both sputum and serum during a viral exacerbation [12, 13] Although all three of these chemokines bind to the CXCR3 receptor, however CXCL11 has increased affinity and CXCL9 the least, implying a hierarchy of activity [9]. The source of these chemokines in the airways of COPD is unclear, however bronchial airway epithelial cells [14–16] and airway smooth muscle cells [17] release these chemokines following stimulation with interferon (IFN)- γ in both the presence and absence of tumour necrosis factor (TNF) α . Classically, binding of IFN γ activates Janus kinases (JAK) 1 and 2 leading to phosphorylation of signal transducer and activation of transcription (STAT)-1 protein, which subsequently dimerizes and binds to genes containing γ -activated sequences [18] including CXCL9, CXCL10 and CXCL11. STAT-1 independent mechanisms may also be invoked and STAT-3 and STAT-5 have been reported to be activated through the IFN γ receptor [19, 20]. Release of CXCL9, CXCL10 and CXCL11 from both airway epithelial cells and airway smooth muscles can be potentiated by the synergistic interactions of TNF α with IFN γ [14, 21]. In the airways of COPD patients, the concentrations of TNF α are elevated [22] and thus the expression of CXCL9, CXCL10 and CXCL11 by structural cells of the airways is likely to be enhanced, driving lymphocyte recruitment.

Previously, we have shown that the epithelial cell line BEAS-2B releases CXCL9, CXCL10 and CXCL11 in response to IFN γ in a manner that is glucocorticosteroid-insensitive but responsive to inhibition via the I κ B kinase, IKK2 [15]. The present study utilized our previous model to assess whether direct inhibition of the JAK pathway could suppress release of CXCL9, CXCL10 and CXCL11 from human lung epithelial cells using two structurally dissimilar compounds. PF956980 ([23]-pyrrolidin-1-yl-methanone hydrate) is a pan-JAK inhibitor with little effect against other kinases, including IKK [24] whereas the novel compound, PF1367550

(4-(3-(1H-benzo[d]imidazol-2-yl)-1H-indazol-6-yl)-3-ethylphenol), is a structurally differentiated, indazole pan-JAK inhibitor with excellent selectivity over non-JAK kinases [25]. Both of these JAK inhibitors were assessed for efficacy and potency to suppress CXCR3 chemokine release from both BEAS-2B cells and human primary airway epithelial cells with a view to understanding the utility of JAK inhibitors as therapeutic agents for inflammation in diseases such as COPD.

Materials and Methods

Materials

All chemicals and reagents were from Sigma Chemical Co (Poole, Dorset, UK) unless indicated otherwise. PF956980 and PF1367550 were kind gifts from Pfizer Inc. (Cambridge, Mass). All drugs were reconstituted at a concentration of 10mM in DMSO and diluted from these stock solutions for use in experiments. All experiments were performed with an appropriate vehicle control which had no effect on any of the responses measured.

Cell culture

BEAS-2B cells (catalogue number CRL-9609) were purchased directly from American Type Culture Collection (Rockville, MD). Keratinocyte serum-free medium (K-SFM), bovine pituitary extract (BPE), and recombinant human epidermal growth factor (EGF) were purchased from Invitrogen (Paisley, UK). BEAS-2B cells were cultured in K-SFM containing 50 µg/ml BPE and 5 ng/ml EGF at 37°C in a humidified atmosphere comprising 5% (v/v) CO₂ in air. Cells were cultured as monolayers in growth factor-free media for 24 h prior to stimulation. Human primary airway epithelial cells were cultured as monolayers from bronchial brushings from both male and female subjects (Table 1) and cultured in LHC-9 media (Invitrogen, Paisley, UK) in collagen (1% w/v) coated flasks. Samples were obtained from non-smokers, smokers and patients with COPD. Smokers had a smoking history of at least 10 pack-years and COPD patients were stable and fulfilled the American Thoracic Society criteria [26]. The subjects were matched for age and smokers and COPD patients for smoking history (Table 1). All subjects gave informed written consent and the study was approved by the NRES London-Chelsea Research Ethics committee, study number 09/H0801/85.

Table 1. Demographic data for subjects providing primary human airway epithelial cells.

	Non-smokersn = 4	Smokersn = 5	COPDn = 7
Age (years)	61 ± 10	53 ± 10	66 ± 4
Gender M:F	2:2	2:3	2:6
FEV ₁ (L)	2.2 ± 0.4	2.3 ± 0.3	0.8± 0.2** ##
FVC (L)	2.8 ± 0.6	2.9 ± 0.3	1.8 ± 0.2*
FEV ₁ (% predicted)	79 ± 2	89 ± 8	38 ± 10**
FVC (% predicted)	83 ± 1	93± 8	65 ± 9
FEV ₁ :FVC	0.83 ± 0.03	0.76 ± 0.05	0.44 ± 0.09*#
Smoking History (Pack Years)	-	23 ± 5	38 ± 6

Data are presented as mean ± S.E.M.

*p<0.05 and

**p<0.01 vs smokers

#p<0.05 and

##p<0.01 vs non-smokers.

doi:10.1371/journal.pone.0128757.t001

Measurement of CXCL9, CXCL10 and CXCL11

Cell-free supernatants were removed 20h post-stimulation and assayed for CXCL9, CXCL10 and CXCL11 using Duoset ELISA kits (R & D Systems Europe, Abingdon, UK) according to the manufacturer's instructions. The detection limits of these assays are 62pg/ml, 31pg/ml, and 7.8 pg/ml respectively.

Cell viability

Cell viability was determined colorimetrically by measuring the reduction of 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide, MTT, to formazan by mitochondrial dehydrogenases, as described previously [27].

Western Blotting

Enhanced chemiluminescence (ECL) reagent and Hybond-ECL nitrocellulose were obtained from GE Healthcare (Little Chalfont, UK). Bis-Tris SDS-PAGE (4–12%) gels and buffers were purchased from Invitrogen Ltd (Paisley, Scotland, UK). Antibodies against phosphorylated and total STAT-1 were purchased from New England Biolabs (Hitchin, UK). Epithelial cells were treated as indicated in six-well plates and lysed in 10 mM Tris-HCl buffer (pH 7.5) containing 150 mM NaCl, 2 mM EDTA, 1 mM sodium orthovanadate, 1% (v/v) Triton X-100, 1 mM phenylmethylsulfonyl fluoride, 10 µg/ml leupeptin, and 10 µg/ml aprotinin for 30 min on ice. The lysates were then centrifuged at 12,000g for 15 min and protein concentration determined using the BCA protein assay (BioRad, Hemel Hempstead, UK). The proteins within the lysates (20 µg) were resolved with 4–12% Bis-Tris gels and transferred onto nitrocellulose membranes. The membranes were then blocked with 5% (w/v) non-fat milk in TBS containing 0.1% (v/v) Tween 20 for 1 h at room temperature and then incubated with antibodies specific for pSTAT-1 overnight at 4°C. The primary antibody was detected with peroxidase-conjugated secondary antibodies and labelled proteins were detected by ECL.

STAT Activation Assays (TransAM)

STAT phosphorylation, nuclear translocation and DNA binding was determined using TransAM assays (Actif Motif, Rixensart, Belgium). Cells were treated with JAK inhibitors for 30 min prior to stimulation for 1h. Nuclear extracts were prepared using the Actif Motif assay reagents. Briefly, the nuclear extracts (10 µg) were incubated in DNA coated plates for 1h. After washing, antibodies supplied with the TransAM kit that were specific for the active forms of STAT1, STAT3, STAT5A and STAT5B were added to the wells and incubated for a further hour. The plates were washed and a secondary HRP-labelled anti-goat Ig antibody was added and the plates incubated for an hour. After washing, the assay was developed using the TMB substrate provided and the reaction stopped using 2N sulphuric acid. The plates were read at 450nm with a reference filter of 610nm. Data were then normalised to non-stimulated cells and calculated as fold change from baseline.

Measurement of mRNA Expression of CXCL9, CXCL10 and CXCL11

Total RNA was extracted from cells and reverse transcribed, as described previously [15]. Gene expression was determined by Taqman real-time PCR on a 7500 Real Time PCR system (Applied Biosystems, Life Technologies Ltd, Paisley, UK) using PCR Master Mix Reagent and 'assays on demand' (Applied Biosystems). HPRT1 gene expression was used as the housekeeping gene and data presented as $\delta\delta$ CT relative to baseline.

Statistical Analyses

Data are presented as mean \pm SEM of 'n' independent determinations. Concentration-response curves were analysed by least-squares, non-linear iterative regression with the 'Graphpad Prism' curve fitting program (GraphPad software, San Diego, USA) and IC₅₀ values were subsequently interpolated from curves of best-fit. Statistical differences were determined using the Kruskal-Wallis test followed by Dunn's multiple comparison test or a Wilcoxon signed rank test as appropriate. Values of $P < 0.05$ were considered significant.

Results

CXCR3 chemokine release from epithelial cells

Initial experiments were devised to compare the release of CXCL9, CXCL10 and CXCL11 from BEAS-2B cells and primary human airway epithelial cells. IFN γ (10ng/ml) in both the absence and presence of TNF α stimulated the release of CXCL9, CXCL10 and CXCL11 from both cell types (Fig 1). IFN γ and TNF α operated synergistically to stimulate release of CXCL9, CXCL10 and CXCL11 from BEAS-2B cells and (Fig 1) primary human airway epithelial cells with no differences in responses of the two cell types (Fig 1). There were no differences in the responses of human primary airway epithelial cells derived from non-smokers, smokers or patients with COPD with respect to any of the stimuli examined (Fig 1B, 1D and 1F) These data suggest that BEAS-2B cells are a good model for the release of CXCL9, CXCL10 and CXCL11 from human airway epithelia and that human primary airway epithelial cells from COPD patients do not release greater levels of CXCL9, CXCL10 and CXCL11 than cells obtained from control subjects.

Effect of JAK inhibitors on CXCR3 chemokine release

The effect of two, distinct JAK inhibitors on the release of CXCL9, CXCL10 and CXCL11 was investigated using both BEAS-2B cells and primary airway epithelial cells and compared to the effect of a glucocorticosteroid, dexamethasone. The selectivity of PF956980 has been reported previously [24] with a reported IC₅₀ of 22nM for JAK/JAK3 and 188 nM for JAK2 in a cellular assay, whereas the selectivity of PF1367550 is reported in Table 2. PF956980 has been reported to be inactive or with an IC₅₀ >30 μ M against AKT, AuroraA, cdk2, cdk6,CHK1, FGFR1, GSK3b, IKKb, IKKi, INSR, MAPK1, MAPKAP-K2, MASK,MET, PAK4, PDK1, PKCb, ROCK1, TaoK3, TrkA [24]. Both JAK inhibitors, PF956980 and PF1367550, attenuated IFN γ -stimulated and IFN γ +TNF α -stimulated CXCR3 chemokine release from BEAS-2B cells in a concentration-dependent manner (Fig 2) (Table 3), with PF1367550 being ~50-fold more potent than PF956980. This contrasts with dexamethasone, which had a very limited effect on CXCL9 and CXCL11 release (Fig 2A, 2C, 2D and 2E) but a maximal ~50% inhibitory effect on IFN γ -stimulated release of CXCL10 (Fig 2B) but none when CXCL10 was measured following stimulation with IFN γ and TNF α together (Fig 2E).

It was possible that concomitant addition of stimulus and JAK inhibitors may elicit a different inhibitory profile, therefore to address this, experiments were performed. Addition of both cytokines concomitantly with PF956980 or PF1367550 showed similar inhibitory profiles to those exhibited when inhibitors were added prophylactically (S1 Fig) with no difference in IC₅₀ values (S1 Table). Experiments were then performed to determine the effects of both JAK inhibitors in human primary airway epithelial cells in order to determine whether their ability to suppress CXCR3 chemokine release from BEAS-2B cells was reflective of their efficacy in primary cells. Again, both PF956980 and PF1367550, inhibited both IFN γ -stimulated and IFN γ + TNF α -stimulated CXCR3 chemokine release from human primary airway epithelial cells in a concentration-dependent manner irrespective of the source of the primary airway epithelial




Fig 1. Effect of IFN γ in the presence and absence of TNF α on the release of CXCL9, CXCL10 and CXCL11 from BEAS-2B and primary human airway epithelial cells. BEAS-2B cells, n = 11 (open bars, panels a, c and e) or human primary airway epithelial cells from non-smokers (n = 4) (closed bars), smokers (n = 5) (horizontal lines) or patients with COPD (n = 7) (hatched bars) (panels b, d and f) were incubated for 20h in the absence or presence of IFN γ (10ng/ml) or IFN γ + TNF α (10ng/ml). Media was harvested and the concentrations of CXCL9 (panels a and b), CXCL10 (panels c and d) and CXCL11 (panels e and f) were measured by ELISA. Data are presented as mean \pm SEM, where * represents p<0.05, **p<0.01 and ***p<0.001.

doi:10.1371/journal.pone.0128757.g001

cells (S2–S4 Figs, Table 4) with cells from non-smokers, smokers and COPD patients responding similarly. Therefore, the data from all primary cells were pooled to increase statistical power (Fig 3) (Table 5) and showed that PF1367550 was ~80-fold more potent than PF956980. In these experimental systems, dexamethasone did not suppress CXCR3 chemokine release.

In order to determine whether any of the inhibitory effects of the JAK inhibitors could be attributed to cell death, MTT assays were performed at the end of every experiment. PF956980 had no effect on cell viability at any of the concentrations used in this study. PF1367550 had no effect on cell viability within 10^3 of the IC₅₀ for this compound and therefore any observed inhibitory effects are driven the pharmacological mechanism, since at 10^{-6} M, and at all concentrations below this, both inhibitors showed no alterations in cell viability compared to untreated cells. Therefore, the effect of the JAK inhibitors to suppress CXCR3 chemokine release is not related to cell toxicity. The limited effect of dexamethasone in these systems suggested that the cells used in this study may be glucocorticosteroid-insensitive. In order to address this issue, the levels of IL-6 released from cells stimulated with IFN γ + TNF α were measured. At baseline, BEAS-2B cells and human primary airway epithelial cells release 41 ± 15 pg/ml, n = 8 and 236 ± 86 pg/ml, n = 7, IL-6 respectively. These levels are increased in the presence of IFN γ + TNF α to 1480 ± 142 pg/ml, n = 4 and 1077 ± 267 pg/ml, n = 7 for BEAS-2B cells and primary airway epithelial cells, respectively. In the presence of dexamethasone, IL-6 release was suppressed

Table 2. Selectivity of PF1367550.

Kinase	IC ₅₀
JAK1 Km	0.8 nM
JAK1 1 mM	0.4 nM
JAK2 Km	1.0 nM
JAK2 1 mM	1.3 nM
JAK3 Km	1.0 nM
JAK3 1 mM	14 nM
TYK2 Km	0.8 nM
TYK2 1 mM	8.4 nM
IL-2-induced P-STAT5 human T lymphocytes (JAK1/3)	11 nM
IL-2-induced IFN γ release by human T lymphocytes (JAK1/3)	14 nM
IL-12-induced IFN γ release by human T lymphocytes (JAK1/2)	45 nM
MLK1 Km	54 nM
MAP4K4 Km	60 nM
TrkA Km	227 nM
GSK3 β Km	617 nM
Protein kinase C β 2 Km	665 nM
MST4 Km	692 nM
TAOK2 Km	781 nM

Calculated IC₅₀ values for PF1367550 against >50 kinases. Data are presented for the kinases that showed >50% inhibition at 1mM ATP or at the Km as indicated (Data from Coe *et al.*, [25]).

doi:10.1371/journal.pone.0128757.t002




Fig 2. Effect of JAK inhibitors, PF956980 and PF1367550 and dexamethasone on IFN γ and IFN γ + TNF α -stimulated release of CXCL9, CXCL10 and CXCL11 from BEAS-2B cells. BEAS-2B cells were pre-treated and incubated for 1h with either PF956980 (\circ), PF1367550 (\bullet), or dexamethasone (\blacktriangle) followed by stimulation for 20h with IFN γ (10ng/ml) (panels a-c) or IFN γ + TNF α (10ng/ml) (panels d-f). Media was harvested and the concentrations of CXCL9 (panels a and d), CXCL10 (panels b and e) and CXCL11 (panels c and f) were measured by ELISA. Data are presented as mean \pm SEM, n = 7–8.

doi:10.1371/journal.pone.0128757.g002

Table 3. IC₅₀ values for PF956980 and PF1367550 on IFN γ and IFN γ +TNF α -stimulated release of CXCL9, CXCL10 and CXCL11 from BEAS-2B cells.

Chemokine	PF956980 IC ₅₀ (μ M)		PF1367550 IC ₅₀ (nM)	
	IFN γ	IFN γ +TNF α	IFN γ	IFN γ +TNF α
CXCL9	0.4 \pm 0.1	0.6 \pm 0.1	3.6 \pm 1.4	13.3 \pm 2.0
CXCL10	0.6 \pm 0.3	2.2 \pm 0.3	6.1 \pm 2.0	50.3 \pm 8.5
CXCL11	0.5 \pm 0.1	0.9 \pm 0.3	11.5 \pm 2.0	13.9 \pm 2.6

BEAS-2B cells were pre-treated for 1h in the presence of either PF956980 or PF1367550 prior to stimulation for 20h in the absence or presence of IFN γ (10ng/ml) or IFN γ +TNF α (10ng/ml). After this time, media was harvested and the concentrations of CXCL9, CXCL10 and CXCL11 measured by ELISA. Data are presented as mean \pm SEM, n = 7–8.

doi:10.1371/journal.pone.0128757.t003

Table 4. IC₅₀ values for PF956980 and PF1367550 on IFN γ and IFN γ +TNF α -stimulated release of CXCL9, CXCL10 and CXCL11 from human primary airway epithelial cells from non-smokers, smokers and COPD patients.

IFN γ	PF956980 EC ₅₀ (μ M)			PF1367550 EC ₅₀ (nM)		
	CXCL9	CXCL10	CXCL11	CXCL9	CXCL10	CXCL11
Non-smokers	0.88 \pm 0.39 n = 4	0.81 \pm 0.15 n = 4	0.66 \pm 0.24 n = 4	9.73 \pm 3.5 n = 3	12.8 \pm 5.6 n = 3	8.5 \pm 0.7 n = 3
Smokers	0.55 \pm 0.09 n = 4	1.28 \pm 0.36 n = 4	0.78 \pm 0.16 n = 4	9.0 \pm 3.00 n = 4	12.3 \pm 0.36 n = 4	6.8 \pm 1.9 n = 4
COPD	0.63 \pm 0.09 n = 5	0.53 \pm 0.01 n = 5	0.82 \pm 0.25 n = 5	7.4 \pm 1.3 n = 6	7.1 \pm 1.3 n = 6	6.3 \pm 1.4 n = 6
IFNγ +TNFα						
Non-smokers	0.61 \pm 0.26 n = 4	2.72 \pm 0.36 n = 4	1.85 \pm 0.67 n = 4	14.1 \pm 7.1 n = 3	20.3 \pm 5.1 n = 3	13.7 \pm 6.7 n = 3
Smokers	0.35 \pm 0.26 n = 4	1.88 \pm 0.41 n = 4	0.87 \pm 0.13 n = 4	15.3 \pm 4.8 n = 4	20.2 \pm 3.6 n = 4	11.8 \pm 2.9 n = 4
COPD	0.59 \pm 0.38 n = 5	1.84 \pm 0.40 n = 5	1.92 \pm 0.67 n = 5	20.8 \pm 4.8 n = 6	29.1 \pm 3.5 n = 6	16.7 \pm 4.1 n = 6

Human primary airway epithelial cells were pre-treated for 1h in the presence of either PF956980 or PF1367550 prior to stimulation for 20h in the absence or presence of IFN γ (10 ng/ml) or IFN γ +TNF α (10 ng/ml). After this time, media was harvested and the concentrations of CXCL9, CXCL10 and CXCL11 measured by ELISA. Data are presented as mean \pm SEM.

doi:10.1371/journal.pone.0128757.t004

by ~50–70% with IC₅₀ values of 16 \pm 8 nM and 11 \pm 7 nM respectively. Therefore, these cells are sensitive to glucocorticosteroid inhibition; however, the release of CXCL9, CXCL10 and CXCL11 following stimulation with either IFN γ or IFN γ in combination with TNF α is not. Both JAK inhibitors also suppressed this response in primary airway epithelial cells but with IC₅₀ values of ~5 and 50-fold greater than those for inhibition of CXCR3 chemokine release (PF956980: 11.4 \pm 9 μ M and PF1367550: 0.8 \pm 0.2 μ M, n = 7) suggesting that these effects may be unrelated to JAK inhibition directly.

Effects of JAK inhibitors on STAT-1

Having ascertained that JAK inhibitors could suppress the glucocorticosteroid-insensitive release of CXCL9, CXCL10 and CXCL11 from airway epithelial cells, it was important to confirm that this was via inhibition of the JAK-STAT pathway and not an off-target effect of these agents. Activation of the JAK-STAT pathway by IFN γ leads to the phosphorylation of STAT-1, therefore experiments were devised to assess whether PF956980 and PF1367550 altered STAT-1 phosphorylation. BEAS-2B cells were stimulated with either IFN γ or IFN γ in combination with TNF α and the phosphorylation of STAT-1 was measured by western blotting as reported previously [15] (Fig 4). Exposure of cells to either IFN γ or IFN γ + TNF α increase STAT-1 phosphorylation and this was inhibited by pre-treatment with either PF956980 or PF1367550 (Fig 4) (IC₅₀ values: PF956980; 0.94 \pm 0.16 μ M, and 0.84 \pm 0.21 μ M, n = 4 for IFN γ and IFN γ + TNF α stimulation respectively; PF1367550; 24 \pm 6nM, and 32 \pm 13nM, n = 4 for IFN γ and IFN γ + TNF α stimulation respectively) with, PF1367550 being ~40-fold more potent at suppressing STAT-1 phosphorylation when compared with PF956980 (Fig 4). These data in BEAS-2B cells were further validated using human primary airway epithelial cells. Both PF956980 and PF1367550 suppressed phosphorylation of STAT-1 by IFN γ and IFN γ + TNF α (Fig 5). The capacity of PF956980 and PF1367550 to suppress JAK-STAT activation was further supported by the observation that both IFN γ and IFN γ + TNF α stimulation of STAT-1 DNA binding could be suppressed by these inhibitors in BEAS-2B cells (Fig 6A). Although, IFN γ failed to promote STAT-3 DNA binding (Fig 6B), IFN γ + TNF α significantly increased STAT-3 binding to DNA in a manner that was suppressed by both PF956980 and PF1367550 (10⁻⁶M). There was no activation of STAT-5A or STAT-5B in this system (data not shown).




Fig 3. Effect of JAK inhibitors, PF956980 and PF1367550 and dexamethasone on IFN γ and IFN γ +TNF α -stimulated release of CXCL9, CXCL10 and CXCL11 from human primary airway epithelial cells. Human primary airway epithelial cells were pre-treated for 1h with either PF956980 (○), PF1367550 (●), or dexamethasone (▲) followed by stimulation for 20h with IFN γ (10 ng/ml) (panels a-c) or IFN γ + TNF α (10 ng/ml) (panels d-f). Media was harvested and the concentrations of CXCL9 (panels a and d), CXCL10 (panels b and e) and CXCL11 (panels c and f) were measured by ELISA. Data are presented as mean \pm SEM, n = 10–13.

doi:10.1371/journal.pone.0128757.g003

Effects of JAK inhibitors on CXCL9, CXCL10, CXCL11 gene transcription

In order to demonstrate that suppression of STAT-1 activation leads to reduced gene expression, experiments were performed using BEAS-2b cells that had been pretreated with JAK inhibitors for 1h prior to stimulation with either IFN γ or IFN γ with TNF α for 4h and RNA isolated (Fig 7). PF956980 suppressed expression of CXCL9, CXCL10 and CXCL11 in a concentration dependent manner irrespective of stimuli (Fig 7). In contrast, PF1367550 completely inhibited expression of all three genes at all concentrations used. Taken together these data suggest that JAK inhibition suppresses release of CXCL9, CXCL10 and CXCL11 by reducing gene transcription which contrasts with the effect of dexamethasone, which had no effect on gene transcription in this system.

Table 5. IC₅₀ values for PF956980 and PF1367550 on IFN γ and IFN γ +TNF α -stimulated release of CXCR3 chemokines CXCL9, CXCL10 and CXCL11 from human primary airway epithelial cells.

Chemokine	PF956980 IC ₅₀ (μ M)			
	IFN γ	IFN γ +TNF α		
CXCL9	0.68 \pm 0.12	0.52 \pm 0.17		
CXCL10	0.96 \pm 0.19	2.12 \pm 0.24		
CXCL11	0.76 \pm 0.12	1.58 \pm 0.33		

Chemokine	PF1367550 IC ₅₀ (nM)	
	IFN γ	IFN γ +TNF α
CXCL9	8.4 \pm 1.3	17.6 \pm 3.0
CXCL10	10.4 \pm 1.7	24.3 \pm 2.4
CXCL11	7.0 \pm 0.9	14.5 \pm 2.4

Human primary airway epithelial cells were pre-treated for 1h in the presence of either PF956980 or PF1367550 prior to stimulation for 20h in the absence or presence of IFN γ (10ng/ml) or IFN γ +TNF α (10ng/ml). After this time, media was harvested and the concentrations of CXCL9, CXCL10 and CXCL11 measured by ELISA. Data are presented as mean \pm SEM, n = 13.

doi:10.1371/journal.pone.0128757.t005




Fig 4. Effect of JAK inhibitors, PF956980 and PF1367550 on IFN γ and IFN γ + TNF α -stimulated phosphorylation of STAT-1 in BEAS-2B cells. BEAS-2B cells were pre-treated for 1h with either PF956980 or PF1367550 followed by stimulation for 10 min with IFN γ (10ng/ml) (\circ) or IFN γ + TNF α (\bullet). Cells were harvested and western blots performed for phosphorylated STAT-1 and total STAT-1 and quantified by densitometry. Data are representative of n = 4 independent experiments (panels a and c) and presented as mean \pm SEM, n = 4 (panels b and d).

doi:10.1371/journal.pone.0128757.g004




Fig 5. Effect of JAK inhibitors, PF956980 and PF1367550 on IFN γ and IFN γ + TNF α -stimulated phosphorylation of STAT-1 in human primary airway epithelial cells. Human primary airway epithelial cells were pre-treated for 1h with either 1 μ M PF956980 (PF956) or 1 μ M PF1367550 (PF136) followed by stimulation for 10 min with either IFN γ (10ng/ml) or IFN γ + TNF α . Cells were harvested and western blots performed for phosphorylated STAT-1 and total STAT-1 and quantified by densitometry. Data are representative of n = 4 independent experiments (panel a) and presented as mean \pm SEM, n = 4 (panel b).

doi:10.1371/journal.pone.0128757.g005

Discussion

Transgenic mice that over-express IFN γ exhibit many of the emphysematous changes observed in COPD [28], thus suggesting that IFN γ is an important molecule driving the development




Fig 6. Effect of JAK inhibitors, PF956980 and PF1367550 on IFN γ and IFN γ + TNF α -stimulated DNA binding of STAT-1 and STAT-3 in BEAS-2B cells. BEAS-2B cells were pre-treated with either PF956980 (PF956) or PF1367550 (PF136) for 1h followed by stimulation with IFN γ (10ng/ml) or IFN γ + TNF α . After 1 h, nuclear extracts of the cells were prepared and TransAM assays for STAT-1 (panel a) and STAT-3 (panel b) were performed. Data are presented as fold change from baseline \pm SEM; n = 4.

doi:10.1371/journal.pone.0128757.g006

and continuing pathophysiology of this disease. IFN γ is produced by CD8⁺ T lymphocytes and these cells specifically accumulate in the lungs of patients with COPD at the sites of small airways disease [3, 29] and are associated with increased expression of the chemokine receptor, CXCR3 [2]. This suggests that recruitment of these cells is via CXCR3 and that attenuation of recruitment of CD8⁺ cells into the airways of these patients could provide an anti-inflammatory treatment for this glucocorticosteroid-insensitive disease. Since the chemokines that bind to and activate CXCR3 are induced by IFN γ , the present study investigated the effect of inhibiting IFN γ signalling in airway epithelial cells using, for the first time in these cells, selective JAK inhibitors and demonstrated reductions in CXCR3 chemokine release. In addition, the synergistic action of TNF α with IFN γ to increase expression and release of CXCR3 chemokines has been reported previously [14, 15, 21] and since TNF α is also elevated in the airways of patients with COPD [22], the effects of JAK inhibition on this stimulus was also evaluated and showed similar attenuation of chemokine output. These data would suggest that JAK inhibition could lead to a reduction in the accumulation of CD8⁺ lymphocytes in COPD, however other chemokines and receptors, possibly CCR5 which is also expressed on these cells [3, 30] could provide an alternative avenue for cell migration and activation. It is of note that in diseases such as COPD, increased levels of proteases including neutrophil elastase and matrix metalloproteinases (MMP) such as MMP-2 and MMP-9 [31, 32], could contribute to chemokine processing. There is evidence that CXCL9, CXCL10 and CXCL11 can all be modified in this way leading to possible alterations in function [33].

PF956980 has been shown to be a selective inhibitor of JAK activation, most prominently against JAK2 and JAK3 with no effect on ~30 other kinase pathways including AKT, GSK3 β , IKK β , IKKi, MAPK1, MAPKAP-K2, MASK, MET, and ROCK1 but with modest effects against PKA and Lck at concentrations >5 μ M [24]. In the present study, this compound completely




Fig 7. Effect of JAK inhibitors, PF956980 and PF1367550 on IFN γ and IFN γ + TNF α -stimulated expression of CXCL9, CXCL10, and CXCL11. BEAS-2B cells were pre-treated with either PF956980 (checked bars), PF1367550 (hashed bars) or dexamethasone (10⁻⁶M white bar) for 1h followed by stimulation with IFN γ (10ng/ml) or IFN γ + TNF α for 6 h. Gene expression of CXCL9 (panels a and b), CXCL10 (panels c and d) and CXCL11 (panels e and f) were quantified by Taqman RT-PCR. Data are presented as fold change from stimulus \pm SEM; n = 3 where 1 indicates the level of mRNA induced by IFN γ or IFN γ + TNF α in the absence of inhibitors.

doi:10.1371/journal.pone.0128757.g007

attenuated IFN γ -stimulated release of CXCL9, CXCL10 and CXCL11. The IC₅₀ reported for PF956980 against JAK3 is ~4nM [24], however this compound has been reported to be effective against IL-4-induced cytotoxic resistance in chronic lymphocytic leukaemia cells at concentrations of ~0.3 μ M [34] in agreement with the potency of PF956980 in the present study. Moreover a second, structurally dissimilar molecule, PF1367550 had a similar profile of activity to PF956980, suggesting that this is not a chemotype-dependent profile, but a function of JAK enzyme inhibition. However, PF1367550 is a more potent molecule in these assays.

Of note, the human airway epithelial cell line, BEAS-2B responded to both stimuli and inhibitors in a similar manner to isolated human primary airway epithelial cells, thus confirming our previous work and that of others that BEAS-2B are a good model for primary cells in this system [14, 15]. The inhibitory effects of both JAK-STAT inhibitors were also observed when TNF α was included with IFN γ to stimulate release of CXCL9, CXCL10, and CXCL11. The synergistic effect of these cytokines to generate CXCL9, CXCL10 and CXCL11 has been reported previously [14, 21] and is thought to be mediated via the action of not only STAT-1 but also NF- κ B and CREB [21]. However, inhibition of the JAK pathway continued to inhibit expression and production of CXCL9, CXCL10 and CXCL11 in the presence of TNF α , indicating that inhibition of STAT-1 phosphorylation is an essential regulatory step in this process and indicates that JAK inhibition may be a good therapeutic approach in COPD where multiple inflammatory mediators interact.

Although IFN γ classically signals through STAT-1 homodimers [35], expression of CXCL11 has also been reported to occur via STAT-3 [36]. The present study was unable to measure a significant increase in STAT-3 DNA binding in response to IFN γ stimulation, in contrast to STAT-1. However, basal DNA binding of STAT-3 was attenuated by both JAK inhibitors. STAT-3 is activated during transformation of cells [37] and since BEAS-2B are a transformed cell line [38], STAT-3 is most likely to be activated basally and can then be inhibited by the JAK inhibitors. Nevertheless, DNA binding of STAT3 increased significantly following stimulation of BEAS-2B cells with the combination of IFN γ and TNF α and again could be suppressed by JAK inhibition. This suppression of STAT signalling by the JAK inhibitors used in this study led to inhibition of gene transcription with again PF1367550 being more effective than PF956980. Taken together, these data indicate that suppression of JAK activation by selective, small molecule inhibitors, such as PF1367550, would be of benefit in inflammatory diseases where recruitment of cells via CXCL9, CXCL10 and CXCL11 is important. Currently, pan-JAK inhibitors are undergoing clinical trials in inflammatory diseases including psoriasis and rheumatoid arthritis [39, 40] with Tofacitinib, which is structurally similar to PF956980 [24] already approved for use in rheumatoid arthritis in the USA. As such, a similar approach could be promising in diseases such as COPD where glucocorticosteroids are not effective.

Supporting Information

S1 Fig. Effect of JAK inhibitors, PF956980 and PF136550 and dexamethasone on IFN γ and IFN γ + TNF α -stimulated release of CXCL9, CXCL10 and CXCL11 from BEAS-2B cells when added concomitantly. BEAS-2B cells were incubated for with either PF956980 (\circ), PF1367550 (\bullet), or dexamethasone (\blacktriangle) and stimulated for 20h with IFN γ (10ng/ml) (panels a-c) or IFN γ + TNF α (10ng/ml) (panels d-f). Media was harvested and the concentrations of CXCL9 (panels a and d), CXCL10 (panels b and e) and CXCL11 (panels c and f) were measured by ELISA. Data are presented as mean \pm SEM, n = 7–8.
(EPS)

S2 Fig. Effect of JAK inhibitors, PF956980 and PF1367550 and dexamethasone on IFN γ and IFN γ + TNF α -stimulated release of CXCL9 from human primary airway epithelial cells derived from non-smokers, smokers and patients with COPD. Human primary airway epithelial cells from non-smokers (panels a and d), smokers (panels b and e) and patients with COPD (panels c and f) were pre-treated and incubated for 1h with either PF956980 (\circ), PF1367550 (\bullet), or dexamethasone (\blacktriangle) followed by stimulation for 20h with either IFN γ (10ng/ml) (panels a-c) or IFN γ + TNF α (10ng/ml) (panels d-f). Media was harvested and the concentrations of CXCL9 were measured by ELISA. Data are presented as mean \pm SEM, n = 5–6.
(EPS)

S3 Fig. Effect of JAK inhibitors, PF956980 and PF1367550 and dexamethasone on IFN γ and IFN γ + TNF α -stimulated release of CXCL10 from human primary airway epithelial cells derived from non-smokers, smokers and patients with COPD. Human primary airway epithelial cells from non-smokers (panels a and d), smokers (panels b and e) and patients with COPD (panels c and f) were pre-treated and incubated for 30 min with either PF956980 (\circ), PF1367550 (\bullet), or dexamethasone (\blacktriangle) followed by stimulation for 20h with either IFN γ (10ng/ml) (panels a-c) or IFN γ + TNF α (10ng/ml) (panels d-f). Media was harvested and the concentrations of CXCL10 were measured by ELISA. Data are presented as mean \pm SEM, n = 5–6. (EPS)

S4 Fig. Effect of JAK inhibitors, PF956980 and PF1367550 and dexamethasone on IFN γ and IFN γ + TNF α -stimulated release of CXCL11 from human primary airway epithelial cells derived from non-smokers, smokers and patients with COPD. Human primary airway epithelial cells from non-smokers (panels a and d), smokers (panels b and e) and patients with COPD (panels c and f) were pre-treated and incubated for 30 min with either PF956980 (\circ), PF1367550 (\bullet), or dexamethasone (\blacktriangle) followed by stimulation for 20h with either IFN γ (10ng/ml) (panels a-c) or IFN γ + TNF α (10ng/ml) (panels d-f). Media was harvested and the concentrations of CXCL10 were measured by ELISA. Data are presented as mean \pm SEM, n = 5–6. (EPS)

S1 Table. IC₅₀ values for PF956980 and PF1367550 when incubated concomitantly with IFN γ and IFN γ +TNF α for the release of CXCL9, CXCL10 and CXCL11 from BEAS-2B cells. BEAS-2B cells were incubated with either PF956980 or PF1367550 for 20h in the absence or presence of IFN γ (10 ng/ml) or IFN γ +TNF α (10 ng/ml). After this time, media was harvested and the concentrations of CXCR3 chemokines measured by ELISA. Data are presented as mean \pm SEM, n = 4 (DOCX)

Acknowledgments

This study was supported by the NIHR Respiratory Disease Biomedical Research Unit at the Royal Brompton and Harefield NHS Foundation Trust and Imperial College London.

Author Contributions

Conceived and designed the experiments: ICK PJB LED. Performed the experiments: PSF PM. Analyzed the data: PSF PM LED. Contributed reagents/materials/analysis tools: ICK. Wrote the paper: PSF PM ICK PJB LED.

References

1. Brusselle GG, Joos GF, Bracke KR. New insights into the immunology of chronic obstructive pulmonary disease. *Lancet*. 2011; 378(9795):1015–26. doi: [10.1016/S0140-6736\(11\)60988-4](https://doi.org/10.1016/S0140-6736(11)60988-4) PMID: [21907865](https://pubmed.ncbi.nlm.nih.gov/21907865/)
2. Saetta M, Mariani M, Panina-Bordignon P, Turato G, Buonsanti C, Baraldo S, et al. Increased expression of the chemokine receptor CXCR3 and its ligand CXCL10 in peripheral airways of smokers with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2002; 165(10):1404–9. Epub 2002/05/23. PMID: [12016104](https://pubmed.ncbi.nlm.nih.gov/12016104/).
3. Saetta M, Baraldo S, Corbino L, Turato G, Braccioni F, Rea F, et al. CD8+ve cells in the lungs of smokers with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 1999; 160(2):711–7. Epub 1999/08/03. PMID: [10430750](https://pubmed.ncbi.nlm.nih.gov/10430750/).
4. Chrysafakis G, Tzanakis N, Kyriakoy D, Tsoumakidou M, Tsiligianni I, Klimathianaki M, et al. Perforin expression and cytotoxic activity of sputum CD8+ lymphocytes in patients with COPD. *Chest*. 2004; 125(1):71–6. Epub 2004/01/14. PMID: [14718423](https://pubmed.ncbi.nlm.nih.gov/14718423/).

5. Siena L, Gjomarkaj M, Elliot J, Pace E, Bruno A, Baraldo S, et al. Reduced apoptosis of CD8+ T-lymphocytes in the airways of smokers with mild/moderate COPD. *Respir Med*. 2011; 105(10):1491–500. Epub 2011/05/27. doi: [10.1016/j.rmed.2011.04.014](https://doi.org/10.1016/j.rmed.2011.04.014) PMID: [21612902](https://pubmed.ncbi.nlm.nih.gov/21612902/).
6. Mannino DM, Buist AS. Global burden of COPD: risk factors, prevalence, and future trends. *Lancet*. 2007; 370(9589):765–73. Epub 2007/09/04. doi: [10.1016/S0140-6736\(07\)61380-4](https://doi.org/10.1016/S0140-6736(07)61380-4) PMID: [17765526](https://pubmed.ncbi.nlm.nih.gov/17765526/).
7. Buist AS, McBurnie MA, Vollmer WM, Gillespie S, Burney P, Mannino DM, et al. International variation in the prevalence of COPD (the BOLD Study): a population-based prevalence study. *Lancet*. 2007; 370(9589):741–50. Epub 2007/09/04. doi: [10.1016/S0140-6736\(07\)61377-4](https://doi.org/10.1016/S0140-6736(07)61377-4) PMID: [17765523](https://pubmed.ncbi.nlm.nih.gov/17765523/).
8. Barnes PJ, Adcock IM. Glucocorticoid resistance in inflammatory diseases. *Lancet*. 2009; 373(9678):1905–17. Epub 2009/06/02. doi: [10.1016/S0140-6736\(09\)60326-3](https://doi.org/10.1016/S0140-6736(09)60326-3) PMID: [19482216](https://pubmed.ncbi.nlm.nih.gov/19482216/).
9. Groom JR, Luster AD. CXCR3 ligands: redundant, collaborative and antagonistic functions. *Immunol Cell Biol*. 2011; 89(2):207–15. Epub 2011/01/12. doi: [10.1038/icc.2010.158](https://doi.org/10.1038/icc.2010.158) PMID: [21221121](https://pubmed.ncbi.nlm.nih.gov/21221121/).
10. Loetscher M, Gerber B, Loetscher P, Jones SA, Piali L, Clark-Lewis I, et al. Chemokine receptor specific for IP10 and MIG: structure, function, and expression in activated T-lymphocytes. *J Exp Med*. 1996; 184(3):963–9. Epub 1996/09/01. PMID: [9064356](https://pubmed.ncbi.nlm.nih.gov/9064356/); PubMed Central PMCID: [PMC2192763](https://pubmed.ncbi.nlm.nih.gov/pmc/PMC2192763/).
11. Costa C, Rufino R, Traves SL, Lapa ESJR, Barnes PJ, Donnelly LE. CXCR3 and CCR5 chemokines in induced sputum from patients with COPD. *Chest*. 2008; 133(1):26–33. Epub 2007/10/11. doi: [10.1378/chest.07-0393](https://doi.org/10.1378/chest.07-0393) PMID: [17925429](https://pubmed.ncbi.nlm.nih.gov/17925429/).
12. Warwick G, Thomas PS, Yates DH. Non-invasive biomarkers in exacerbations of obstructive lung disease. *Respirology*. 2013; 18(5):874–84. Epub 2013/03/26. doi: [10.1111/resp.12089](https://doi.org/10.1111/resp.12089) PMID: [23521049](https://pubmed.ncbi.nlm.nih.gov/23521049/).
13. Quint JK, Donaldson GC, Goldring JJ, Baghai-Ravary R, Hurst JR, Wedzicha JA. Serum IP-10 as a biomarker of human rhinovirus infection at exacerbation of COPD. *Chest*. 2010; 137(4):812–22. Epub 2009/10/20. doi: [10.1378/chest.09-1541](https://doi.org/10.1378/chest.09-1541) PMID: [19837822](https://pubmed.ncbi.nlm.nih.gov/19837822/); PubMed Central PMCID: [PMC2851557](https://pubmed.ncbi.nlm.nih.gov/pmc/PMC2851557/).
14. Sauty A, Dziejman M, Taha RA, Iarossi AS, Neote K, Garcia-Zepeda EA, et al. The T cell-specific CXC chemokines IP-10, MIG, and I-TAC are expressed by activated human bronchial epithelial cells. *J Immunol*. 1999; 162(6):3549–58. PMID: [10092813](https://pubmed.ncbi.nlm.nih.gov/10092813/)
15. Tudhope SJ, Catley MC, Fenwick PS, Russell RE, Rumsey WL, Newton R, et al. The role of I κ B kinase 2, but not activation of NF- κ B, in the release of CXCR3 ligands from IFN- γ -stimulated human bronchial epithelial cells. *J Immunol*. 2007; 179(9):6237–45. PMID: [17947699](https://pubmed.ncbi.nlm.nih.gov/17947699/)
16. Pechkovsky DV, Goldmann T, Ludwig C, Prasse A, Vollmer E, Muller-Quernheim J, et al. CCR2 and CXCR3 agonistic chemokines are differently expressed and regulated in human alveolar epithelial cells type II. *Respiratory Research*. 2005; 6(1):75. Epub 2005/07/22. doi: [10.1186/1465-9921-6-75](https://doi.org/10.1186/1465-9921-6-75) PMID: [16033640](https://pubmed.ncbi.nlm.nih.gov/16033640/); PubMed Central PMCID: [PMC1185567](https://pubmed.ncbi.nlm.nih.gov/pmc/PMC1185567/).
17. Hardaker EL, Bacon AM, Carlson K, Roshak AK, Foley JJ, Schmidt DB, et al. Regulation of TNF-alpha and IFN-gamma-induced CXCL10 expression: participation of the airway smooth muscle in the pulmonary inflammatory response in chronic obstructive pulmonary disease. *FASEB J*. 2004; 18(1):191–3. Epub 2003/11/05. doi: [10.1096/fj.03-0170fje](https://doi.org/10.1096/fj.03-0170fje) PMID: [14597565](https://pubmed.ncbi.nlm.nih.gov/14597565/).
18. Murray PJ. The JAK-STAT signaling pathway: input and output integration. *J Immunol*. 2007; 178(5):2623–9. Epub 2007/02/22. PMID: [17312100](https://pubmed.ncbi.nlm.nih.gov/17312100/).
19. Meinke A, Barahmand-Pour F, Wohrl S, Stoiber D, Decker T. Activation of different Stat5 isoforms contributes to cell-type-restricted signaling in response to interferons. *Mol Cell Biol*. 1996; 16(12):6937–44. Epub 1996/12/01. PMID: [8943349](https://pubmed.ncbi.nlm.nih.gov/8943349/); PubMed Central PMCID: [PMC231697](https://pubmed.ncbi.nlm.nih.gov/pmc/PMC231697/).
20. Gough DJ, Levy DE, Johnstone RW, Clarke CJ. IFN γ signaling—Does it mean JAK-STAT? *Cytokine Growth Factor Reviews*. 2008; 19(5):383–94. doi: [10.1016/j.cytogfr.2008.08.004](https://doi.org/10.1016/j.cytogfr.2008.08.004) PMID: [18929502](https://pubmed.ncbi.nlm.nih.gov/18929502/)
21. Clarke DL, Clifford RL, Jindarat S, Proud D, Pang L, Belvisi M, et al. TNF α and IFN γ Synergistically Enhance Transcriptional Activation of CXCL10 in Human Airway Smooth Muscle Cells via STAT-1, NF- κ B, and the Transcriptional Coactivator CREB-binding Protein. *J Biol Chem*. 2010; 285(38):29101–10. doi: [10.1074/jbc.M109.0999952](https://doi.org/10.1074/jbc.M109.0999952) PMID: [20833730](https://pubmed.ncbi.nlm.nih.gov/20833730/)
22. Keatings VM, Collins PD, Scott DM, Barnes PJ. Differences in interleukin-8 and tumor necrosis factor-alpha in induced sputum from patients with chronic obstructive pulmonary disease or asthma. *Am J Respir Crit Care Med*. 1996; 153(2):530–4. Epub 1996/02/01. PMID: [8564092](https://pubmed.ncbi.nlm.nih.gov/8564092/).
23. Cox MA, Jenh CH, Gonsiorek W, Fine J, Narula SK, Zavodny PJ, et al. Human interferon-inducible 10-kDa protein and human interferon-inducible T cell alpha chemoattractant are allotypic ligands for human CXCR3: differential binding to receptor states. *Mol Pharmacol*. 2001; 59(4):707–15. PMID: [11259614](https://pubmed.ncbi.nlm.nih.gov/11259614/)
24. Changelian PS, Moshinsky D, Kuhn CF, Flanagan ME, Munchhof MJ, Harris TM, et al. The specificity of JAK3 kinase inhibitors. *Blood*. 2008; 111(4):2155–7. Epub 2007/12/21. doi: [10.1182/blood-2007-09-115030](https://doi.org/10.1182/blood-2007-09-115030) PMID: [18094329](https://pubmed.ncbi.nlm.nih.gov/18094329/).

25. Coe JWD, D.M.; Jones, P.; Kortum, S.W.; Sabnis, Y.A.; Wakenhut, F.M.; Whitlock, G.A., inventor/Indazole 2013 2013.
26. ATS. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease. American Thoracic Society. *Am J Respir Crit Care Med.* 1995; 152(5 Pt 2):S77–121. PMID: [7582322](#)
27. Catley MC, Sukkar MB, Chung KF, Jaffee B, Liao SM, Coyle AJ, et al. Validation of the anti-inflammatory properties of small molecule I kappa B kinase (IKK)-2 inhibitors by comparison with adenoviral-mediated delivery of dominant-negative IKK1 and IKK2 in human airways smooth muscle. *Mol Pharmacol.* 2006; 70(2):697–705. doi: [10.1124/mol.106.023150](#) PMID: [ISI:000239117900030](#).
28. Wang Z, Zheng T, Zhu Z, Homer RJ, Riese RJ, Chapman HA Jr., et al. Interferon gamma induction of pulmonary emphysema in the adult murine lung. *J Exp Med.* 2000; 192(11):1587–600. Epub 2000/12/06. PMID: [11104801](#); PubMed Central PMCID: PMC2193095.
29. Saetta M, Di Stefano A, Maestrelli P, Ferrarresso A, Drigo R, Potena A, et al. Activated T-lymphocytes and macrophages in bronchial mucosa of subjects with chronic bronchitis. *Am Rev Respir Dis.* 1993; 147(2):301–6. Epub 1993/02/01. PMID: [8430952](#).
30. Freeman CM, Curtis JL, Chensue SW. CC Chemokine Receptor 5 and CXC Chemokine Receptor 6 Expression by Lung CD8+ Cells Correlates with Chronic Obstructive Pulmonary Disease Severity. *Am J Pathol.* 2007; 171(3):767–76. PMID: [17640964](#)
31. Culpitt SV, Rogers DF, Traves SL, Barnes PJ, Donnelly LE. Sputum matrix metalloproteinases: comparison between chronic obstructive pulmonary disease and asthma. *Respir Med.* 2005; 99(6):703–10. PMID: [15878486](#)
32. Burnett D, Chamba A, Hill SL, Stockley RA. Neutrophils from subjects with chronic obstructive lung disease show enhanced chemotaxis and extracellular proteolysis. *Lancet.* 1987; 2(8567):1043–6. PMID: [2889963](#)
33. Van den Steen PE, Husson SJ, Proost P, Van Damme J, Opdenakker G. Carboxyterminal cleavage of the chemokines MIG and IP-10 by gelatinase B and neutrophil collagenase. *Biochem Biophys Res Commun.* 2003; 310(3):889–96. Epub 2003/10/11. PMID: [14550288](#).
34. Steele AJ, Prentice AG, Cwynarski K, Hoffbrand AV, Hart SM, Lowdell MW, et al. The JAK3-selective inhibitor PF-956980 reverses the resistance to cytotoxic agents induced by interleukin-4 treatment of chronic lymphocytic leukemia cells: potential for reversal of cytoprotection by the microenvironment. *Blood.* 2010; 116(22):4569–77. Epub 2010/08/19. doi: [10.1182/blood-2009-09-245811](#) PMID: [20716767](#).
35. Hu X, Ivashkiv LB. Cross-regulation of Signaling Pathways by Interferon- γ : Implications for Immune Responses and Autoimmune Diseases. *Immunity.* 2009; 31(4):539–50. doi: [10.1016/j.immuni.2009.09.002](#) PMID: [19833085](#)
36. Yang CH, Wei L, Pfeffer SR, Du Z, Murti A, Valentine WJ, et al. Identification of CXCL11 as a STAT3-Dependent Gene Induced by IFN. *J Immunol.* 2007; 178(2):986–92. PMID: [17202361](#)
37. Aggarwal BB, Kunnumakkara AB, Harikumar KB, Gupta SR, Tharakan ST, Koca C, et al. Signal transducer and activator of transcription-3, inflammation, and cancer: how intimate is the relationship? *Ann N Y Acad Sci.* 2009; 1171(1):59–76. Epub 2009/09/03. doi: [10.1111/j.1749-6632.2009.04911.x](#) PMID: [19723038](#); PubMed Central PMCID: PMC3141289.
38. Ke Y, Reddel RR, Gerwin BI, Miyashita M, McMenamin M, Lechner JF, et al. Human bronchial epithelial cells with integrated SV40 virus T antigen genes retain the ability to undergo squamous differentiation. *Differentiation.* 1988; 38(1):60–6. Epub 1988/06/01. PMID: [2846394](#).
39. Hsu L, Armstrong AW. JAK inhibitors: treatment efficacy and safety profile in patients with psoriasis. *Journal of immunology research.* 2014; 2014:283617. Epub 2014/06/03. doi: [10.1155/2014/283617](#) PMID: [24883332](#); PubMed Central PMCID: PMC4027021.
40. Vyas D, O'Dell KM, Bandy JL, Boyce EG. Tofacitinib: The First Janus Kinase (JAK) inhibitor for the treatment of rheumatoid arthritis. *Ann Pharmacother.* 2013; 47(11):1524–31. Epub 2013/11/29. doi: [10.1177/1060028013512790](#) PMID: [24285764](#).