

Effect of Multisensory Stimulation on Analgesia in Term Neonates: A Randomized Controlled Trial

CARLO VALERIO BELLINI, FRANCO BAGNOLI, SERAFINA PERRONE, ANNA NENCI,
DUCCIO MARIA CORDELLI, MARA FUSI,
SIMONA CECCARELLI, AND GIUSEPPE BUONOCORE

Department of Pediatrics, Obstetrics, and Reproductive Medicine, University of Siena, Italy

ABSTRACT

Many attempts have been made to obtain safe and effective analgesia in newborns. Oral glucose-water has been found to have analgesic properties in neonates. We investigated whether other sensory stimulation added to oral glucose provided more effective analgesia than oral glucose alone. In a randomized prospective double-blind trial, we studied 120 term newborns during heel prick. The babies were divided randomly into six groups of 20, and each group was treated with a different procedure during heel prick: *A*) control; *B*) 1 mL 33% oral glucose given 2 min before the heel prick; *C*) sucking; *D*) 1 mL 33% oral glucose plus sucking; *E*) multisensory stimulation including 1 mL 33% oral glucose (sensorial saturation); *F*) multisensory stimulation without oral glucose. Sensorial saturation consisted in massage, voice, eye contact, and perfume smelling during heel prick. Each heel prick was filmed and

assigned a point score according to the Douleur Aiguë du Nouveau-né (DAN) neonatal acute pain scale. Camera recording began 30 s before the heel prick, so it was impossible for the scorers to distinguish procedure *A* (control) from *B* (glucose given 2 min before), *C* (sucking water) from *D* (sucking glucose), and *E* (multisensory stimulation and glucose) from *F* (multisensory stimulation and water) from the video. Procedure *E* (multisensory stimulation and glucose) was found to be the most effective procedure, and the analgesia was even more effective than that produced by procedure *D* (sucking glucose). We conclude that sensorial saturation is an effective analgesic technique that potentiates the analgesic effect of oral sugar. It can be used for minor painful procedures on newborns. (*Pediatr Res* 51: 460–463, 2002)

Newborns feel pain (1, 2). Repeated painful stimuli lower their pain threshold (3, 4) by overstimulation of NMDA receptors, which may lead to excitotoxic brain damage (5). Until a few years ago, it was claimed that the word pain was inappropriate for newborns, as pain is a subjective experience that newborns, because of their age, cannot have (6). Until the 1980s, analgesics were rarely administered to newborns even in the case of surgery (7). Now we know that anesthesia reduces brain damage due to hypoxemia, hypertension, tachycardia, variations in heart rate, and increased intracranial pressure (8, 9), all of which are particularly dangerous because of immature cerebral vasoregulation in the premature (10).

The number of painful stimuli needs to be kept to a minimum, and every effort should be made to render them less painful. Guidelines for neonatal analgesia have been suggested (11–15), especially for the most routine type of pain, blood sampling, which is usually performed by heel prick. To avoid the drawbacks of general and local analgesics (16–21), types of nonpharmacologic analgesia have been proposed, including

nonnutritional sucking and instillation of glucose or other sweet liquids on the newborn's tongue (22). The analgesic effect of glucose is thought to stimulate an increase in plasma concentrations of β -endorphin (23–27) by a preabsorptive mechanism (28).

Very recently, we observed that sensory stimuli combined with oral glucose during heel prick greatly reduced manifestations of pain in preterm babies (29). This combination was even more effective in premature babies than oral glucose. We called it sensorial saturation (SS) because it works through competition between nonpainful and painful stimuli, not to indicate that nonpainful stimuli cause saturation of sensorial pathways. In the present study, we investigate whether SS promotes analgesia in term babies and whether it is more effective than that obtained with sugar alone. We also studied the various components of SS (sucking, oral glucose, other sensory stimuli) to determine which components were the most effective.

METHODS

Subjects. This prospective randomized double-blind clinical study was designed to study the reaction of normal term neonates to pain in the Siena hospital. The study was approved

Received June 25, 2001; accepted August 10, 2001.

Correspondence: Giuseppe Buonocore, M.D., Department of Pediatrics, Obstetrics, and Reproductive Medicine, University of Siena, Policlinico "Le Scotte," V. le Bracci 36, 53100 Siena, Italy; e-mail: buonocore@unisi.it

by the local ethics committee, and the parents gave their written informed consent. Five parents refused. Inclusion criteria were the following: Apgar score at least 9 at 5 min; gestational age 38–41 wk; more than 48 h after delivery; more than 2 h after last meal; availability of the physiotherapist and a researcher to film the heel prick. The blood sampling studied was that carried out at 96 h of age for phenylketonuria and hypothyroid screening. This particular sampling was chosen because it is compulsory for all babies, has a standard procedure, is always carried out 3 h after a meal, and is long enough after delivery to permit stabilization after cesarean section. None of the babies had ever been given analgesic or sedative drugs.

Protocol. A total of 120 babies were studied. Heel prick was carried out in a quiet room barred to other persons. Parents could be present, but they had to be silent and abstain from interfering during the preparatory phase and for 30 s after the heel prick. The sampling was performed by experienced nurses; they held the baby's heel in their hands before the prick to warm it; after the preparatory period, they disinfected the heel and lanced it with a sterile blood lancet, waited 2 s, and began sampling. In all cases, the analgesic procedure was carried out by the same experienced physiotherapist (A.N.).

The babies were randomly assigned to one of six groups on the basis of a random number table. Calculation of sample size with mean and SD of 2.5 showed that to achieve 80% power and 0.02 type I error (α) in detecting a 2-point difference in Douleur Aiguë du Nouveau-né (DAN) scale between groups, at least 17 subjects were required in each group.

Each group underwent a different procedure for the heel prick:

A Control group. No analgesic procedure.

B Glucose before heel prick. The tip of a 1-mL syringe without needle was placed in the baby's mouth, and 1 mL of 33% glucose solution was instilled with gentle movements of the syringe to stimulate sucking for 30 s; heel prick was performed 120 s later.

C Water during heel prick (sucking). The tip of a 1-mL syringe without needle was placed in the baby's mouth, and 1 mL of distilled water was instilled with gentle movements of the syringe to stimulate sucking for 30 s before, during, and after heel prick.

D Glucose and sucking. As for C, but the syringe contained 33% glucose solution.

E Sensorial saturation. As for D, but with a series of sensory stimuli before, during, and after the heel prick (see below).

F Sensorial saturation without sugar. As for E, but the syringe contained distilled water.

We used 33% glucose/water solution because of its availability in our ward.

The physiotherapist did not know whether the syringe contained water or glucose solution. The baby's reaction was filmed with a SONY video camera from 30 s before to 30 s after the heel prick. The films were seen by observers who did not know the method used each time. It was impossible for them to distinguish the following pairs of groups in the films: A (control) from B (glucose given 2 min before), C (sucking water) from D (sucking glucose), and E (multisensory stimu-

lation and glucose) from F (multisensory stimulation and water).

The observers attributed a state of alertness according to the observational rating system of Prechtl and Beintema (30), which distinguishes five types of behavioral state: 1) eyes closed, regular breathing, no movements; 2) eyes closed, irregular breathing, no movements; 3) eyes open, no gross movements; 4) eyes open, continual gross movements, no crying; 5) eyes open or closed, fussing, or crying. They then assigned a score on the DAN scale (Table 1) (31, 32). This scale scores pain from 0 to 10, where 0 is no pain and 10 is maximal pain. It evaluates three items: facial expression, limb movements, and vocal expression.

SS. This procedure consisted of the following: 1) laying the infant on its side with legs and arms flexed but free to move; 2) looking the infant in the face, close up, to attract its attention and, simultaneously, 3) massaging the infant's face and back; 4) speaking to the infant gently, but firmly; 5) letting the infant smell the fragrance of baby perfume (Colonia Analcolica-Humana) on the physiotherapist's hands; 6) instilling 33% glucose on the infant's tongue with the timing and method described in C.

Statistical analysis was performed with GBStat v 6.5 PC software, using multiple *t* test comparisons with Bonferroni adjustment to α (cry data) and the Kruskal-Wallis ANOVA test for comparison of median DAN scores between groups.

RESULTS

From February to June 2001, we studied 120 newborns, all of white Caucasian race. (Table 2 shows the characteristics of the six groups of infants enrolled in the study.

Figure 1 shows mean DAN scores obtained with the five analgesic procedures and controls. We observed no statistically

Table 1. DAN*: behavioral acute pain-rating scale for neonates

Measure	Score
Facial expressions	
Calm	0
Snivels and alternates gentle eye opening and closing	1
Determine intensity of one or more of eye squeeze, brow bulge, nasolabial furrow	
Mild, intermittent with return to calm	2
Moderate	3
Very pronounced, continuous	4
Limb movements	
Calm or gentle movements	0
Determine intensity of one or more of the following signs: pedals, toes spread, legs tensed and pulled up, agitation of arms, withdrawal reaction	
Mild, intermittent with return to calm	1
Moderate	2
Very pronounced, continuous	3
Vocal expression	
No complaints	0
Moans briefly; for intubated child, looks anxious or uneasy	1
Intermittent crying; for intubated child, gesticulations of intermittent crying	2
Long-lasting crying, continuous howl; for intubated child, gesticulations of continuous crying	3

* See Ref. 31.

Table 2.

	A Control	B Glucose before heel prick	C Sucking water during heel prick	D Sucking glucose during heel prick	E SS	F SS
Mean birth weight (range) grams	3320 (3900–2950)	3160 (3790–2820)	3150 (3570–2840)	3570 (3900–3120)	3610 (3840–3470)	3430 (3900–2690)
Boys/girls	11/9	12/8	9/11	10/10	9/11	11/9
Median arousal state score (1–5)	3	3	3	3	3	4
Range	1–4	1–4	1–4	1–4	1–4	2–5




Figure 1. Comparison of median DAN scores of the five analgesic treatments and controls. (A) controls, (B) oral 33% glucose, (C) sucking, (D) glucose and sucking, (E) SS with glucose, (F) sensorial stimulation without oral glucose.

significant reduction of score in procedures B (glucose) and F (SS without sugar) with respect to A (controls). Procedure C (sucking) disclosed a reduction of DAN score ($p = 0.001$), but the maximum of analgesia with respect to controls was obtained with procedures D (glucose plus sucking) ($p < 0.0001$) and E (SS) ($p < 0.0001$). Procedure E was even more effective than procedure D ($p = 0.004$).

Figure 2 shows the mean duration of crying in the first 30 s after lancing. We observed a decrease from a mean value of 25 s with procedure A (control) to 7 s with procedure D and 2.8 s with procedure E (SS).

DISCUSSION

The methods used in previous studies of nonpharmacologic analgesia in neonates have led to a considerable reduction but not elimination of signs of pain perception (21, 33–39). The best results have been obtained with simultaneous administration of oral sugar and pacifier (40), which also has been used




Figure 2. Cry (seconds) during the 30 s after the heel prick. (A) controls, (B) oral 33% glucose, (C) sucking, (D) glucose and sucking, (E) SS, (F) multi-sensorial stimulation without oral glucose.

during venous blood sampling (31). Nevertheless, most of these studies were performed with a waiting time of 2 min between instillation of sweet solution and heel prick, sometimes too long a time for a busy ward. Moreover, analgesia limited to administration of analgesic substances leaves out human presence, which is most important in moments of pain (41, 42). SS includes human presence and requires only a few seconds.

Our results confirm the analgesic effect of orally administered glucose: analgesia was statistically greater in the group of babies who sucked 33% glucose (group D) than that of babies who sucked water (group C).

When glucose was given 2 min before heel prick (B), pain score was not significantly different from control (A). Signs of pain, nevertheless, decreased when sugar was offered immediately before and during heel prick to stimulate sucking (D). This is in line with Johnston *et al.* (43) who advised giving sugar 2 min before and during heel prick in preterm babies for maximal analgesic effect and with those authors proposing pacifiers plus glucose during heel prick (31). Sugar, therefore, does not seem to create analgesia by a pharmacologic effect alone but seems rather to have an effect on the central control of painful stimuli (44) according to the gate control theory (45): the brain is not a passive receiver of nociceptive input but can influence the information received, deciding whether it is important enough to record. Stimulation of sensory channels prevents nociceptive nerve impulses from getting through (46, 47).

So we increased with SS the number of channels engaged during heel prick by adding auditory, tactile, visual, olfactory, and vestibular stimuli to the gustatory one. Alone (F), these stimuli did not produce an analgesic effect, or rather they increased awareness and irritated the baby (Table 2). When combined with glucose (E), however, they increased the analgesic effect. To function, they evidently require a favorable background situation such as that of a baby already intent on sucking a sweet liquid.

The preeminence of the gustatory stimulus may be explained by the fact that the chemoreceptor system is the first to become active in all species of animals and is already functional in the fetus (48–50).

Different sensory stimuli have been used to activate gate-control mechanisms to block nociceptive transmission: massage (51) or water mattresses (52), tactile (53) and acoustic (54) stimuli or both (55), sucking (39), oral sucrose or glucose (28, 35, 36, 38, 40, 56–58), and other sweet liquids (22, 34). Nevertheless, the technique used in the present study is the first to obtain an evident and significant decrease in the response to

pain in term neonates by stimulating various sensory channels simultaneously and in a codified manner. Simultaneous stimulation of various channels had already been used or in a noncodified manner, namely, general consolation by parents (33) or unsuccessfully combining sucrose with rocking (59).

With SS, we found an almost complete absence of pain reaction during heel prick; the babies rarely cried, and if they did, only for a few seconds (Fig. 2). Other researchers (33–37) succeeded in reducing crying of premature babies with oral sucrose, but the infants nevertheless cried for many seconds and in some cases for more than 1 min, much more than with our technique, which reduced the mean crying time in the 30 s after heel prick to 2.8 s.

Haouari *et al.* (37) were on the right track when they wrote, “We don’t know whether simply cuddling an infant after heel prick is as effective in reducing crying as 50% sucrose.” In actual fact, concentrated sugar solution is much more effective than cuddling, but the latter or at least its sensory component, *i.e.* massage, voice, rocking, eye fixing, adds further analgesia to the analgesic effect of glucose and provides a human factor in the form of company to a baby facing pain. SS is a technique that can be used for all newborns undergoing blood samples or other minor painful procedures.

REFERENCES

- Anand KJ, Hickey PR 1987 Pain and its effects in the human neonate and fetus. *N Engl J Med* 317:1321–1326
- Fitzgerald M, Millard C, MacIntosh N 1988 Hyperalgesia in premature infants. *Lancet* 1:292
- McIntosh N 1997 Pain in the newborn, a possible new starting point. *Eur J Pediatr* 156:173–177
- Menon G, Anand KJ, McIntosh N 1998 Practical approach to analgesia and sedation in the neonatal intensive care unit. *Semin Perinatol* 22:417–424
- Anand KJ, Scalzo FM 2000 Can adverse neonatal experiences alter brain development and subsequent behavior? *Biol Neonate* 77:69–82
- Richard T 1985 Can a fetus feel pain? *BMJ* 291:1220–1221
- Beyer JE, DeGood DE, Ashley LC, Russell GA 1983 Patterns of postoperative analgesic use with adults and children following cardiac surgery. *Pain* 17:71–81
- Anand KJ 1998 Clinical importance of pain and stress in preterm neonates. *Biol Neonate* 73:1–9
- Stevens BJ, Johnston CC 1994 Physiological responses of premature infants to a painful stimulus. *Nurs Res* 43:226–231
- Tsuji M, Saul P, du Plessis A, Eichenwald E, Sobh J, Crocker R, Volpe JJ 2000 Cerebral intravascular oxygenation correlates with mean arterial pressure in critically ill premature infants. *Pediatrics* 106:625–632
- Canadian Pediatric Society. Fetus and Newborn Committee 2000 Prevention and management of pain and stress in the neonate. *Pediatrics* 104:454–458
- Spaeth JP, O’Hara IB, Kurth CD 1998 Anesthesia for the micropreemie. *Semin Perinatol* 22:390–401
- Stevens B, Gibbins S, Franck LS 2000 Treatment of pain in the neonatal intensive care unit. *Pediatr Clin North Am* 47:633–640
- Carbajal R, Simon N 1995 Sédation et analgésie chez l’enfant. *Arch Pediatr* 2:1089–1096
- Anand KJ 2001 Consensus statement for the prevention and management of pain in the newborn. *Arch Pediatr Adolesc Med* 155:173–180
- Suresh S, Anand KJ 1998 Opioid tolerance in neonates: mechanisms, diagnosis, assessment, and management. *Semin Perinatol* 22:425–433
- Jacqz-Aigrain E, Burtin P 1996 Clinical pharmacokinetics of sedatives in neonates. *Clin Pharmacokinet* 31:423–443
- Levene M 1995 Pain relief and sedation during neonatal intensive care. *Eur J Pediatr* 154:S22–S23
- Law RMT, Halpern S, Martins RF, Reich H, Innanen V, Ohlsson A 1996 Measurement of methemoglobin after EMLA analgesia for newborn circumcision. *Biol Neonate* 70:213–217
- Gourrier E, Karoubi P, El Hanache A, Merbouche S, Mouchino G, Dhabhi S, Leraillez J 1995 Utilisation de la crème EMLA chez le nouveau-né à terme et prématuré. Etude d’efficacité et de tolérance. *Arch Pediatr* 2:1041–1046
- Lemmen RJ, Semmekrot BA 1996 Muscle rigidity causing life-threatening hypercapnia following fentanyl administration in a premature infant. *Eur J Pediatr* 155:1067
- Blass EM 1997 Milk-induced hypoalgesia in human newborns. *Pediatrics* 99:825–829
- Balon-Perin S, Kolanowski J, Berbinschi A, Franchimont P, Ketelslegers JM 1991 The effects of glucose ingestion and fasting on plasma immunoreactive beta-endorphin, adrenocorticotrophic hormone, and cortisol in obese subjects. *J Endocrinol Invest* 14:919–925
- Tropeano G, Lucisano A, Liberale I, Barini A, Vuolo IP, Martino G, Menini E, Dell’Acqua S 1994 Insulin, C-peptide, androgens, and beta-endorphin response to oral glucose in patients with polycystic ovary syndrome. *J Clin Endocrinol Metab* 78:305–309
- Blass EM, Fitzgerald E 1988 Milk-induced analgesia and comforting in 10-day-old rats: opioid mediation. *Pharmacol Biochem Behav* 29:9–13
- Shide DJ, Blass EM 1989 Opioidlike effects in intraoral infusions of corn oil and polycose on stress reactions in 10-day-old rats. *Behav Neurosci* 103:1168–1175
- Blass E, Fitzgerald E, Kehoe P 1987 Interactions between sucrose pain and isolation distress. *Pharmacol Biochem Behav* 26:483–489
- Ramenghi LA, Evans DJ, Levene MI 1999 “Sucrose analgesia”: absorptive mechanism or taste perception? *Arch Dis Child Fetal Neonatal Ed* 80:F146–F147
- Belliemi CV, Buonocore G, Nenci A, Franci N, Cordelli DM, Bagnoli F 2001 Sensorial saturation: an effective analgesic tool for heel-prick in preterm infants. A prospective randomized trial. *Biol Neonate* 80:15–18
- Prechtl HF, Beintema DJ 1964 The neurological examination of the full term newborn infant. In: *Clinics in Developmental Medicine* N. 12 Heinemann Ed., London
- Carbajal R, Chauvet X, Courdec S, Olivier-Martin M 1999 Randomized trial of analgesic effects of sucrose, glucose, and pacifiers in term neonates. *BMJ* 319:1397–1402
- Carbajal R, Paupe A, Hoenn E, Lenclen R, Olivier-Martin M 1997 DAN: une échelle comportementale d’évaluation de la douleur aigue du nouveau-né. *Arch Pediatr* 4:623–628
- Overgaard C, Knudsen A 1999 Pain relieving effect of sucrose in newborns during heel prick. *Biol Neonate* 75:279–284
- Bucher HU, Baumgartner R, Bucher N, Seiler M, Fauchère JC 2000 Artificial sweetener reduces nociceptive reaction in term newborn infants. *Early Hum Dev* 59:51–60
- Ramenghi LA, Griffith GC, Wood CM, Levene MI 1996 Effect of non-sucrose sweet tasting solution on neonatal heel prick responses. *Arch Dis Child* 74:F129–F131
- Skogsdal Y, Eriksson M, Schollin J 1997 Analgesia in newborns given oral glucose. *Acta Paediatr* 86:217–220
- Haouari N, Wood C, Griffiths G, Levene M 1995 The analgesic effect of sucrose in full term infants: a randomised controlled trial. *BMJ* 310:1498–1500
- Rushforth JA, Levene MI 1993 Effect of sucrose on crying in response to heel stab. *Arch Dis Child* 69:388–389
- Blass EM, Hoffmeyer LB 1991 Sucrose as an analgesic for newborn infants. *Pediatrics* 87:215–218
- Blass EM, Watt LB 1999 Suckling and sucrose-induced analgesia in human newborns. *Pain* 83:611–623
- Als H, Duffy FH, McAnulty GB 1996 Effectiveness of individualised neurodevelopmental care in the newborn intensive care unit. *Acta Paediatr* 416:S21–S30
- Als H, Gilkerson L 1997 The role of relationship-based developmentally supportive newborn intensive care in strengthening outcome of preterm infants. *Semin Perinatol* 21:178–189
- Johnston CC, Stremler R, Horton L, Friedman A 1999 Effect of repeated doses of sucrose during heel stick procedure in preterm neonates. *Biol Neonate* 75:160–166
- Lindahl S 1997 Calming minds or killing pain in newborn infants? *Acta Paediatr* 86:787–788
- Melzack R, Wall PD 1965 Pain mechanisms: a new theory. *Science* 150:971–979
- Wall PD 1978 The gate control theory of pain mechanism. A re-examination and re-statement. *Brain* 101:1–18
- Melzack R 1999 From the gate to the neuromatrix. *Pain* 6:S121–S126
- Relier JP 1996 Importance of fetal perceptions in the organization of the mother-fetus interactions. *Biol Neonate* 69:165–212
- Robin M 2000 Premier regards, premiers échanges. In: Herbinet E, Busnel MC (eds) *L’Aube des Sens*. Stock, Paris, pp 55–64
- Herbinet E, Busnel MC 2000 *L’Aube des Sens*. Stock, Paris
- Franck LS, Lawhon G 1998 Environmental and behavioral strategies to prevent and manage neonatal pain. *Semin Perinatol* 22:434–443
- Korner AF, Kraemer HC, Haffner ME, Cospser LM 1975 Effects of waterbed flotation on premature infants: a pilot study. *Pediatrics* 56:361–367
- Gray L, Watt L, Blass EM 2000 Skin-to-skin contact is analgesic in healthy newborns. *Pediatrics* 105:e14
- Locsin RG 1981 The effect of music on the pain of selected post-operative patients. *J Adv Nurs* 6:19–25
- McIntosh N, van Veen L, Brameyer H 1994 Alleviation of the pain of heel prick in preterm infants. *Arch Dis Child* 70:F177–F181
- Stevens B, Ohlsson A 2001 Sucrose for analgesia in newborn infants undergoing painful procedures (Cochrane Review). In: *The Cochrane Library*, Issue 4, Oxford
- Abad F, Diaz NM, Domenech E, Robayna M, Rico J 1996 Oral sweet solution reduces pain-related behaviour in preterm infants. *Acta Paediatr* 85:854–858
- Bucher HU, Moser T, Von Siebental K, Keel M, Wolf M, Duc G 1995 Sucrose reduces pain reaction to heel lancing in preterm infants: a placebo-controlled, randomized and masked study. *Pediatr Res* 38:332–335
- Johnston CC, Stremler RL, Stevens BJ, Horton LJ 1997 Effectiveness of oral sucrose and simulated rocking on pain response in preterm neonates. *Pain* 72:193–199