



Effect of polluted water on soil and plant contamination by heavy metals in El-Mahla El-Kobra, Egypt

Esawy Kasem Mahmoud¹ and Adel Mohamed Ghoneim^{2,3}

¹Soil and Water Sciences Department, Faculty of Agriculture at Tanta, Tanta University, Tanta, Egypt

²Soil Science Department, College of Food and Agricultural Sciences, King Saud University, P.O. Box 2460, Riyadh 11451, Saudi Arabia

³Agricultural Research Center, Field Crops Research Institute, Rice Research and Training Center, Sakha, Kafr El-Sheikh, 33717, Egypt

Correspondence to: Adel Mohamed Ghoneim (aghoneim@ksu.edu.sa)

Received: 24 March 2015 – Published in Solid Earth Discuss.: 8 February 2016

Revised: 19 April 2016 – Accepted: 19 April 2016 – Published: 29 April 2016

Abstract. The discharge of untreated waste water in Zefta drain and drain no. 5 is becoming a problem for many farmers in the El-Mahla El-Kobra area, Egypt. The discharged water contains high levels of contaminants considered hazardous to the ecosystem. Some plants, soil, water, and sediment samples were collected from the El-Mahla El-Kobra area to evaluate the contamination by heavy metals. The results showed that the heavy metals, pH, sodium adsorption ratio (SAR), biochemical oxygen demand (BOD), and chemical oxygen demand (COD) in the water of Zefta drain and drain no. 5 exceeded permissible limits for irrigation. In rice and maize shoots grown in soils irrigated by contaminated water from Zefta drain and drain no. 5, the bioaccumulation factors for Cd, Pb, Zn, Cu, and Mn were higher than 1.0. The heavy metals content of irrigated soils from Zefta drain and drain no. 5 exceeded the upper limit of background heavy metals. In this study, the mean contaminant factor values of the drain no. 5 sediments revealed that Zn, Mn, Cu, Cd, Pb, and Ni > 6, indicating very high contamination. The bioaccumulation coefficient values of *Cynodon dactylon*, *Phragmites australis*, and *Typha domingensis* aquatic plants growing in Zefta drain are high. These species can be considered as hyperaccumulators for the decontamination of contaminated water.

1 Introduction

Contamination of soils by heavy metals, such as Cd, Ni, Zn, Pb, and Cu, has increased dramatically during the last few decades (Chibuike and Obiora, 2014) due to mining, smelting, manufacturing, use of agricultural fertilizers and pesticides, municipal waste, traffic emissions, and industrial effluents (Morgan, 2013; Chibuike and Obiora, 2014). Contamination of soils by heavy metals is now widespread (Al-Nagger et al., 2013). Land degradation caused by heavy metals has significant adverse effects on the environment and ecosystem worldwide (Li et al., 2013; Chen et al., 2015). Dispersion of heavy metals in irrigated soils and the plants that are growing results in the contamination of food that may be hazardous to humans and animals (Jolly et al., 2013). Heavy metals in effluents are poorly soluble in water, and cannot be degraded; they tend to accumulate in soils and subsequently accumulate in plants (Ghoneim et al., 2014). In addition, heavy metals persist in soil which then leach down into the groundwater and may induce enhanced antioxidant enzymatic activities in plants or become adsorbed with solid soil particles (Iannelli et al., 2002). According to Roy and McDonald (2013), carrots grown in soils contaminated by Cd have the potential to cause toxicological problems in men, women, and young children. Cd uptake by carrot roots was about 5 times more than the regulatory limits for men, 8 times more for women, and 12 times more for children. High levels of Cd in soil were identified as causing itai-itai disease in Toyama Prefecture, Japan; however, soil solution levels



Figure 1. Location of study area.

similarly high in Cd do not seem to cause health problems for people living in Shipham, England (Morgan, 2013). For the Cu-contaminated soils planted with tomato (*Solanum lycopersicum* L.), these values would range between 32.9 and 1696.5 mg kg⁻¹, depending on soil properties (Sacristán et al., 2015). Accumulation of toxic heavy metals in living plant cells results in various deficiencies, reduction of cell activities, and inhibition of plant growth (Farooqi et al., 2009).

Transfer of heavy metals from soils to plants is one of the key pathways for exposure of humans via the food chain. In order to assess risks to health associated with metals in soils, it is necessary to predict transfer of heavy metals from soil to tissues of plants for subsequent use in phytoremediation (Roy and McDonald, 2013; Ye et al., 2014). Heavy metal pollution is persistent, covert, and irreversible (Wang et al., 2011). This kind of pollution not only degrades the quality of the food crops, atmosphere, and water, but also threatens the health of human and animals (Dong et al., 2011). Excessive intake of the Pb to human body can damage the nervous, skeletal, endocrine, enzymatic, circulatory, and immune system (Zhang et al., 2012). The chronic effects of Cd consist of lung cancer, pulmonary adenocarcinomas, prostatic prolifer-

ative lesions, kidney dysfunction, bone fractures, and hypertension (Brevik et al., 2015).

Soil is a key part of the Earth system as it control the hydrological, erosional, biological, and geochemical cycles. The soil system also offers goods, services, and resources to humankind (Berendse et al., 2015; Brevik et al., 2015; Decock et al., 2015; Smith et al., 2015). This is why it is necessary to research how soils are affected by societies. Pollution is one of these damaging human activities, and we need more information and assessment of soil pollution (Mahmoud and El-Kader, 2015; Riding et al., 2015; Roy and McDonald, 2015; Wang et al., 2015). Soil degradation is now considered a challenge of a global dimension and is included in environmental policy frameworks. A prime example is the United Nations Convention to Combat Desertification (UNCCD), which recognizes the important role of soils in sustainable development and has anticipated the ambitious aim to achieve zero net land degradation by 2030 (UNCCD, 2012). Soils have been used to detect the deposition, accumulation, and distribution of heavy metals in different locations (Alirzayeva et al., 2006; Onder et al., 2007), but little quantitative information is available on the contamination of agri-

cultural soils in El-Mahla El-Kobra, Egypt, by heavy metals. To close this knowledge gap, this study investigated the contamination of agricultural soil and plants by heavy metals in residential and industrial areas in El-Mahla El-Kobra, Egypt.

2 Materials and methods

2.1 Site description, samples, and analysis

El-Mahla El-Kobra (Fig. 1) is located at 30°34' N, 30°45' E. The dominant sources of heavy metal pollution are waste water irrigation, manure, and sediment applications for metallic ores. The El-Mahla El-Kobra area is densely populated (4.5 million 462 683 km⁻²) and contains 183 industrial factories such as for textiles, food, oil, and other industries. The quantity of industrial and municipal waste water is around 243 500 m³ day⁻¹ (107 500 m³ day⁻¹ municipal sewage and 136 000 m³ day⁻¹ industrial waste water), which discharges into Zefra drain (flow, 354 240 m³ day⁻¹) and drain no. 5 (flow, 265 248 m³ day⁻¹) without treatment, except 63 627 m³ day⁻¹ municipal waste water which can be treated at Dawakhlia plant.

Seventy representative soil samples (0–30 cm) in summer 2012 were collected from cultivated lands of El-Mahla El-Kobra, Egypt, which are irrigated with drainage water from Zefra drain and drain no. 5, and 15 samples of soil were collected which is irrigated with water from Baher El Mlah canal (fresh water). The soil samples were air-dried and ground to pass through a 2 mm screen for chemical analysis. The soils' pH was determined in a saturated soil paste extract (Richards, 1954). Calcium and magnesium levels were determined titrimetrically using versenate (Jackson, 1973). The level of sodium was determined using a flame photometer (Richards, 1954). The level of total carbonate was determined using the calcimeter as a CaCO₃ percentage according to Loeppert and Suarez (1996). The total heavy metals (Cd, Pb, Zn, Fe, Mn, Cu, and Ni) were measured by the atomic absorption spectrophotometer after the soil samples had digested concentrated mixtures of HNO₃ and HClO₄ acids (Page, 1982). Samples of rice and maize cultivated crops (age 65 days in summer 2012) that are grown in the studied soils, and three other aquatic plant species (*Cynodon dactylon*, *Phragmites australis*, and *Typha domingensis*) which are grown in Zefra drain, were also collected at different times. The plant samples were dried in an oven at 75 °C for 72 h. The total heavy metals content in plant shoots was measured using the atomic absorption spectrophotometer after the plant samples had digested concentrated H₂SO₄ and H₂O₂ (Chapman and Pratt, 1961).

2.2 Transfer of heavy metals

The bioconcentration factor (BF) of each metal in plants was calculated by dividing the total content in the plants by the total content in soil (Brooks, 1998). In addition, 17 water sam-




Figure 2. Concentration of heavy metals in maize and rice shoots grown in soils irrigated by water from Zefra drain and drain no. 5.

ples were collected from Zefra drain and drain no. 5 at different times (March 2012 to March 2013) at about 20 cm below the water surface and were chemically analysed for pH, electrical conductivity (EC), sodium adsorption ratio (SAR), biological oxygen demand (BOD), chemical oxygen demand (COD), and heavy metals content (APHA, 2005). The bioaccumulation coefficients of each heavy metal in aquatic plants were calculated by dividing the total heavy metal content in aquatic plants by the concentration in water.

The contaminant factor (Cf) for soil is the ratio obtained by dividing the concentration of each heavy metal in the sediment by the background values (Håkanson, 1980).

$$Cf = C_{\text{Heavy metal}} / C_{\text{Background}} \quad (1)$$

According to Håkanson (1980), the values of Cf < 1 indicate low contamination; 1 < Cf < 3 indicates moderate contamination; 3 < Cf < 6 indicates considerable contamination; and Cf > 6 indicates very high contamination.

3 Results and discussion

3.1 Effect of contaminated water on plant and soil contamination

Heavy metals content was higher in rice and maize shoots grown in the soil around the Zefra drain than the same crops in soil irrigated by water from drain no. 5 (Fig. 2). This was due to the high total heavy metal contents in these soils (Table 1). Maize shoots contained more Fe, Cd, Mn, and Pb than rice shoots, and this may be attributed to the planting of rice under flooded conditions. Under flooded conditions, Fe, Cd, Mn, and Pb could be precipitate as FeS₂, CdS, MnS, and PbS, respectively, due to the reducing conditions. Heavy metals content in rice and maize shoots exceeded the defined limits reported by Kabata-Pendias and Pendias (1992) and were above the levels acceptable for elemental composition of uncontaminated plant tissue. Alloway (1990) reported that in angiosperms, uncontaminated plant tissue contains 0.64,

Table 1. Total concentrations of heavy metals in soils irrigated by contaminated water from Zefta drain, drain no. 5, and Baher El Mlah canal.

Parameters	Unit	Surrounding soils			Upper limit of total heavy metals in soils (Chen et al., 1992)
		Drain no. 5	Zefta drain	Baher El Mlah	
pH	–	7.80–8.30	7.80–8.50	7.30	–
CaCO ₃	%	4.10–8.20	3.28–5.74	4.10	–
Fe	mg kg ⁻¹	1226–4989	1790–4757	933	–
Zn	mg kg ⁻¹	102–187	184–449	54	120
Mn	mg kg ⁻¹	341–800	172–853	264	–
Cu	mg kg ⁻¹	82–167	123–386	60	35
Cd	mg kg ⁻¹	13–28	21–33	11	3
Pb	mg kg ⁻¹	48–92	55–80	53	120
Ni	mg kg ⁻¹	55–133	104–164	31	60

2.4, 160, and 14 mg kg⁻¹ of Cd, Pb, Zn, and Cu, respectively. It is clear from Fig. 2 and Table 1 that the concentrations of Cd in rice and maize shoots are higher than other heavy metals compared with the maximum limits according to Kabata-Pendias and Pendias (1992). Li et al. (1994) found that plants absorb Cd more readily than other heavy metals and levels are often reached that are hazardous to human health before any stress symptoms appear. Chitdeshwari et al. (2002) reported that the use of sewage water increased the uptake of Cd and Cr in *Amaranthus* crops. Phosphate fertilizers were sources of Cd used in fertilization of rice and maize plants in this study area. Phosphate fertilizers were even measured to contain 200 mg Cd kg⁻¹ (Nziguheba and Smolders, 2008). The city of El-Mahla El-Kobra is densely populated and is the capital of the local textile industry. Large amounts of industrial and contaminated water are discharged directly into irrigation canals without treatment, which often contain heavy metals that contribute to metals' enrichment in soil (Fakayode and Onianwa, 2002). In addition, rice and wheat ash fertilization is carried out in El-Mahla El-Kobra on a large scale (Abou-Sekkina et al., 2010). Application of ash to agricultural soils contributes significantly to the greater concentration of Cd in agricultural soil from El-Mahla El-Kobra. The higher concentration of Zn observed might be due to abrasion of tyres, barks, and Zn-containing compounds, which are used in some manufactured goods, such as paints, cosmetics, automobile tyres, and batteries (Imperato et al., 2003).

The range of pH, EC, and heavy metal contents in soil samples which are irrigated by water from Zefta drain, drain no. 5, and Baher El Mlah are compared with background limits (Table 3). The soils irrigated by contaminated water from Zefta drain and drain no. 5 show an increase of soil pH with comparison to soils irrigated by water from Baher El Mlah (fresh water). Similar results were noticed by Gupta et al. (2010) and Saffari and Saffari (2013), who reported that after irrigation with sewage water, pH increased signifi-

Table 2. Bioconcentration factors of heavy metals in maize and rice shoots grown in soils irrigated by waste water from Zefta drain, drain no. 5, and limits of heavy metals.

Elements	Drain no. 5		Zefta drain		Limits of heavy metals ¹ mg kg ⁻¹
	Rice	Maize	Rice	Maize	
Fe	0.29	0.35	0.54	0.34	–
Mn	3.40	2.97	1.71	1.51	300–500
Cu	0.59	1.54	1.75	1.83	20–100
Zn	2.82	1.71	1.44	1.32	100–400
Pb	6.73	6.83	5.26	5.66	30–300
Cd	6.14	6.45	6.55	2.26	5–30

¹ Kabata-Pendias and Pendias (1992).

cantly. The reason for an increasing soil pH may be attributed to high pH values in Zefta drain and drain no. 5 (Table 3). The soils irrigated by contaminated water from Zefta drain and drain no. 5 affect the total dissolved solids (TDSs) (Table 3). Indeed, in comparison to soils which are irrigated by water from Baher El Mlah, the TDS value is greater in Zefta drain and drain no. 5. These results were in agreement with Mollahoseini (2013) and Khaskhoussy et al. (2013), who reported that irrigating with sewage water increased soil salinity, exchangeable Na, K, Ca, Mg, and available P. In general, the concentrations of heavy metals in soils irrigated by water from Zefta drain and drain no. 5 exceeded the upper background limit of total heavy metals (Chen et al., 1992). Contents of Mn, Cd, and Ni in soils at Zefta drain were higher than in soils at drain no. 5, due to the high concentration of heavy metals in Zefta drain water (Table 3). The level of heavy metals of soils irrigated by water from Zefta drain and drain no. 5 was higher than the levels of the surrounding soils of Baher El Mlah canal. Similar results were reported by Chen et al. (1992), who found high levels of heavy metals in soils which are irrigated by polluted industrial waste water. These results coincided with El-Gendi et al. (1997) who indi-

cated that irrigating sandy soil in the Abou-Rawash area with drainage water increased total Cu, Zn, and Fe, which reached 125, 170, and 5 times that of the virgin soil in the same area. Both Cd and Pb levels in soils measured during this study were higher than those reported by Nassef et al. (2006) and Suci et al. (2008). These differences might be related to different anthropogenic activities and concentrations of urbanization at each site.

3.2 Bioconcentration factors (BFs)

The BF values in the rice and maize shoots are presented in Table 2. In rice and maize grown in soils irrigated by water from Zefta drain and drain no. 5, the BF values for Cd, Pb, Zn, Cu, and Mn were higher than 1. This indicates that bioconcentrations of Cd, Pb, Cu, and Mn were high in the plants studied. Fe was an exception because its BF value was lower than 1, indicating low accumulation in studied plants. The BF values for Zn and Cu of rice shoots were higher than maize shoots grown in the same soils irrigated by water from Zefta drain and drain no. 5. These concentrations were attributed to using $ZnSO_4$ as Zn fertilizer in rice cultivation. Zhao et al. (2010) reported that BF values tend to decrease with increasing soil heavy metal concentration, and values lower than 0.20 are considered normal when plants are grown on polluted soils (McGrath and Zhao, 2003). The differences in the obtained BF values depend on the heavy metals and the plant species. The higher BF value for Cd indicates that Cd accumulated in rice and maize shoots. These differences might be related to heavy metal-binding capacity to roots (Toth et al., 2009), available metals, interactions between physicochemical parameters, and the plant species grown in these soils (Bose and Bhattacharyya, 2008).

3.3 Quality of drainage water

Concentrations of BOD and COD ranged from 442 and 978 to 632 and 2445 mgL^{-1} in Zefta drain, while the BOD and COD value ranged from 540 and 882 to 723 and 2301 mgL^{-1} in drain no. 5 (Table 3). This water would be classified as high strength (Metcalf and Eddy, 2003). These results were in agreement with Pescond (1992), who reported that chemical properties of water include total dissolved solids (TDSs); BOD and COD showed higher values in untreated sewage water compared to groundwater. The BOD / COD ratio in Zefta drain and drain no. 5 ranged from 0.25 and 0.31 to 0.45 and 0.61, respectively. The waste water with a BOD / COD ratio below 0.50 contains some toxic components such as dyes and heavy metals (Linsley et al., 1992).

The average TDSs was 1016 in drain no. 5, 1130 in Zefta drain, and 334 mgL^{-1} in Baher El Mlah canal. The SARs in water of Zefta drain and drain no. 5 were above 12, which is considered a potential level for aggregate slaking, soil swelling, and clay dispersion, and thus reduction in hydraulic




Figure 3. Bioaccumulation coefficients of heavy metals in *Typha domingensis*, *Phragmites australis*, and *Cynodon dactylon* plants grown in Zefta drain.

conductivity (Mace and Amrhein, 2001). The heavy metals in the two drains were higher than in the water of Baher El Mlah canal which could be attributed to discharge of industrial waste water into the two drains without treatment. The level of heavy metals exceeded the criteria limits for irrigation water (FAO, 2010; E.C.S, 48/1992). Similar results reported by Matloub and Mehana (1998) showed that sewage often has high values of temperature, pH, hardness, alkalinity, COD, TDS, NO_3^- , NO_2^- , Na, K, Ca, and Mg. Chitdeshwari et al. (2002) reported that increased levels of sewage water increased the uptake of Cd and Cr in *Amaranthus* crops.

3.4 Heavy metal concentrations in sediments

The high heavy metal concentrations in sediments of drain no. 5 (Table 4) can be attributed to higher pH in water which can form ions of insoluble precipitates. The measured concentrations of heavy metals are higher than US Environmental Protection Agency (EPA) toxicity values (US EPA, 1999). Similar findings were reported by Thuy et al. (2007) who found that heavy metals in sediments of five canals which received untreated industrial waste water exceeded the US EPA toxicity levels. The partitioning of heavy metals between sediment and water can be expressed by distribution coefficient (K_d) values ($L kg^{-1}$). K_d values of sediment samples were the highest for Zn, Cd, and Mn, and lowest for Pb, Cu, and Ni. The higher K_d value indicates that the sorption of heavy metals by sediments was strong (Salomons and Forstner, 1980). Sediments are both carriers and potential sources of contaminants in aquatic systems, and these materials also affect groundwater quality and agricultural products when disposed of on land. In this study, the mean Cf values of the drain no. 5 sediments revealed that Zn, Mn, Cu, Cd, Pb, and Ni > 6 , indicating very high contamination due to the direct discharge of waste water from the residential and industrial areas.

Table 3. The chemical analysis of water collected from the Zefta drain, drain no. 5, and Baher El Mlah canal.

Parameters	Units	Drain no. 5	Zefta drain	Baher El Mlah	Water criteria for irrigation water
pH	–	9.80	12.2	7.20	6.5–8.4
TDS	mg L ⁻¹	1016	1130	334	2000
SAR	–	17.3	18.2	6.00	6–12
BOD	mg L ⁻¹	540–723	442–632	0.00	40 ¹
COD	mg L ⁻¹	882–2301	978–2445	0.00	60 ¹
Fe	mg L ⁻¹	0.09	0.56	0.01	5.0
Zn	mg L ⁻¹	0.02	0.037	0.00	2.00
Mn	mg L ⁻¹	0.68	2.91	0.03	0.20
Cu	mg L ⁻¹	0.15	0.28	0.12	0.20
Cd	mg L ⁻¹	0.03	0.07	0.001	0.01
Pb	mg L ⁻¹	1.05	0.18	0.05	5.00
Ni	mg L ⁻¹	0.12	0.31	0.02	0.20

¹ Egyptian Chemical Standards (48/1992).

Table 4. Average heavy metal concentrations, contaminant factor, and distribution coefficients (Kd) in sediments of drain no. 5 compared with toxicological reference values (US EPA, 1999).

Element	Conc. (mg kg ⁻¹)		Et	Cf	Kd (L kg ⁻¹)
	mean	± SD			
Zn	647.5	± 36.7	110	6.25	32375.0
Mn	2125.0	± 74.3	–	12.7	3125.0
Cu	425.0	± 12.4	16	4.25	2833.3
Cd	97.5	± 4.6	0.6	9.55	3250.0
Pb	145.0	± 4.5	31	4.80	138.10
Ni	195.0	± 9.8	–	7.33	1625.0

Et: US EPA toxicity reference value; Cf: contaminant factor; Kd: distribution coefficients (L kg⁻¹).

3.5 Bioaccumulation coefficients of aquatic plants

The bioaccumulation coefficients of heavy metals in plants of *Cynodon*, *dactylon*, *Phragmites australis*, and *Typha domingensis* grown in Zefta drain are shown in Fig. 3. The bioaccumulation coefficients of metals in *Cynodon dactylon* were higher than in *Phragmites australis* and *Typha domingensis*. These plant species can be considered as hyperaccumulators, and are used for the decontamination of contaminated water. The use of plants for decontamination of polluted waters has been described as rhizofiltration (Brooks, 1998). The three species would be useful for phytoremediation of contaminated water in a particular area. Bonanno (2013) showed that *Phragmites australis* and *Typha domingensis* species may be used as biomonitors of trace element contamination in sediment. Overall, *T. domingensis* and *P. australis* showed a greater capacity of bioaccumulation as well as a greater efficiency of element removal than *A. donax*. In particular, *T. domingensis* and *P. australis* may be used for Hg phy-

tostabilization; the former also acted as a hyperaccumulator for trace elements' phytoextraction and phytostabilization. In contaminated wetlands, the presence of *T. domingensis* and *P. australis* may increase the general retention of trace elements. Wafaa et al. (2010) demonstrated that *Phragmites australis* and *Tamarix aphillya* species are significant as vegetation filters and for cleaning the soils from contamination by heavy metals by phytoextraction. Antioxidant thiol compounds were probably involved in the mechanisms used by *P. australis* to alleviate metal toxicity. As *P. australis* is considered suitable for phytostabilizing metal-contaminated sediments, understanding its tolerance mechanisms to toxic metals is important to optimize the conditions for applying this plant in phytoremediation (Rocha et al., 2014).

4 Conclusions

Delta drains often receive high amounts of organic and inorganic pollutants from industrial and domestic waste water as well as diffuse agricultural drainage. High priority should be given to Zefta drain and drain no. 5 sites which receive high loads of pollutants. This was confirmed by the lower water quality and soils polluted by heavy metals in the El-Mahla El-Kobra area. Industrial and municipal waste water sources in El-Mahla El-Kobra area must be treated before being discharged in Zefta drain and drain no. 5. Using agricultural soils contaminated by heavy metals in the cultivation of rice and maize crops for human consumption may result in health hazards.

Acknowledgements. The authors appreciate and thank the College of Food and Agricultural Sciences, Research Center, and Deanship of Scientific Research, King Saud University, Saudi Arabia, for the financial support of this research study. We also acknowledge Tanta University, governorate of Tanta, Egypt, for the funds and support in carrying out this research.

Edited by: A. Cerdà

References

- Abou-Sekkina, M. M., Issa, R. M., Bastawisy, A. E., and El-Helece, W. A.: Characterization and evaluation of thermodynamic parameters for Egyptian heap fired rice straw ash (RSA), *Int. J. Chem.*, 2, 81–88, 2010.
- Alirzayeva, E. G., Shirvani, T. S., Yazici, M. A., Alverdiyeva, S. M., Shukurov, E. S., Ozturk, L., Ali-Zade, V. M., and Cakmak, I.: Heavy Metal accumulation in Artemisia and Foliaceous Lichen species from the Azerbaijan flora, *Forest Snow and Landscape Research*, 80, 339–348, 2006.
- Alloway, B. J.: *Heavy Metals in Soils*, John Wiley & Sons, Inc. New York, ISBN 0470215984, 1990.
- Al Naggar, Y. A., Naiem, E. A., Seif, A. I., and Mona, M. H.: Honey bees and their products as a bio-indicator of environmental pollution with heavy metals, *Mellifera*, 13, 10–20, 2013.
- APHA, AWWA and WEF: *Standard methods for the examination of water and wastewater*, XX Edn., American Public Health Association, Washington, DC, 2005.
- Berendse, F., van Ruijven, J., Jongejans, E., and Keesstra, S.: Loss of plant species diversity reduces soil erosion resistance, *Ecosystems*, 18, 881–888, 2015.
- Bonanno G.: Comparative performance of trace element bioaccumulation and biomonitoring in the plant species *Typha domingensis*, *Phragmites australis* and *Arundo donax*, *Ecotox. Environ. Safe*, 97, 124–130, 2013.
- Bose, S. and Bhattacharyya, A. K.: Heavy metal accumulation in wheat plant grown in soil amended with industrial sludge, *Chemosphere*, 70, 1264–1272, 2008.
- Brevik, E. C., Cerdà, A., Mataix-Solera, J., Pereg, L., Quinton, J. N., Six, J., and Van Oost, K.: The interdisciplinary nature of SOIL, *SOIL*, 1, 117–129, doi:10.5194/soil-1-117-2015, 2015.
- Brooks, R.: *Plants that hyper accumulated heavy metals* CAB international, London, UK, 1998.
- Chapman, H. D. and Pratt, P. F.: *Methods of Analysis for Soils, Plant and Water*, Univ. of California, Division of Agric. Sci., USA, Chapter 2, 56–64, 1961.
- Chen, X. W., Tsz-Fung Wong, J., Mo, W. Y., Man, Y. B., Wang-Wai Ng, C., and Wong, M. H.: Ecological Performance of the Restored South East New Territories (SENT) Landfill in Hong Kong (2000–2012), *Land Degrad. Dev.*, 1, 1–13, doi:10.1002/ldr.2366, 2015.
- Chen, Z. S., Lee, D. Y., Wong, D., and Wang, Y.: Effect of various treatments on the uptake of Cd from polluted soils by vegetable crops, in: *Proceedings of 3rd work shop of soil pollution and prevention*, National Chuny-Hsing University Taiwan, ROC, 277–292, 1992.
- Chibuikwe, G. U. and Obiora, S. C.: Heavy Metal Polluted Soils: Effect on plants and bioremediation methods, *Appl. Environ. Soil Sci.*, 12 pp., doi:10.1155/2014/752708, 2014.
- Chitdeshwari, T., Savithri, P., and Mahimai Raja, S.: Effect of sewage bio-solids compost on biomass yield of Amaranths and heavy metal availability, *J. Indian Soc. Soil Sci.*, 50, 480–484, 2002.
- Decock, C., Lee, J., Necpalova, M., Pereira, E. I. P., Tendall, D. M., and Six, J.: Mitigating N₂O emissions from soil: from patching leaks to transformative action, *Soil*, 1, 687–694, 2015.
- Dong, J., Yang, Q. W., Sun, L. N., Zeng, Q., Liu, S. J., and Pan, J.: Assessing the concentration and potential dietary risk of heavy metals in vegetables at a Pb/Zn mine site, China, *Environ. Earth Sci.*, 64, 1317–1321, 2011.
- Egyptian Chemical Standards: Protection of the Nile River and water stream from pollution, Ministry of Irrigation, Cairo, Egypt, low No. 48/1992.
- El-Gendi, S. A., Badawy, S. H., and Helal, M. I.: Mobility of some heavy metal nutrients in sandy soils irrigated with sewage effluent, *J. Agric. Sci. Mansoura Univ.*, 22, 3535–3552, 1997.
- Fakayode, S. and Onianwa, P.: Heavy metal contamination of soil and bioaccumulation in Guinea grass (*Panicum maximum*) around Ikeja industrial estate, Lagos, Nigeria. *Environ. Geology*, 43, 145–150, 2002.
- FAO.: The wealth of waste: The economics of wastewater use in agriculture, in: *FAO Water Report No. 35*, Water Development and Management Unit, Food and Agriculture Organization of the United Nations, ISBN 978-92-5-106578-5, 2010.
- Farooqi, Z. R., Iqbal, M. Z., Kabir, M., and Shafiq, M.: Toxic effects of lead and cadmium on germination and seedling growth of *Albezia lebbek (L.) Benth*, *Pak. J. Bot.* 41, 27–33, 2009.
- Ghoneim, A. M., Al-Zahrani, S., El-Maghraby, S., and Al-Farraj, A.: Heavy metal distribution in *Fagonia indica* and *Cenchrus ciliaris* native vegetation plant species, *J. Food Agric. Environ.*, 12, 320–324, 2014.
- Gupta, S., Satpati, S., Nayek, S., and Garai, D.: Effect of wastewater irrigation on vegetables in relation to bioaccumulation of heavy metals and biochemical changes, *Environ. Monit. Assess.*, 165, 169–177, 2010.
- Håkanson, L.: Ecological risk index for aquatic pollution control: sediment logical approach, *Water Res.*, 14, 975–1001, 1980.
- Hernandez, T., Moreno, J., and Costa, F.: Influence of sewage sludge application on crop yields and heavy metal availability, *Soil Sci. Plant Nutr.*, 37, 201–210, 1991.
- Iannelli, M.A., Pietrini, F., Flore, L., Petrilli, L., and Massacci, A.: Antioxidant response to cadmium in *Phragmites australis* plants, *Plant Physiol. Biochem.*, 40, 977–982, 2002.
- Imperato, M., Adamo, P., Naimo, D., Arienzo, M., Stanzione, D., and Violante, P.: Spatial distribution of metals in urban soils of Naples city (Italy), *Environ. Poll.*, 124, 247–256, 2013.
- Jackson, M. L.: *Soil Chemical Analysis*, Constable and Company Ltd, London, 1973.
- Jolly, Y. N., Islam, A., and Akbar, S.: Transfer of metals from soil to vegetables and possible health risk assessment, *Springer Plus*, 2, 1–8, 2013.
- Kabata-Pendias, A. and Pendias, H.: *Trace Elements in Soils and Plants*, Boca Raton, FL: CRC Press, 1992.
- Khaskhoussy, K., Hachicha, M., Kahlaoui, B., Messoudi-Nefzi, B., Rejeb, A., Jouzdan, O., and Arselan, A.: Effect of treated

- wastewater on soil and corn crop in the Tunisian area, *J. Appl. Sci. Res.*, 9, 132–140, 2013.
- Li, G. C., Haw-Tarn, L., and Chi-Sen, L.: Uptake of heavy metals by plants in Taiwan, paper from conference title: Biogeochemistry of trace elements, *Environ. Geochem. Hlth.*, 2, 153–160, 1994.
- Li, X. L., Gao, J., Brierley, G., Qiao, Y. M., Zhang, J., and Yang, Y. W.: Rangeland degradation on the Qinghai–Tibet Plateau: implications for rehabilitation, *Land Degrad. Dev.*, 24, 72–80, 2013.
- Linsley, R. K., Joseph, B., Franzini, D. L., and Freyberg, G. T.: *Water Resources Engineering, Fourth Edn.*, McGraw-Hill, Inc., New York, 1992.
- Loeppert, R. H. and Suarez, D. L.: Carbonate and gypsum, in: *Methods of Soil Analysis*, edited by: Sparks, D. L., Part3, SSSA Book Ser. 5., SSSA, Madison, WI, 437–474, 1996.
- Lone, M. I. and Rizwan, M.: Evolution of industrial effluents for irrigation and their effect on soil and chemical properties, in: *Proc. NSMTCC 97, Environ. Poll., Islamabad, Pakistan*, 30, 269–280, 1997.
- Mace, J. E. and Amrhein, C.: Leaching and reclamation of a soil irrigated with moderate SAR waters, *Soil Sci. Soc. Am. J.*, 65, 199–204, 2001.
- Mahmoud, E. and Abd El-Kader, N.: Heavy metal immobilization in contaminated soils using phosphogypsum and rice straw compost, *Land Degrad. Dev.*, 26, 819–824, 2015.
- Matloub, M. A. and Mehana, T. A.: Utilization of treated sewage effluent for reclaiming a salt-affected soil, *Egypt, J. Appl. Sci.*, 13, 298–316, 1998.
- McGrath, S. P. and Zhao, F. J.: Phytoextraction of metals and metalloids from contaminated soils, *Current Opinion in Biotechnology*, 14, 277–282, 2003.
- Metcalf and Eddy (Eds.): *Wastewater Engineering: Treatment, Disposal, Reuse*, 4th Edn., McGraw-Hill, Inc. New York, 2003.
- Mollahoseini, H.: Long term effects of municipal wastewater irrigation on some properties of a semiarid region soil of Iran, *Int. J. Agr. Plant Pro.*, 4, 1023–1028, 2013.
- Morgan, R.: Soil, heavy metals, and human health, in: *Soils and human health*, edited by: Brevik, E. C. and Burgess, L. C., Boca Raton, FL, USA, CRC Press, 59–82, 2013.
- Nassef, M., Hannigan, R., EL Sayed, K. A., and Tahawy, M. S.: Determination of some heavy metals in the environment of Sadat industrial city, *Proceedings of the 2nd Environmental Physics Conference*, 14–152, 2006.
- Nziguheba, G. and Smolders, A.: Inputs of trace elements in agricultural soils via phosphate fertilizers in European countries, *Sci. Total Environ.*, 390, 53–57, 2008.
- Onder, S., Dursun, S., Gezgin, S., and Demirbas, A.: Determination of Metal Pollution in Grass and Soil of City Centre Green Areas (Konya, Turkey), *Polish J. Environ. Studies*, 16, 145–154, 2007.
- Page, M. A. (Ed.): *Methods of soil analysis, Part 2*, Academic press, New York, 1982.
- Pescod, M. B.: *Wastewater treatment and use in agriculture*, Bull. FAO 47, 125, Rome, Italy, 1992.
- Richards, R. L.: *Diagnosis and improvement of saline and alkali soils*, USDA, Agriculture Handbook, No. 60US Gov., Printing office Washington, 1954.
- Riding, M. J., Martin, F. L., Jones, K. C., and Semple, K. T.: Carbon nanomaterials in clean and contaminated soils: environmental implications and applications, *SOIL*, 1, 1–21, doi:10.5194/soil-1-1-2015, 2015.
- Rocha, A. S., Marisa, R., Almeida, M., Clara, P., Bastoa, M., and Teresa, S. D.: Antioxidant response of *Phragmites australis* to Cu and Cd contamination, *Ecotox. Environ. Safe.*, 109, 152–160, 2014.
- Roy, M. and McDonald, L. M.: metal uptake in plants and health risk assessments metal-contaminated smelter soils, *Land Degrad. Dev.*, 26, 785–792, doi:10.1002/ldr.2237, 2013.
- Roy, M. and McDonald, L. M.: Metal uptake in plants and health risk assessments in metal-contaminated smelter soils, *Land Degrad. Dev.*, 26, 785–792, 2015.
- Sacristán, D., Peñarroya, B., and Recatalá, L.: Increasing the knowledge on the management of Cu-contaminated agricultural soils by cropping tomato (*Solanum Lycopersicum L.*), *Land Degrad. Dev.*, 26, 587–595, doi:10.1002/ldr.2319, 2015.
- Saffari, V. R. and Saffari, M.: Effect of treated municipal wastewater on bean growth, soil chemical properties, and chemical fractions of zinc and copper, *Arab. J. Geol. Sci.*, 6, 4475–4485, 2013.
- Saleemi, M. A.: Environmental assessment and management of irrigation and drainage scheme for sustainable agriculture growth, *Environ. Protection Agency Bull.*, (Lahore), 64 pp., 1993.
- Salomons, W. and Forstner, V.: Trace metal analysis on pollution sediments 11, Evaluation of environmental impact *Environ. Technol. Lett.*, 1, 506–517, 1980.
- Salwa, F. Shalaby, E., Bahy, A., and El Deen, A.: Water management problems associated with urban sprawl in Gharbia Governorate, Egypt using remote sensing and GIS, *Int. J. Adv. Rem. Sens. & GIS*, 2, 243–259, 2013.
- Smith, P., Cotrufo, M. F., Rumpel, C., Paustian, K., Kuikman, P. J., Elliott, J. A., McDowell, R., Griffiths, R. I., Asakawa, S., Bustamante, M., House, J. I., Sobocká, J., Harper, R., Pan, G., West, P. C., Gerber, J. S., Clark, J. M., Adhya, T., Scholes, R. J., and Scholes, M. C.: Biogeochemical cycles and biodiversity as key drivers of ecosystem services provided by soils, *SOIL*, 1, 665–685, doi:10.5194/soil-1-665-2015, 2015.
- Suciu, I. C., Cosma, M., Todici, S. D., Bolboacă, S. D., and Jantschi, L.: Analysis of soil metal pollution and pattern in central Transylvania, *Int. J. Molecul. Sci.*, 9, 434–453, 2008.
- Thuy, L. T., Nguyen, N. V., and Tu, T. L.: Anthropogenic input of selected heavy metals (Cu, Cr, Pb, Zn and Cd) in aquatic sediments of Hochiminh city, Vietnam, 2007.
- Toth, T., Tomas, J., Lazor, P., Bajcan, D., and Jomova, D.: The Transfer of metals from contaminated soils into agricultural plants in high Tatars region, *Czech J. Food Sci.*, 27, 390–393, 2009.
- UNCCD: Zero net land degradation: A sustainable development goal for Rio C20, Bonn, Germany, United Nations Convention to Combat Desertification, 2012.
- US Environmental Protection Agency: Screening level ecological risk assessment protocol for hazardous waste combustion facilities, Vol. 3, Appendix E: Toxicity reference values, EAP530-D99-001C, 1999.
- Wafaa, A., Al-Taisan, A. A., Al-Qarawi, A., and Moodi, S.: Effect of water stress by polyethylene glycol 8000 and sodium chloride on germination of Ephedra Alata Decne seeds, *Biol. Sci. J.*, 17, 253–257, 2010.
- Wang, H.-Q., Zhao, Q., Zeng, D.-H., Hu, Y.-L., and Yu, Z.-Y.: Remediation of a magnesium-contaminated soil by chemical amendments and leaching, *Land Degrad. Dev.*, 26, 613–619, 2015.

- Wang, M., Song, H., Chen, W., Lu, C., Hu, Q., and Ren Z.: Cancer mortality in a Chinese population surrounding a multi-metal sulphide mine in Guangdong province: an ecologic study, *BMC Public Health*, 16, 319, doi:10.1186/1471-2458-11-319, 2011.
- Ye, M., Sun, M., Liu, Z., Ni, N., Chen, Y., Gu, C., Kengara, F. O., Li, H., and Jiang, X.: Evaluation of enhanced soil washing process and phytoremediation with maize oil, carboxymethyl-cyclodextrin, and vetiver grass for the recovery of organochlorine pesticides and heavy metals from a pesticide factory site, *J. Environ. Manage.*, 141, 161–168, 2014.
- Zhang, H., Lin, Y. H., Zhang, Z., Zhang, X., Shaw, S. L., Knipping, E. M., Weber, R. J., Gold, A., Kamens, R. M., and Surratt, J. D.: Secondary organic aerosol formation from methacrolein photooxidation: Roles of NO_x level, relative humidity, and aerosol acidity, *Environ. Chem.*, 9, 247–262, 2012.
- Zhao, K., Liu, X., Xu, J., and Selim, H.: Heavy metal contaminations in a soil – rice system: identification of spatial dependence in relation to soil properties of paddy fields, *J. Hazard Mater.*, 181, 778–787, 2010.