



Original Research

Effect of pomegranate (*Punica granatum* L.) juice on kidney, liver, heart and testis histopathological changes, and the tissues lipid peroxidation and antioxidant status in lead acetate-treated rats

D. S. Aksu^{1*}, Y. S. Sağlam², S. Yildirim², T. Aksu³

¹ Department of Physiology, Faculty of Veterinary Medicine, Yuzuncu Yıl University, 65080, Van, Turkey

² Department of Pathology, Faculty of Veterinary Medicine, Ataturk University, 25100, Erzurum, Turkey

³ Department of Animal Science, Faculty of Veterinary Medicine, Yuzuncu Yıl University, 65080, Van, Turkey

Correspondence to: daksu@yyu.edu.tr

Received July 24, 2017; Accepted October 4, 2017; Published October 31, 2017

Doi: <http://dx.doi.org/10.14715/cmb/2017.63.10.5>

Copyright: © 2017 by the C.M.B. Association. All rights reserved.

Abstract: Pomegranate juice (PJ) contains relevant amounts of active biological compounds which alleviate the detrimental effects of chronic heavy metal exposure. This study investigated the protective potential of PJ against lead-induced oxidative stress. A total of forty adult male Sprague Dawley rats were divided into four experimental groups. The animals were fed a standard pellet diet and tap water *ad libitum*. The rats were divided into four groups (n=10 for each group): control, lead acetate (2000 ppm), low-treated PJ- a daily dose of 2.000 ppm lead plus 30µl pomegranate juice (included 1.050 µmol total polyphenols, gallic acid equivalent), and high-treated PJ- a daily dose of 2.000 ppm lead plus 60µl pomegranate juice (included 2.100 µmol total polyphenols, gallic acid equivalent). The treatments were delivered for 5 weeks. After the treatment period, the tissues samples (kidney, liver, heart and testis) were collected. Tissue lead (Pb) and mineral amounts (copper, zinc, and iron), tissues lipid peroxidation level and antioxidant status, and tissues histopathological changes were determined. The results showed that the highest rate lead loading was in the kidney and the testis. Pomegranate juice was decreased the lead levels of soft tissues examined; increased Zn amounts in tissues of which the lead accumulation was higher (kidney and the testis); decreased the copper, zinc and the iron levels of the liver and heart tissues, without creating a weakness in antioxidant capacity of these tissues, restricted the oxidative stress by decreasing lipid peroxidation, improved both of the activities of antioxidant enzymes such as superoxide dismutase (SOD) and catalase (CAT), and the level of glutathione (GSH) in all the tissues examined in lead-treated groups. As histopathological findings, the cellular damage induced by lead in the tissues of the kidney, liver and the heart were observed to have been partially prevented by PJ treatment. The protective effect of PJ was more pronounced in the testis compared to those others.

Key words: Lead; Phenolic compounds; Oxidative stress.

Introduction

Chronic exposure to lead results in its accumulation in vital organs, with serious pathological effects apparent in these organs after a short time of intestinal absorption (1,2). Pathological damages can occur in vital organs depending on duration and severity of lead exposure. Lead exposure causes renal dysfunction, liver cirrhosis, damage to the cardiovascular system and anemia (3). It can also cause some pathological damages in the ovaries and testes, causes by poor reproductive performance (4). Elimination of lead from tissues is slowly. Prolonged exposure to a low level of lead can hereby cause detrimental effects and disruptions occurs in membrane-related process such as the activities of membrane enzymes, endo- and exo-cytosis, transport of solutes across the bilayer and signal transduction processes (5,6). Lead accumulation in tissues causes oxidative stress by inducing the generation of reactive oxygen species (ROS), thereby reducing the anti-oxidant defence system of cells (1). Heavy metals can impair the metabolism of essential minerals (7). Minerals are defined as metal, and they can cause cellular damage and toxicity by interacting with biological

macromolecules in the presence of heavy metals such as lead (8,9). From the other side, the cells need trace minerals such as copper (Cu), zinc (Zn) and iron (Fe) to perform their normal biological functions. Therefore, detection and prevention of lead toxicity is a major international public health priority. The current therapeutic strategy for heavy metal toxicity is usually chelating agents. These agents are generally non-specific with respect to metals, and they have many side effects. The lead chelate complex can remain in tissue, and re-distribute to other tissues over time (10). In recent years, studies have begun to focus on the use of anti-oxidants as protectants against the harmful effects of lead (11-13). The pomegranate (*Punica granatum*, Punicaceae) is consumed fresh or can be processed into juice, jams, or a type of wine. Pomegranate juice contains in a satisfactory amounts of ascorbic acid, and the minerals include Fe, Ca, Mg, Se, Zn, as well as considerable amounts of phenolic compounds (14,15). Pomegranate juice has higher antioxidant capacity than other polyphenol-rich beverages (16). The antioxidant capacity of pomegranate juice in terms of free radical scavenging and iron reduction capacity has been shown to be three times higher than that of red wine and green tea (17). Com-

mercial pomegranate juice is obtained by pressing the whole pomegranate fruit and its peels. Pomegranate juice has considerable antioxidant potential with a high amount of phenolic compounds in comparison to commonly consumed fruit juices, such as grape, cranberry, grapefruit, or orange juice (18,19). Phenolic compounds are natural substances found in plants, fruits, vegetables and plant-derived consumables. These compounds show their mainly antioxidant effects by different mechanisms like inducing expression of protective genes against oxidative stress, regulation of reactive oxygen species by interacting with oxidative pathways and scavenging metal ions as pathogenic free radicals (20). Guo *et al.* (21) reported that 250 mL of pomegranate pulp juice (PPJ) daily for four weeks given to healthy elderly persons increased the plasma antioxidant capacity from 1.33 mmol to 1.46 mmol, while the other's consuming apple juice experienced no significant increase in antioxidant capacity. In addition, plasma carbonyl content (biomarker for oxidant/antioxidant barrier impairment in various inflammatory diseases) significantly decreased in the persons consuming the PPJ compared to the person consuming apple juice. Plasma vitamin E, ascorbic acid, and reduced glutathione values did not differ significantly between the groups, leading researchers to conclude that pomegranate phenolics may be responsible for the observed results. The molecular interactions of phenolic compounds with biological systems continue to be mostly speculative. The free-radical scavenging capability of polyphenols has been primarily tested with *in vitro* studies. However, phenolic compounds are structurally altered *in vivo*. As far as we know, inadequate *in vivo* studies have examined the lead binding activity of PJ, its interaction with minerals (copper, zinc and iron) and the protective effect of PJ against lead-induced oxidative damage. Therefore, the aim of the present study was to evaluate the lead binding activity of PJ, its interactions with the minerals (copper, zinc and iron), and the protective effect on the liver, kidney, heart and the testis of rats exposed to lead.

Materials and Methods

Chemicals

All chemicals were obtained from Sigma Chemical Inc. (St. Louis, MO, USA) and Merck Chemical Inc. (Darmstadt, Germany).

Pomegranate Juice (PJ) preparation

Fresh pomegranate fruit (*Punica granatum*) was purchased from a local retailer. Fresh fruits were washed and then squeezed to remove whole pomegranate juice.

Determination of Total Sugars (TS), Ascorbic Acid (AA) and Total Anthocyanins (TAs) of PJ

In order to estimate TS amount, the method described by Ranganna (22) was used, and the amount was expressed as g L⁻¹ of PJ. Amount of AA was determined using the method described by Ruck (23) and expressed as mg L⁻¹ of PJ. The total anthocyanins were estimated by the pH differential method using two buffer systems: potassium chloride buffer, pH 1.0 (25 mM), and sodium acetate buffer, pH 4.5 (0.4 M) (24). The total anthocyanins content was calculated as follows:

$$TAs = ((A \times MW \times DF \times 100) / MA)$$

where A = (A510 - A700) pH 1.0 - (A510 - A700) pH4.5; MW: molecular weight; DF: dilution factor; MA: molar absorptive coefficient of cyanidin-3-glucosid. Results were expressed as cyanidin-3-glucoside mg L⁻¹ of PJ.

Determination of Total Phenolic Content (TPC) and Antioxidant Activity (AA) of PJ

TPC was measured using the Folin-Ciocalteu method described by Ough and Amerine (25) using gallic acid as reference. The antioxidant capacity of pomegranate juice was evaluated based on free antioxidant activity radical scavenging capacity by the DPPH radical (expressed as inhibition percentage)(26).

Determination of copper, zinc, manganese and iron amounts of PJ

The fruit juice sample was analysed for minerals (copper, zinc, manganese and iron) using atomic absorption spectrophotometer (AAS, model AAS-200SN Rotalab).

Animals and experimental design

In this experimental study, forty adult male Sprague Dawley rats (300 ±10 gr) were purchased from an Experimental Research Institute. The animals were housed in plastic cages at 12-light/dark cycle, 21 ± 2°C with water and food available *ad libitum*. Prior to use, the animals were acclimatized under these conditions for 15 days. The rats were divided into four groups (n=10 for each group): 1) control, 2) lead - a daily dose of 2000 ppm lead acetate alone, 3) low-treated PJ- a daily dose of 2.000 ppm lead plus 30µl pomegranate juice (included 1.050 µmol total polyphenols, gallic acid equivalent), and 4) high-treated PJ- a daily dose of 2.000 ppm lead plus 60µl pomegranate juice (included 2.100 µmol total polyphenols, gallic acid equivalent). The experiments were approved by the local ethical committee at Mustafa Kemal University (B.30.2.M.K.Ü.0.00/05). Lead acetate was dissolved in water and delivered by drinking water, and pomegranate juice was applied by oral gavage. The experiment lasted for five weeks. The dose of PJ treatments was used based on results of previous study (27,28). At the end of the treatments, the rats were anesthetized and sacrificed following an overnight fasting. The organs (kidneys, livers, heart and testis) were dissected and washed in cold ice saline (0.9%) for lipid peroxidation, antioxidant enzyme and molecules, lead and micronutrient analysis. For histopathological examination, tissue sections were obtained from the tissues samples and fixed in 10% neutral buffered formalin, embedded in paraffin wax, sectioned at 5 µm thick, and stained with haematoxylin and eosin (HE). The sections were then examined and photographed under an optical microscope (29). For histopathological examination, tissue sections (kidneys, livers, heart and testis) were quickly removed and processed for histopathology analyses. Tissue sections were fixed in 10% neutral buffered formaldehyde for 48 hours and washed under tap water overnight. Following routine tissue preparation procedures, tissue samples were dehydrated through graded series of alcohol and xylene and embedded in paraffin blocks. Paraffin serial sections were cut at a thickness

of 5 and stained with haematoxylin and eosin (HE). The sections were then examined and photographed under an optical microscope (29). The sections examined on the light microscope were scored as no (-), mild (+), moderate (++) and severe (+++) according to the lesion severity (30). Other parts of tissues homogenate was prepared in an ice cold homogenization buffer (0.32 M sucrose, 1 mM EDTA, 10 mM Tris HCl, pH 7.4), and cytosolic samples of tissues homogenate were obtained by centrifuging at 10.000 g for 10 min at +4°C.

Lipid peroxidation assay

The tissues lipid peroxidation (as malondialdehyde, MDA) levels were determined using the method described by Yoshiko *et al* (31), based on the thiobarbituric-acid (TBA) reaction. The optical density was measured at 535 nm by spectrophotometer (Shimadzu UV 1208).

Lead, copper, zinc and iron estimation in tissues

Tissues were digested with concentrated nitric acid-hydrogen peroxide (2:3) using a microwave digestion system (speedwave MWS-2 Berghof products + Instruments Harresstr.1. 72800 Enien Germany) (32). The samples were brought to a constant volume and the tissue mineral contents were measured using an inductively couple plasma spectrophotometer (Perkin-Elmer, optima 2100 DV, ICP/OES, Shelton, CT 06484-4794, USA) (33).

Antioxidant enzyme and molecules

The method described by Sedlak and Lindsay (34) was used in order to determine the glutathione (GSH) concentrations of tissues examined. All of the non-protein sulfhydryl groups of cells are in the form of reduced GSH. The chromogen 5,5'-Dithiobis 2-nitrobenzoic acid (DTNB) is a disulfide compound that is readily reduced by sulfhydryl compounds to an intensely yellow compound. The absorbance of the reduced chromogen was measured at 412 nm. Catalase activity in the tissue homogenate was assayed by the decrease in absorbance of hydrogen peroxide at 240 nm as per the method of Aebi (35). Superoxide dismutase activity in the tissue homogenate was assayed spectrophotometrically as described by Sun *et al* (36). This method is based on inhibition of nitroblue tetrazolium (NBT) reduction by superoxide anions. The resulting reduction of NBT was measured at 560 nm by spectrophotometry. We determined all the tissues' protein levels with the Bradford reagent using a spectrophotometer at 595 nm.

Statistical analysis

The appropriate Sum of Squares Method was selected for ANOVA (37). The data were analyzed by the one-factor ANOVA using the general linear models procedure of SAS (38). Software for the main effect of treatments. The differences between the means were determined by the Duncan's multiple range test at a significance level of $P < 0.05$.

Table 1. Estimated chemical parameters of pomegranate juice.

Total sugars, g L ⁻¹	189.41
Ascorbic acid, mg L ⁻¹	124.76
Total anthocyanins, mg L ⁻¹	88.45
Total phenolics, mg L ⁻¹ gallic acid equivalent	6645
Total antioxidant activity, %	28.78
Copper, mg L ⁻¹	0.28
Zinc, mg L ⁻¹	0.39
Manganese, mg L ⁻¹	1.96
Iron, mg L ⁻¹	2.38

ted for ANOVA (37). The data were analyzed by the one-factor ANOVA using the general linear models procedure of SAS (38). Software for the main effect of treatments. The differences between the means were determined by the Duncan's multiple range test at a significance level of $P < 0.05$.

Results

Total Sugars (TS), Total Anthocyanins (TAs), Ascorbic Acid (A), Total Phenolics (TPs), Antioxidant Activity (AA) and mineral (Cu, Zn, Mn, and Fe) concentrations of pomegranate Juice

TS, TAs, A, TPs, AA, and mineral concentrations of pomegranate juice have been presented in Table 1. The results were in accordance with previous reports on other pomegranate cultivars (39,40). The pomegranate juice appears to be a good source of nutrients and variation in the mineral composition could originate from the pomegranate cultivar as well as agro-climatic conditions, handling practices and manufacturing conditions.

Histopathological examination of the tissues

The histopathological changes scores of tissues and examination have been displayed in Table 2-5, and Figure 1-4, respectively. Histopathological examination revealed normal histology of kidney in control group (Figure 1A). Lead administration produced severe hyperemia and infiltration of mononuclear cells in glomerular capillaries and veins in intertubular, dilatation of

Table 2. Histopathological changes scored in kidney tissue.

	Groups			
	Control	Lead	Low-treated PJ	High-treated PJ
Tubul epithelia degeneration	-	+++	+++	+
Tubul epithelia necrosis	-	+++	++	-
Mononuclear cell infiltration	-	+++	++	+
Hyperemia and hemorrhage	-	+++	++	+
Tubul dilatation	-	+++	+++	++

+++; severe; ++:middle; +:little; -: no.

Table 3. Histopathological changes scored in liver tissue.

	Groups			
	Control	Lead	Low-treated PJ	High-treated PJ
Hydropatic degeneration	-	+++	++	+
Hepatocyte necrosis	-	+++	++	-
Mononuclear cell infiltration	-	++	++	-
Hyperemia and hemorrhage	-	+++	+	+

+++; severe; ++:middle; +:little; -: no.

Table 4. Histopathological changes scored in heart tissue.

	Groups			
	Control	Lead	Low-treated PJ	High-treated PJ
Hyaline degeneration	-	+++	++	++
Zenker's degeneration	-	+++	++	-
Mononuclear cell infiltration	-	+++	++	+
Hyperemia and hemorrhage	-	++	+	-

+++: severe; ++:middle; +:little; -: no.

Table 5. Histopathological changes scored in testis tissue.

	Groups			
	Control	Lead	Low-treated PJ	High-treated PJ
Atrophy in seminiferous tubules	-	+++	++	-
Intertubular edema	-	++	+++	+
Slimming on the tubul wall	-	+++	++	-
Active spermium	+++	-	+	++

+++: severe; ++:middle; +:little; -: no.

tubules, degeneration and necrosis of tubular epithelium (Figure 1-B). In low-treated PJ group, hyperemia in tubular glomerular capillary and intertubular range, dilatation in the tubule lumen, degeneration in the tubular epithelium were observed in kidney tissues (Figure 1-C). In high-treated PJ group, dilatation in tubules and very few degeneration in tubular epithelium vessels with nearly normal histological structures were determined (Figure 1-D). For liver tissue, normal histological structures were observed in control group (Figure 2-A). In the group receiving lead alone, diffuse hydropic degeneration, necrosis, hyperemia in sinusoidal and acinar veins were detected (Figure 2-B) whilst degeneration in acinar region, few necrotic hepatocytes, sinusoidal dilatation and hyperemia were determined in low-treated PJ (Figure 2-C). In high-treated PJ group, a few number of degenerated hepatocytes in the acinar region, mild hyperemia (Figure 2-D) were observed in liver tissues. Also, histopathological examination of the heart sections of control group showed normal cardiac myocytes (Figure 3-A). In the group receiving lead alone, the common hyperemia and perivascular mononuclear cell infiltration, hyaline degeneration and Zenker's necrosis in the myocardium were detected (Figure 3-B). In low-treated PJ group, mononuclear cell infiltration, hyaline degeneration and a small number of Zenker's necrosis were determined (Figure 3-C). In high-treated PJ group, a few number of degenerated cells and interstitial hyperemia were observed (Figure 3-D). For testis tissue, normal histological structure were observed in control group (Figure 4-A). Lead administration produced atrophy of seminiferous tubular, thinning of the tubule wall due to




Figure 1. A. normal histological appearance of kidney tissue of control group, Bar: 50µm; B: kidney tissue of the rat received 2000 ppm lead alone; severe hyperemia and infiltration of mononuclear cells in glomerular capillaries and veins in intertubular, dilatation of tubules, degeneration (thin arrows) and necrosis of tubular epithelium (thick arrows), Bar: 50µm; C: kidney tissue of the rat received 2000 ppm lead plus 30µl pomegranate juice (low-treatment); hyperemia (thick arrows), hyperemia in tubular glomerular capillary and intertubular range, dilatation in the tubule lumen, degeneration in the tubular epithelium (thin arrows), Bar: 50µm. D: kidney tissue of the rat received 2000 ppm lead plus 60µl pomegranate juice (high-treatment); dilatation in tubules and very few degeneration in tubular epithelium (arrows) Bar: 50µm.




Figure 2. A. normal histological appearance of liver tissue of control group, Bar: 50µm; B: liver tissue of the rat received 2000 ppm lead alone; diffuse hydropic degeneration, necrosis, sinusoidal and central veins hyperemia (arrows), Bar: 50µm; C: liver tissue of the rat received 2000 ppm lead plus 30µl pomegranate juice (low-treatment); degeneration (thin arrow) in central region and few necrotic hepatocytes, sinusoidal dilatation (thick arrow) and hyperemia, Bar: 50µm; D: liver tissue of the rat received 2000 ppm lead plus 60µl pomegranate juice (high-treatment); a few number of central degenerated hepatocytes (arrows), mild central hyperemia, Bar: 50µm.

reduction in germ cells, the absence of spermatazoa in some tubule lumen. A small number of spermatazoa in some tubule lumen were detected in the group receiving lead alone (Figure 4-B). In low-treated PJ group, degeneration of germ cell and intertubular edema, a small number of spermatazoa in tubule lumen were observed (Figure 4-C). In high-treated PJ group, mild edema in intertubular range was determined whilst many sperme-




Figure 3. A. normal histological appearance of heart tissue of control group, Bar: 20µm; B: heart tissue of the rat received 2000 ppm lead alone; common heart hyperemia and perivascular mononuclear cell infiltration (Thick arrow), hyaline degeneration and necrosis (thin arrow), Bar: 20µm; C: heart tissue of the rat received 2000 ppm lead plus 30µl pomegranate juice (low-treatment); mononuclear cell infiltration, hyaline degeneration and in a small number of necrosis (arrows), Bar: 20µm; D: heart tissue of the rat received 2000 ppm lead plus 60µl pomegranate juice (high-treatment); a few number of degenerated cells (arrows) and vascular hyperemia. Bar: 20µm.

tazoa were also found (Figure 4-D).

Tissues' lead (Pb) and mineral levels

Table 6 shows the lead, copper, zinc and the iron levels of the kidney, liver, heart and the testis tissues. The lead levels of the tissues in the group that received a daily dose of 2.000 ppm lead (as lead acetate) were highest compared to those others ($P < 0.001$). In the present study, the highest lead accumulation was observed in the kidney. The rate of lead accumulation was 13.7 times higher in the kidneys of the lead-exposed rats compared to the controls. It was 8.6 times higher in the testis and 2.5 times higher in the heart and liver of the lead-exposed rats compared to the controls. Experimental application of lead increased the zinc accumulation in the kidney; it also increased the iron accumulation in the liver, heart and the testis tissues ($P < 0.001$). Contrarily, the lead application decreased the copper and zinc levels in the testis ($P < 0.001$). Furthermore, the copper and iron levels of the kidney, and the copper and zinc levels of the liver and the heart did not change by the experimental application of lead. The levels of both phenolic compounds decreased the lead amount of examined tissues compared to the group that received lead alone ($P < 0.001$). Additionally, both phenolic compounds levels further increased the zinc amount of kidney tissue compared to the the group that received lead alone. Phenolic compounds at 60 µl level significantly decreased the copper, zinc and iron amounts of liver tissue, even below the value of the control group ($P < 0.001$). Both phenolic compounds levels decreased the copper and zinc amounts of heart tissue compared to those of the controls and the group that received lead alone ($P < 0.001$). On the other hand, phenolic compounds at 30 µl level did not affect the iron accumulation in heart tissue, while phenolic compounds at 60 µl level significantly decreased the iron amount, even down to




Figure 4. A. normal histological appearance of tubules seminiferi of rat testis tissue of control group, Bar: 50µm; B: testis tissue of the rat received 2000 ppm lead alone; atrophy of seminiferus tubular, thinning of the tubule wall due to reduction in germ cells that make up tubular wall, the absence of spermatzoa in some tubule lumen (thick arrow), in a small number of spermatzoa in some tubule lumen (thin arrow), Bar: 50µm; C: testis tissue of the rat received 2000 ppm lead plus 30µl pomegranate juice (low-treatment); degeneration of germ cell and intertubular edema,(thick arrow) fewer making spermatozoa (thin arrow), Bar: 50µm; D: testis tissue of the rat received 2000 ppm lead plus 60µl pomegranate juice (high-treatment); mild edema in intertubular range (arrows), Bar: 50µm.

the iron level of the control group ($P < 0.01$). In the testis, the reduced levels of copper in the group receiving lead alone were increased to the control level in both groups that received PJ. The increased levels of iron in the group receiving lead alone were also decreased to that of control level in both of groups that received PJ ($P < 0.001$). Furthermore, the amount of zinc was increased in both PJ levels, even higher than that of the controls ($P < 0.001$).

Lipid peroxide levels and antioxidant parameters

Lipid peroxidation, determined by MDA formation, was increased in all examined tissues of the rats exposed to lead while PJ administration significantly decreased the level of lipid peroxidation in all tissues. When the group receiving lead alone was compared with the controls, it was observed that the GSH level was significantly decreased in all tissues while SOD and CAT activities were significantly increased. The GSH levels in both of the groups receiving PJ were significantly lower than that of the controls, although its level was significantly higher than the group receiving lead alone ($P < 0.001$). In particular, the GSH levels of heart muscle in the group receiving 60 µl PJ was very close to that of the controls. When examining the CAT and the SOD activities of the groups receiving both levels of pomegranate juice, it was observed that CAT activities in all tissues were lower than the group receiving lead alone, while it was higher than the controls' level. The SOD activities in the heart and the testis tissues of the group receiving 60 µl PJ were close to that of the control group ($P < 0.001$).

Table 6. Tissue lead (Pb) and mineral amounts of experimental groups.

	Control	Lead	Low-treated PJ	High-treated PJ	P
Items, mg/kg	Kidney				
Lead	0.03 ^d ±0.002	0.41 ^a ±0.034	0.18 ^b ±0.016	0.09 ^c ±0.001	0.001
Copper	5.11±0.295	5.45±0.505	5.56±0.465	4.52±0.223	0.249
Zinc	9.05 ^d ±0.091	11.87 ^c ±0.145	16.64 ^a ±0.799	15.01 ^b ±0.374	0.001
Iron	125.19±3.776	119.18±1.455	123.28±4.918	121.74±4.380	0.732
	Liver				
Lead	0.11 ^d ±0.002	0.26 ^a ±0.012	0.19 ^b ±0.003	0.14 ^c ±0.003	0.001
Copper	2.88 ^a ±0.091	2.64 ^a ±0.138	2.82 ^a ±0.153	2.20 ^b ±0.035	0.001
Zinc	41.66 ^a ±0.844	41.65 ^a ±1.260	38.72 ^{ab} ±1.192	37.77 ^b ±0.427	0.013
Iron	36.37 ^b ±0.631	39.32 ^a ±0.717	39.87 ^a ±0.536	33.01 ^c ±0.490	0.001
	Heart				
Lead	0.03 ^d ±0.001	0.08 ^a ±0.002	0.06 ^b ±0.001	0.04 ^c ±0.001	0.001
Copper	0.50 ^a ±0.011	0.49 ^a ±0.012	0.42 ^b ±0.021	0.32 ^c ±0.008	0.001
Zinc	35.59 ^a ±1.814	38.67 ^a ±0.727	31.35 ^b ±0.819	30.81 ^b ±0.927	0.001
Iron	6.06 ^b ±0.339	7.54 ^a ±0.274	7.11 ^a ±0.187	5.85 ^b ±0.210	0.001
	Testis				
Lead	0.17 ^d ±0.011	1.47 ^a ±0.095	0.57 ^b ±0.021	0.41 ^c ±0.025	0.001
Copper	3.62 ^a ±0.091	3.06 ^b ±0.084	3.90 ^a ±0.180	3.64 ^a ±0.202	0.003
Zinc	53.54 ^b ±2.108	47.31 ^c ±1.625	64.38 ^a ±2.312	63.42 ^a ±2.505	0.001
Iron	151.58 ^b ±4.65	180.99 ^a ±3.14	157.21 ^b ±5.211	162.30 ^b ±4.648	0.001

^{abcd}: Mean values within the same row labelled different letters are statistically significant (P<0.001).

Table 7. Lipid peroxide levels and antioxidant status of experimental groups.

	Control	Lead	Low-treated PJ	High-treated PJ	P
Tissues	MDA (µmol/mg protein)				
Kidney	12.51 ^d ±0.28	17.72 ^a ±0.33	15.95 ^b ±0.14	14.72 ^c ±0.18	0.001
Liver	12.01 ^d ±0.39	17.15 ^a ±0.19	15.28 ^b ±0.15	14.42 ^c ±0.06	0.001
Heart	22.65 ^c ±0.37	44.66 ^a ±2.30	29.14 ^b ±2.94	25.51 ^c ±0.22	0.001
Testis	7.25 ^b ±0.43	13.20 ^a ±0.99	12.13 ^a ±0.79	7.35 ^b ±0.41	0.001
	CAT (U/mg protein)				
Kidney	1.34 ^c ±0.02	1.90 ^a ±0.04	1.59 ^b ±0.07	1.48 ^b ±0.03	0.001
Liver	0.26 ^c ±0.02	1.22 ^a ±0.08	0.70 ^b ±0.05	0.60 ^b ±0.03	0.001
Heart	0.28 ^b ±0.01	0.57 ^a ±0.07	0.38 ^b ±0.08	0.33 ^b ±0.03	0.001
Testis	0.11 ^b ±0.01	0.33 ^a ±0.04	0.14 ^b ±0.01	0.10 ^b ±0.01	0.001
	SOD (U/mg protein)				
Kidney	0.88 ^c ±0.03	1.28 ^a ±0.03	1.11 ^b ±0.03	1.06 ^b ±0.04	0.001
Liver	1.17 ^c ±0.05	1.78 ^a ±0.07	1.56 ^b ±0.08	1.41 ^b ±0.08	0.001
Heart	0.22 ^c ±0.02	0.34 ^a ±0.01	0.27 ^b ±0.01	0.25 ^{bc} ±0.01	0.001
Testis	0.76 ^b ±0.04	1.01 ^a ±0.06	1.01 ^a ±0.04	0.87 ^{ab} ±0.04	0.001
	GSH (µmol/mg protein)				
Kidney	0.77 ^a ±0.02	0.53 ^c ±0.02	0.63 ^b ±0.01	0.68 ^b ±0.02	0.001
Liver	6.45 ^a ±0.27	2.77 ^d ±0.05	3.92 ^c ±0.21	5.09 ^b ±0.28	0.001
Heart	0.63 ^a ±0.02	0.37 ^c ±0.01	0.53 ^b ±0.03	0.58 ^{ab} ±0.03	0.001
Testis	0.35 ^a ±0.01	0.24 ^c ±0.02	0.28 ^b ±0.02	0.31 ^b ±0.02	0.001

^{abcd}: Mean values within the same row labelled different letters are statistically significant (P<0.001).

Discussion

To our knowledge, an inadequate number of studies has been carried out to determine the protective effects of phenolic compounds in pomegranate juice on mammalian tissues against lead accumulation and lead induced oxidative damage although the structure and mode of action of phenolic compounds and pome-

granate juice have been discussed. We therefore investigated the ameliorative effects of pomegranate juice against lead accumulation, the lead binding activity, interactions with minerals (copper, zinc and iron) and the protective effect on the liver, kidney, heart and testis of rats exposed to lead.

In the current study, the highest lead accumulation was observed in the kidney. Kidney cells can concen-

trate the lead in intranuclear inclusion bodies which are storage sites that may have a protective effect during excessive lead exposure (42). The histopathological findings in kidney tissue in this study are also supported by this statement. Lead exposure has affected the mineral amounts in tissues examined. According to this; lead exposure increased the zinc loading in the kidney; it increased the Fe loading in the liver, heart and the testis tissues. On the other hand, lead exposure decreased the loading of copper and zinc of the testis whilst it did not affect the amounts of copper and iron of the kidney, and the amounts of copper and zinc of the liver and the heart, respectively. It is known that zinc homeostasis is regulated by the kidney and that a small amount is excreted through bile (42). We also thought that the lead-induced kidney damage observed as pathological results may have resulted in a reduction in the urinary excretion of zinc; thus, Zn deposition may be increased in the kidneys. Another important finding of this study is that lead exposure resulted in the accumulation of iron in the liver, heart and the testis (Table 6). Iron is an essential element, and it is necessary for normal cellular functioning (43). Iron and lead occupy similar niches within the body. Therefore, they compete for similar binding sites, particularly during absorption (44). In doing so, the increased lead absorption in the gastrointestinal system may cause a limitation in iron absorption. When iron is deficient, hepatocytes produce less or no hepcidin (45). Iron loading in the liver and other tissues develops in animals with reduced hepcidin expression (46). In the present study, exposure to lead decreased the testicular copper and zinc amounts. The testes requires a high zinc concentration to maintain their normal physiology. Zinc maintains the redox balance by modulating several Zn-dependent enzyme metallothioneins (MT) (47). Metallothioneins are metal-binding proteins that are cysteine-rich and of low molecular weight. In fact, metallothioneins have a protective effect against toxic metals. Although the specific region for heavy metals-binding of metallothioneins is lower in number, their expressions are increased in the presence of heavy metals such as cadmium, lead and mercury (48). The metal-binding regions of metallothioneins also have specific areas for copper and zinc. The metal binding capacity of metallothioneins is directly related to the metal density in cells (45).

The reduction in the tissues' lead amounts in the rats that received PJ could be attributed to the lead-fixing activity of polyphenolic compounds. As far as we know, there is no previously study expressing these results, but the present findings showed that phenolic compounds in PJ decreased the tissue lead amounts. Two attachment sites for metal ions have been suggested within the molecular structure of flavanols, which are chemical compounds of polyphenols; the *o*-diphenolic groups in the 30,40-dihydroxy positions in the B ring and the keto structure 4-keto, 3-hydroxy or 4-keto and 5-hydroxy in the C ring. These functional groups bind to Cu and Fe ions. With a closer look at the molecular structures, the same chemical and molecular features can also be found in other chemical components of phenolic compounds (49). We can therefore speculate that polyphenols in PJ may have the ability of binding the lead as well as copper and iron. Although the binding of metals, such

as copper and iron, by phenolic compounds can reduce the bioavailability of these metals, this negative effect can also be advantageous in organisms. Copper and iron participate in Fenton and Heber–Weiss reactions, generating highly reactive hydroxyl radicals, which are associated with many pathological conditions (50). The binding of these metals may decrease the hydroxyl radical-induced cell toxicity. As a remarkable finding, the zinc amount also increased in the kidney and the testis, to which the most lead was loaded when pomagranate juice were applied. This finding has drawn attention onto the protective effects of pomagranate juice in lead exposure once again. High Zn concentration is essential for the normal physiology of the testes and Zn homeostasis is regulated by the kidney (51). A previous study indicated that zinc acts as an anti-oxidant and possibly as a chelator agent in lead toxicity, and PJ is rich in Zn (52). The reason for the zinc loading in the kidney and the testis tissues can be attributed to this mechanism.

As expected, the lead exposure enhanced the generation of ROS and lipid peroxidation, and caused cell damage. Lipid peroxidation is a well-known mechanism of oxidative damage caused by ROS and it has been used a potential marker of oxidative stress. In this study, the increases in lipid peroxidation (indicated by a high MDA level) in the tissues of lead-exposed rats were altered in the animals' antioxidant defence systems, including decreased GSH levels in all tissues examined and increased SOD and CAT activity (Table 7). The increase in SOD and CAT activities in all the tissues of the rats exposed to lead alone could be a compensation mechanism to counteract the decreasing level of GSH. Glutathione (GSH) contains reactive sulfhydryl groups (-SH), which play an important role in protecting the cell membrane lipids from ROS attack. However, GSH is rapidly oxidized by oxidants (53). Lead causes an elongation in fatty acids by increasing the number of double bonds, thereby increasing the lipid peroxidation in the cell membranes. Additionally, the affinity of lead for sulfhydryl groups (-SH) adversely affects the integrity of cell membranes (54). Peroxidation of the phospholipid structure in cells leads to deterioration of cell membrane integrity and ultimately to cell death (5). Thus, the histopathological findings such as cell damage in the glomerulus, liver hepatocytes, heart muscle cells and seminiferous tubules observed in the group receiving lead alone indicated the cell damage. These histopathological findings were supported by previous studies (55,56). Superoxide dismutase and CAT are important enzymatic antioxidants involved in protecting the cells against the detrimental effects of free radicals. SOD is a well-known major antioxidant enzyme containing copper and zinc (57). In the current study, increasing the supplementation level of PJ decreased the copper and zinc amounts, even to a level lower than those of controls, but this reduction did not create a weakness in SOD activity of the liver and the heart (Table 6 and 7). CAT is a heme-containing antioxidant enzyme, which acts sequentially to SOD in the conversion of hydrogen peroxide to water. Fe+3 protoporphyrin is a major component of CAT, and CAT activity is related to the copper level (58). Copper is necessary for adequate utilization of iron, which is an important component of CAT. Similarly, increasing the supplementation level of PJ decreased the copper and

iron levels of the liver and the heart, but this reduction did not create a weakness in CAT activity in these tissues (Table 6 and 7). According to these findings, reduction of the amounts of copper-, zinc- and iron caused by phenolic compounds in the PJ did not create any negative impact on either of the antioxidant activities SOD and CAT in the liver and heart tissues. The chelation of these micronutrients may be beneficial in living organisms by inhibiting the oxidative damage in these tissues. The close link between these minerals (copper and iron) and Fenton and Heber–Weiss reactions that are generating highly reactive hydroxyl radicals, which are associated with many pathological conditions is well known (50). In this study, PJ could alleviate the oxidative damage by decreasing the lipid peroxidation (indicated by low MDA level), improving the activities of antioxidant enzyme (CAT and SOD) and GSH level (Table 7). This contribution could be attributed to the presence of ascorbic acid, total anthocyanins, total phenolics and total antioxidant activity in PJ (Table 1). In agreement with the present study, Wie *et al.*, (59) found that treatment with EPP (the extracts of pomegranate peels, 150mg/kg body weight) and EPS (the extracts of pomegranate seeds, 100mg/kg body weight) markedly decreased the hepatic MDA levels and efficiently increased the hepatic SOD and GSH-Px activities compared with that of the carbon tetrachloride (1mL/kg body weight) injected groups in rats. Treatment with EPP and EPS obviously alleviated the collagen deposition and liver injury. In another study conducted by Reckziegel *et al.*, (60), the antioxidant protection of gallic acid (GA) against toxicity induced by Pb in rats were investigated. It was observed that an increase in Pb levels in blood, liver and kidney ($P < 0.001$) in Pb-exposed rats in relation to control group. This affect was not reverted by GA treatment. SOD activity was decreased in liver and kidney of Pb-exposed rats in relation to control group, and the events were partially reverted in liver and completely in kidney by GA. In liver CAT activity decreased in Pb-exposed rats compared with control group, and this affect reversed by GA treatment ($P < 0.05$). No changes were observed in kidney CAT activity (60). These compounds have protective effects against the generation of high reactive species such as hydroxyl radicals ($\text{OH}\cdot$), hydrogen peroxide (H_2O_2) and superoxide anions ($\text{O}_2^{\cdot-}$) (16). On the other side, the polyphenolic compounds in PJ did not exactly repair the tissue damage compared to the normal cell structure, but showed an ameliorative effect against lead-induced pathological damages on the rat tissues examined, especially in the testes. This effect was especially pronounced in the group that received $2100\mu\text{mol}$ polyphenols (high treated group) by pomegranate juice, such that lead-induced pathological damages were mild or of low intensity in this group compared to those of others. Oyagbemi *et al.* (61) observed the histopathological changes like degeneration and coagulation necrosis in tubul epithelium, mononuclear cell infiltration in inter tubular spaces, and severe hyperemia and hemorrhage in the kidneys of Pb-exposed rats (0.25, 0.50, and 1.0 mg/mL of lead acetate). They reported an increase in the severity of lesions with increasing Pb-doses. Similarly, Abdou and Hassan (3) reported hepatic histopathological changes like degeneration and necrosis in hepatocytes, dilatation in portal veins,

hyperemia and haemorrhage, mononuclear cell infiltration in portal region in the liver of cadmium-exposed rats. Kolawole *et al.*(4) observed a significant reduction in the number of sperm and motility, and histopathological changes like significant thinning in the wall of the testis tubules, deterioration of the tubulosic structure in the rats that received 2.25mg/kg body weight of lead acetate (4). Muhammed (62) reported that the phenolic compound extract of green tea protected kidney tissues against toxicity effect of cadmium sulphate by improving the histopathological alterations and normalizing the kidney biochemical parameters. A group which orally gavage with phenolic compound extract of green tea (at a dose 400mg/kg of body weight) showed high cellularity glomeruli and normal renal convoluted tubules which lining with normal endothelial cells (62). In a new study (63) where avacado fruit extract (AFE) and avacado seed extract (ASE) were used polyphenol-rich ingredients in diethylnitrosamine/2-acetylaminofluorene (DEN/2AAF)-exposed rats, it was observed that the DEN/2AAF-administration created histopathological changes in both of kidney and heart like mononuclear leucocytic aggregation and perivascular oedema, severe vacuolar degenerative changes of the epithelial lining of renal tubules, brown pigments accumulation and few karyomegalic nuclei, perivascular oedema, intramuscular oedema, degeneration of certain cardiomyocytes and intramuscular haemorrhage, respectively (63). The treatment of AFE and ASE showed congestion in the blood vessels with nearly normal histological structure in kidney, and no histopathological changes in heart (63). The histopathological findings of the previously mentioned studies are consistent with our findings.

It was concluded that the ameliorative effect of PJ on metal detoxification was carried out with ascorbic acid, minerals including Fe, Ca, Mg, Se, Zn, as well as considerable amounts of phenolic compounds together. As a result, regular consumption of pomagranate juice could be useful in protecting from chronic lead exposure, which is an important public health problem.

Author agreement

The authors have no conflicts of interest to identify. All of the authors have reviewed and approved this manuscript prior to submission. This manuscript describes the original work of the authors and has not been previously considered for publication.

Acknowledgments

This work was supported by Mustafa Kemal University (MKUBAP-08G0101).

References

1. Mansouri MT, Cauli O. Motor alterations induced by chronic lead exposure. *Environ Toxicol Phar* 2009; 27: 307–13
2. Navarro-Moreno LG, Quintanar-Escorza MA, González S, Mondragón R, Cerbón-Solorzano J, Valdés J, Calderón-Salinas JV. Effects of lead intoxication on intercellular junctions and biochemical alterations of the renal proximal tubule cells. *Toxicol in Vitro* 2009; 23: 1298–1304.
3. Abdou HM, Hassan MA. Protective role of omega-3 polyunsaturated fatty acid against lead acetate-induced toxicity in liver and kidney of female rats. *Biomed Res Int*. 2014; 1-11

4. Kolawole TA, Dapper DV, Ojeka SO. Ameliorative effects of the methanolic extract of the rind of *Citrullus lanatus* on lead acetate induced toxicity on semen parameters and reproductive hormones of male albino wistar rats. *European J Med Plants* 2014; 4 (9): 1125-37
5. Ercal N, Gurer-Orhan H, Aykin-Burns N. Toxic metals and oxidative stress. Part I: mechanisms involved in metal-induced oxidative damage. *Curr Top Med Chem* 2001; 1: 529-39
6. Noriega GO, Tomaro ML, del Batlle AM. Bilirubin is highly effective in preventing in vivo δ -aminolevulinic acid-induced oxidative cell damage. *Biochim Biophys Acta* 2003; 638: 173-8
7. Alonso ML, Montaña FP, Miranda M, Castillo C, Hernández J, Benedito JL. Interactions between toxic (As, Cd, Hg and Pb) and nutritional essential (Ca, Co, Cr, Cu, Fe, Mn, Mo, Ni, Se, Zn) elements in the tissues of cattle from NW Spain. *Biometals* 2004; 17: 389-97.
8. Flora SJS, Mittal M, Mehta A. Heavy metal induced oxidative stress and its possible reversal by chelation therapy. *Indian J Med Res* 2008; 128: 501-23.
9. Leonard SS, Harris GK, Shi XL. Metal-induced oxidative stress and signal transduction. *Free Rad Biol Med* 2004; 37: 1921-42
10. Lowry JA. Oral chelation therapy for patients with lead poisoning. *American Academy of Pediatrics*. http://www.who.int/selection_medicine/committees/expert/18/applications/4_2Lead_Oral_Chelators.pdf. (accessed 16.25.11)
11. Saxena G, Flora SJS. Lead induced oxidative stress and hematological alterations and their response to combined administration of calcium disodium EDTA with a Thiol chelator in rats. *J Biochem Mol Toxic* 2004; 18 (4): 221-33
12. Aksu DS, Didin M, Kayıkçı F. The protective role of polyphenols on blood cells in rats exposed to lead. *Rev Romana Med Lab* 2012; 20 (3/4):47-57
13. Aksu DS, Sağlam YS, Aksu T. The investigation of neuroprotective effects of pomegranate juice against low level lead induced oxidative stress in rats brain. *Eurasian J Vet Sci* 2016; 2(4): 255-59
14. Espin JC, García-Conesa MT, Tomás-Barberán FA. Nutraceuticals: facts and fiction. *Phytochemistry* 2007; 68: 2896–3008
15. Lansky EP, Newman RA. *Punica granatum* (pomegranate) and its potential for prevention and treatment of inflammation and cancer. *J Ethnopharmacol* 2007; 109:177–206
16. Seeram NP, Aviram M, Zhang Y, Henning SM, Feng L, Dreher M, Heber D. Comparison of antioxidant potency of commonly consumed polyphenol-rich beverages in the United States. *J Agric Food Chem* 2008; 56: 1415–22
17. Gill MI, Tomás-Barberán FA, Hess-Pierce B, Holcroft DM, Kader AA. Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing. *J Agric Food Chem* 2000; 48: 4581–89
18. Azadzi KM, Schulman RN, Aviram M, Siroky MB. Oxidative stress in arteriogenic erectile dysfunction: prophylactic role of antioxidants. *J Urol* 2005; 174:386–93
19. Rosenblat M, Aviram M. In: *Pomegranates: Ancient Roots to Modern Medicine, Antioxidative Properties of Pomegranate: In vitro studies*. Seeram NP, Heber D. (eds.) Taylor and Francis Group, New York, 2006, pp 31–43
20. Kelsey NA, Wilkins HM, Linseman DA. Nutraceutical antioxidants as novel neuroprotective agents. *Molecules* 2010; 15: 7792-814
21. Guo C, Wei J, Yang J, Xu J, Pang W, Jiang Y. Pomegranate juice is potentially better than apple juice in improving antioxidant function in elderly subjects. *Nutr Res* 2008; 28: 72-7
22. Ranganna S. Sugar estimation. In: *Handbook of analysis and quality control for fruit and vegetable products*. Ranganna S. (ed.), 2nd edn. Tata McGraw-Hill, New Delhi, 2001, pp 12-17
23. Ruck JA. *Chemical methods of analysis of fruits and vegetables*. Dep Agri Canada Publication No. 1154, Canada, 1963.
24. Giusti MM and Wrolstad RE. Characterization and measurement of anthocyanins by UV-visible spectroscopy. In: *Current protocols in food analytical chemistry*. Wrolstad RE and Schwartz SJ (eds.) John Wiley and Sons, New York, 2001, pp 1–13
25. Ough CS and Amerine MA *Methods for analysis of musts and wines*. A Wiley Inter-Science Publication, New York, 1998.
26. Cuendet M, Hostettmann K, Potterat O. Iridoid glucosides with free radical scavenging properties from *Fragaria bluei*. *Helvetica Chim Acta* 1997; 80: 1144-52
27. Aviram M, Dornfeld L, Rosenblat M, Volkova N, Kaplan M, Coleman R, Hayek T, Presser D, Fuhrman B. Pomegranate juice consumption reduces oxidative stress, atherogenic modifications to LDL, and platelet aggregation: studies in humans and in atherosclerotic apolipoprotein E-deficient mice. *Am J Clin Nutr* 2000; 71: 1062–76.
28. Kaplan M, Hayek T, Raz A, Coleman R, Dornfeld L, Vaya J, Aviram M. Pomegranate juice supplementation to atherosclerotic mice reduces macrophage lipid peroxidation, cellular cholesterol accumulation and development of atherosclerosis. *J Nutr* 2001; 131(8): 2082-89.
29. İlhan F, Vural SA, Yıldırım S, Sözdutalmaz İ, Alcigir ME. Expression of p53 protein, Jaagsiekte sheep retrovirus matrix protein, and surfactant protein in the lungs of sheep with pulmonary adenomatosis. *J Vet Diagn Invest* 2016; 28 (3): 249-56
30. Hoshino J, Mise K, Ueno T, Imafuku A, Kawada M, Sumida K, Hiramatsu R, Hasegawa E, Yamanouchi M, Hayami N, Suwabe T, Sawa N, Hara S, Fujii T, Ohashi K, Ubara Y, Takaichi K. A pathological scoring system to predict renal outcome in diabetic nephropathy. *Am J Nephrol* 2015; 41: 337-344
31. Yoshiko T, Kawada K, Shimada T. Lipid peroxidation in maternal and cord blood and protective mechanism against active-oxygen toxicity in the blood. *American J Obstet Gynecol* 1979; 135:372-376
32. Mertens D. AOAC Official Method 922.02. In: Horwitz W and Latimer GW (eds) *Preparation of laboratory sample. Official methods of analysis*, Chapter 3. Eighteenth ed. AOAC-International Suite 500,481. Maryland: North Frederick Avenue, 20877-2417, Gaithersburg, (2005a) pp 1-2
33. Mertens D. AOAC official method 975.03. In: Horwitz W and Latimer GW (eds) *Metal in plants and pet foods. Official methods of analysis* Chapter 3. Eighteenth ed., AOAC-International Suite 500, 481. Maryland: North Frederick Avenue, 20877-2417, Gaithersburg, (2005b) pp 3-4
34. Sedlak J, Lindsay RH. Estimation of total protein-bound and nonprotein sulfhydryl groups in tissue with Ellman's reagent. *Anal Biochem* 1967; 25: 192-205
35. Aebi H. Catalase in vitro. *Methods Enzymol* 1984; 105:121-6.
36. Sun Y, Oberley LW, Ying L. A simple method for clinical assay of superoxide dismutase. *Clin Chem* 1988; 34: 497-500.
37. Ergün G, Aktaş S. Comparisons of sum of squares methods in ANOVA Models. *Kafkas Univ Vet Fak Derg* 2009; 15 (3):481-4
38. SAS: SAS/STAT. User's Guide. Release 6.08 ed., SAS. Institute Inc, Cary, North Carolina, USA, 1994.
39. Cam M, Hisil Y, Durmaz G. Classification of eight pomegranate juices based on antioxidant capacity measured by four methods. *Food Chem* 2009; 112: 721–6
40. Akhavan H, Barzegar M, Weidlich H, Zimmermann BF. Phenolic compounds and antioxidant activity of juices from ten Iranian pomegranate cultivars depend on extraction. *Chemist* 2015; 1-7
41. Arslan H, Saripinar-Aksu D, Ozdemir S, Yavuz O, Or ME, Barutcu UB. Evaluation of the relationship of blood heavy metal, trace element levels and antioxidative metabolism in cattle which are living near the trunk roads. *Kafkas Univ Vet Fak Derg* 17 (Suppl A) 2011; 77-82
42. Gupta RC. *Veterinary Toxicology. Basic and clinical principles*,

- 2nd edn. Academic Press in an Imprint of Elsevier, London, 2012.
43. Merle U, Fein E, Gehrke SG, Stremmel W, Kulaksiz H. The iron regulatory peptide hepcidin is expressed in the heart and regulated by hypoxia and inflammation. *Endocrinology* 2007; 148 (6):2663-8
44. Kwong WT, Friello P, Semba RD. Interactions between iron deficiency and lead poisoning: epidemiology and pathogenesis. *Sci Total Environ* 2004; 330 (1-3): 21-37
45. Ganz T, Nemeth E (2012) Hepcidin and iron homeostasis. *Biochimica et Biophysica Acta* 2012; 1823:1434-43
46. Ganz T. Hepcidin and iron regulation, 10 years later. *Blood* 2011; 117 (17): 4425-33
47. Oteiza PI. Zinc and the modulation of redox homeostasis. *Free Radic Biol Med* 2012; 53(9): 1748-59.
48. Carpena E, Andreani G, Isani G. Metallothionein functions and structural characteristics. *J Trace Elem Med Biol* 2007; 21: 35-9
49. David MP, Valente P, Pereira JA, Paula B. Andrade PB. Phenolics: from chemistry to biology. *Molecules* 2009; 14: 2202-11
50. Prousek J. Fenton chemistry in biology and medicine. *Pure Appl Chem* 2007; 79: 2325-38
51. Kambe T, Tsuji T, Hashimoto A, Itsumura N. The physiological, biochemical, and molecular roles of zinc transporters in zinc homeostasis and metabolism. *Phys Rev* 2015; 95: 749-84
52. Jomova K, Valko M. Advances in metal-induced oxidative stress and human disease. *Toxicol* 2011; 283: 65-87
53. Cremers CM, Jacob U. Oxidant sensing by reversible disulfide bond formation. *J Biol Chem* 2013; 288 (37): 26489-96
54. Yuan G, Dai S, Yin Z, Lu H, Jia R, Xu J et al. Sub-chronic lead and cadmium co-induce apoptosis protein expression in liver and kidney of rats. *Int J Clin Exp Pathol* 2014; 7 (6): 2905-14.
55. El-Neweshy MS, El-Sayed YS. Influence of vitamin C supplementation on lead-induced histopathological alterations in male rats. *Exp Toxicol Pathol* 2011; 63 (3):221-7
56. Patra RC, Swarup D, Dwivedi SK. Antioxidant effects of α -tocopherol, ascorbic acid and L-methionine on lead induced oxidative stress to the liver, kidney and brain in rats. *Toxicol* 2001; 162 (2): 81-8
57. Gurer H, Ercal N. Can antioxidants be beneficial in the treatment of lead poisoning? *Free Radic Biol Med* 2000; 29 (10):927-45
58. Aksu DS, Aksu T, Özsoy B, Baytok E. The effects of replacing inorganic with a lower level of organically complexed minerals (Cu, Zn and Mn) in broiler diets on lipid peroxidation and antioxidant defense systems. *Asian-Aust J Anim Sci* 2010; 23 (8): 1066-72
59. Wei X, Fang R, Yang Y, Bi X, Ren G, Luo A et al. , Zhao M, Zang W. Protective effects of extracts from Pomegranate peels and seeds on liver fibrosis induced by carbon tetrachloride in rats. *BMC Complement Altern Med* 2015; 15: 389
60. Reckziegel P, Dias VT, Benvegenu DM, Bouffleur N, Barcelos RCS, Segat HJ et al. Antioxidant protection of gallic acid against toxicity induced by Pb in blood, liver and kidney of rats. *Toxicol Reports* 2016; 3: 351-6
61. Oyagbemi AA, Omobowale TO, Akinrinde AS, Saba AB, Ogundolu BS, Daramola O. Lack of reversal of oxidative damage in renal tissues of lead acetate-treated rats. 2015; 30 (11):1235-43
62. Muhammed ZI. Effect of phenolic compound extract of green tea to ameliorate the Cadmium Sulphate toxicity on the female rat kidneys. *J Pharm Biol Sci* 2014; 9 (2):44-50.
63. Abdel-Moneim AA, Osama MA, Hanaa IF, Eman EM. The preventive effects of avocado fruit and seed extracts on cardio-nephrotoxicity induced by diethylnitrosamine/2-acetylaminoflurine in wistar rats. *Basic Sci Med J* 2017; 6(1): 4-13