

Effect of Potassium Nitrate and Calcium Chloride priming on germination and seedling growth of Honesty (*Lunaria annua* L.)

Reyaz Ahmad Bhat¹, F U Khan¹, Z. A. Bhat¹, F. A. Khan², Javaid Ahmad Wani³, Abu Manzar⁴, I. T Nazki¹, Neelofar¹, Qazi Altaf¹ and Nasir Hamid Masoodi¹

¹*Division of Floriculture and Landscape Architecture, Faculty of Horticulture, SKUAST-K, Shalimar Campus, Jammu and Kashmir, India*

²*Division of Basic Sciences and Humanities, SKUAST-K, Shalimar Campus, Jammu and Kashmir, India*

³*Division of Soil Science, SKUAST-K, Shalimar Campus, Jammu and Kashmir, India*

⁴*Division of Entomology, SKUAST-K, Shalimar Campus, Jammu and Kashmir, India*

Corresponding author's email id: bhatriaz.2012@gmail.com

Abstract

Seed priming improves seed performance under environmental conditions. The study was designed to evaluate the effect of different priming treatments on the germination behaviour of Honesty (*Lunaria annua* L.). The experiment was conducted under a complete randomized design (CRD) with three replications. Seed priming was done with different concentrations of calcium chloride (CaCl₂), potassium nitrate (KNO₃) and hydro priming. All the treatments except T₁ (Control) had a significant effect on seedling establishment and seedling vigour. Results show that maximum invigoration was observed in seeds primed at 1% KNO₃ while minimum invigoration was observed in T₁ (Control). It was concluded that the germination percentage could be increased by using lower concentrations of KNO₃ and CaCl₂.

Keywords: Seed priming, Honesty, Potassium nitrate, Calcium chloride.

INTRODUCTION

Honesty (*Lunaria annua* L.), that is also known as silver dollar, satin flower, money plant, and moonwort, belongs to family Brassicaceae. It's a tall biennial ornamental with hairy stems that's native to southeastern Europe and western Asia. It has unstalked, coarsely serrated leaves with huge scentless flowers of 25-30 mm in diameter that bloom from April to June (Fitter *et al.*, 1980) if the Vernalization conditions have been met throughout the previous fall and winter (Pierik, 1967). a biennial crop that starts growing in the spring and stays vegetative for the first year. Lunaria is a spring flowering plant with toothed leaves and odorless flowers that grows without petioles (April to June). This plant, in addition to its attractiveness, produces lovely violet and white flowers that may be used as decorative plants at home for several years after the fruits (coins) have reached full maturity and the fruit inflorescence has been harvested. The skeletal appearance of the Lunaria, which is most stunning in its dried condition, is most eye-catching in the winter garden. Lunaria gets its common and botanical names from the oval seed pods and the gleaming silique that remains once the pod splits to release the seed.

Seed germination is considered as critical stage in the whole life of a plant (Yang *et al.*, 2008). Poor seed germination and uneven seedling emergence results into low quality products which leads to financial losses to growers. Several techniques and strategies have been developed to improve seed germination and uniform seedling growth. Seed priming is a presoaking seed treatment to improve seed germination and uniform seedling emergence in a wide range of crops (McDonald, 2000). In seed priming, seeds are partially hydrated so that it activates metabolic activities but prevents radicle protrusion. After seed priming, seeds are re-dried to initial moisture content prior to sowing

Osmotic adjustment or priming of seeds prior to sowing is known as an efficient way to increase germination and emergence rate in some species with for example small embryo or species with stepwise seed development (Sivritepe, 2000). The beneficial effects of priming have also been demonstrated for many field crops (Mehmet *et al.*, 2006). Seed

priming has been successfully demonstrated to improve germination and emergence in seeds of many crops, particularly seeds of vegetables and small seeded grasses (Arif *et al.*, 2007). The purpose of these treatments is to shorten the emergence and to protect seed from biotic and abiotic factor during critical phase of seedling establishment so as to synchronize emergence, which lead to uniform stand and improved yield. These priming treatments which enhance seed germination include hydro-priming (Basra *et al.*, 2003; Afzal *et al.*, 2004), osmopriming and hormonal priming (Afzal *et al.*, 2006). It was indicated that primed seed emerged one to three days earlier than non primed seeds and quickly became apparent. Similarly, improvement in germination, reduction in germination time and enhanced emergence in hydro-primed seeds were reported by Harris *et al.* (2001). Priming is the enhancement of physiological and biochemical events in seeds during suspension of germination by low osmotic potential and negligible matric-potential of the imbibing medium. Salts or non penetrating organic solutes in liquid medium (osmo-conditioning) or matrix solutions (matri-conditioning) are used to establish equilibrium water potential between seed and the osmotic medium needed for conditioning (Khan, 1993). The present study was planned to evaluate the effects of KNO₃ and CaCl₂ on seed germination of *Lunaria annua* L.

2. MATERIALS AND METHODS

Experiment was conducted in laboratory of division of floriculture and landscape architecture faculty of Horticulture, Sher-e-Kashmir university of Agriculture Sciences Srinagar Kashmir, during the year 2021 To improve the rate of germination and reduce the time required for germination, seed priming was done with different concentration of calcium chloride (CaCl₂) potassium nitrate (KNO₃) and hydro-priming to determine its effect on seed germination (Table 1 & 2). For priming, seeds were soaked in distilled water and different concentration of potassium nitrate solution for 48 h. After treatment, the seeds were placed on wet filter paper treated with fungicide Dithane M-45 @ 1 g/L in growth room. The seeds were kept in growth chamber at 24 ± 2°C. The seeds were watered at three days interval and kept under close observation. Treatments were designed with three replications and 10 seeds per replication in complete randomized design (CRD). Experimental data was analyzed statistically adopting the techniques of analysis of variance (ANOVA) for Completely

Randomized Design (CRD). The level of significance of the treatment mean square at 5% probability was tested against F calculate value for Completely Randomized Design (CRD). The data were recorded on days to 50 % germination, germination percentage, seedling fresh weight, seedling dry weight, root length, shoot length, seedling vigour index and root/shoot ratio.

3. RESULTS

3.1 Seedling Establishment

Seedling establishment of *Lunaria annua* L. includes number of days required by seeds to 50% germination (%) and Germination percentage. The results of the present study regarding days to 50% germination & germination percentage presented in the Table 1 shows statistically significant differences among different treatments but the seed priming with KNO₃ improved the stand establishment of Honesty (*Lunaria annua* L.) grown in growth chamber. Honesty (*Lunaria annua* L.) seeds primed with 1% KNO₃ had maximum emergence rate (76.3 %), so they were showing better performances when compared to the other treatments. Minimum germination (40%) was observed in non treated I.e. T₁ (control) seeds of Honesty (*Lunaria annua* L.), while, in the case of days to 50% germination, the maximum number of days (14.6) was observed in non treated seeds of *Lunaria annua* L. The seeds treated with 1% KNO₃ germinated earlier than all other treatments.

3.2 Seedling Vigor

Statistical analysis of data about seedling vigor revealed that all priming treatments significantly improved the seedling shoot length in Honesty (*Lunaria annua* L.), whereas the maximum seedling root and shoot length was achieved in Lunaria seed primed with 1% (6.9 & 7.1 cm), followed by 2% KNO₃ solution (6.4 & 6.3 cm), whereas the lowest values for seedling root length and shoot length was observed in control (3.5 & 3 cm), plants raised from seeds treated with 1% KNO₃ showed higher values for seedling fresh weight (116.5 mg) and dry weight (25.7 mg) respectively as compared to other treatments (Table 2).

Table 1. Effect of different priming treatments on days to 50% germination, germination percentage, seedling fresh weight & seedling dry weight of *Lunaria annua* L.

Treatment combination	Days to 50% germination	Germination percentage	Seedling Fresh weight	Seedling dry weight
T1: Control	14.6	40.0	102.8	19.1
T2: CaCl ₂ (1 %)	12.9	56.2	108.6	21.2
T3: CaCl ₂ (2 %)	11.7	58.3	110.0	22.9
T4: CaCl ₂ (3 %)	11.6	61.1	114.2	24.6
T5: CaCl ₂ (4 %)	12.8	54.0	109.0	22.7
T6: KNO ₃ (1 %)	10.0	72.4	116.5	25.7
T7: KNO ₃ (2 %)	11.0	69.3	114.6	25.6
T8: KNO ₃ (3 %)	11.3	68.1	111.9	24.3
T9: KNO ₃ (4 %)	12.3	67.3	112.3	24.1
C.D.	1.39	4.09	7.48	2.92
SE(m)	0.46	1.36	2.50	0.97

Table2. Effect of seed priming treatments on Seedling dry weight, root length, shoot length, Seedling vigour index and root/shoot ratio

Treatment combination	Root length (cm)	Shoot length (cm)	Seedling vigour index	Root/shoot ratio
T1: Control	3.5	3.0	332.4	0.33
T2: CaCl₂ (1 %)	5.4	3.9	419.1	1.02
T3: CaCl₂ (2 %)	5.6	4.0	516.6	0.59
T4: CaCl₂ (3 %)	6.1	4.1	550.4	0.73
T5: CaCl₂ (4 %)	5.7	4.3	510.0	0.91
T6: KNO₃ (1 %)	6.9	7.1	670.0	1.09
T7: KNO₃ (2 %)	6.4	6.3	636.6	1.01
T8: KNO₃ (3 %)	5.8	4.7	590.3	1.02
T9: KNO₃ (4 %)	5.7	4.4	603.3	0.82
C.D	1.19	0.89	128.3	0.36
SE(m)	0.40	0.29	42.86	0.12

Figure1. Graphical representation of effect of different priming treatments on various seedling establishment parameters





Figure2. Graphical representation of effect of different priming treatments on various seedling vigor parameters



4. DISCUSSION

4.1 Seedling establishment

Honesty (*Lunaria annua* L.) seed priming with different concentrations of KNO₃ and CaCl₂ affected the emergence of seedling and the speed of seed germination. Major events in other literature on priming includes metabolic changes, such as repair of DNA and increases in the biosynthesis of RNA (Bray 2017), and enhancement in the respiration process of seed (Singh *et al.*, 2013). This indicates that the time of seed imbibition is very important for seed priming. For the study of seed priming of Honesty (*Lunaria annua* L.) with different levels of KNO₃ and CaCl₂, it is important to know about the emergence percentage and emergence time.

The data shown in Table 1 indicated that priming of Honesty (*Lunaria annua* L.) seeds with 1% KNO₃ was better than other treatments in terms of days to 50% germination and germination percentage. The mechanism of action of KNO₃ on the improvement of seed germination and/or early growth is far from being completely understood. The complexity about nitrate effects on seed germination and early seedling growth could be due to its dual role as a nutrient and a signaling molecule (Duermeyer *et al.*, 2018). Nitrate stimulation of seed germination is often associated with plant species whose seeds require light for germination (Hilhorst *et al.*, 1986; Footitt *et al.*, 2013). Our study is in correspondence with another study that revealed that the emergence percentage of wheat seeds was decreased when they were primed with >1% KNO₃ (Shafiei *et al.*, 2012). This indicates that KNO₃ concentration above a certain threshold may not be appropriate to boost seed germination. Seed priming with 1% KNO₃ was found useful in terms of emergence percentage in sorghum (Shehzad *et al.*, 2012) and rice (Ruttanaruangboworn 2017). Besides, soybean seed primed with 1% KNO₃ for 1 day enhanced the emergence percentage as compared to non-treated seeds, both in laboratory and field experiments (Mohammadi 2018).


4.2 Seedling Vigor

Seedling vigor is the combined result of the emerged seeds under a wide range of biotic and abiotic stresses. Seedling vigor is not a single measurable entity, but it is a sum of many growth parameters, such as seedling length, seedling fresh weight, and seedling dry weight (ISTA 2015). Maximum vigor was observed when seed priming with 1% KNO₃ was done. Our study is in line

with another study in which seedling vigor of wheat was improved by priming with KNO₃ (Shafiei *et al.*, 2012). Similar results were found in corn when the priming of seed was done with 1% KNO₃ (Hadinezhad *et al.*, 2013). Our findings are similar to other studies, in which the shoot length of watermelon and tomato were increased by the seed priming with KNO₃ (Demir *et al.*, 1999 & Mirabi 2012). Seed priming with 1% KNO₃ improved the vegetative growth of watermelon (Oliveira *et al.*, 2019) and tomato (Vaktabhai *et al.*, 2017), respectively, under salt stress. Seed priming with KNO₃ can cause a significant increase in seedling vigor of the wheat crop as compared to hydro-priming or dry broadcasting (Basra *et al.*, 2006)

5. Conclusion

The performance of honesty is diminished by the poor quality of seed. Therefore, the present study was conducted to improve the quality of honesty seed by priming with different concentrations of KNO₃ & CaCl₂. The results presented in this paper revealed that honesty seeds primed with 1% KNO₃ proved to be successful for improving seedling establishment and vigor. The present study provides the direction towards further molecular investigation related to the seed priming of honesty.



of Agriculture & Biology. 8: 23-28

Afzal I., Aslam N., Mahmood F., Hussain A., Irfan S. (2004). Enhancement of germination and emergence of canola seeds by different priming techniques. *Caderno de Pesquisa Serie Biologia. 16: 19-33.*

Arif M., Jan M T., Marwat K B., Khan M A. (2007). Seed priming improves emergence and yield of soybean. *Pakistan Journal of Botany. 40 (3):1169-1177.*

- Basra SMA, Farooq M, Khaliq A (2003). Comparative study of presowing seed enhancement treatment in indica rice (*Oryza sativa* L.). *Pakistan Journal of Life & Social Science*. 1(1): 5-9.
- Bray. C. M. (2017). Biochemical processes during the osmopriming of seeds. In *Seed Development and Germination*; Jaime Kigel; CRC Press: Boca Raton, FL, USA,
- Demir, I., Van De Venter, H.A. (1999). The effect of priming treatments on the performance of watermelon (*Citrullus lanatus* (Thunb.) Matsum. and Nakai) seeds under temperature and osmotic stress. *Seed Science & Technology*. 27: 871–876.
- Fitter, R., Fitter, A., Blamey, M., 1980. The Wild Flowers of Britain and Northern Europe. William Collins Sons, Glasgow, pp. 90-91.
- Hadinezhad, P., Payamenur, V., Mohamadi, J., Ghaderifar, F. (2013). The effect of priming on seed germination and seedling growth in *Quercus castaneifolia*. *Seed Science & Technology*. 41:121–124.
- Harris D, Pathan AK, Gothkar P, Joshi A, Chivasa W, Nyamudeza P (2001). On-farm seed priming: using participatory methods to revive and refine a key technology. *Agricultural Systems*. 69: 151-164.
- ISTA. International Rules for Seed Testing (2015). *The International Seed Testing Association* (ISTA): Bassersdorf, Switzerland, 1–276.
- Khan AA (1993). Preplant physiological seed conditioning. *Horticultural Reviews*. 13: 131–81.
- McDonald MB. Seed priming (2000). *Seed technology and its biological basis*, 287-325.
- Mehmet D., Okcu K G, Atak M., Cikili Y and Kolsarici O (2006). Seed treatment to overcome salt and drought stress during germination in sunflower (*Helianthus annus* L.). *European Journal of Agronomy*. 3: 291-295.
- Mirabi, E., Hasanabadi, M. (2012). Effect of seed priming on some characteristic of seedling and seed vigor of tomato (*Lycopersicon esculentum*). *Journal of Advanced Laboratory Research in Biology*. 3:237–240.

- Mohammadi, G.R. (2009). The effect of seed priming on plant traits of late-spring seeded soybean (*Glycine max* L.). *American-Eurasian Journal of Agricultural & Environmental Sciences*. **5**: 322–326.
- Oliveira, C.E.D.S., Steiner, F., Zuffo, A.M., Zoz, T., Alves, C.Z., Aguiar, V.C.B. (2019). Seed priming improves the germination and growth rate of melon seedlings under saline stress. *Ciência Rural* **49**.
- Pierik, R.L.M., 1967. Regeneration, vernalization and flowering in *Lunaria annua* L. in vivo and in vitro. *Meded. Landb. Hogesch. Wageningen* 67 (6), 1 -71
- Ruttanaruangboworn, A., Chanprasert, W., Tobunluepop, P., Onwimol, D. (2017). Effect of seed priming with different concentrations of potassium nitrate on the pattern of seed imbibition and germination of rice (*Oryza sativa* L.). *Journal of Integrated Agriculture*. **16**: 605–613.
- Shafiei, A.M.; Ghobadi, M. (2012). The effects of source of priming and post-priming storage duration on seed germination and seedling growth characteristics in wheat (*Triticum aestivum* L.). *Journal of Agriculture Science*. **4**, 256.
- Shehzad, M., Ayub, M., Ahmad, A. U. H., Yaseen, M. (2012). Influence of priming techniques on emergence and seedling growth of forage sorghum (*Sorghum bicolor* L.). *Journal of Animal and Plant Sciences*. **22**:154–158.
- Singh, G., Gill, S.S., Sandhu, K.K. (2013). Improved performance of muskmelon (*Cucumis melo*) seeds with osmoconditioning. *Acta Agrobot*. **52**, 121–137.
- Sivritepe, H O. (2000). The effects of osmotic conditioning treatments on salt tolerance of onion seeds. III. *National symposium on vegetable production*, Isparta, Turkey, pp. 475-481.
- Steel, R G D., Torrie J H., Dickey DA (1997). Principles and procedures of statistics, a biological approach. 3rd Edition. McGraw Hill Book Co., Inc., NY, USA, pp. 352-358
- Vaktabhai, C.K., Kumar, S. (2017) Seedling invigouration by halo priming in tomato against salt stress. *Journal of Pharmacology and Phytochemistry*., **6**, 716–722.

- Yang, Q. H, Wei X., Zeng X. L, Ye W. H., Yin X. J., ZhangMing W. (2008). Seed biology and germination ecophysiology of *Camellia nitidissima*. *Forest Ecology and Management*. 255(1):113-118.
- Duermeyer, L.; Khodapanahi, E.; Yan, D.; Krapp, A.; Rothstein, S.J.; Nambara, E. (2018). Regulation of seed dormancy and germination by nitrate. *Seed Science Research*. 28, 150–157.
- Hilhorst, H.W.M.; Smitt, A.I.; Karssen, C.M. (1986). Gibberellin-biosynthesis and -sensitivity mediated stimulation of seed germination of *Sisymbrium officinale* by red light and nitrate. *Plant Physioly*. 67, 285–290.
- Footitt, S.; Huang, Z.; Clay, H.A.; Mead, A.; Finch-Savage, W.E. (2013). Temperature, light and nitrate sensing coordinate Arabidopsis seed dormancy cycling, resulting in winter and summer annual phenotypes. *Plant Journal*, 74, 1003–1015.

Figure1. Graphical representation of effect of different priming treatments on various seedling establishment parameters





Figure2. Graphical representation of effect of different priming treatments on various seedling vigor parameters






Fig.3 a) Primed honesty seed in Petri-plates b) 50% seed germination c, d) measuring different seedling parameters