

Original Research Article

<https://doi.org/10.20546/ijcmas.2018.710.230>

Effect of Time of Pruning on Fruit Production and Berry Quality of Grapes (*Vitis vinifera*) under Bankura Condition of West Bengal

Ranjit Pal^{1*} and S.N. Ghosh²

¹Department of Fruit Science, Rani Lakshmi Bai Central Agricultural University, Jhansi, Utter Pradesh Pin-284003, India

²Department of Fruits and Orchard Management, Faculty of Horticulture, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia, West Bengal, Pin-741252, India

*Corresponding author

ABSTRACT

In India, viticulture is one of the most important horticultural industries in views of its area, production, productivity, value addition and employment creation in both rural and urban areas. The grape in West Bengal is now a new fruit crop for utilizing the up and medium fallow lands in western part of the state. In this part, the soil is red and laterite and acidic in nature and this area received very low rainfall and experience a long dry spell during month of January to May. In such climatic condition pruning is often considered as emergency practice. Correct time and proper pruning very important for productive and healthy over the life of grapevine. As there was no attempt was made to find out proper time of pruning on grape vines under this climatic condition. The aim of this research was to standardize the proper time of pruning on cultivar Arka Trishna under this condition. The investigation was therefore under taken in this direction. The study was conducted at Taldangra in Bankura district of West Bengal (India) during 2012 and 2014. The vine was pruned on four different dates i.e. 20th January, 30th January, 10th February and 20th February. The experiment was laid out in a Randomized Block Design with six replications. The vines were trained on 'Y'-trellis system. The results indicated that maximum yield and yield attributes character per vine was recorded when vines were pruned on 30th January. The berry quality was increased when vines were pruned on 20th February. On the basis of present findings, it can be concluded that 30th January appeared to be the appropriate time for pruning the vines under Bankura conditions with regard to yield attributes and quality.

Keywords

Grape, Arka Trishna, Pruning, Yield, Quality

Article Info

Accepted:
15 September 2018
Available Online:
10 October 2018

Introduction

In India, viticulture is one of the most important horticultural industries in views of its area, production, productivity, value addition and employment creation in both rural and urban areas. Traditionally, grapes are

important fruit crops of temperate region of the world, but in India, grape is grown under temperate, subtropical and tropical climatic condition (Chadha, 2008). The area under grape cultivation is increasing day by day in India. The grape in West Bengal is now a new fruit crop for utilizing the up and medium

fallow lands in western part of the state. In this part, the soil is red and laterite and acidic in nature and this area received very low rainfall and experience a long dry spell during month of January to May. In grape cultivation, the most important and expensive cultural operation is pruning and it plays an important role in grape which affects yield and quality of grapes (Chadha and Shikhamany, 1999). In such climatic condition pruning is often considered as emergency practices. Correct time and proper pruning very important for productive and healthy over the life of grapevine. As there was no attempt was made to find out proper time of pruning on Arka Trishna cultivar of grapes under this climatic condition. The cultivar Arka Trishna is hybrid between Bangalore Blue x Convent Large Black. It is very high yield potential, less susceptible to insects and diseases (Radha and Mathew, 2007). It is possible to take two crops in a year in Bangalore condition but in West Bengal condition only one crop is possible. The aim of this research was to standardize the proper time of pruning on cultivar Arka Trishna under this climatic condition.

Materials and Methods

The investigation was carried out at the Horticultural Research and Development Farm, Govt. of West Bengal, Taldandra district Bankura during 2012-2014.

The site is situated at 23° 01'60" N latitude and 87°06'00 'E longitude with an altitude of 88.0 m above mean sea level. The site is sub-tropical with little rainfall during summer. The average precipitation (June to October) in the area about 1100 mm. The maximum and minimum temperature during summer months varies between 40°C and 24°C respectively while during winter it ranges between 25.5°C and 11.2°C. The soil of the orchard was laterite having pH 6.5, Organic Carbon 0.55%.

Available (natural) soil content of nitrogen (N), phosphorus (P₂O₅) and potassium (K₂O) was 310.30, 40.27 and 155.73 kg per hectare respectively.

The experiment was laid out in Randomized Block Design with four treatments with six replications. The cultivar Arka Trishna grapes with four pruning dates, i.e., 20th January, 30th January, 10th February and 20th February are details of the experiments. The vines were trained on 'Y' trellis system and planting distance at 2m x 3m. All vines were kept under uniform cultural practices. Vine yield and physical characteristics of fruits were recorded when the fruits were ready for harvest. Incidence of thrips was measured by regular field observation. Mortality of renewal shoots were measured after nine months of pruning and expressed in percentage. For chemical analysis of the fruits, the methods were followed as described by A.O.A.C. (1990).

Method of pruning

The method of pruning consisted of removal of not only the past season's shoots at the level indicated but also the removal of unwanted old woods, dried, dead shoots and unthrifty growth during pruning dormant condition.

Time of pruning

The cultivar Arka Trishna grapes with four pruning dates, i.e., 20th January, 30th January, 10th February and 20th February and fruits were harvested during the month of May – June.

Statistical analysis

The data were analyzed statistically and test of significance were done by following the statistical method RBD as described by Pance

and Sukhatme (1985). The significance of the mean difference between the treatments was determined by computing the standard error and critical difference.

Results and Discussion

Yield and yield attribute characteristics of grapes

The yield attributes characters of fruits have been presented in Table 1 and 2. Yield per vine was significantly varied due to different date of pruning during period of investigation. The average yield per vine was highest when vines were pruned on 30th January (3.36 kg) followed by 10th February (2.95 kg) and lowest was recorded when vines were pruned on 20th January (1.79 kg).

Maximum number of bunches per vine was recorded when vines were pruned on 30th January (32.83) closely followed by 10th February (30.67).

Highest bunch weight (102.40 g), longest bunch length (12.5 cm) and bunch breadth (8.2 cm) was recorded when vines were pruned on 30th January. Highest yield and yield attributes characters was recorded from 30th January pruned vines may be due to higher fruitfulness percentage of the spur and increasing size of bunches after berry set. Increasing in fruit yield with 30th January pruning was noted by Ghosh *et al.*, (2012) in cultivar Arka Neelamani under Jhargram condition. Chadha and Mand (1969) found that the yield was great when vines were pruned from 26th January to 5th February under North India condition.

The data presented in Table 3 and 4 showed that highest ten berries weight (33.07 g), average berry size (length x breadth) was recorded when vines were pruned on 10th February. Highest juice content was measured when vines were pruned on 10th February (78.88 %) closely followed by 20th February (78.01 %).

Table.1 Effect of different date of pruning on yield and number of bunches of grapes cv. Arka Trishna

Treatments	Yield per vine (Kg)	Number of bunches per vine
Pruning on 20 th January	1.79	20.16
Pruning on 30 th January	3.36	32.83
Pruning on 10 th February	2.95	30.67
Pruning on 20 th February	1.82	21.16
CD at 5%	0.51	7.40

Table.2 Effect of different date of pruning on bunch weight, length and breadth of grapes cv. Arka Trishna

Treatments	Bunch weight (g)	Bunch length (cm)	Bunch breadth (cm)
Pruning on 20 th January	89.21	10.8	6.2
Pruning on 30 th January	102.40	12.5	8.2
Pruning on 10 th February	96.25	10.3	6.5
Pruning on 20 th February	86.36	10.2	6.2
CD at 5%	6.91	0.98	1.02

Table.3 Effect of different date of pruning on ten berries weight and average berry length of grapes cv. Arka Trishna

Treatments	10 berries weight (g)	Average berry length (cm)
Pruning on 20 th January	29.18	2.82
Pruning on 30 th January	31.57	3.05
Pruning on 10 th February	33.07	3.17
Pruning on 20 th February	32.02	3.07
CD at 5%	2.08	0.17

Table.4 Effect of different date of pruning on average berries breadth and juice content of grapes cv. Arka Trishna

Treatments	Average berry breadth (cm)	Juice content (%)
Pruning on 20 th January	2.82	76.89
Pruning on 30 th January	3.05	76.64
Pruning on 10 th February	3.17	78.88
Pruning on 20 th February	3.07	78.01
CD at 5%	0.17	0.94

Table.5 Effect of different date of pruning on mortality of shoots and incidence of thrips of grapes cv. Arka Trishna

Treatments	Mortality percentage of renewal shoots	Incidence of thrips
Pruning on 20 th January	14.20	Negligible
Pruning on 30 th January	12.50	Negligible
Pruning on 10 th February	11.69	Negligible
Pruning on 20 th February	11.41	Medium
CD at 5%	2.64	-

Table.6 Effect of different date of pruning on date of panicle emergence and harvesting period of grapes cv. Arka Trishna

Treatments	Date of panicle emergence	Date of harvest	
		1 st harvest	Last harvest
Pruning on 20 th January	8 th February	18 th May	31 st May
Pruning on 30 th January	19 th February	23 rd May	31 st May
Pruning on 10 th February	28 th February	28 th May	7 rd June
Pruning on 20 th February	9 th March	30 th May	10 th June
CD at 5%	-	-	-

Table.7 Effect of different date of pruning on TSS, Acidity and TSS/acidity ratio of grapes cv. Arka Trishna

Treatments	TSS (° Brix)	Acidity (%)	TSS/acid ratio
Pruning on 20 th January	18.7	0.57	32.80
Pruning on 30 th January	19.6	0.46	42.60
Pruning on 10 th February	20.9	0.45	46.45
Pruning on 20 th February	22.2	0.45	49.34
CD at 5%	1.10	0.14	8.63

Table.8 Effect of different date of pruning on Total sugar, reducing sugar and Ascorbic acid contain of grapes cv. Arka Trishna

Treatments	Total sugar (%)	Reducing sugar (%)	Ascorbic acid (mg/100 ml juice)
Pruning on 20 th January	15.45	13.66	4.00
Pruning on 30 th January	14.76	13.54	4.45
Pruning on 10 th February	15.75	14.25	3.50
Pruning on 20 th February	16.15	14.80	3.86
CD at 5%	0.87	0.92	0.94

It was evident from the data presented in Table 5 and 6 observed that pruning on 20th January recorded increasing mortality percentage of renewal shoots. It was further noted that delay in pruning (20th February) resulted more incidence of thrips, at present is considered as serious ones. The data showed that pruning on 30th January resulted in completion of harvesting process within the month of May i.e., before on set of rainy season while pruning on 20th February, harvesting time extended up to 10th June when the rainy season had already started.

Quality characteristics of grapes

The chemical composition of berries of different treatments has been presented in Table 7 and 8. The data showed that different date of pruning exhibit significant variation with respect to total soluble solids of Arka Trishna grapes. Highest total soluble solids content of the berry was measured when vines were pruned on 20th February (22.2 ° brix)

and lowest when vines were pruned on 30th January (18.7 ° brix). This kind of similar phenomena was reported by earlier worker Ghosh *et al.*, (2012) reported that highest TSS content in the berries when vines were pruned on 20th January in Arka Neelamani cultivar under Jhargram condition. The highest acidity percentage (0.57 %) of berry was measured when vines were pruned on 20th January and lowest when vines were pruned on 10th February and 20th February (0.45 %).

Most important quality character TSS/acid ratio was highest when vines were pruned on 20th February (49.34). The total sugar (16.15 %) and reducing sugar (14.80 %) content was highest when vines were pruned on 20th February. Increasing TSS content and decreasing acid content of berry under this date of pruning may be due to favorable temperature and relative humidity during fruit maturity period. Ascorbic acid content was recorded highest when vines were pruned on 30th January (4.45 mg/100 ml juice).

Considering the above study in view it can be concluded that proper time of pruning is very essential for improving production of grapes under this climatic condition. On the basis of present findings, it can be concluded that 30th January appeared to be the appropriate time for pruning the vines under Bankura conditions with regard to yield attributes and quality.

References

- A.O.A.C. 1990. Official method of analysis. Association of analytical chemists. 15th edn. Washington, D.C.
- Chadha, K, L, 2008. Indian viticulture scenario. *Acta Horticulture*. 785: 59-68.
- Chadha, K.L., and Shikhamany, S.D. 1999. The Grapes. Improvement, Production and Post-Harvest Management. Malhotra Publishing House, New Delhi.
- Chadha, K.L., Mand, A.S. 1969. Effect of time and severity of pruning on maturity, yield and quality of Anab-e-Shahi grape. *Journal of Research*. Punjab Agricultural University. 6: 808-20.
- Ghosh, S.N., Bera, B, Roy, S and Kundu, A. 2012. Adaptation and commercialization of viticulture in West Bengal - a new area in India. *Acta Horticulture*. 931: 389-399.
- Panse, V.G., and Sukhatme, P.V. 1985. Design of Experiments. VIII Randomized Blocks and Latin Square. In: Statistical methodology for Agricultural workers. Indian Council of Agricultural Research, New Delhi.
- Radha, T., and Mathew, L. 2007. Fruit Crops: Sub-Tropical Fruits. New India Publishing Agency. Pritam Pura, New Delhi.

How to cite this article:

Ranjit Pal and Ghosh, S.N. 2018. Effect of Time of Pruning on Fruit Production and Berry Quality of Grapes (*Vitis vinifera*) under Bankura Condition of West Bengal. *Int.J.Curr.Microbiol.App.Sci*. 7(10): 1994-1999. doi: <https://doi.org/10.20546/ijemas.2018.710.230>