



Published in final edited form as:

*J Biol Rhythms*. 2019 April ; 34(2): 195–204. doi:10.1177/0748730419830013.

## Effect of white light devoid of “cyan” spectrum radiation on nighttime melatonin suppression over a 1-h exposure duration

Rohan Nagare, Mark S. Rea, Barbara Plitnick, Mariana G. Figueiro\*

Lighting Research Center, Rensselaer Polytechnic Institute, Troy, NY

### Abstract

The intrinsically photosensitive retinal ganglion cells are the main conduit of the light signal emanating from the retina to the biological clock located in the suprachiasmatic nuclei of the hypothalamus. Lighting manufacturers are developing white light sources that are devoid of wavelengths around 480 nm (“cyan-gap”) to reduce their impact on the circadian system. The present study was designed to investigate whether exposure to a “cyan-gap,” 3000 K white light source spectrally tuned to reduce radiant power between 475 and 495 nm (reducing stimulation of the melanopsin-containing photoreceptor) would suppress melatonin less than a conventional 3000 K light source.

The study’s 2 phases employed a within-subjects experimental design involving the same 16 adult participants. In Phase 1, participants were exposed for 1 h to 3 experimental conditions over the course of 3 consecutive weeks: (1) dim light control (<5 lux); (2) 800 lux at the eyes of a 3000 K light source; and (3) 800 lux at the eyes of a 3000 K “cyan-gap” modified (3000 K mod) light source. The same protocol was repeated in Phase 2, but light levels were reduced to 400 lux at the eyes. As hypothesized, there were significant main effects of light level ( $F_{1,12} = 9.1$ ,  $p < 0.05$ ,  $\eta_p^2 = 0.43$ ) and exposure duration ( $F_{1,12} = 47.7$ ,  $p < 0.05$ ,  $\eta_p^2 = 0.80$ ), but there was no significant main effect of spectrum ( $F_{1,12} = 0.16$ ,  $p > 0.05$ ,  $\eta_p^2 = 0.01$ ). There were no significant interactions with spectrum. Contrary to our model predictions, our results showed that short-term exposures to “cyan-gap” light sources suppressed melatonin similarly to conventional light sources of the same CCT and photopic illuminance at the eyes.

### Keywords

Light at night; circadian; melatonin suppression; spectrum; cyan-gap lighting; cyan-deficient lighting

### INTRODUCTION

The human circadian clock “free runs” (i.e., cycles independently, without external cues) at a period that is slightly greater than 24 h. The 24-h, light–dark pattern reaching the retina is the major synchronizer of the circadian clock to one’s local position on Earth (Czeisler et al., 1989; Zeitzer et al., 2000a; Khalsa et al., 2003). A person whose circadian system is

\*To whom all correspondence should be addressed: Mariana Figueiro, Lighting Research Center, 21 Union Street, Troy, NY 12180-3352. [figuem@rpi.edu](mailto:figuem@rpi.edu).

synchronized to local time is said to be entrained. Asynchrony or misalignment between the endogenous biological clock and the external light–dark cycle, as is likely for shift-workers or people experiencing rapid transmeridian travel, leads to circadian disruption. It is important to address disruption of circadian rhythms, as it has been associated with a series of decrements such as poor sleep, depression, increased risk for metabolic and cardiovascular disease, and a higher incidence of cancer in workers whose schedules involve evening and night shifts (Davis and Mirick, 2006; Maemura et al., 2007; Stevens et al., 2007; Young and Bray, 2007; Costa et al., 2013; Karatsoreos, 2014; Potter et al., 2016).

Nocturnal melatonin suppression is a primary outcome for characterizing the impact of light on the biological clock (Brainard et al., 2001; Thapan et al., 2001). Published psychophysical studies of nocturnal melatonin suppression using narrowband light sources have shown that the human circadian system is maximally sensitive to short-wavelength (“blue”) light, peaking close to 460 nm (Brainard et al., 2001; Thapan et al., 2001). In regard to nighttime exposures to broadband sources, past studies have demonstrated that filtering out short-wavelength light < 530 nm from polychromatic white light sources limits the suppression of melatonin secretion in humans (Kayumov et al., 2005; Sasseville et al., 2006; Van de Werken et al., 2013; Rahman et al., 2017). Studies by Rahman et al. (2011) have further shown that filtering out short wavelength energy < 480 nm from a broadband light source prevents nocturnal melatonin suppression; however, significant suppression was recorded for the same light source when energy at only wavelengths < 460 nm was filtered out.

It has been demonstrated that the intrinsically photosensitive retinal ganglion cells (ipRGCs) are the main conduit of the light signal emanating from the retina to the biological clock located in the suprachiasmatic nuclei (SCN) of the hypothalamus. The ipRGCs express melanopsin, which makes them intrinsically photosensitive, and *in situ* have a peak spectral sensitivity close to 480 nm (Bailes and Lucas, 2013). The peak spectral sensitivity to short-wavelength light observed for acute melatonin suppression at 460 nm cannot be exclusively attributed to this single type of photoreceptor. Rather, the ipRGCs receive input from the more distal photoreceptors (i.e., rods and cones) in the process of circadian phototransduction (Foster et al., 1991; Ruby et al., 2002; Hattar et al., 2003).

Nevertheless, lighting manufacturers have begun developing white light sources that are devoid of wavelengths around 480 nm in an effort to provide white light while avoiding circadian-effective light, categorizing these newly developed light sources as “cyan-deficient” or “cyan-gap”. A “cyan-gap” spectrum type has, for example, been employed in a recent study by Souman et al. (2018), who exposed 15 participants to 3 h of the same amount of photopic illuminance of white light with either low or high radiant power between 450 nm and 500 nm, which is close to the maximal sensitivity of the ipRGCs. Their results showed that melatonin suppression, calculated using area under the curve for 3 h, was significantly reduced for the “cyan-gap” source relative to the white light source without selective attenuation of short wavelengths between 450 nm and 500 nm. In another recent nighttime study involving 11 participants (aged 21–30 years), Allen et al. (2018) reported an increase in subjective sleepiness and salivary melatonin levels following exposure to a

customized visual display unit (VDU) providing lower “melanopic stimulation” compared to a standard VDU, without changing the visual appearance of the display.

In the wake of these developments, the present study was designed to test the effectiveness of a newly developed “cyan-gap” LED light source on nighttime melatonin suppression. Unlike the white light source used in the Souman et al. studies (2700K), which had enhanced radiant power at very short wavelengths (i.e., 410 nm), the light source used in the current study (3000 K mod) had enhanced radiant power at 450 nm and reduced radiant power between around 475 nm and 495 nm.

To predict the impact of this newly developed light source on melatonin suppression, this study employed the Rea et al. model of human circadian phototransduction (Rea et al., 2005; Rea et al., 2012; Rea and Figueiro, 2018), which is based in part on the light-induced nocturnal melatonin suppression data from Brainard et al. (2001) and Thapan et al. (2001). The model is constrained by the fundamental knowledge of retinal neurophysiology and neuroanatomy and provides a framework for depicting how classical photoreceptors (i.e., rods and cones) provide input to the ipRGCs. Mathematically, for any light source, the model converts the spectral irradiance at the eyes into circadian light ( $CL_A$ ), reflecting the spectral sensitivity of the circadian system, and then transforms it into a circadian stimulus (CS) value reflecting the absolute sensitivity of the circadian system. Thus, CS is a measure of the effectiveness of the retinal light stimulus for the human circadian system from threshold ( $CS \approx 0.1$ ) to saturation ( $CS \approx 0.7$ ) after a 1-h exposure. Using the model calculations, it was hypothesized that the “cyan-gap” light source would induce a small ( $\approx 8\%$  at both light levels) but significant reduction in melatonin suppression compared to a conventional, 3000 K light source. It was also hypothesized that melatonin suppression would be significantly greater at higher light levels and longer exposure durations.

## MATERIALS AND METHODS

### Participant Selection

Sixteen adult participants (12 females, 4 males; mean  $\pm$  standard deviation [SD] age = 37.8 years  $\pm$  13.9) were recruited for the study via personal referrals, word of mouth, Rensselaer Polytechnic Institute’s social media platform, and lists of participants from previous studies. All participants completed the protocol. Depicting the clock hour of mid-sleep times on free days, the participants’ mean  $\pm$  SD Munich Chronotype Questionnaire score (Roenneberg et al., 2003) was  $2.94 \pm 1.29$ , indicating that they were neither extreme larks (early persons) nor extreme owls (late persons). The study was conducted in 2 phases (see Experimental Conditions) of 3 weeks each, with Phase 1 running from 10 February 2017 to 3 March 2017 and Phase 2 running from 7 April 2017 to 5 May 2017. All participants were pre-screened for major health problems such as bipolar disorder, seasonal depression, cardiovascular disease, diabetes, and high blood pressure. Participants were excluded from the experiment if they were taking any over-the-counter melatonin or prescription medications such as blood pressure medicine, antidepressants, sleep medicine, or beta-blockers. Participants’ reporting eye diseases such as cataracts or glaucoma were also excluded, as were participants who were identified by the study team as having color vision deficiency according to Ishihara

color blindness tests (Ishihara, 1960). No participants were scheduled to undertake transmeridian travel over the course of each study phase.

In order to maintain a stable circadian melatonin rhythm, all participants were required to follow a consistent sleep–wake schedule during the week leading up to each experimental session, with bedtimes no later than 23:00 and wake times no later than 07:30. Participants were also requested to refrain from caffeine consumption 12 h prior to the beginning of each experimental session. Given that all of the participants were full-time workers, their sleep schedules were presumed to be regular throughout the duration of the study.

This study conformed to the Code of Federal Regulations (CFR) document Protection of Human Subjects, 45 CFR 46, (2016), and international ethical standards (Portaluppi et al., 2010). It was reviewed, approved, and monitored by Rensselaer Polytechnic Institute’s Institutional Review Board. Informed consent was obtained from all study participants.

### Experimental Conditions

The study’s 2 phases employed a within-subjects experimental design involving the same 16 participants to minimize the effects of potential confounding factors such as age and individual differences. During each phase, the participants reported to the Lighting Research Center’s laboratory in Troy, NY, on 3 nights that were separated by at least 1 week to allow for a washout period between the conditions. The participants were randomly divided into 3 groups (two groups of 5 and one group of 6 participants), and each group was exposed to a single experimental condition per session in a planned, counter-balanced order to avoid order effects. Each experimental session began with a 30-min dim light adaptation period that preceded presentation of the experimental conditions.

For each study phase, participants were exposed to 3 experimental conditions comprised of one dim light control condition and 2 lighting interventions derived from spectrally distinct white light sources: 3000 K and 3000 K “cyan-gap” modified (3000 K mod). On random allocation, the study phases delivered either high (Phase 1) or low (Phase 2) levels of the 2 intervention spectra at the participants’ eyes during separate sessions. The higher light level employed in Phase 1 (approximately 800 lux at the eyes), although not typically found in the built environment, was selected to maximize the signal-to-noise ratio and, therefore, reliably detect melatonin suppression from all participants. The lower light level employed in Phase 2 (approximately 400 lux at the eyes) was also selected because it would still maximize the signal-to-noise ratio and minimize the impact of individual differences in the response while being representative of light levels one can potentially experience in healthcare and commercial applications. These higher light levels would also minimize the impact of photic history on the experimental results. The dim-light control condition was employed to collect participants’ baseline melatonin levels for analysis.

Spectral irradiance measurements for the lighting interventions were performed using a spectrometer (Model USB650, Ocean Optics, Winter Park, FL, USA), and their relative spectral power distributions are shown in Figure 1. The photometric characteristics (i.e., illuminance,  $CL_A$ , melanopic illuminance, and CS) of the light stimulus delivered, based on the Daysimeter data (see Recording Equipment) and calculated following the Rea et al. (CS)

(Rea et al., 2005; Rea et al., 2012; Rea and Figueiro, 2018) and Enezi et al. (2011) (melanopic illuminance) models, are provided in Table 1. The corresponding  $\alpha$ -opic irradiances of the lighting interventions, following the SI-compliant approach recommended by the Commission Internationale de l'Éclairage (CIE) (2015) are provided in Table 2.

### Lighting Apparatus

The lighting interventions were delivered via custom-built luminaires that were mounted on laboratory tables. Each luminaire housed an LED light bar composed of 48 low-power white LEDs (Lumileds, San Jose, CA, USA) operating in a constant current mode (200 mA max). White, spectrally neutral diffusers were installed on the luminaires to eliminate potential discomfort glare and provide a uniform light distribution. The luminaires were mounted on adjustable stands, and their height and distance from participants' eyes were individually adjusted for each participant to ensure delivery of the desired stimulus of 800 lux at the eyes for Phase 1, and 400 lux at the eyes for Phase 2 (Figure 2). For a fixed width of 18 in. and a standard viewing distance of 30 in., the luminaires delivered an image that was approximately 33° in width at the participants' eyes. Consistency of the light stimulus throughout the 60-min exposure was ensured through continual monitoring of the light levels through spot measurements taken with an illuminance meter every 10 min. The luminaires were not employed for the dim light control condition.

### Recording Equipment

The study did not employ chin rests to control the participants' gaze angle. Potential variability within the individual spectral and absolute characteristics of the stimulus delivered to the participants was addressed by continual data collection from the Daysimeter (Model 12, Lighting Research Center, Troy, NY USA). The Daysimeter was positioned near each participant's eyes, mounted on lensless eyeglasses frames, and recorded the light stimulus at 30-s intervals throughout the 60-min light exposure.

### Study Protocol

Participants arrived at the laboratory at 23:30 and remained in dim light (< 5 lux at the eyes) for 30 min, followed by a 60-min exposure to one of the 3 experimental conditions (Figure 3). During the experiment, the participants were instructed to not remove or block the Daysimeter's sensor. The participants were not required to directly look at the light emitting fixtures but were instructed to face them to ensure minimum variability with respect to the target stimulus (see Figure 2). Participants were free to operate their personal electronic devices (i.e., computers, tablets, cell phones, etc.) and were required to perform a similar task (e.g., browse the Internet, watch a video or movie, read an e-book, etc.) on all study nights. All displays were covered with orange-tinted media (Roscolux #21 golden amber, Rosco Laboratories, Stamford, CT, USA) that filtered out radiation < 525 nm to prevent participants from receiving additional circadian-effective light from their self-luminous devices. In a previous study, photometric measurements of the stimulus emitted by similarly filtered electronic displays revealed increased light levels of < 5 lux, which translated to an additional CS of < 0.001 (Rea and Figueiro, 2018). During each session, the participants refrained from consuming any food and were allotted a 10-min window to drink water, following each saliva sample time. Periodic visual monitoring was carried out to ensure

compliance with the experimental protocol and confirm that none of the participants closed their eyes.

Three saliva samples were collected from each participant over the course of each experimental session. The first sample was taken immediately before the beginning of the lighting intervention after a 30-min dim light exposure and the 2 remaining samples were taken thereafter at 30-min intervals. At 01:00, after the final saliva sample was collected, the participants were released from the laboratory. Saliva samples (1 ml) were collected using the Salivette system (Sarstedt, Nümbrecht, DE) wherein the participant chews on a plain cotton cylinder for 1–2 min (not timed individually) that was immediately placed in a test tube, centrifuged for 5 min at 1000 g, and frozen (–20 ° C). The frozen samples from each session were assayed in a single batch using melatonin radioimmunoassay kits (Direct Melatonin RIA, ALPCO, Salem, NH, USA). The sensitivity of the saliva sample assay was reported to be 0.3 pg/mL and the intra- and inter-assay coefficients of variability were 11–14% and 14–17%, respectively.

Melatonin data were acquired from all participants for the low light level (Phase 2). For the high light level (Phase 1), melatonin data could not be processed from one participant for the “3000 K mod” condition (S2), one participant for the 3000 K condition (S2), and from one participant for both the 3000 K and “3000 K mod” conditions (all samples), due to a lack of sufficient saliva within the samples collected.

## Data Analysis

Circadian-effective light levels (Rea et al., 2010) were calculated from the Daysimeter data and transformed to the CS metric following the Rea et al. model (Rea et al., 2005; Rea et al., 2012; Rea and Figueiro, 2018). Melatonin suppression was determined by comparing melatonin levels collected during the dim light control night to those collected at the corresponding times on each lighting intervention night (see Figure 3). For each session, melatonin concentrations for S2 and S3 were first normalized to S1, and the melatonin suppression at each time was then calculated using Equation 1.

$$\left(1 - \frac{M_n}{M_d}\right) \times 100 \quad (1)$$

where  $M_n$  is the normalized melatonin concentration at each time on the respective intervention nights and  $M_d$  is the normalized melatonin concentration at each corresponding time on the dim light control night.

Primary statistical analyses were performed using repeated measures analysis of variance (ANOVA) with spectrum, light level, and exposure duration employed as within-subject factors. Baseline melatonin levels across the 2 study phases with 3 experimental conditions each were also subjected to repeated measures ANOVA. The ANOVA was performed with SPSS statistical software (SPSS version 25, IBM, Armonk, NY, USA). Further evaluation for the main effects and interactions was performed using *post hoc*, 1-sample *t*-tests. The test results were considered statistically significant if the *p* value was <0.05.

## RESULTS

### Absolute melatonin levels

The ANOVA revealed that the absolute baseline melatonin levels recorded at the beginning of each study night (S1) were not significantly different across the 2 study phases ( $F_{1,14} = 1.24$ ,  $p = 0.28$ ,  $\eta_p^2 = 0.08$ ) and the 3 experimental conditions ( $F_{2,28} = 1.51$ ,  $p = 0.24$ ,  $\eta_p^2 = 0.10$ ), suggesting that the light treatment was delivered at a similar circadian phase for all participants on all study nights (Table 3). Figure 4 shows the absolute melatonin data for the dim light control and the 2 intervention spectra for both light levels at baseline and after 30-min and 60-min exposures.

### Effect of lighting characteristics on melatonin suppression

The ANOVA revealed significant main effects of light level ( $F_{1,12} = 9.1$ ,  $p < 0.05$ ,  $\eta_p^2 = 0.43$ ) and exposure duration ( $F_{1,12} = 47.7$ ,  $p < 0.05$ ,  $\eta_p^2 = 0.80$ ), wherein, as expected, greater melatonin suppression was observed following exposure to higher light levels and longer durations during the participants' biological night (Figure 5).

The ANOVA did not reveal a significant main effect of intervention spectrum ( $F_{1,12} = 0.16$ ,  $p = 0.69$ ,  $\eta_p^2 = 0.01$ ), whereby the mean melatonin suppression between the 2 spectra was not statistically discernible (Figure 6) for both phases of the study light levels. The mean  $\pm$  SD melatonin suppression following a 1-h exposure to the high light level was  $49\% \pm 13$  and  $50\% \pm 13$ , for the 3000 K and 3000 K mod interventions, respectively. The mean  $\pm$  SD melatonin suppression following a 1-h exposure to the low light level was  $34\% \pm 11$  and  $36\% \pm 11$ , for the 3000 K and 3000 K mod interventions, respectively. Statistical analysis did not reveal any outliers within the entire data set, suggesting that the results were not skewed by any particular participant(s). The median (IQR) melatonin suppression following a 1-h exposure to the low light level was 34% (15%) and 32% (17%), for the 3000 K and 3000 K mod interventions, respectively. The median (IQR) melatonin suppression following a 1-h exposure to the high light level was 44% (19%) and 53% (19%), for the 3000 K and 3000 K mod interventions, respectively. No statistically significant interactions between the independent variables were identified ( $p > 0.05$ ).

*Post hoc* statistical analyses showed that melatonin suppression following a 1-h exposure was significantly greater than zero for both the "3000 K mod" (Phase 1:  $t_{14} = 15.0$ ,  $p < 0.05$ ; Phase 2:  $t_{15} = 12.9$ ,  $p < 0.05$ ) and the conventional 3000 K (Phase 1:  $t_{14} = 14.1$ ,  $p < 0.05$ ; Phase 2:  $t_{15} = 12.4$ ,  $p < 0.05$ ) sources. Furthermore, melatonin suppression following exposure to the 3000 K intervention was not significantly different ( $p > 0.05$ ) from the *a priori* CS levels predicted for both the high ( $t_{14} = -0.06$ ,  $p > 0.05$ ) and low ( $t_{15} = 0.07$ ,  $p > 0.05$ ) light levels. However, melatonin suppression was significantly greater than the predicted CS levels for the 3000 K mod intervention at the high ( $t_{14} = 2.59$ ;  $p < 0.05$ ) and low ( $t_{14} = 3.47$ ;  $p < 0.05$ ) light levels, following a 1-h exposure.

## DISCUSSION

The present study was designed to investigate whether exposures to a "cyan-gap" (3000 K mod) white light source that is spectrally tuned to reduce stimulation of the melanopsin-

containing photoreceptor suppressed less melatonin than a conventional 3000 K light source. Contrary to our model predictions and conventional wisdom, the results showed that participants exposed to the “cyan-gap” light source experienced similar levels of melatonin suppression (50% at high light levels, 36% at low light levels) compared to those experienced after exposure to the conventional 3000 K intervention (49% at high light levels, 34% at low light levels; see Tables 1 and 2). The measured levels of light-induced nocturnal melatonin suppression after 1-h exposures closely matched the predicted CS values for the conventional 3000 K light source; however, measured suppression levels for the “cyan-gap” source were statistically equal to those for the conventional source. Therefore, suppression was greater for the “cyan-gap” source than would be expected if the ipRGC, melanopsin-containing photoreceptor were solely responsible for nocturnal melatonin suppression by both light sources.

Although there was no statistical difference between the “cyan-gap” source and the conventional source at either light level, the results are consistent with our hypothesis and with those from many previous studies showing that higher light levels and longer exposure durations result in significantly greater melatonin suppression (Zeitzer et al., 2000b; Chang et al., 2012; Nagare et al., 2018b). The present data, as well as those reported by Nagare et al. (2018b) for light levels commonly experienced in indoor environments with white light sources, suggest that lighting dose (amount  $\times$  duration) is of greater relative importance than spectrum for the same CCT when describing the effect of photic stimulus on the human circadian system.

At first glance, the results of the present study may appear to contradict those of Souman et al. (2018), who demonstrated reduced melatonin suppression with the reduction of spectral power between 450 nm and 500 nm (Souman et al., 2018). Several methodological differences between the 2 studies are noteworthy, however. First, the Souman et al. study used an exposure duration of 3 h rather than the 1 h duration employed here. Importantly, Souman et al. showed no significant differences in melatonin suppression for their “cyan-gap” source and their conventional source after 1-h exposure, consistent with the present study. Second, the melanopsin stimulation ratio of the “high MEF” to “low MEF” light sources from the Souman et al. study was almost twice ( $\approx 3.5$ ) the melanopsin stimulation ratio ( $\approx 1.6$ ) of the 2 light sources employed in the present study. A third key difference is that the absolute levels of irradiance at the eyes provided by Souman et al. were much lower than those employed in the present study. As such, it is possible that differential habituation to the 2 low-light stimuli may have occurred in the Souman et al. study. A recent study by Rahman et al. (2017) also examined melatonin suppression following sustained low-level light exposures ( $\approx 50$  lux at the eyes) to a conventional, 4100 K fluorescent light source (FL) and a “blue-depleted” LED light source (C-LED). Although nocturnal melatonin suppression by the C-LED source was significantly lower than that from the fluorescent source after 8 h of exposure, there was no difference between the 2 sources initially following collection of melatonin data early in the night. Thus, even though the results from Rahman et al. are consistent with the results presented here as well as those from Souman et al., it is nevertheless important to examine the temporal dynamics of light exposures in the future to determine why 2 light stimuli of different spectral composition would have equal nocturnal melatonin suppressing effects for short durations but not for long durations.



A limitation of the present study is that we did not monitor and control participants' photic history leading up to the experimental sessions. Since all participants were employed adults or students with regular schedules, it was presumed that they would receive consistent daily light exposures over the course of the two 3-week study phases. The study's protocol also did not employ chin rests to control the participants' gaze angle, although any potential variability within the individual spectral and absolute characteristics of the stimulus was addressed by continual data collection from the Daysimeter positioned near the participants' eyes. Again, there is no evidence that the absolute light levels were confounded with light source spectrum. Finally, while we acknowledge a sex disparity among the study's participants (75% females), it should be noted that past studies have reported no differences in measures of circadian preference and sleep with respect to sex (Giannotti et al., 2002; Russo et al., 2007).

The findings from the present research should be of particular interest to lighting manufacturers who are interested in producing light sources that can have a differential effect on the circadian system without affecting photopic light levels and CCTs, which are key lighting parameters considered by light specifiers. The results of the present study suggest that for short-term exposures "cyan-gap" light sources affect the human circadian system similarly to conventional light sources of the same CCT and photopic illuminance at the eyes.

## ACKNOWLEDGEMENTS


The present study was funded by LumiLeds, the Light and Health Alliance (Acuity Brands, CREE, Current Powered by GE, Ketra, Osram, Philips, and USAI Lighting) and the NIH Training Program in Alzheimer's Disease Clinical and Translational Research (NIA 5T32AG057464).

## REFERENCES

- Allen AE, Hazelhoff EM, Martial FP, Cajochen C, and Lucas RJ (2018) Exploiting metamerism to regulate the impact of a visual display on alertness and melatonin suppression independent of visual appearance. *Sleep* 8:10.1093/sleep/zsy1100.
- Bailes HJ, and Lucas RJ (2013) Human melanopsin forms a pigment maximally sensitive to blue light ( $\lambda_{max} \approx 479$  nm) supporting activation of Gq/11 and Gi/o signalling cascades. *Proc Biol Sci* 280:20122987. [PubMed: 23554393]
- Brainard GC, Hanifin JP, Greeson JM, Byrne B, Glickman G, Gerner E, and Rollag MD (2001) Action spectrum for melatonin regulation in humans: evidence for a novel circadian photoreceptor. *J Neurosci* 21:6405–6412. [PubMed: 11487664]
- Chang AM, Santhi N, St Hilaire M, Gronfier C, Bradstreet DS, Duffy JF, Lockley SW, Kronauer RE, and Czeisler CA (2012) Human responses to bright light of different durations. *J Physiol* 590:3103–3112. [PubMed: 22526883]
- Commission Internationale de l'Éclairage (2015) Technical Note: Report on the First International Workshop on Circadian and Neurophysiological Photometry, 2013. In, Commission Internationale de l'Éclairage, Vienna, Austria.
- Costa IC, Carvalho HN, and Fernandes L (2013) Aging, circadian rhythms and depressive disorders: a review. *Am J Neurodegener Dis* 2:228–246. [PubMed: 24319642]
- Czeisler CA, Kronauer RE, Allan JS, Duffy JF, Jewett ME, Brown EN, and Ronda JM (1989) Bright light induction of strong (type 0) resetting of the human circadian pacemaker. *Science* 244:1328–1333. [PubMed: 2734611]
- Davis S, and Mirick DK (2006) Circadian disruption, shift work and the risk of cancer: a summary of the evidence and studies in Seattle. *Cancer Causes Control* 17:539–545. [PubMed: 16596308]

- Enezi Ja, Revell V, Brown T, Wynne J, Schlangen L, and Lucas R (2011) A “melanopic” spectral efficiency function predicts the sensitivity of melanopsin photoreceptors to polychromatic lights. *J Biol Rhythms* 26:314–323. [PubMed: 21775290]
- Foster R, Provencio I, Hudson D, Fiske S, De Grip W, and Menaker M (1991) Circadian photoreception in the retinally degenerate mouse (rd/rd). *J Comp Physiol A* 169:39–50. [PubMed: 1941717]
- Giannotti F, Cortesi F, Sebastiani T, and Ottaviano S (2002) Circadian preference, sleep and daytime behaviour in adolescence. *J Sleep Res* 11:191–199. [PubMed: 12220314]
- Hattar S, Lucas RJ, Mrosovsky N, Thompson SH, Douglas RH, Hankins MW, Lem J, Biel M, Hofmann F, Foster RG, and Yau K-W (2003) Melanopsin and rod-cone photoreceptive systems account for all major accessory visual functions in mice. *Nature* 424:76–81. [PubMed: 12808468]
- Ishihara S (1960) Tests for colour-blindness. H. K. Lewis & Co., London.
- Karatsoreos IN (2014) Links between circadian rhythms and psychiatric disease. *Front Behav Neurosci* 8:162. [PubMed: 24834040]
- Kayumov L, Casper RF, Hawa RJ, Perelman B, Chung SA, Sokalsky S, and Shapiro CM (2005) Blocking low-wavelength light prevents nocturnal melatonin suppression with no adverse effect on performance during simulated shift work. *J Clin Endocrinol Metab* 90:2755–2761. [PubMed: 15713707]
- Khalsa SB, Jewett ME, Cajochen C, and Czeisler CA (2003) A phase response curve to single bright light pulses in human subjects. *J Physiol* 549:945–952. [PubMed: 12717008]
- Lucas RJ, Peirson SN, Berson DM, Brown TM, Cooper HM, Czeisler CA, Figueiro MG, Gamlin PD, Lockley SW, O’Hagan JB, Price LL, Provencio I, Skene DJ, and Brainard GC (2014) Measuring and using light in the melanopsin age. *Trends Neurosci* 37:1–9. [PubMed: 24287308]
- Maemura K, Takeda N, and Nagai R (2007) Circadian rhythms in the CNS and peripheral clock disorders: role of the biological clock in cardiovascular diseases. *J Pharmacol Sci* 103:134–138. [PubMed: 17299249]
- Nagare R, Plitnick B, and Figueiro MG (2018a) Does the iPad Night Shift mode reduce melatonin suppression? *Light Res Technol*:10.1177/1072211177153517748189.
- Nagare R, Plitnick B, and Figueiro MG (2018b) Effect of exposure duration and light spectra on nighttime melatonin suppression in adolescents and adults. *Light Res Technol*: 10.1177/1072211177153518763003.
- Portaluppi F, Smolensky MH, and Touitou Y (2010) Ethics and methods for biological rhythm research on animals and human beings. *Chronobiol Int* 27:1911–1929. [PubMed: 20969531]
- Potter GD, Skene DJ, Arendt J, Cade JE, Grant PJ, and Hardie LJ (2016) Circadian rhythm and sleep disruption: Causes, metabolic consequences, and countermeasures. *Endocr Rev* 37:584–608. [PubMed: 27763782]
- Rahman SA, Marcu S, Shapiro CM, Brown TJ, and Casper RF (2011) Spectral modulation attenuates molecular, endocrine, and neurobehavioral disruption induced by nocturnal light exposure. *Am J Physiol Endocrinol Metab* 300:E518–527. [PubMed: 21177289]
- Rahman SA, St Hilaire MA, and Lockley SW (2017) The effects of spectral tuning of evening ambient light on melatonin suppression, alertness and sleep. *Physiol Behav* 177:221–229. [PubMed: 28472667]
- Rea MS, Figueiro MG, Bierman A, and Bullough JD (2010) Circadian light. *J Circadian Rhythms* 8:2. [PubMed: 20377841]
- Rea MS and Figueiro MG (2018) Light as a circadian stimulus for architectural lighting. *Light Res Technol* 50:497–510.
- Rea MS, Figueiro MG, Bierman A, and Hamner R (2012) Modelling the spectral sensitivity of the human circadian system. *Light Res Technol* 44:386–396.
- Rea MS, Figueiro MG, Bullough JD, and Bierman A (2005) A model of phototransduction by the human circadian system. *Brain Res Rev* 50:213–228. [PubMed: 16216333]
- Roenneberg T, Wirz-Justice A, and Mrosovsky M (2003) Life Between Clocks: Daily Temporal Patterns of Human Chronotypes. *J Biol Rhythms* 18:80–90. [PubMed: 12568247]
- Ruby N, Brennan T, and Xie X (2002) Role of melanopsin in circadian responses to light. *Science* 298:2211–2213. [PubMed: 12481140]

- Russo PM, Bruni O, Lucidi F, Ferri R, and Violani C (2007) Sleep habits and circadian preference in Italian children and adolescents. *J Sleep Res* 16:163–169. [PubMed: 17542946]
- Sasseville A, Paquet N, Sévigny J, and Hébert M (2006) Blue blocker glasses impede the capacity of bright light to suppress melatonin production. *J Pineal Res* 41:73–78. [PubMed: 16842544]
- Souman JL, Borra T, de Goijer I, Schlangen LJM, Vlaskamp BNS, and Lucassen MP (2018) Spectral tuning of white light allows for strong reduction in melatonin suppression without changing illumination level or color temperature. *J Biol Rhythms* 33:420–431. [PubMed: 29984614]
- Stevens RG, Blask DE, Brainard GC, Hansen J, Lockley SW, Provencio I, Rea MS, and Reinlib L (2007) Meeting report: the role of environmental lighting and circadian disruption in cancer and other diseases. *Environ Health Perspect* 115:1357–1362. [PubMed: 17805428]
- Thapan K, Arendt J, and Skene DJ (2001) An action spectrum for melatonin suppression: evidence for a novel non-rod, non-cone photoreceptor system in humans. *J Physiol* 535:261–267. [PubMed: 11507175]
- Van de Werken M, Gimenez MC, de Vries B, Beersma DG, and Gordijn MC (2013) Short-wavelength attenuated polychromatic white light during work at night: limited melatonin suppression without substantial decline of alertness. *Chronobiol Int* 30:843–854. [PubMed: 23705821]
- Wyszecki G, and Stiles WS (1982) *Color Science: Concepts and Methods, Quantitative Data and Formulae*. John Wiley & Sons, New York, NY.
- Young ME, and Bray MS (2007) Potential role for peripheral circadian clock dyssynchrony in the pathogenesis of cardiovascular dysfunction. *Sleep Med* 8:656–667. [PubMed: 17387040]
- Zeitler JM, Dijk DJ, Kronauer R, Brown E, and Czeisler C (2000a) Sensitivity of the human circadian pacemaker to nocturnal light: melatonin phase resetting and suppression. *J Physiol* 526:695–702. [PubMed: 10922269]
- Zeitler JM, Dijk DJ, Kronauer RE, Brown EN, and Czeisler CA (2000b) Sensitivity of the human circadian pacemaker to nocturnal light: melatonin phase resetting and suppression. *J Physiol* 526:695–702. [PubMed: 10922269]




**Figure 1.**

The relative spectral power distributions of the lighting interventions employed in this study and the melanopsin function adjusted to account for crystalline lens transmission according to Wyszecki and Stiles (1982). The 3000 K light is the conventional source and the 3000 K mod is the “cyan-gap” source.




**Figure 2.**

The experimental setup, showing participants performing their choice of task with the desktop luminaires in place. The luminaires were mounted on adjustable stands, and their height and distance from participants' eyes were individually adjusted for each participant to ensure delivery of the desired stimulus. For a fixed width of 18 in. and a standard viewing distance of 30 in., the luminaires delivered an image that was approximately  $33^\circ$  in width at the participants' eyes.




**Figure 3.**

The experimental protocol used for the study, showing the relationship between experimental conditions, saliva sample times (S1, S2, and S3), and Daysimeter use. The participants were exposed to all 3 experimental conditions in both phases of the study, with high levels for the lighting interventions being delivered during Phase 1 and low light levels for the lighting interventions being delivered during Phase 2 (see Tables 1 and 2). Light levels were maintained at < 5 lux at the eyes for the dim light period (represented by the hatched pattern) preceding the lighting conditions and the dim light control.



**Figure 4.**


Raw melatonin data for the dim light control and the 2 intervention spectra for both light levels at baseline and after 30-min and 60-min exposures. The data for the high light level are shown on the left, and those for the low light level are shown on the right. The error bars represent standard error of the mean.



**Figure 5.**

The graph on the left depicts the main effect of light level on nocturnal melatonin suppression compared to the dim-light control condition, combined across the 2 spectra and the 2 exposure durations. The graph on the right depicts the main effect of exposure duration on nocturnal melatonin suppression compared to the dim-light control condition, combined across the 2 spectra and the 2 light levels. The error bars represent SD. The asterisks represent statistical significance ( $p < 0.05$ ).





**Figure 6.** Mean  $\pm$  SD melatonin suppression for the 2 light levels after a 1-h exposure, normalized to the control (dim-light condition), compared to the a priori CS predictions derived following Rea and Figueiro, 2018. The error bars represent SD.

**Table 1.**

Photometric characteristics of the lighting interventions, based on the Daysimeter data and calculated following the Rea and Figueiro (2018) (CS) and Enezi et al. (2011) (Melanopic illuminance)

Study phase (light level)	Metric	3000 K	3000 K mod
Phase 1 (high light level)	Mean $\pm$ SD photopic illuminance (lux)	897 $\pm$ 96.8	787 $\pm$ 119.4
	CL <sub>A</sub> <sup>a</sup>	790	482
	Melanopic illuminance <sup>b</sup>	436	265
	Predicted CS <sup>a</sup>	0.49	0.41
Phase 2 (low light level)	Mean $\pm$ SD photopic illuminance (lux)	384 $\pm$ 50.3	361 $\pm$ 34.1
	CL <sub>A</sub> <sup>a</sup>	338	221
	Melanopic illuminance <sup>b</sup>	187	122
	Predicted CS <sup>a</sup>	0.34	0.26

(a) Rea and Figueiro (2018).

(b) Enezi et al. (2011)

**Table 2.**

Calculated  $\alpha$ -opic irradiances for all experimental conditions, following the SI-compliant approach recommended by the CIE

Study phase (light level)	Lighting intervention	Cyanopic irradiance $\mu\text{W. cm}^{-2}$	Melanopic irradiance $\mu\text{W. cm}^{-2}$	Rhodopic irradiance $\mu\text{W. cm}^{-2}$	Chloropic irradiance $\mu\text{W. cm}^{-2}$	Erythroptic irradiance $\mu\text{W. cm}^{-2}$
Phase 1 (high light level)	3000 K	21.5	51.4	73.4	118.4	154.5
	3000 K mod	27.9	29.2	51.9	102.7	129.6
Phase 2 (low light level)	3000 K	9.2	22.0	31.4	50.7	66.2
	3000 K mod	12.8	13.4	23.8	47.1	59.4

Notes: The  $\alpha$ -opic irradiances are calculated using the CIE's (Commission Internationale de l'Éclairage, 2015) proposed SI-compliant version of the Lucas et al. (2014) toolbox, which is available as a freely downloadable electronic document ([http://files.cie.co.at/784\\_TN003\\_Toolbox.xls](http://files.cie.co.at/784_TN003_Toolbox.xls)).

**Table 3.**

Mean absolute baseline melatonin levels at S1

Study variable	Study phase					
	Phase 1 (high light level)			Phase 2 (low light level)		
Experimental condition	Dim light	3000 K	3000 K mod	Dim light	3000 K	3000 K mod
Mean $\pm$ SD melatonin level (pg/mL)	9.0 $\pm$ 5.1	10.5 $\pm$ 6.2	10.2 $\pm$ 5.1	12.0 $\pm$ 9.1	11.8 $\pm$ 8.1	11.6 $\pm$ 9.0

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript