

Effects of 2,2-dichloroacetamide (DCAcAm), an emerging disinfection by-product in drinking water, on the intestinal microbiota of adult zebrafish

Bin Xue, Chenyu Li, Shang Wang, Chen Zhao, Kun Dai, Wenhong Li, Zhuge Xi, Jingfeng Wang, Zhigang Qiu and Zhiqiang Shen

ABSTRACT

The presence of disinfection by-products (DBPs) increases the mutagenicity of water and may pose adverse health effects. Gut microbiota exerts a fundamental role on host physiology, and how extrinsic perturbations influence its composition has been increasingly examined. However, the effect of DBPs on gut microbiota is still poorly understood. In the present study, adult zebrafish were exposed to different concentrations of dichloroacetamide (DCAcAm, an emerging nitrogenous DBP) for 30 days. Sequencing of 16S rRNA amplicons revealed a significant change in the richness and diversity of microbiota in the gut of DCAcAm-exposed zebrafish. At the phylum level, the abundance of Proteobacteria decreased and the abundance of Fusobacteria and Firmicutes increased significantly in the gut after exposure to 100 and 500 µg/L DCAcAm. At the genus level, the abundances of several bacteria which are considered pathogens or opportunistic pathogens in fish and closely related to fish metabolism, disease and inflammation (*Aeromonas*, *Stenotrophomonas*, *Bacteroides* and *Ralstonia*) increased in the DCAcAm-treated groups. Our results reveal that DBPs in drinking water potentially affect gut microbiota composition, which may contribute to the toxicity assessment of DBPs in future and provide new insight into the complex interactions between the DBPs in drinking water and host health.

Key words | dichloroacetamide, disinfection by-products, gut microbiota, zebrafish

Bin Xue
Chenyu Li
Shang Wang
Chen Zhao
Wenhong Li
Zhuge Xi
Jingfeng Wang
Zhigang Qiu
Zhiqiang Shen (corresponding author)
Tianjin Institute of Environmental & Operational
Medicine,
Key Laboratory of Risk Assessment and Control for
Environment & Food Safety,
Tianjin 300050,
China
E-mail: tianjinszq922@sohu.com

Kun Dai
Tianjin Rehabilitation Center,
Tianjin, 300191,
China

INTRODUCTION

The disinfection of drinking water has been hailed as one of the most important advances ever for protecting public health (Ngwenya *et al.* 2013). While the efficacy of chemical disinfectants in controlling pathogens is well recognized, one of the main concerns is the reactivity of disinfectants with organic and inorganic substances to generate harmful disinfection by-products (DBPs) (Yang *et al.* 2019). Research into DBPs dates back to the discovery of trichloromethane in the 1970s and since then, numerous studies have been conducted to increase the understanding of DBPs in terms of their formation processes and exposure levels in drinking water distribution systems and the associated human health

effects (Li & Mitch 2018; Mian *et al.* 2018). To date, around 700 DBPs have been identified in drinking waters, and the possible population health impacts associated with exposure to emerging DBPs are of particular interest due to their high cytotoxicity, genotoxicity, developmental toxicity and potential carcinogenicity (Cortes & Marcos 2018; Han & Zhang 2018). Accumulating evidence demonstrates that exposure to DBPs is associated with the increased risks of birth defects and a number of diseases such as colon, colorectal, rectal and bladder cancer (Rahman *et al.* 2014; Hrudey *et al.* 2015; Han *et al.* 2017; Jiang *et al.* 2017). In recent years, haloacetamides (HAcAms) in drinking water have

raised wide concerns due to their high toxicity compared with the regulated carbonaceous DBPs (Bond *et al.* 2011; Yang *et al.* 2014). Dichloroacetamide (DCAcAm), the most frequently detected member of HAcAms, were approximately two orders of magnitude more cytotoxic than dichloroacetic acid (Huang *et al.* 2012). Thus, the risk of DCAcAm for humans via long-term water drinking cannot be ignored.

Gut microbiota, the large number of microbial species residing in the gastrointestinal tract, is increasingly known to play a pivotal role in the maintenance of host health (Abdollahi-Roodsaz *et al.* 2016; Cani 2018). It has been established that changes in the gut microbiota are associated with not only obesity, diabetes and liver diseases but also cancer and even neurodegenerative diseases (Barichella *et al.* 2018; Yoshida *et al.* 2018). Furthermore, gut microbiota dynamics are highly sensitive to exogenous stressors. A few recent studies have demonstrated that exposure to different kinds of environmental pollutants, including antibiotics, nanomaterials, pesticides, heavy metals and persistent organic pollutants, can effectively induce gut microbiota dysbiosis in different experimental models (Kan *et al.* 2015; Jin *et al.* 2017; Chen *et al.* 2018a, 2018b). However, information about the effects of DBPs on the intestinal microbiota remains limited.

Considering the prevalence and potential adverse human health effects of DBPs in disinfected drinking water, we wondered whether the 'DBPs' associated with the 'gut microbiota'. For this purpose, we exposed adult zebrafish to various concentrations of DCAcAm, an emerging nitrogenous DBP that was widely detected in disinfected drinking water, and determined whether or not they could induce microbiota dysbiosis in the gut by using 16S rRNA gene deep sequencing. The results acquired in the present study may contribute to the current knowledge of the link between DBPs and intestinal microbiota, and provide new information regarding DBPs-induced toxicity.

METHODS

Chemicals and reagents

The DCAcAm was obtained from Alfa Aesar (Karlsruhe, Germany). A stock solution of DCAcAm was prepared in

dimethyl sulfoxide of high-performance liquid chromatography-grade (DMSO; Sigma-Aldrich, USA). All other reagents were of analytic grade.

Fish maintenance and exposure

Adult zebrafish were purchased from the China Zebrafish Resource Center. Each fish was raised in a separate tank with 300 mL charcoal-filtered and 24-h aerated tap water at a constant ambient temperature of 28.0 ± 2.0 °C under a light/dark cycle of 14/10 h. The fish were fed twice daily with commercial flake food (Sera Vipran, Petco, USA). After 2 weeks' acclimation, fish were randomly divided into four groups (10 fish in each group), including a control group (0 µg/L DCAcAm) and the three DCAcAm-exposure groups (10, 100 and 500 µg/L DCAcAm). The control group received an equal volume of DMSO alone. The final DMSO concentrations in the control and exposure groups were <0.001%. The water was renewed daily, to maintain the appropriate concentrations of DCAcAm. After 30 days of exposure, the zebrafish were anesthetized in 0.03% MS-222 (Sigma-Aldrich) and dissected for their intestines, which were immediately snap-frozen in liquid nitrogen and stored at -80 °C for the following microbiota analyses.

Measurement of DCAcAm

The actual concentrations of DCAcAm in exposure media after renewal were measured according to a previous method with a small modification (Yu *et al.* 2015; Lin *et al.* 2016). Briefly, the extraction of DCAcAm was performed using liquid-liquid extraction with ethyl acetate. The detection was carried out using an Agilent 7890A capillary gas chromatograph coupled with a micro-electron capture detector. DCAcAm was separated via DB-5MS column (30 mm × 0.25 mm × 0.25 µm, Agilent, USA).

16S rRNA amplicon sequencing and bioinformatic analyses

There are a series of procedures from DNA extraction to final data acquisition including total DNA extraction,

polymerase chain reaction and product purification, library construction, sequencing and data analysis. All procedures were conducted using Novogene Bioinformatics Technology Co. Ltd (Beijing, China).

Briefly, after each exposure phase, nine zebrafish in each group were randomly chosen for gut microbial community analysis. From each group, three intestines excised from three individual zebrafish were pooled together and considered a biological replicate ($n = 3$ per treatment group). Total bacteria DNA was extracted from intestinal samples using the cetyltrimethylammonium bromide method. The purity and concentration of DNA extraction were determined by electrophoresis on 1% agarose gels. Genomic DNA were then used for the amplification of 16S rRNA using the primer pair 341F (50-CCTAYGGGRBGCAS CAG-30) and 806R (50-GGACTACNNGGGTATCTAAT-30), targeting the V3–V4 hypervariable regions. Then, the amplicons were sequenced on an Ion S5™ XL platform and 400 bp/600 bp single-end reads were generated. The raw reads were filtered according to the cut adapt quality controlled process (V1.9.1, <http://cutadapt.readthedocs.io/en/stable/>) and chimera sequences by use of UCHIME algorithm (UCHIME Algorithm, http://www.drive5.com/usearch/manual/uchime_algo.html) (Edgar *et al.* 2011; Haas *et al.* 2011). The effective tags were then clustered by UPARSE (Uparse v7.0.1001 <http://drive5.com/uparse/>) to operational taxonomic units according to 97% similarity. After taxonomic annotation based on the RDP classifier (version 2.2) and the Green Genes Database, the number of sequences was calculated and summed for each classification level (e.g., phylum and genus) to compare the gut microbial abundance and diversity among exposure groups. Alpha- and beta-diversity analyses were conducted

with QIIME (Version 1.7.0) and displayed with R software (Version 2.15.3). After rarefying the number of reads, observed-species, Chao1, Shannon, Simpson, ACE and Good's coverage were calculated, and statistical significance was determined using Student's *t*-tests and a 5% false-discovery rate. Principal coordinate analysis (PCoA) was displayed by WGCNA package, stat packages and ggplot2 package in R software (Version 2.15.3) to get principal coordinates and visualize from complex, multidimensional data.

RESULTS AND DISCUSSION

Concentration of DCaAm in the exposure media

Nitrogenous DBPs in drinking water, especially the emerging HAcAms which have demonstrated higher cytotoxicity and genotoxicity than carbonaceous DBPs and other nitrogenous DBPs (e.g., haloacetonitriles), have limited study in terms of their formation, removal and toxicity properties and thus are a cause of wide concern (Bond *et al.* 2011). Among the detected HAcAms, DCaAm is present at the highest concentration and caused significant chronic cytotoxicity and acute genotoxicity (Huang *et al.* 2012). In this study, DCaAm-induced changes in gut microbiota composition of adult zebrafish were characterized to provide a new perspective on the investigation of DBPs' toxicity. A low treatment concentration of DCaAm (10 µg/L) was selected to represent DCaAm exposure from oral intake of drinking water (Cortes & Marcos 2018), while the medium treatment (100 µg/L) represents those of fish with high contents of organic nitrogen which can serve as

Table 1 | Alpha diversity in zebrafish intestines after DCaAm exposure (0, 10, 100 and 500 µg/L)^a

Group	Observed species	Shannon ^b	Simpson ^b	Chao1 ^c	ACE ^c	Good's coverage (%) ^d
0	717 ± 154	4.38 ± 0.76	0.78 ± 0.10	765.37 ± 182.24	772.69 ± 179.85	99.87 ± 0.06
10	412 ± 85*	2.70 ± 0.38*	0.66 ± 0.07	463.02 ± 86.14	468.24 ± 90.80	99.90 ± 0.01
100	425 ± 65*	3.13 ± 0.22	0.74 ± 0.06	483.94 ± 71.48	488.76 ± 39.38	99.87 ± 0.06
500	515 ± 98	3.48 ± 0.37	0.72 ± 0.07	576.46 ± 91.75	566.81 ± 89.90	99.90 ± 0.00

^aValues represent the mean ± SD of replicates.

^bIndicative of bacterial community diversity.

^cIndicative of bacterial community richness.

^dIndicative of bacterial sequencing coverage.

* $P < 0.05$ indicates significant difference between DCaAm exposure groups and the control group.

a source of DCaAm precursors, and an additional high DCaAm exposure (500 $\mu\text{g/L}$) for fish in polluted systems. Concentrations of DCaAm in each DCaAm exposure group were detected to be $9.82 \pm 0.21 \mu\text{g/L}$, $100.43 \pm 0.82 \mu\text{g/L}$ and $498.96 \pm 1.48 \mu\text{g/L}$, respectively. No DCaAm was found in the control group. Concentration used in the present study was an environmentally realistic one; hence, there is a guarantee of greater ecological relevance from the results based on the current data.

Overall characterization of the sequencing data

Given the essential role of the gut microbiome in a variety of aspects of human health coupled with the potential health risk of DCaAm, there is a need to elucidate the effects of DCaAm exposure on the gut microbiome. Although DCaAm has been demonstrated to cause the acute metabolism and DNA damage in zebrafish (Lin et al. 2016), toxicological literature concerning the effects of DCaAm on gut microbiota of zebrafish is scarce. To the best of our knowledge, only one similar study to date has demonstrated that the diversity of the gut microbiome in mice was significantly reduced due to trichloroacetamide exposure. Consistent with our findings, a number of perturbed gut microbes such as Firmicutes, Bacteroidetes and Proteobacteria were significantly changed (Zhang et al. 2015).

In the present study, adult zebrafish were exposed to different concentrations of DCaAm for 30 days. The summary information of pyrosequencing of the V3–V4 region of 16S rRNA genes is shown in Table 1. The values of Good's coverage were nearly 100% for all sequences in the four groups, suggesting sufficient sequencing depth for the investigation of gut microbiota. There were differences between the control and DCaAm-treated groups (10 and 100 $\mu\text{g/L}$) regarding the number of species detected (Figure 1(a)). The Shannon diversity index also suggested that the diversity of the gut microbiota was influenced by 10 $\mu\text{g/L}$ DCaAm exposure (Figure 1(b)). No significant differences were observed among DCaAm treatment groups. In addition, PCoA results showed that communities from different DCaAm exposure groups were clearly more distant, particularly from those of the control group (Figure 1(c)).




Figure 1 | Effects of DCaAm on the composition of microbiota in the gut, as detected by 16S rRNA gene sequencing. (a) Observed species and (b) Shannon index of the diversity of gut microbiota after DCaAm exposure. (c) Gut microbiota patterns in controls and DCaAm-treated groups differentiated by principal coordinate analysis.




Figure 2 | Relative abundance of: (a) top 10 phyla and (b) top 30 genera after exposure to 0, 10, 100 and 500 µg/L DCACAm, respectively.

Dysbiosis of gut microbiota induced by DCACAm

Presented here is the first observation that DCACAm can induce microbiota dysbiosis in the gut of adult zebrafish.

At phylum level, Proteobacteria were dominant in all groups, representing 81.48%, 93.96% and 87.44% in the control, 10 and 500 µg/L DCACAm-treated groups, respectively (Figure 2(a)). Previous studies using zebrafish also

identify the dominance of gut microbiota by Proteobacteria (Chen *et al.* 2018a; Jin *et al.* 2018), although Davis *et al.* (2016) observed highest abundance of Fusobacteria in zebrafish intestines. These geographical differences in gut microbial community of domesticated zebrafish have historical connections with laboratory aquaculture facilities (Chen *et al.* 2018a). Interestingly, the relative abundance of Proteobacteria was decreased from 81.48% in the control group to 48.89% after 100 µg/L DCACAm exposure for 30 days (Figure 2(a)). In contrast, the composition of Fusobacteria increased from 5.8% in the control group to 47.22% in the 100 µg/L DCACAm-treated group. Furthermore, the composition of Firmicutes in the control group was approximately 2.92%, but it increased to approximately 8.21% in the 500 µg/L DCACAm-treated group. Similar results were obtained in the study of Jin *et al.* (2018) that showed the abundance of Proteobacteria decreased significantly and the abundance of Fusobacteria and Firmicutes increased in the gut of zebrafish after exposure to 1,000 mg/L 0.5 and 50 mm polystyrene microplastic for 14 days.

At the genus level, *Stenotrophomonas* was the most abundant genus in all groups, with the exception of the 100 µg/L DCACAm-treated group, which was dominated by *Cetobacterium* (Figure 2(b)). The abundance of *Cetobacterium* slightly decreased in 10 and 500 µg/L DCACAm-treated groups, and significantly increased in 100 µg/L DCACAm exposure group (Figure 2(b)). This increased abundance of *Cetobacterium* is consistent with a previous study, which showed that fish had higher abundance of *Cetobacterium* after administration with dexamethasone (Qi *et al.* 2019). *Cetobacterium*, with a role in sulfur metabolism, has been shown to produce vitamin B12 with high efficiency and to ferment peptides and carbohydrates (Bridges *et al.* 2018; Chang *et al.* 2019). Therefore, we assume that the changes in composition of the fish gut microbiome may suggest an adaptive response to increase DCACAm detoxification; however, further research should be targeted toward elucidating the underlying mechanisms of these perturbations. Furthermore, although the functions of some of the altered bacteria (Figure 2(b)), such as *Phyllobacterium* and *Acinetobacter*, remain unclear in fish, the functions of several other bacteria, including *Stenotrophomonas*, *Bacteroides* and

Ralstonia, are considered pathogens or opportunistic pathogens in fish and closely related to fish metabolism, disease and inflammation (Smith *et al.* 2011; Janda *et al.* 2016; Jin *et al.* 2018). In addition, the *Aeromonas*, present only in the 100 and 500 µg/L DCACAm-treated group, are disease-related bacteria and their reproduction has been demonstrated to cause inflammation of the intestinal tract and infection of soft tissue in a previous study (Chen *et al.* 2018b).

CONCLUSIONS

In conclusion, we have observed that different concentrations of DCACAm can induce microbiota dysbiosis in the adult zebrafish gut after a 30-day exposure. The results may extend our knowledge regarding the correlations of gut microbial community shift with DCACAm exposure and provide new information concerning DBPs-induced toxicity in zebrafish. Clearly, the dysbiosis of gut microbiota by chronic DCACAm exposure in this study is only preliminary. Further research should be targeted toward exploring the relationship between gut microbes and host health, and elucidating the underlying mechanisms of these perturbations.

ACKNOWLEDGEMENTS

This work was supported by the National Natural Science Foundation of China (Grants 51678565, 31470234), AWS18J004, BWS17J025 and AWS16J004, and the Natural Science Foundation of Tianjin, China (Grants 17JCZDJC39100). The authors declare no competing financial interests related to the publication of this study. Bin Xue and Chenyu Li contributed equally to this work.

REFERENCES

- Abdollahi-Roodsaz, S., Abramson, S. B. & Scher, J. U. 2016 [The metabolic role of the gut microbiota in health and rheumatic disease: mechanisms and interventions](#). *Nat. Rev. Rheumatol.* **12** (8), 446–455.

- Barichella, M., Severgnini, M., Cilia, R., Cassani, E., Bolliri, C., Caronni, S., Ferri, V., Cancellò, R., Ceccarani, C., Faierman, S., Pinelli, G., De Bellis, G., Zecca, L., Cereda, E., Consolandi, C. & Pezzoli, G. 2018 Unraveling gut microbiota in Parkinson's disease and atypical parkinsonism. *Mov. Disord.* **34**, 396–405.
- Bond, T., Huang, J., Templeton, M. R. & Graham, N. 2011 Occurrence and control of nitrogenous disinfection by-products in drinking water – a review. *Water Res.* **45** (15), 4341–4354.
- Bridges, K. N., Zhang, Y., Curran, T. E., Magnuson, J. T., Venables, B. J., Durrer, K. E., Allen, M. S. & Roberts, A. P. 2018 Alterations to the intestinal microbiome and metabolome of *Pimephales promelas* and *Mus musculus* following exposure to dietary methylmercury. *Environ. Sci. Technol.* **52** (15), 8774–8784.
- Cani, P. D. 2018 Human gut microbiome: hopes, threats and promises. *Gut* **67** (9), 1716–1725.
- Chang, X., Li, H., Feng, J., Chen, Y., Nie, G. & Zhang, J. 2019 Effects of cadmium exposure on the composition and diversity of the intestinal microbial community of common carp (*Cyprinus carpio* L.). *Ecotoxicol. Environ. Saf.* **171**, 92–98.
- Chen, L., Guo, Y., Hu, C., Lam, P. K. S., Lam, J. C. W. & Zhou, B. 2018a Dysbiosis of gut microbiota by chronic coexposure to titanium dioxide nanoparticles and bisphenol A: implications for host health in zebrafish. *Environ. Pollut.* **234**, 307–317.
- Chen, L., Zhang, W., Hua, J., Hu, C., Lok-Shun Lai, N., Qian, P. Y., Lam, P. K. S., Lam, J. C. W. & Zhou, B. 2018b Dysregulation of intestinal health by environmental pollutants: involvement of the estrogen receptor and aryl hydrocarbon receptor. *Environ. Sci. Technol.* **52** (4), 2323–2330.
- Cortes, C. & Marcos, R. 2018 Genotoxicity of disinfection byproducts and disinfected waters: a review of recent literature. *Mutat. Res.* **831**, 1–12.
- Davis, D. J., Doerr, H. M., Grzelak, A. K., Busi, S. B., Jasarevic, E., Ericsson, A. C. & Bryda, E. C. 2016 *Lactobacillus plantarum* attenuates anxiety-related behavior and protects against stress-induced dysbiosis in adult zebrafish. *Sci. Rep.* **6**, 33726.
- Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C. & Knight, R. 2011 UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* **27** (16), 2194–2200.
- Haas, B. J., Gevers, D., Earl, A. M., Feldgarden, M., Ward, D. V., Giannoukos, G., Ciulla, D., Tabbaa, D., Highlander, S. K., Sodergren, E., Methe, B., DeSantis, T. Z., Petrosino, J. F., Knight, R. & Birren, B. W. 2011 Chimeric 16S rRNA sequence formation and detection in Sanger and 454-pyrosequenced PCR amplicons. *Genome Res.* **21** (3), 494–504.
- Han, J. & Zhang, X. 2018 Evaluating the comparative toxicity of DBP mixtures from different disinfection scenarios: a new approach by combining freeze-drying or rotoevaporation with a marine polychaete bioassay. *Environ. Sci. Technol.* **52** (18), 10552–10561.
- Han, J., Zhang, X., Liu, J., Zhu, X. & Gong, T. 2017 Characterization of halogenated DBPs and identification of new DBPs trihalomethanols in chlorine dioxide treated drinking water with multiple extractions. *J. Environ. Sci. (China)* **58**, 83–92.
- Hrudey, S. E., Backer, L. C., Humpage, A. R., Krasner, S. W., Michaud, D. S., Moore, L. E., Singer, P. C. & Stanford, B. D. 2015 Evaluating evidence for association of human bladder cancer with drinking-water chlorination disinfection by-products. *J. Toxicol. Environ. Health B Crit. Rev.* **18** (5), 213–241.
- Huang, H., Wu, Q. Y., Hu, H. Y. & Mitch, W. A. 2012 Dichloroacetonitrile and dichloroacetamide can form independently during chlorination and chloramination of drinking waters, model organic matters, and wastewater effluents. *Environ. Sci. Technol.* **46** (19), 10624–10631.
- Janda, J. M., Abbott, S. L. & McIver, C. J. 2016 *Plesiomonas shigelloides* revisited. *Clin. Microbiol. Rev.* **29** (2), 349–374.
- Jiang, J., Zhang, X., Zhu, X. & Li, Y. 2017 Removal of intermediate aromatic halogenated DBPs by activated carbon adsorption: a new approach to controlling halogenated DBPs in chlorinated drinking water. *Environ. Sci. Technol.* **51** (6), 3435–3444.
- Jin, Y., Wu, S., Zeng, Z. & Fu, Z. 2017 Effects of environmental pollutants on gut microbiota. *Environ. Pollut.* **222**, 1–9.
- Jin, Y., Xia, J., Pan, Z., Yang, J., Wang, W. & Fu, Z. 2018 Polystyrene microplastics induce microbiota dysbiosis and inflammation in the gut of adult zebrafish. *Environ. Pollut.* **235**, 322–329.
- Kan, H., Zhao, F., Zhang, X. X., Ren, H. & Gao, S. 2015 Correlations of gut microbial community shift with hepatic damage and growth inhibition of *Carassius auratus* induced by pentachlorophenol exposure. *Environ. Sci. Technol.* **49** (19), 11894–11902.
- Li, X. F. & Mitch, W. A. 2018 Drinking water disinfection byproducts (DBPs) and human health effects: multidisciplinary challenges and opportunities. *Environ. Sci. Technol.* **52** (4), 1681–1689.
- Lin, T., Zhou, D., Yu, S. & Chen, W. 2016 The removal process of 2,2-dichloroacetamide (DCAcAm), a new disinfection by-product, in drinking water treatment process and its toxicity on zebrafish. *Chemosphere* **159**, 403–411.
- Mian, H. R., Hu, G., Hewage, K., Rodriguez, M. J. & Sadiq, R. 2018 Prioritization of unregulated disinfection by-products in drinking water distribution systems for human health risk mitigation: a critical review. *Water Res.* **147**, 112–131.
- Ngwenya, N., Ncube, E. J. & Parsons, J. 2013 Recent advances in drinking water disinfection: successes and challenges. *Rev. Environ. Contam. Toxicol.* **222**, 111–170.
- Qi, X. Z., Tu, X., Zha, J. W., Huang, A. G., Wang, G. X. & Ling, F. 2019 Immunosuppression-induced alterations in fish gut microbiota may increase the susceptibility to pathogens. *Fish Shellfish Immunol.* **88**, 540–545.
- Rahman, M. B., Cowie, C., Driscoll, T., Summerhayes, R. J., Armstrong, B. K. & Clements, M. S. 2014 Colon and rectal

- cancer incidence and water trihalomethane concentrations in New South Wales, Australia. *BMC Cancer* **14**, 445.
- Smith, B., Bode, S., Petersen, B. L., Jensen, T. K., Pippert, C., Kloppenborg, J., Boye, M., Krogfelt, K. A. & Molbak, L. 2011 Community analysis of bacteria colonizing intestinal tissue of neonates with necrotizing enterocolitis. *BMC Microbiol.* **11**, 73.
- Yang, F., Zhang, J., Chu, W., Yin, D. & Templeton, M. R. 2014 Haloactamides versus halomethanes formation and toxicity in chloraminated drinking water. *J. Hazard. Mater.* **274**, 156–163.
- Yang, M., Zhang, X., Liang, Q. & Yang, B. 2019 Application of (LC)/MS/MS precursor ion scan for evaluating the occurrence, formation and control of polar halogenated DBPs in disinfected waters: a review. *Water Res.* **158**, 322–337.
- Yoshida, N., Emoto, T., Yamashita, T., Watanabe, H., Hayashi, T., Tabata, T., Hoshi, N., Hatano, N., Ozawa, G., Sasaki, N., Mizoguchi, T., Amin, H. Z., Hirota, Y., Ogawa, W., Yamada, T. & Hirata, K. I. 2018 *Bacteroides vulgatus* and *Bacteroides dorei* reduce gut microbial lipopolysaccharide production and inhibit atherosclerosis. *Circulation.* **138** (22), 2486–2498.
- Yu, S., Lin, T., Chen, W. & Tao, H. 2015 The toxicity of a new disinfection by-product, 2,2-dichloroacetamide (DCAcAm), on adult zebrafish (*Dani rerio*) and its occurrence in the chlorinated drinking water. *Chemosphere* **139**, 40–46.
- Zhang, Y., Zhao, F., Deng, Y., Zhao, Y. & Ren, H. 2015 Metagenomic and metabolomic analysis of the toxic effects of trichloroacetamide-induced gut microbiome and urine metabolome perturbations in mice. *J. Proteome Res.* **14** (4), 1752–1761.

First received 11 April 2019; accepted in revised form 26 June 2019. Available online 29 July 2019