

# Effects of daily iron supplementation in primary-school-aged children: systematic review and meta-analysis of randomized controlled trials

Michael Low MBBS BMedSci, Ann Farrell MBBS, Beverley-Ann Biggs PhD, Sant-Rayn Pasricha PhD

See related commentary by Gray-Donald at [www.cmaj.ca/lookup/doi/10.1503/cmaj.131249](http://www.cmaj.ca/lookup/doi/10.1503/cmaj.131249)

## ABSTRACT

**Background:** Anemia is an important public health and clinical problem. Observational studies have linked iron deficiency and anemia in children with many poor outcomes, including impaired cognitive development; however, iron supplementation, a widely used preventive and therapeutic strategy, is associated with adverse effects. Primary-school-aged children are at a critical stage in intellectual development, and optimization of their cognitive performance could have long-lasting individual and population benefits. In this study, we summarize the evidence for the benefits and safety of daily iron supplementation in primary-school-aged children.

**Methods:** We searched electronic databases (including MEDLINE and Embase) and other sources (July 2013) for randomized and quasi-randomized controlled trials involving daily iron supplementation in children aged 5–12 years. We combined the data using random effects meta-analysis.

**Results:** We identified 16 501 studies; of these, we evaluated 76 full-text papers and included 32 studies including 7089 children. Of the included studies, 31 were conducted in low- or middle-income settings. Iron supplementation improved global cognitive scores (standardized mean difference 0.50, 95% confidence interval [CI] 0.11 to 0.90,  $p = 0.01$ ), intelligence quotient among anemic children (mean difference 4.55, 95% CI 0.16 to 8.94,  $p = 0.04$ ) and measures of attention and concentration. Iron supplementation also improved age-adjusted height among all children and age-adjusted weight among anemic children. Iron supplementation reduced the risk of anemia by 50% and the risk of iron deficiency by 79%. Adherence in the trial settings was generally high. Safety data were limited.

**Interpretation:** Our analysis suggests that iron supplementation safely improves hematologic and nonhematologic outcomes among primary-school-aged children in low- or middle-income settings and is well-tolerated.

### Competing interests:

Michael Low has received travel expenses from Gilead Sciences. Sant-Rayn Pasricha is co-investigator on an unrestricted research grant from Vifor Pharma. No competing interests declared by Ann Farrell and Beverley-Ann Biggs.

This article has been peer reviewed.

### Correspondence to:

Sant-Rayn Pasricha,  
[sant-rayn.pasricha@unimelb.edu.au](mailto:sant-rayn.pasricha@unimelb.edu.au)

*CMAJ* 2013, DOI:10.1503/cmaj.130628

An estimated 25% of school-aged children worldwide are anemic.<sup>1</sup> Iron deficiency is thought to account for about half of the global cases of anemia<sup>2</sup> and is associated with inadequate dietary iron and, in developing settings, hookworm and schistosomiasis.<sup>3</sup> In developed settings, anemia is prevalent among disadvantaged populations, including newly arrived refugees, indigenous people<sup>4</sup> and some ethnic groups (e.g., Hispanic people in the United States).<sup>5,6</sup> About 3% of primary-school-aged children in Canada are anemic.<sup>7</sup> Programs to address anemia are constrained by concerns that iron supplements cause adverse effects, including an increased risk of infections such as malaria in endemic areas.<sup>8</sup>

In observational studies, iron deficiency has been associated with impaired cognitive and physical development. It has been estimated that

each 10 g/L decrement in hemoglobin reduces future intelligence quotient (IQ) by 1.73 points.<sup>9</sup> However, observational data are susceptible to confounding,<sup>10</sup> and a causal relation between iron deficiency and cognitive impairment has not been confirmed.<sup>11</sup> Randomized controlled trials should overcome confounding, but results of trials examining this question have not agreed.

Optimizing cognitive and physical development in primary-school-aged children could have life-long benefits.<sup>12</sup> However, anemia-control recommendations must balance safety and efficacy. We performed a systematic review of the effects of daily iron supplementation, a commonly used strategy to combat anemia,<sup>2</sup> in primary-school-aged children. We examined cognitive, growth and hematologic outcomes and adverse effects across all settings.

## Methods

The protocol for our systematic review and meta-analysis is registered with PROSPERO ([www.crd.york.ac.uk/PROSPERO/display\\_record.asp?ID=CRD42011001208](http://www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42011001208)). We included outcomes that could be used to inform guidelines for anemia control.<sup>13</sup>

### Information sources

We searched the following electronic databases: Scopus, MEDLINE, Embase, Cochrane Central Register of Controlled Trials, ProQuest Digital Theses, the Australian Digital Theses Program database, WHO regional databases and OpenGrey. We reviewed the references of the identified articles and of previous systematic reviews. There was no language restriction applied. The search strategy for Scopus is presented in Appendix 1 (available at [www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.130628/-/DC1](http://www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.130628/-/DC1)). We performed the searches on July 4, 2013, for all databases except Embase (searched in June 2011). We searched the World Health Organization's International Clinical Trials Registry Platform in July 2013 for potentially eligible ongoing trials.

### Inclusion criteria

We included randomized controlled trials that included primary-school-aged children (5–12 yr) who were randomly assigned to daily ( $\geq 5$  d/wk) oral iron supplementation or control. We included studies that did not specifically recruit participants from this age range if the mean or median age of participants was between 5 and 12 years, if more than 75% of participants were aged 5–12 years, or if most of the study's recruitment age range overlapped 5–12 years. We excluded studies that included only children with a known developmental disability or a condition that substantially altered iron metabolism, including severe anemia. We included trials involving participants from all countries and socioeconomic backgrounds.

### Outcomes

We included the following primary outcomes: cognitive performance (as measured by study authors); physical growth; safety (gastrointestinal adverse effects, infection [including malaria], morbidity and, if measured, mortality); and hemoglobin, iron indices, prevalence of anemia (hemoglobin  $< 120$  g/L or as defined by study authors) and iron deficiency.

The secondary outcomes were the effects of iron on other micronutrients (including zinc) and physical performance.

### Data extraction

Three authors (M.L., A.F., S.-R.P.) screened the titles and abstracts identified by the search and excluded clearly ineligible references. Two authors (M.L., A.F.) independently assessed the remaining full-text studies against the inclusion criteria; their assessments were checked by a third author (S.-R.P.). Data were independently extracted by 2 authors (M.L., A.F.) and checked by a third author (S.-R.P.). One author (M.L.) entered the data into Review Manager software (RevMan 5.1, The Cochrane Collaboration) and a second author (S.-R.P.) checked entries for accuracy. Discrepancies were resolved through discussion.

### Assessment of risk of bias

We assessed the risk of bias by use of the Cochrane risk of bias assessment tool, which addresses selection, performance, attrition, detection and reporting bias through the evaluation of reported sequence generation, allocation concealment, blinding, incomplete outcome data and selective outcome reporting.<sup>14</sup> We used sensitivity analysis to examine the effects of removing studies with a high risk of bias (studies with poor or unclear randomization or allocation concealment, and poor or unclear blinding or high or imbalanced losses to follow-up) from the analysis.<sup>15</sup> For outcomes that had more than 10 included trials, we examined funnel plots for evidence of publication bias.

### Statistical analysis

Meta-analysis was performed for outcomes for which at least 2 studies provided data. We determined the mean difference (MD) for the difference between outcome means or the change from baseline for continuous data measured on the same scale. We calculated standardized mean differences (SMDs; difference of means is standardized according to standard deviation) for continuous data that were reported using different scales. If for a particular outcome, some studies reported outcome data only and others reported change from baseline data only, these data were combined using MD if the same scale had been used. If both outcome and change data were reported in the same study for the same outcome, the outcome data were used. For dichotomous data, the results are presented as risk ratios (RRs). For rates, the results are presented as rate ratios.

We included studies that provided iron supplementation along with a second intervention (cointervention), provided that the cointervention was applied identically in the control arm. For studies with a  $2 \times 2$  factorial design (e.g., participants randomized to iron or control, then a second randomization to another intervention or

control), we extracted the data and entered them into the meta-analysis separately.

We quantified heterogeneity using  $I^2$ . We considered substantial heterogeneity to exist if  $I^2$  was above 50%. We performed predefined subgroup analyses for outcomes that had 5 or more included studies. We included the following subgroups: baseline anemia status; baseline iron status; baseline iron deficiency anemia status; dose of iron supplementation; duration of iron supplementation; iron combination (iron alone or with a cointervention); and reported malaria endemicity of the setting. Differences between subgroups were tested using  $\chi^2$  and  $I^2$  statistics.<sup>16</sup> We created forest plots to present the effect size and relative weights for each study and the overall calculated effect size. Random-effects meta-analysis was used with calculation of  $\tau^2$  because of potential statistical, clinical and methodologic heterogeneity.<sup>8</sup>

## Results

The selection of studies for inclusion is illustrated in Figure 1. Our search strategy identified 16 501 potentially eligible titles and abstracts. We did not identify any potentially eligible unpublished trials in the International Clinical Trials Registry Platform. Following screening, 32 studies were deemed eligible; this included 7089 children (3837 randomized to iron supplements and 3252 to control). All studies except 1 were performed in low- or middle-income countries. Table 1 presents the characteristics of the included studies. Only 4 studies were considered at low overall risk of bias (Table 2).

The primary outcome findings are presented in Table 3. A complete table of findings including secondary outcomes is presented in Appendix 2, available at [www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.130628/-/DC1](http://www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.130628/-/DC1).

## Cognition

No studies reporting cognitive outcomes were at low overall risk of bias.

## Global cognitive performance

Global cognitive performance was reported in 9 studies: 5 studies used IQ (tests: Raven Progressive Matrices, Wechsler Intelligence Scale for Children [WISC], Test of Non-verbal Intelligence 2nd Edition),<sup>28,43,44,55–58</sup> 3 studies used author-adapted scales of global cognitive performance (visual memory, digit span, mazes test, clerical task tests with maximum scores of 40),<sup>28–31,34</sup> and 1 study used overall school performance.<sup>17,18</sup>


Data from 2 other studies could not be included because global cognitive scores were not presented.

Sen and Kanani<sup>52,53</sup> found significant improvements from iron supplementation in digit span, maze testing and visual memory testing but not clerical tasks in 110 girls aged 9–13 years. Rico and colleagues<sup>47–50</sup> found no difference with iron supplementation on various cognitive domains measured with an adapted WISC in 590 children aged 6–8 years other than a small significant improvement in responses on the stimulus discrimination task.

Children who received iron supplementation had higher global cognitive scores at the end of the intervention compared with children who received the control (SMD 0.50, 95% confidence interval [CI] 0.11 to 0.90;  $p = 0.01$ ,  $I^2 = 93%$ , 9 studies,  $n = 2355$ ) (Figure 2). This beneficial effect was seen among children who were anemic at baseline (SMD 0.29, 95% CI 0.07 to 0.51;  $p = 0.01$ ,  $I^2 = 22%$ , 6 studies,  $n = 487$ ) but not among those without anemia (SMD 0.01, 95% CI –0.10 to 0.11;  $p = 0.9$ ,  $I^2 = 0%$ , 4 studies,  $n = 1361$ ) (test for subgroup difference:  $p = 0.05$ ). Although fewer than 10 studies reported this outcome, a Funnel plot did not indicate publication bias (Appendix 3, available at [www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.130628/-/DC1](http://www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.130628/-/DC1)).

## Intelligence quotient

The most commonly reported global cognitive score was age-adjusted IQ. There was no overall



**Figure 1: Flow diagram of identification of included studies.** In total, 16 501 titles were identified, of which 43 references pertaining to 32 studies were eligible for inclusion in the systematic review.

**Table 1 (part 1 of 3):** Studies included in the systematic review\*

Study	Country	Age at recruitment, yr	Baseline anemia, iron deficiency status	Intervention	Control	No. randomized	Duration	Outcomes
Ayoya et al. <sup>17,18</sup>	Mali	7–12	Anemic, iron status unknown	Ferrous sulfate 100 mg + anthelmintic therapy	Anthelmintic therapy	Total: 202; iron: 105; placebo: 97	12 wk	Anemia, hemoglobin, malaria, ferritin, soluble transferrin receptor, cognitive score
Berger et al. <sup>19</sup>	Bolivia	3.3–8.3	Anemic, iron status unknown	Ferrous sulfate calculated to 3–4 mg/kg/d as closest amount to 20 mg or 36 mg/d	Placebo (once per wk)	Total: 115; iron: 58; placebo: 57	16 wk (4 mo)	Anemia, hemoglobin, zinc protoporphyrin
Boivin et al. <sup>20</sup>	Zaire	Mean age 7.7	Anemia, iron status unknown	20 mL of iron supplementation (formulation not stated) ± levamisole	No intervention ± levamisole	Total: 47; iron: 7; placebo: 15; iron + anthelmintic: 17; placebo + levamisole: 8	4 wk, 30 d	Data not extractable (all data presented as levamisole v. no levamisole) (not included in analysis)
Bradfield et al. <sup>21</sup>	Peru	7–13	Anemia and iron status unknown	Ferrous sulfate 5 mg ± anthelmintic therapy	Placebo ± anthelmintic therapy	Total: 87; iron: 51; placebo: 10; iron + anthelmintic: 16; placebo + anthelmintic: 10	10 wk	Hemoglobin, hematocrit, no. with helminth infection
Charoenlarp <sup>22</sup>	Thailand	5–14	Anemia and iron status unknown	Ferrous gluconate (300 mg) ± folate	Placebo ± folate	Total: 437; iron: 214; placebo: 223	3 mo	Hematocrit
Charoenlarp et al. <sup>23</sup>	Thailand	6–12	Anemia and iron status unknown	Ferrous sulfate (200 mg) ± mebendazole (if hookworms present)	Placebo ± mebendazole (if hookworms present)	Total: 72; iron: 33; placebo: 39	5 mo	Hemoglobin, hematocrit
Chwang et al. <sup>24</sup>	Indonesia	8.2–13.5	Subgroups: anemic, not anemic, iron deficient, not iron deficient, iron deficiency anemia, nonanemic noniron deficient	Ferrous sulfate (10 mg)	Placebo	Total: 139; iron: 59; placebo: 60	12 wk	Hemoglobin, serum iron, transferrin saturation, total iron binding capacity, height, weight, arm circumference, morbidity score; z scores for height, weight, arm circumference
de Silva et al. <sup>25</sup>	Sri Lanka	5–10	Anemia and iron status unknown	Ferrous sulfate (60 mg elemental iron)	Placebo	Total: 363; iron: 261; placebo: 102	8 wk	Anemia, iron deficiency, hemoglobin, ferritin, no. of URTI episodes, URTI episodes with fever, total no. of days sick with URTI, constipation, vomiting, GI upset
Fallahi et al. <sup>26</sup>	Iran	Grade 5 (about age 11 yr)	Anemia and iron status unknown	Ferrous sulfate (20 mg) + zinc	Zinc	Total: 54; iron: 26; placebo: 28	4 mo	Hemoglobin, ferritin, zinc, retinol
Gebreselassie <sup>27</sup>	Ethiopia	5–14	Anemic and iron deficient	Ferrous sulfate (300 mg)	Placebo	Total: 480; iron: 239; placebo: 241	3 mo	Anemia, iron deficiency, hemoglobin, ferritin, malaria, splenomegaly
Gopaldas et al. <sup>28–31</sup>	India	8–15	Iron status unknown. Subgroups: anemic, nonanemic	Ferrous sulfate (30 mg or 40 mg)	Placebo	Total: 210; iron: 140; placebo: 70	60 d	Hemoglobin, cognitive scores, exercise tests
Greisen <sup>32</sup>	Zambia	5–15	Anemic, iron status unknown	Ferrous fumarate (300 mg) ± folate	Placebo ± folate	Total: 430; iron: 218; placebo: 212	7 wk	Hemoglobin
Harvey et al. <sup>33</sup>	Papua New Guinea	8–12	Anemic, iron status unknown	Ferrous sulfate (200 mg)	Placebo	Total: 300; iron: 156; placebo: 144	16 wk	Hemoglobin, gastrointestinal upset, malaria

Continued

**Table 1 (part 2 of 3):** Studies included in the systematic review\*

Study	Country	Age at recruitment, yr	Baseline anemia, iron deficiency status	Intervention	Control	No. randomized	Duration	Outcomes
Kashyap et al. <sup>34</sup>	India	8–15	Anemia and iron status unknown	Ferrous sulfate (200 mg)	Placebo	Total: 130; iron: 65; placebo: 65	8 mo	Hemoglobin, cognitive scores
Konofal et al. <sup>35</sup>	France	5–8	Anemia status unknown, iron deficient	Ferrous sulfate (300 mg)	Placebo	Total: 23; iron: 18; placebo: 5	3 mo	Hemoglobin, ferritin, serum iron, transferrin saturation, constipation and abdominal pain
Latham et al. <sup>36</sup>	Kenya	Mean age control: 8.1; iron: 7.9	Anemia and iron status unknown	Ferrous sulfate (400 mg) + albendazole	Placebo + albendazole	Total: 55; iron: 29; placebo: 26	15 wk	Hemoglobin, height, weight, arm circumference, malaria, <i>Ascaris</i> infection
Lawless et al. <sup>37</sup>	Kenya	6–11	Anemia and iron status unknown	Ferrous sulfate (150 mg)	Placebo	Total: 86; iron: 42; placebo: 44	14 wk	Hemoglobin, ferritin, height, weight, z scores for height and weight
Nchito et al. <sup>38–40</sup>	Zambia	7–15 (mean 10.2)	Anemia and iron status unknown	Ferrous dextran (60 mg elemental iron) ± multivitamin	Placebo ± multivitamin	Total: 215; iron: 61; placebo: 50; Iron + multivitamin: 57; placebo + multivitamin: 47	10 mo	<i>Ascaris</i> infection
Paracha et al. <sup>41</sup>	Pakistan	11	Subgroups: anemic, not anemic, iron deficient, not iron deficient, iron deficient anemia, nonanemic noniron deficient	Ferrous gluconate (76 mg elemental iron) + multivitamin	Placebo + multivitamin	Total: 119; iron: 61; placebo: 58	11 wk	Hemoglobin, hematocrit, MCV, MCH, MCHC, ferritin, transferrin saturation
Perrone et al. <sup>42</sup>	Italy	4–11	Not anemic, iron status unknown	Iron polymaltose (12.5 mg elemental iron)	Placebo	Total: 16; iron: 8; placebo: 8	1 yr	Ferritin, zinc, serum copper, z scores for weight and height
Pollitt et al. <sup>43,44</sup>	Thailand	9–11	Subgroups: anemic, not anemic, iron deficient, not iron deficient, iron deficient anemia, iron deficient not anemic, nonanemic noniron deficient	Ferrous sulfate (50 mg/100 mg) + albendazole	Placebo + albendazole	Total: 1358; iron: 679; placebo: 679	16 wk	Hemoglobin, ferritin, transferrin saturation, cognitive scores
Powers et al. <sup>45</sup>	Gambia	4–12	Anemia and iron status unknown	Ferrous sulfate (200 mg)	Placebo	Total: 50; iron: 25; placebo: 25	6 wk	Hemoglobin, hematocrit, MCV, MCH, serum iron
Richard et al. <sup>46</sup>	Peru	0.5–15	Anemia and iron status unknown	Ferrous sulfate (15 mg elemental iron) ± zinc	Placebo ± zinc	Total: 836; iron: 208; placebo: 209; iron + zinc: 210; placebo + zinc: 209	7 mo	Hemoglobin, zinc, malaria, lower respiratory tract infection, diarrhea
Rico et al. <sup>47–50</sup>	Mexico	6–8	Anemia and iron status unknown	Ferrous fumarate (10 mg elemental iron) ± zinc	Placebo or zinc	Total: 590; iron: 141; placebo: 151; iron + zinc: 152; placebo + zinc: 146	6 mo	Hemoglobin, ferritin, zinc, lead, height, weight, cognitive scores
Rosales et al. <sup>51</sup>	Guatemala	8–11	Anemia and iron status unknown	Iron citrate (20 mg elemental iron) ± zinc	Placebo ± zinc	Total: 64; iron: 15; placebo: 15; iron + zinc: 19; placebo + zinc: 15	8 wk	Iron deficiency, zinc deficiency, hemoglobin, ferritin, zinc, CRP
Sen et al. <sup>52,53</sup>	India	9–13	Anemia and iron status unknown	Unknown (100 mg elemental iron) + folic acid	Control	Total: 110; iron: 59; placebo: 41	1 yr	Hemoglobin, cognitive scores, maximum steps climbed, recovery time
Sen et al. <sup>54</sup>	India	9–13	Iron status unknown. Subgroups: anemic, not anemic	Unknown (elemental iron 100 mg)	Control	Total: 100; iron: 59; placebo: 41	1 yr	Hemoglobin

Continued

**Table 1 (part 3 of 3):** Studies included in the systematic review\*

Study	Country	Age at recruitment, yr	Baseline anemia, iron deficiency status	Intervention	Control	No. randomized	Duration	Outcomes
Seshadri et al. (study 1) <sup>28,55</sup>	India	5–8	Anemia and iron status unknown	Unknown (20 mg elemental iron) + folic acid	Control	Total: 97; iron: 63; placebo: 34	60 d	Hemoglobin, IQ (performance, verbal total)
Seshadri et al. (study 2) <sup>28,55</sup>	India	5–6	Anemic, iron status unknown	Unknown (40 mg elemental iron) + folic acid	Placebo	Total: 28; iron: 14; placebo: 14	60 d	Hemoglobin, IQ (performance, verbal, total)
Seshadri et al. (study 4) <sup>28</sup>	India	8–15	Iron status unknown. Subgroups: anemic, nonanemic	Unknown (60 mg elemental iron)	Placebo	Total: 130; iron: 65; placebo: 65	12 mo	Anemia, iron deficiency, hemoglobin, IQ (performance, verbal, total)
Soemantri et al. <sup>56</sup>	Indonesia	8.1–11.6	Subgroups: anemic, not anemic, iron deficient, not iron deficient, iron deficient anemia, iron deficient not anemic, nonanemic noniron deficient	Ferrous sulfate (10 mg/kg/d)	Placebo	Total: 130; iron: 71; placebo: 59	3 mo	Hemoglobin, transferrin saturation, cognitive scores
Sungthong et al. <sup>57,58</sup>	Thailand	6–13	Anemia and iron status unknown	Ferrous sulfate (300 mg) + albendazole	Placebo + albendazole	Total: 263; iron: 140; placebo: 123	16 wk	Hemoglobin, ferritin, height, weight, z scores for height and weight, cognitive scores

Note: CRP = C-reactive protein, GI = gastrointestinal, IQ = intelligence quotient, MCH = mean cell hemoglobin, MCHC = mean cell hemoglobin concentration, MCV = mean cell volume, URTI = upper respiratory tract infection.  
 \*Two included studies<sup>52,53,54</sup> used clustered randomization at the school level but the reported data were not clearly adjusted for clustering. We imputed the design effect for hemoglobin from a recent Cochrane review,<sup>8</sup> and the intra-cluster correlation coefficient for physical activity and cognitive assessment from other cluster RCTs reporting these outcomes;<sup>59,60</sup> sample size was then adjusted by dividing by the design effect.

benefit of iron supplementation on IQ scores (MD 5.47, 95% CI –3.24 to 4.18;  $p = 0.2$ ,  $I^2 = 97\%$ , 5 studies,  $n = 1874$ ). However, our subgroup analysis showed that children who had anemia at baseline had significant improvements in their IQ after iron supplementation compared with children who received the control (MD 4.55, 95% CI 0.16 to 8.94;  $p = 0.04$ ,  $I^2 = 28\%$ , 3 studies,  $n = 186$ ) (Appendix 3).

#### Other cognitive tests

Compared with children in the control group, children who received iron supplementation had superior performance on the maze test (MD 1.30, 95% CI 0.9 to 1.7;  $p < 0.001$ ,  $I^2 = 0\%$ , 4 studies,  $n = 288$ ) and clerical task scores (SMD 0.44, 95% CI 0.14 to 0.75;  $p = 0.005$ ,  $I^2 = 30\%$ , 4 studies,  $n = 290$ ). There was no difference between iron supplementation and control for other measures including digit span, visual memory score, math scores and language test scores (Appendix 2).

#### Growth

##### Height

Iron supplementation was not associated with a difference in absolute height. However, children who received iron supplements experienced a

small advantage in height-for-age (z score) (Figure 3). When we included only studies with a low risk of bias, the benefit of iron supplementation was still apparent (MD 0.12, 95% CI 0.00 to 0.23;  $p = 0.05$ , 2 studies,  $n = 922$ ).

#### Weight

There was no difference in absolute weight or weight-for-age (z score) between children who received iron or control. One study included subgroups of children with iron deficiency anemia and children without iron deficiency or anemia.<sup>24</sup> The authors reported that the children with iron deficiency anemia who received iron had a significant improvement in weight-for-age (MD 0.13, 95% CI 0.04 to 0.22;  $p = 0.005$ ,  $n = 78$ ); this effect was not seen among children without anemia (MD 0.18, 95% CI –0.24 to 0.60;  $p = 0.4$ ,  $n = 57$ ).<sup>24</sup>

#### Safety

##### Malaria infection

The prevalence of malaria parasitemia was similar between iron and control groups (RR 1.10, 95% CI 0.94 to 1.29;  $p = 0.9$ ,  $I^2 = 93\%$ , 4 studies,  $n = 919$ ) (Appendix 4, available at [www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.130628/-DC1](http://www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.130628/-DC1)). There were no differences in the rate of positive

tests for all malaria infections (rate ratio 1.05, 95% CI 0.90 to 1.21;  $p = 0.6$ ,  $F = 0\%$ , 4 studies,  $n = 1818$ ), *Plasmodium falciparum* infections (rate ratio 1.02, 95% CI 0.81 to 1.29;  $p = 0.9$ ,  $F = 0\%$ , 2 studies,  $n = 1136$ ) and *Plasmodium vivax* infections (rate ratio 1.07, 95% CI 0.85 to 1.34;  $p = 0.6$ ,  $F = 0\%$ , 2 studies,  $n = 1136$ ).

#### Other infections

De Silva and colleagues<sup>25</sup> reported fewer respiratory tract infections (MD  $-0.56$ , 95% CI  $-0.82$  to  $-0.30$ ;  $p < 0.001$ ,  $n = 363$ ) and fewer days off sick with respiratory tract infections (MD  $-7.04$ ,

95% CI  $-9.75$  to  $-4.33$ ;  $p < 0.001$ ,  $n = 363$ ) among children who received iron supplementation. Richard and colleagues<sup>46</sup> reported no difference in the rate of acute respiratory tract infections between groups (rate ratio 0.91, 95% CI 0.65–1.27;  $p = 0.6$ ,  $n = 836$ ).

#### Gastrointestinal adverse effects

There was no difference in the number of children with gastrointestinal upset, constipation, vomiting or diarrhea or the rates of diarrhea between children who received iron supplementation or control (Appendix 4).

**Table 2:** Analysis of the risk of bias among the included studies

Study	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding (participants and personnel; performance bias)	Blinding (outcome assessment; detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)
Ayoya et al. <sup>17,18</sup>	Yes	Yes	Yes	Yes	Yes	Unknown
Berger et al. <sup>19</sup>	Unknown	Unknown	Yes	Yes	Yes	Unknown
Boivin et al. <sup>20</sup>	Unknown	Unknown	Unknown	Unknown	Yes	Unknown
Bradfield et al. <sup>21</sup>	Unknown	Yes	Yes	Yes	Unknown	Unknown
Charoenlarp <sup>22</sup>	Unknown	Unknown	Yes	Yes	Yes	Unknown
Charoenlarp et al. <sup>23</sup>	Unknown	Unknown	Yes	Yes	Unknown	Unknown
Chwang et al. <sup>24</sup>	Unknown	Unknown	Unknown	Yes	Unknown	Unknown
de Silva et al. <sup>25</sup>	Unknown	Unknown	Yes	Yes	Yes	Unknown
Fallahi et al. <sup>26</sup>	Unknown	Unknown	Yes	Yes	Yes	Unknown
Gebreselassie <sup>27</sup>	Yes	Yes	Yes	Yes	Yes	Yes
Gopaldas et al. <sup>28-31</sup>	Unknown	Unknown	Yes	Yes	Unknown	Unknown
Greisen <sup>32</sup>	No	Unknown	Unknown	Yes	Yes	Unknown
Harvey et al. <sup>33</sup>	Unknown	Unknown	Unknown	Unknown	Yes	Unknown
Kashyap et al. <sup>34</sup>	Unknown	Unknown	Unknown	Unknown	Yes	Unknown
Konofal et al. <sup>35</sup>	Yes	Unknown	Yes	Yes	Yes	Unknown
Latham et al. <sup>36</sup>	Unknown	Unknown	Unknown	Yes	Yes	Unknown
Lawless et al. <sup>37</sup>	Yes	Yes	Yes	Yes	Unknown	Unknown
Nchito et al. <sup>38-40</sup>	Unknown	No	Unknown	Yes	No	Unknown
Paracha et al. <sup>41</sup>	Unknown	Unknown	Unknown	Yes	Yes	Unknown
Perrone et al. <sup>42</sup>	Unknown	Unknown	Unknown	Yes	Yes	Unknown
Pollitt et al. <sup>43,44</sup>	Unknown	Unknown	Yes	Yes	No	No
Powers et al. <sup>45</sup>	Unknown	Unknown	Yes	Yes	Yes	Unknown
Richard et al. <sup>46</sup>	Yes	Yes	Yes	Yes	Yes	Unknown
Rico et al. <sup>47-50</sup>	Yes	Unknown	Yes	Yes	Yes	Yes
Rosales et al. <sup>51</sup>	Yes	Yes	Yes	Yes	Yes	Unknown
Sen et al. <sup>52,53</sup>	No	No	No	No	Unknown	Unknown
Sen et al. <sup>54</sup>	Unknown	Unknown	No	No	Unknown	Unknown
Seshadri et al. (study 1) <sup>28,55</sup>	Unknown	Unknown	No	No	Unknown	Unknown
Seshadri et al. (study 2) <sup>28,55</sup>	Yes	Unknown	Unknown	Unknown	Unknown	Unknown
Seshadri et al. (study 4) <sup>28</sup>	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown
Soemantri et al. <sup>56</sup>	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown
Sungthong et al. <sup>57,58</sup>	Yes	Yes	Unknown	Unknown	Yes	Unknown

### Morbidity and mortality

Chwang and colleagues<sup>24</sup> reported that children who received iron supplementation had a lower overall morbidity score than children in the control group (MD -2.73, 95% CI -3.56 to -1.90;  $p < 0.001$ ,  $n = 119$ ). Mortality, admission to hospital, nonregular clinic visits and morbidity from iron poisoning were not reported in any study.

### Hematologic and iron indices

Iron supplementation improved hemoglobin and ferritin levels, anemia and iron deficiency (Table 3, Appendix 2, Appendix 5 [available at [www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.130628/-/DC1](http://www.cmaj.ca/lookup/suppl/doi:10.1503/cmaj.130628/-/DC1)]). Funnel plots for hemoglobin and ferritin levels did not show any evidence of publication bias (Appendix 5). When we included only the 4 studies at low risk of bias, the beneficial effects of iron supplementation were no longer observed for hemoglobin, ferritin or iron deficiency.

Only 2 studies reported iron deficiency anemia.<sup>17,18,57,58</sup> Iron supplementation reduced the prevalence of anemia (RR 0.12, 95% CI 0.02 to 0.66;  $p = 0.01$ , 334 children).

### Other micronutrients

There were no differences in the prevalence of zinc deficiency<sup>51</sup> or in the serum levels of zinc,<sup>51</sup> lead,<sup>47-50</sup> retinol<sup>26</sup> or copper.<sup>42</sup>

### Exercise performance

Gopaldas and colleagues<sup>28-31</sup> reported a reduction in pulse rate during the Harvard Step Test among children who received iron supplementation. However, Sen and Kanani<sup>52,53</sup> found no difference between the iron and control groups in the number of steps climbed or recovery time.

### Adherence

Twenty-one studies reported adherence. We were unable, however, to perform a meta-analysis because of differences in reporting. Five studies (835 children) used fully or partially supervised iron supplementation. Of the remaining studies, 4 (820 children) reported adherence of more than 90% in the control and iron supplementation groups,<sup>21,25,27,57,58</sup> and 7 studies (2071 children) reported more than 80% compliance in both groups.<sup>19,21,25,27,33,46,57,58</sup> Only 3 studies reported lower

**Table 3:** Effect of iron supplementation on primary outcomes among children aged 5–12 years

Outcome	No. of trials	No. of participants	Effect size (95% CI)	<i>p</i> for effect	<i>I</i> <sup>2</sup> , %
<b>Cognitive performance</b>					
Global cognitive performance*	9	2355	0.50 (0.11 to 0.90)	0.01	93
Intelligence quotient†	5	1874	5.47 (-3.24 to 14.18)	0.2	97
<b>Growth</b>					
Height†	5	1111	-0.37 (-1.14 to 0.40)	0.3	0
Z score for height†	5	1318	0.09 (0.01 to 0.17)	0.03	10
Weight†	5	1111	0.12 (-0.50 to 0.75)	0.7	0
Z score for weight†	5	1318	0.10 (-0.03 to 0.23)	0.1	70
<b>Hematology and iron indices</b>					
Hemoglobin, g/L†	28	6545	8.38 (6.21 to 10.56)	< 0.001	97
Anemia‡	7	1763	0.50 (0.39 to 0.64)	< 0.001	85
Iron deficiency‡	4	1020	0.21 (0.07 to 0.63)	0.006	85
Ferritin, g/L†	14	3612	28.45 (18.03 to 38.86)	< 0.001	98
<b>Other micronutrients</b>					
Serum zinc, mol/L †	5	1553	0.16 (-0.46 to 0.79)	0.6	76
<b>Safety</b>					
Malaria prevalence‡	4	919	1.10 (0.94 to 1.29)	0.2	0
Rates of positive malaria test (any species)§	4	1818	1.05 (0.90 to 1.21)	0.6	0
Gastrointestinal upset‡	4	576	1.30 (0.89 to 1.91)	0.2	0
Constipation‡	2	202	3.44 (0.66 to 19.68)	0.2	6
Vomiting‡	2	202	0.86 (0.13 to 5.67)	0.9	0
*Standardized mean difference. †Mean difference. ‡Risk ratio. §Rate ratio.					



adherence rates: Sen and Kanani<sup>52-54</sup> reported an adherence rate of 72%, and Nchito and colleagues<sup>38-40</sup> adherence of 50%. Six studies (2651 children) excluded participants who failed to take a predefined percentage of supplements (ranging

from 66% to 90% of total tablets).<sup>22,23,44,47-51,55</sup> Five studies (1894 children) compared adherence between children taking iron and control; they reported similar proportions of children taking at least a specified percentage of tablets.<sup>19,21,25,27,33</sup>




Figure 2: Forest plots for global cognitive performance and cognitive performance by anemia status. \*Inverse variance, random effects. †Author-adapted scales included clerical task test, visual memory test, digit span test and maze test. Note: CI = confidence interval, TONI-II = Test of Non-verbal Intelligence, 2nd edition, SD = standard deviation, SMD = standardized mean difference, WISC = Wechsler Intelligence Scale for Children.

## Interpretation

Daily supplementation provides the highest dose of iron of any nonparenteral approach and is a commonly recommended clinical and public health strategy for the prevention and treatment of anemia.<sup>2</sup> Potential strategies for anemia control include improved dietary iron by improving dietary diversity or fortification of staple foods and condiments, home-based fortification with multiple micronutrients, iron supplementation and control of helminthic infection.<sup>6</sup> Our data are chiefly extrapolatable to iron supplementation interventions; alternative approaches to improving iron stores such as food fortification are under evaluation by other systematic reviews.<sup>61</sup>

In this review, we found evidence of a benefit of iron supplementation on cognitive performance among primary-school-aged children, including on IQ among children with anemia. Iron may also improve growth. Daily iron supplementation decreased the prevalence of anemia by about 50% and reduced the prevalence of iron deficiency by 79%. Iron supplements appear to be well-tolerated, but safety data are limited.

Previous systematic reviews have evaluated the efficacy of iron delivered by multiple approaches (i.e., supplementation, parenteral administration, fortification) among children of broad age ranges; outcomes included hemoglobin level,<sup>62</sup> mental and physical development,<sup>63</sup> growth,<sup>64</sup> exercise capacity,<sup>65</sup> malaria<sup>8</sup> and infectious diseases<sup>66</sup> (but not other adverse effects). Iron supplementation among iron-deficient children without anemia has also been reviewed.<sup>67</sup> A Cochrane review of intermittent iron supplementation in children found

insufficient studies reporting cognitive outcomes or adverse effects.<sup>15</sup>

Two previous reviews have evaluated the effects of iron on cognitive outcomes among primary-school-aged children. Sachdev and colleagues<sup>63</sup> reported improvements in overall cognitive performance (SMD 0.30) and IQ (SMD 0.41) among children who received parenteral or enteral iron (including fortification). Falkingham and colleagues<sup>68</sup> found no improvement in overall IQ from iron supplementation, but they reported improvements in IQ among participants with anemia (MD 2.5) and improvements in attention and concentration among adults and children older than 6 years of age.

Our search strategy enabled us to identify and include more eligible studies than either Falkingham and colleagues<sup>68</sup> (5 additional studies) or Sachdev and colleagues<sup>63</sup> (7 additional studies). We have improved on previous reviews by performing a broader search, providing a comprehensive evaluation of the benefits and safety, and addressing a specific, widely used strategy of iron administration among children at an age at which developmental benefits are critical.

## Limitations

Our conclusions are limited by poor reporting in the included studies: many studies did not report the randomization method (20 studies), allocation concealment (25 studies) or blinding (18 studies). Although biochemically measured outcomes (e.g., hematologic and iron indices) are unlikely to be affected by the blinding of outcome assessors, cognitive scores have the potential for performance and detection bias.

Our review is also limited by significant het-




Figure 3: Forest plot for height. Forest plot of the z scores for height. \*Inverse variance, random effects. Note: CI = confidence interval, SD = standard deviation.

erogeneity, which we addressed by use of random effects meta-analysis and predefined subgroup analyses. This heterogeneity appears to be mainly because of differences in baseline anemia status. Unfortunately, several important outcomes (particularly physical performance and growth) were reported by relatively few studies, and only 6 studies reported safety outcomes. Further randomized trials evaluating the nonhematologic effects of iron, especially outcomes such as cognitive performance, growth and safety, are needed.

## Conclusion

In this review, we found evidence that iron supplementation benefits global cognitive performance. Anemia and iron deficiency were prevalent in the included studies (69% and 59%, respectively, in the control groups); thus, routine daily iron supplementation is likely to benefit cognitive performance in primary-school-aged children in developing settings where anemia is prevalent and testing hemoglobin before iron supplementation may not be feasible. In developed settings, our data support ensuring that anemia is detected and treated. Reductions in the prevalence of anemia and iron deficiency reported here should guide the targets of anemia-control programs. Cognitive performance is associated with educational achievement,<sup>69</sup> future income and productivity.<sup>70</sup> Daily iron supplementation could benefit educational attainment and economic potential at the individual level and, in settings where anemia is prevalent, population level.

## References

1. Benoist B, McLean E, Egli I, et al. *Worldwide prevalence of anaemia 1993–2005*. Geneva (Switzerland): World Health Organization; 2008.
2. *Iron deficiency: assessment, prevention, and control: a guide for programme managers*. Geneva (Switzerland): World Health Organization and United Nations Children's Fund/United Nations University; 2001.
3. Stoltzfus RJ, Chwya HM, Tielsch JM, et al. Epidemiology of iron deficiency anemia in Zanzibari schoolchildren: the importance of hookworms. *Am J Clin Nutr* 1997;65:153-9.
4. Christofides A, Schauer C, Zlotkin SH. Iron deficiency and anemia prevalence and associated etiologic risk factors in First Nations and Inuit communities in Northern Ontario and Nunavut. *Can J Public Health* 2005;96:304-7.
5. Lawson MS, Thomas M, Hardiman A. Iron status of Asian children aged 2 years living in England. *Arch Dis Child* 1998;78:420-6.
6. Pasricha SR, Drakesmith H, Black J, et al. Control of iron deficiency anemia in low and middle-income countries. *Blood* 2013; 121:2607-17.
7. Cooper M, Greene-Finestone L, Lowell H, et al. *Iron sufficiency of Canadians*. Ottawa (ON): Statistics Canada; 2012. Cat. no. 82-003-X.
8. Okebe JU, Yahav D, Shbita R.M.P. Oral iron supplements for children in malaria-endemic areas. *Cochrane Database Syst Rev* 2011;(10):CD006589.
9. Stoltzfus RJ, Mullany L, Black RE. Iron deficiency anaemia. In: Ezzati M, Lopez AD, Rodgers A, et al., editors. *Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors*. Geneva (Switzerland): World Health Organization; 2004.
10. Grantham-McGregor S, Ani C. A review of studies on the effect of iron deficiency on cognitive development in children. *J Nutr* 2001;131:949S-666S.
11. McCann JC, Ames BN. An overview of evidence for a causal relation between iron deficiency during development and deficits in cognitive or behavioral function. *Am J Clin Nutr* 2007;85:931-45.
12. Goswami U, Bryant P. *Children's cognitive development and learning: primary review research survey 2/1a*. Cambridge (UK): University of Cambridge, Faculty of Education; 2007.
13. Pena-Rosas JP, De-Regil LM, Rogers LM, et al. Translating research into action: WHO evidence-informed guidelines for safe and effective micronutrient interventions. *J Nutr* 2012;142: 197S-204S.
14. Higgins JP, Altman DG, Sterne JA. Assessing risk of bias in included studies. In: Higgins JPT, Green S, editors. *Cochrane handbook for systematic reviews of interventions*. Oxford (UK): The Cochrane Collaboration; 2011.
15. De-Regil LM, Jefferds ME, Sylvestry AC, et al. Intermittent iron supplementation for improving nutrition and development in children under 12 years of age. *Cochrane Database Syst Rev* 2011;(12):CD009085.
16. *Introduction to meta-analysis*. Borenstein M, Hedges L, Rothstein H. Chichester (UK): John Wiley & Sons; 2007.
17. Ayoya MA, Spiekermann-Brouwer GM, Traoré AK, et al. Effect on school attendance and performance of iron and multiple micronutrients as adjunct to drug treatment of *Schistosoma*-infected anemic schoolchildren. *Food Nutr Bull* 2012;33:235-41.
18. Ayoya MA, Spiekermann-Brouwer GM, Traoré AK, Stoltzfus RJ, et al. Multiple micronutrients including iron are not more effective than iron alone for improving hemoglobin and iron status of Malian school children. *J Nutr* 2009;139:1972-9.
19. Berger J, Aguayo VM, Tellez W, et al. Weekly iron supplementation is as effective as 5 day per week iron supplementation in Bolivian school children living at high altitude. *Eur J Clin Nutr* 1997;51:381-6.
20. Boivin MJ, Giordani B. Improvements in cognitive performance for schoolchildren in Zaire, Africa, following an iron supplement and treatment for intestinal parasites. *J Pediatr Psychol* 1993;18:249-64.
21. Bradfield RB, Jensen MV, Gonzales L, et al. Effect of low-level iron and vitamin supplementation on a tropical anemia. *Am J Clin Nutr* 1968;21:57-67.
22. Charoenlarp P. Effect of iron and folate supplementation on haematocrit levels of school children in a rural area of Central Thailand. *Southeast Asian J Trop Med Public Health* 1973;4:588-92.
23. Charoenlarp P, Pholpothi T, Chatpunyaporn P, et al. The effect of riboflavin on the hematologic changes in iron supplementation of schoolchildren. *Southeast Asian J Trop Med Public Health* 1980; 11:97-103.
24. Chwang LC, Soemantri AG, Pollitt E. Iron supplementation and physical growth of rural Indonesian children. *Am J Clin Nutr* 1988; 47:496-501.
25. de Silva A, Atukorala S, Weerasinghe I, et al. Iron supplementation improves iron status and reduces morbidity in children with or without upper respiratory tract infections: a randomized controlled study in Colombo, Sri Lanka. *Am J Clin Nutr* 2003;77:234-41.
26. Fallahi E, Kimiagar M, Nazari A, et al. Effect of zinc and iron supplementation on indicators of iron, zinc and vitamin A status of primary school children. *Pak J Biol Sci* 2007;10:1088-92.
27. Gebreselassie H. *Iron supplementation and malaria infection: results of a randomized controlled field trial* [dissertation]. Montreal (QC): McGill University, 1996.
28. Seshadri S, Gopaldas T. Impact of iron supplementation on cognitive functions in preschool and school-aged children: the Indian experience. *Am J Clin Nutr* 1989;50:675-84, discussion 85-6.
29. Gopaldas T, Kale M. Prophylactic iron supplementation for underprivileged school boys. I. Two levels of dosing and efficacy of teacher-distributions. *Indian Pediatr* 1985;22:731-6.
30. Gopaldas T, Kale M, Bhardwaj P. Prophylactic iron supplementation for underprivileged school boys. III. Impact on submaximal work capacity. *Indian Pediatr* 1985;22:745-52.
31. Gopaldas T, Kale M, Bhardwaj P. Prophylactic iron supplementation for underprivileged school boys. II. Impact on selected tests of cognitive function. *Indian Pediatr* 1985;22:737-43.
32. Greisen G. Mild anaemia in African school children: effect on running performance and an intervention trial. *Acta Paediatr Scand* 1986;75:662-7.
33. Harvey PWJ, Heywood PF, Nesheim MC, et al. The effect of iron therapy on malarial infection in Papua New Guinean schoolchildren. *Am J Trop Med Hyg* 1989;40:12-8.
34. Kashyap P, Gopaldas T. Impact of hematinic supplementation on cognitive function in underprivileged school girls (8–15 yrs of age). *Nutr Res* 1987;7:1117-26.
35. Konofal E, Lecendreux M, Deron J, et al. Effects of iron supplementation on attention deficit hyperactivity disorder in children. *Pediatr Neurol* 2008;38:20-6.
36. Latham MC, Stephenson LS, Kinoti SN, et al. Improvements in growth following iron supplementation in young Kenyan school children. *Nutrition* 1990;6:159-65.

37. Lawless JW, Latham MC, Stephenson LS, et al. Iron supplementation improves appetite and growth in anemic Kenyan primary school children. *J Nutr* 1994;124:645-54.
38. Nchito M, Friis H, Michaelsen KF, et al. Iron supplementation increases small intestine permeability in primary schoolchildren in Lusaka, Zambia. *Trans R Soc Trop Med Hyg* 2006;100:791-4.
39. Nchito M, Geissler PW, Mubila L, et al. Effects of iron and multi-micronutrient supplementation on geophagy: a two-by-two factorial study among Zambian schoolchildren in Lusaka. *Trans R Soc Trop Med Hyg* 2004;98:218-27.
40. Nchito M, Geissler PW, Mubila L, et al. The effect of iron and multi-micronutrient supplementation on *Ascaris lumbricoides* reinfection among Zambian schoolchildren. *Trans R Soc Trop Med Hyg* 2009;103:229-36.
41. Paracha PI, Khan SM, Ahmad I, et al. Effect of iron supplementation on biochemical indices of iron status in selected pre-adolescent schoolgirls in North West Frontier Province, Pakistan. *Asia Pac J Clin Nutr* 1993;2:177-81.
42. Perrone L, Salerno M, Gialanella G, et al. Long-term zinc and iron supplementation in children of short stature: effect of growth and on trace element content in tissues. *J Trace Elem Med Biol* 1999;13:51-6.
43. Hathirat P, Valyasevi A, Kotchabhakdi NJ, et al. Effects of an iron supplementation trial on the Fe status of Thai schoolchildren. *Br J Nutr* 1992;68:245-52.
44. Pollitt E, Hathirat P, Kotchabhakdi NJ, et al. Iron deficiency and educational achievement in Thailand. *Am J Clin Nutr* 1989;50:687-96, discussion 96-7.
45. Powers HJ, Bates CJ, Prentice AM, et al. The relative effectiveness of iron and iron with riboflavin in correcting a microcytic anaemia in men and children in rural Gambia. *Hum Nutr Clin Nutr* 1983;37:413-25.
46. Richard SA, Zavaleta N, Caulfield LE, et al. Zinc and iron supplementation and malaria, diarrhea, and respiratory infections in children in the Peruvian Amazon. *Am J Trop Med Hyg* 2006;75:126-32.
47. Kordas K. *Effects of iron and zinc supplementation on cognitive function, achievement and behavior of lead-exposed Mexican school children*. Baltimore (MD): Johns Hopkins University; 2003.
48. Kordas K, Stoltzfus RJ, Lopez P, et al. Iron and zinc supplementation does not improve parent or teacher ratings of behavior in first grade Mexican children exposed to lead. *J Pediatr* 2005;147:632-9.
49. Rico JA, Kordas K, Lopez P, et al. Efficacy of iron and/or zinc supplementation on cognitive performance of lead-exposed Mexican schoolchildren: a randomized, placebo-controlled trial. *Pediatrics* 2006;117:e518-27.
50. Rosado JL, Lopez P, Kordas K, et al. Iron and/or zinc supplementation did not reduce blood lead concentrations in children in a randomized, placebo-controlled trial. *J Nutr* 2006;136:2378-83.
51. Rosales FJ, Kang Y, Pfeiffer B, et al. Twice the recommended daily allowance of iron is associated with an increase in plasma alpha-1 antichymotrypsin concentrations in Guatemalan school-aged children. *Nutr Res* 2004;24:875-87.
52. Sen A, Kanani SJ. Impact of iron-folic acid supplementation on cognitive abilities of school girls in Vadodara. *Indian Pediatr* 2009;46:137-43.
53. Sen A, Kanani SJ. Physical work capacity of young underprivileged school girls impact of daily vs intermittent iron folic acid supplementation: a randomized controlled trial. *Indian Pediatr* 2009;46:849-54.
54. Sen A, Kanani S. Intermittent iron folate supplementation: impact on hematinic status and growth of school girls. *ISRN Hematol* 2012:482153.
55. Seshadri S, Hirode K, Naik P, et al. Behavioural responses of young anaemic Indian children to iron-folic acid supplements. *Br J Nutr* 1982;48:233-40.
56. Soemantri AG. Preliminary findings on iron supplementation and learning achievement of rural Indonesian children. *Am J Clin Nutr* 1989;50:698-701; discussion -2.
57. Sungthong R, Mo-Suwan L, Chongsuvivatwong V, et al. Once weekly is superior to daily iron supplementation on height gain but not on hematological improvement among schoolchildren in Thailand. *J Nutr* 2002;132:418-22.
58. Sungthong R, Mo-suwan L, Chongsuvivatwong V, et al. Once-weekly and 5-days a week iron supplementation differentially affect cognitive function but not school performance in Thai children. *J Nutr* 2004;134:2349-54.
59. Hedges L, Hedberg EC. Intraclass correlations for planning group randomized experiments in rural education. *J Res Rural Educ* 2007;22.
60. Kriemler S, Schindler C, Meyer U, et al. Effect of school based physical activity programme (KISS) on fitness and adiposity in primary schoolchildren: cluster randomized controlled trial. *BMJ* 2010;340:c785.
61. Pasricha S-R, De-Regil LM, Garcia-Casal MN, et al. Fortification of maize flour with iron for preventing anaemia and iron deficiency in populations. *Cochrane Database Syst Rev* 2012;(11)CD010187.
62. Gera T, Sachdev HP, Nestel P, et al. Effect of iron supplementation on haemoglobin response in children: systematic review of randomised controlled trials. *J Pediatr Gastroenterol Nutr* 2007;44:468-86.
63. Sachdev H, Gera T, Nestal P. Effect of iron supplementation on mental and motor development in children: systematic review of randomized controlled trials. *Public Health Nutr* 2005;8:117-32.
64. Sachdev H, Gera T, Nestel P. Effect of iron supplementation on physical growth in children: systematic review of randomized controlled trials. *Public Health Nutr* 2006;9:904-20.
65. Gera T, Sachdev S, Nestal P. Effect of iron supplementation on physical performance in children and adolescents: systematic review of randomized controlled trials. *Indian Pediatr* 2007;44:15-24.
66. Gera T, Sachdev HP. Effect of iron supplementation on incidence of infectious illness in children: systematic review. *BMJ* 2002;325:1142.
67. Abdullah K, Kendzerska T, Shah P, et al. Efficacy of oral iron therapy in improving the developmental outcome of pre-school children with non-anaemic iron deficiency: a systematic review. *Public Health Nutr* 2013;16:1497-506.
68. Falkingham M, Abdelhamid A, Curtis P, et al. The effects of oral iron supplementation on cognition in older children and adults: a systematic review and meta-analysis. *Nutr J* 2010;9:4.
69. Taubman P, Wales T. *Mental ability and higher education attainment in the 20th century*. New York (NY): McGraw-Hill; 1972.
70. Horton S, Ross J. The economics of iron deficiency. *Food Policy* 2003;28:51-75.

**Affiliations:** Department of Clinical Haematology (Low), The Alfred Hospital, Prahran; Thalassaemia Service (Farrell, Pasricha), Southern Health, Clayton; Department of Medicine (Biggs, Pasricha), The Royal Melbourne Hospital, Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne, Carlton; Nossal Institute for Global Health (Pasricha), Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne, Carlton, Victoria, Australia.

**Contributors:** Sant-Rayn Pasricha and Michael Low conceived the systematic review design, developed the search strategy, undertook the search, screened papers for eligibility, extracted data, analyzed the results and wrote the manuscript. Ann Farrell screened papers for eligibility, extracted data and wrote the manuscript. Beverley-Ann Biggs conceived the systematic review design, interpreted the data and revised the manuscript critically for important intellectual content. All of the authors approved the final version submitted for publication.

**Funding:** Sant-Rayn Pasricha was supported by a Victoria Fellowship from the Government of Victoria; a CRB Blackburn Scholarship from the Royal Australasian College of Physicians; an Overseas Research Experience Scholarship from the University of Melbourne; and a National Health and Medical Research Council CJ Martin Early Career Fellowship. Michael Low was supported by an Alfred Research Trust Grant. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Acknowledgements:** The authors thank the staff in the Department of Nutrition for Health and Development and the Library and Information Networks for Knowledge, World Health Organization, Geneva, for their guidance and assistance in planning and implementing this study. They also thank Patrick Condron, librarian at The University of Melbourne, for assistance with the electronic search strategies, and Maggie Low, University of Adelaide, and Dr. Denise Lee, Alfred Hospital, for assistance with acquiring the publications.