



Effects of H_2O_2 , Fe^{2+} and Fe^{3+} on curcumin-induced chromosomal aberrations in CHO cells

Lusânia Maria Greggi Antunes¹, Maria Cristina Paiva Araújo^{2†}, Francisca da Luz Dias¹ and Catarina Satie Takahashi^{2,3}

¹Faculdade de Medicina do Triângulo Mineiro, Departamento de Ciências Biológicas, Uberaba, MG, Brazil.

²Universidade de São Paulo, Faculdade de Medicina de Ribeirão Preto, Departamento de Genética, Ribeirão Preto, SP, Brazil.

³Universidade de São Paulo, Faculdade de Filosofia Ciências e Letras de Ribeirão Preto, Departamento de Biologia, Ribeirão Preto, SP, Brazil.

Abstract

The effects of H_2O_2 , Fe^{2+} and Fe^{3+} on curcumin-induced clastogenicity were evaluated in CHO cells. Curcumin combined with H_2O_2 did not increase the chromosomal aberrations more than expected based on a simple additive effect. In contrast, the combination of curcumin-Fe significantly decreased the total number of chromosomal aberrations and the number of abnormal metaphases. The clastogenicity of curcumin may be related to its pro-oxidant properties and its ability to generate free radicals.

Key words: CHO, chromosomal aberrations, curcumin, hydrogen peroxide, mutagenesis.

Received: April 26, 2004; Accepted: August 18, 2004.

Experiments *in vitro* and epidemiological studies have shown that some compounds present in the diet have antimutagenic and anticarcinogenic properties (Surh and Ferguson, 2003). Turmeric, a spice obtained from the rhizome of *Curcuma longa* Linn (Zingiberaceae), and its active principle curcumin, have been studied for their ability to protect cells from DNA damage (Polasa *et al.*, 2004). Curcumin, widely used as a coloring for foods, has antioxidant, antimutagenic and anticarcinogenic properties (Nagabhushan and Bhide, 1992; Aggarwal *et al.*, 2003).

Curcumin reduces chromosomal aberrations in rodent bone marrow cells exposed to gamma-radiation or treated with the antitumor drug cisplatin (Abraham *et al.*, 1993; Antunes *et al.*, 2000). However, clastogenic effects of curcumin have also been reported. An increase in the frequency of chromosomal damage has been seen in mice and rats treated acutely and chronically with curcumin (Giri *et al.*, 1990; Mukhopadhyay *et al.*, 1998), and curcumin is mutagenic in cultured Chinese hamster fibroblasts (Ishidate *et al.*, 1984). At concentrations up to 10 $\mu\text{g}/\text{mL}$, curcumin is

mutagenic in cultured CHO cells and, when combined with the antitumor drugs bleomycin or doxorubicin, the frequency of chromosomal aberrations is markedly increased (Araújo *et al.*, 1999; Antunes *et al.*, 1999).

The clastogenicity of curcumin in mammalian cells *in vitro* has been attributed to a pro-oxidant action of this compound (Araújo *et al.*, 1999). In agreement with this proposed mechanism, thiourea, a hydroxyl radical scavenger, significantly inhibited the increase in chromosomal aberrations in CHO cells treated with curcumin (Araújo *et al.*, 2001). Thus, hydroxyl radicals generated by curcumin could contribute to clastogenicity *in vitro*.

A combination of Fe accessibility and high production of H_2O_2 and O_2^- results in a pro-oxidant state within cells (Meneghini, 1997). The molecular structure of curcumin is susceptible to auto-oxidation in the presence of oxygen and transition metal ions (Sahu and Washington, 1992). Indeed, curcumin is a good scavenger of H_2O_2 at high concentrations, but at low concentrations it activates the Fenton reaction to increase the production of H_2O_2 (Kuchandy and Rao, 1990).

Although, there is evidence indicating that metal ions are involved in curcumin-induced mutagenicity, there is still little on this subject in the literature. There have been no studies of the effects of H_2O_2 on curcumin-induced chro-

Send correspondence to Lusânia Maria Greggi Antunes. Faculdade de Medicina do Triângulo Mineiro, Departamento de Ciências Biológicas, Praça Manoel Terra 330, 38015-050 Uberaba, MG, Brazil. E-mail: lusaantunes@yahoo.com.br.

†Deceased June 21, 2001.

mosomal aberrations in CHO cells. To examine the hypothesis that the generation of free radicals is involved in curcumin-induced clastogenicity, CHO cells were treated with a pre-determined clastogenic concentration of curcumin alone or in combination with H₂O₂, Fe²⁺ or Fe³⁺.

Curcumin (diferuloylmethane; CAS No. 458-37-7) was purchased from Sigma Chemicals Co. (St. Louis, MO). DMSO and FeCl₃ were purchased from Merck (Darmstadt, F.R.G.). FeSO₄ and H₂O₂ were obtained from Reagen (Brazil). Dulbecco's modified Eagle's medium (DMEM) and HAM-F10 were purchased from Gibco (Invitrogen Corporation, USA). Fetal calf serum (FCS) was obtained from Cultilab (Campinas, SP, Brazil). Chinese hamster ovary cells (CHO-9) were grown as monolayers at 37 °C in 25-cm² flasks containing HAM-F10 plus DMEM (1:1 ratio), supplemented with 10% FCS, penicillin (0.06 mg/mL) and streptomycin (0.1 mg/mL). For all experiments, exponentially growing cells were seeded at a density of 1 x 10⁶/5 mL flask. Curcumin was dissolved in 0.5% DMSO. CHO cells were incubated for 14 h with H₂O₂ (1.7, 3.4 or 6.8 µg/mL), FeCl₃ (1.25, 2.5 or 5.0 µg/mL) or FeSO₄ (1.25, 2.5 or 5.0 µg/mL) in the absence or presence of curcumin.

Colcemid (0.1 µg/mL) was added to the culture medium 2 h before fixation of the cells. Each experiment was repeated three times and 300 metaphases (100 in each experiment) were analyzed per treatment to assess the frequencies of chromosomal aberrations. The mitotic index (MI) was defined as the percentage of metaphases in 3000 cells analyzed per treatment. The differences in the number of abnormal metaphases, total number of chromosomal aberrations, and mitotic indices in the absence and presence of curcumin were compared by analysis of variance (ANOVA).

The effects of H₂O₂, Fe²⁺ and Fe³⁺ on curcumin-induced chromosomal aberrations in CHO cells are shown in Tables 1 and 2. Curcumin was not cytotoxic at up to

15 µg/mL. However, as expected, there was a significant (p < 0.05) increase in the total number of chromosomal aberrations and in the number of abnormal metaphases after treatment with curcumin.

Curcumin readily penetrates into the cytoplasm and can accumulate in membrane structures (Joe *et al.*, 2004), primarily because of the molecule's structure, which consists of two isomers, *i.e.* the β-diketone and enol forms. The diketone form is a potent ligand for metals such as iron (Sun *et al.*, 2002). In the presence of copper or chromium, curcumin becomes a pro-oxidant and damages DNA (Ahsan and Hadi, 1998). The antioxidant/pro-oxidant action of phenolic compounds depends on factors such as metal reducing potential and chelating behavior (Decker, 1997).

Cultured CHO cells treated with different concentrations of H₂O₂ alone (Table 1) showed a significant (p < 0.05) increase in chromosomal aberrations at all concentrations tested. There was a concentration-dependent response in the total number of chromosomal aberrations and in the number of abnormal metaphases in cultures treated with H₂O₂. The most frequently detected aberrations were chromatid breaks followed by chromosomal breaks. At the highest H₂O₂ concentration, a significant (p < 0.05) decrease in the mitotic index was also observed when compared to control cultures. In combined treatments of curcumin and H₂O₂, all of the concentrations of H₂O₂ tested increased the total number of chromosomal aberrations induced by curcumin (Table 1). However, this combination did not increase the frequency of chromosomal aberrations beyond what was expected from a simple additive effect when compared to H₂O₂ and curcumin alone (H₂O₂: 144 at the highest concentration; curcumin: 68, and curcumin + H₂O₂: 225).

The influence of iron on the effects of curcumin has been investigated because of this metal's important role in

Table 1 - Effects of different concentrations of H₂O₂ on chromosomal aberrations induced by curcumin (CMN) in CHO cells.

Concentrations (µg/mL)		MI (%)	Gaps	Chromosomal aberrations		Total	Abnormal cells (Mean ± SD)
H ₂ O ₂	CMN			B'	B''		
0	0	11.0	1	5	0	5	1.6 ± 1.0
1.7	0	10.6	2	41	8	49 ^a	13.3 ^a ± 2.8
3.4	0	10.3	5	101	16	117 ^a	21.3 ^a ± 3.2
6.8	0	7.0 ^a	3	122	22	144 ^a	24.6 ^a ± 2.5
0	15	10.4	11	56	12	68 ^a	17.0 ^a ± 4.0
1.7	15	8.1	5	117	11	128 ^a	29.6 ^a ± 3.2
3.4	15	9.6	4	153	17	170 ^a	35.6 ^a ± 3.0
6.8	15	8.6	8	198	27	225 ^a	46.0 ^a ± 4.9

Three hundred cells per treatment were analyzed for chromosomal aberrations, and 3000 cells were scored for MI. Gaps were not included in the total number of abnormal metaphases or in the total number of chromosomal aberrations.

B' - chromatid break; B'' - chromosomal break; SD - standard deviation.

^ap < 0.05 for treated vs. control cultures.

Table 2 - Effects of different concentrations of Fe²⁺ and Fe³⁺ on chromosomal aberrations induced by curcumin (CMN) in CHO cells.

Concentrations (µg/mL)		MI (%)	Gaps	Chromosomal aberrations		Total	Abnormal cells (Mean ± SD)
Fe ²⁺	CMN			B'	B''		
0	0	11.7	1	2	1	3	1.0 ± 0.5
1.25	0	12.6	1	2	5	7	2.3 ± 0.5
2.5	0	11.7	3	4	2	6	2.0 ± 1.0
5.0	0	10.8	1	3	2	5	1.6 ± 0.6
0	15	9.2	11	51	12	63 ^a	16.6 ^a ± 4.0
1.25	15	8.5	16	47	8	55 ^a	13.6 ^a ± 4.6
2.5	15	8.2	3	21	8	29 ^b	9.0 ^b ± 4.0
5.0	15	9.7	2	8	0	8 ^b	2.6 ^b ± 2.8
Fe ³⁺	CMN						
0	0	11.6	3	3	2	5	1.6 ± 1.1
1.25	0	8.9	3	5	2	7	2.3 ± 1.0
2.5	0	10.2	2	3	0	3	1.0 ± 0.5
5.0	0	9.9	1	4	3	7	2.3 ± 1.1
0	15	7.8	12	71	7	78 ^a	18.3 ^a ± 6.0
1.25	15	7.1	4	35	4	39 ^b	10.0 ^b ± 3.5
2.5	15	10.5	0	11	4	15 ^b	5.3 ^b ± 2.5
5.0	15	7.8	0	9	4	13 ^b	4.3 ^b ± 1.5

Three hundred cells per treatment were analyzed for chromosomal aberrations, and 3000 cells were scored for MI. Gaps were not included in the total number of abnormal metaphases or in the total number of chromosomal aberrations.

B' - chromatid break; B'' - chromosomal break; SD - standard deviation.

^ap < 0.05 for treated vs. control cultures.

^bp < 0.05 for combined treatment vs. curcumin alone.

biological processes such as oxygen transfer and DNA synthesis (Bernabé-Pineda *et al.*, 2004). Table 2 shows that neither Fe²⁺ nor Fe³⁺ alone was clastogenic in any of the concentrations tested. The mitotic indices were also unaffected when compared to control cultures. The combined treatments with curcumin and Fe caused a significant (p < 0.05), concentration-dependent decrease in the total number of chromosomal aberrations and in the number of abnormal metaphases. At 2.5 or 5.0 µg of Fe²⁺/mL, there was a significant reduction in the total number of chromosomal aberrations induced by curcumin from 63 to 29 (54%) and 8 (87%), respectively. A similar response was seen in CHO cells treated with Fe³⁺ (Table 2).

The simultaneous addition of Fe²⁺ or Fe³⁺ significantly protected CHO cells against curcumin-induced chromosomal damage in a concentration-dependent manner compared to treatment with curcumin alone. This suggested the possibility that curcumin bound strongly to Fe. The reaction between the complexes curcumin-Fe²⁺ and curcumin-Fe³⁺ studied in aqueous media using UV spectrophotometry and cyclic voltammetry also showed a similar behavior, and indicated that a chemical reaction had occurred between the curcumin and Fe before the formation of the complexes (Bernabé-Pineda *et al.*, 2004).

Verma and Goldin (2003) recently reported that in the presence of proteins Cu²⁺ ions may not react with curcumin

to generate DNA damaging species, in contrast to the data obtained with a curcumin-Cu²⁺ combination in cell-free systems (Ahsan and Hadi, 1998). These authors suggested that the inhibitory effect of Cu²⁺ ions could be partly attributed to the presence of protein in the medium since the binding of copper to proteins would influence the activity of curcumin (Verma and Goldin, 2003). In the experiments described here, the presence of serum proteins in the medium could have influenced the activity of the curcumin-Fe complex. Although the exact mechanism by which curcumin induces chromosomal aberrations remains to be elucidated, the clastogenic activity of this compound is apparently related to its pro-oxidant properties and to its ability to generate free radicals.

Acknowledgments

The authors thank Dr. Sérgio N. Kronka (UNESP - Campus Jaboticabal) for helping with the statistical analysis, and Sueli A. Neves and Luis Augusto da Costa Jr. for technical assistance. This work was supported by CAPES and CNPq.

References

Abraham SK, Sarma L and Kesavan PC (1993) Protective effects of chlorogenic acid, curcumin and beta-carotene against

- gamma-radiation-induced *in vivo* chromosomal damage. *Mutat Res* 303:109-112.
- Aggarwal BB, Kumar A and Bharti AC (2003) Anticancer potential of curcumin: Preclinical and clinical studies. *Anticancer Res* 23:363-398.
- Ahsan H and Hadi SM (1998) Strand scission in DNA induced by curcumin in the presence of Cu(II). *Cancer Lett* 124:23-30.
- Antunes LMG, Araújo MCP, Darin JDC and Bianchi MLP (2000) Effects of the antioxidants curcumin and vitamin C on cisplatin-induced clastogenesis in Wistar rat bone marrow cells. *Mutat Res* 465:131-137.
- Antunes LMG, Araújo MCP, Dias FL and Takahashi CS (1999) Modulatory effects of curcumin on the chromosomal damage induced by doxorubicin in Chinese hamster ovary cells. *Teratogen Carcinogen Mutagen* 19:1-8.
- Araújo MCP, Antunes LMG and Takahashi CS (2001) Protective effect of thiourea, a hydroxyl-radical scavenger, on curcumin-induced chromosomal aberrations in an *in vitro* mammalian cell system. *Teratogen Carcinogen Mutagen* 21:175-180.
- Araújo MCP, Dias FL, Kronka SN and Takahashi CS (1999) Effects of turmeric and its active principle, curcumin, on bleomycin-induced chromosome aberrations in Chinese hamster ovary cells. *Genet Mol Biol* 22:407-413.
- Bernabé-Pineda M, Ramirez-Silva MT, Romero-Romo MA, González-Vergara E and Rojas-Hernández A (2004) Spectrophotometric and electrochemical determination of the formation constants of the complexes curcumin-Fe(III)-water and curcumin-Fe(II)-water. *Spectrochim Acta Part A* 60:1105-1113.
- Decker EA (1997) Phenolics: Prooxidants or antioxidants? *Nutr Rev* 55:396-407.
- Giri AK, Das SK, Talukder G and Sharma A (1990) Sister-chromatid exchange and chromosome aberrations induced by curcumin and tartrazine on mammalian cells *in vivo*. *Cytobios* 62:111-118.
- Ishidate M, Sofuni T, Yoshikawa K, Hayashi M, Nohmi T, Sawada M and Matsuoka A (1984) Primary mutagenicity screening of food additives currently used in Japan. *Food Chem Toxicol* 22:623-636.
- Joe B, Vijaykumar M and Lokesh BR (2004) Biological properties of curcumin-cellular and molecular mechanisms of action. *Crit Rev Food Sci Nutr* 44:97-111.
- Kuchandy E and Rao MNA (1990) Oxygen radical scavenging activity of curcumin. *Int J Pharm* 58:237-240.
- Meneghini R (1997) Iron homeostasis, oxidative stress, and DNA damage. *Free Radic Biol Med* 23:783-792.
- Mukhopadhyay MJ, Saha A and Mukherjee A (1998) Studies on the clastogenic effect of turmeric and curcumin on cyclophosphamide and mitomycin *in vivo*. *Food Chem Toxicol* 36:73-76.
- Nagabhushan M and Bhide SV (1992) Curcumin as an inhibitor of cancer. *J Am Coll Nutr* 11:192-198.
- Polasa K, Naidu NA, Ravindranath I and Krishanaswamy K (2004) Inhibition of B(a)P induced strand breaks in presence of curcumin. *Mutat Res* 557:203-213.
- Sahu SC and Washington MC (1992) Effect of ascorbic acid and curcumin on quercetin-induced nuclear DNA damage, lipid peroxidation and protein degradation. *Cancer Lett* 63:237-241.
- Sun Y-M, Zhang H-Y, Chen D-Z and Liu C-B (2002) Theoretical elucidation on the antioxidant mechanism of curcumin: A DFT study. *Org Lett* 4:2909-2911.
- Surh Y-J. and Ferguson LR (2003) Dietary and medicinal antimutagens and anticarcinogens: Molecular mechanisms and chemopreventive potential - highlights of a symposium. *Mutat Res* 523-524:1-8.
- Verma SP and Goldin BR (2003) Copper modulates activities of genistein, nitric oxide, and curcumin in breast tumor cells. *Biochem Biophys Res Commun* 310:104-108.

Associate Editor: Carlos F.M. Menck