

Effects of imperfect noise correlations on decoherence-free subsystems: SU(2) diffusion model



Rafał Demkowicz-Dobrzański

Center for Theoretical Physics of the Polish Academy of
Sciences, Warszawa, Poland
Nicolaus Copernicus University, Toruń, Poland

joint work with

Piotr Kolenderski, Konrad Banaszek

Nicolaus Copernicus University, Toruń, Poland



Depolarizing channel

- **Random unitary rotation of a qubit:**

$$|\psi\rangle = \cos\theta|\leftrightarrow\rangle + \sin\theta e^{i\phi}|\updownarrow\rangle$$

$$\mathcal{E}(|\psi\rangle\langle\psi|) = \int dU U|\psi\rangle\langle\psi|U^\dagger = \mathbb{1}/2$$

- **In long fibers the output polarization of a photon is completely random**



Collectively depolarizing channel

- **N qubit depolarizing channel, where each qubit experience the same disturbance**

$$\mathcal{E}(\rho_N) = \int dU U^{\otimes N} \rho_N U^{\dagger \otimes N}$$

$SU(2)$ Haar measure

N qubit state

- **The model applies e.g. to:**

- photons transmitted through a long fiber
- spins $\frac{1}{2}$ being sent through a slowly varying magnetic field
- communication in the absence of reference frames

Structure of the output state

- Irreducible subspaces under the action of $U^{\otimes N}$:

$$\mathcal{H}^{\otimes N} = \bigoplus_{j=0}^{N/2} \underbrace{\mathcal{H}_j \oplus \dots \oplus \mathcal{H}_j}_{d_j \text{ times}} = \bigoplus_{j=0}^{N/2} \mathcal{H}_j \otimes \mathbb{C}_{d_j} \quad \begin{array}{l} \text{multiplicity subspace} \\ \text{(decoherence free subsystem)} \end{array}$$

$$\mathcal{T}(\rho) := \mathcal{E}(\rho) = \int dU U^{\otimes N} \rho U^{\dagger \otimes N} = \bigoplus_{j=0}^{N/2} \frac{p_j}{2j+1} \mathbb{1}_{\mathcal{H}_j} \otimes \rho_j$$


$$p_j = \text{Tr}(P_j \rho) \quad \rho_j = \frac{1}{p_j} \text{Tr}_{\mathcal{H}_j}(P_j \rho P_j) \quad P_j - \text{projection on } \mathcal{H}_j \otimes \mathbb{C}_{d_j}$$

- Faithfully transmitted states - allow for noiseless classical and quantum communication

$$\rho = \bigoplus_{j=0}^{N/2} \frac{p_j}{2j+1} \mathbb{1}_{\mathcal{H}_j} \otimes \rho_j \quad \text{twirled structure}$$

Photons and salt bags

K. Banaszek, A. Dragan, W. Wasilewski, and G. Radzewicz, Phys. Rev. Lett. **92**, 257901 (2004)



$$|\psi\rangle = \frac{1}{\sqrt{2}} (|\uparrow\leftrightarrow\rangle - |\leftrightarrow\downarrow\rangle)$$

**Salt bags =
integrating with
Haar measure**

**singlet state
remains intact!**

What happens if noise is not perfectly correlated?

In other words:

what happens if oscillation time of salt bags is comparable with photon separation time

Imperfectly correlated noise model

- Consecutive qubits experience slightly different rotations

$$\mathcal{E}(\rho) = \int dU_1 \dots dU_N p(U_1, \dots, U_N) U_1 \otimes \dots \otimes U_N \rho U_1^\dagger \dots \otimes U_N^\dagger$$

- The action described via a stationary Markov process

$$p(U_1, \dots, U_N) = p(U_N | U_{N-1}) \dots p(U_2 | U_1)$$

What is the natural choice for conditional probability?

$$p(U_i | U_{i-1}) = ?$$

Diffusion on the SU(2) group

- **Isotropic diffusion on SU(2)**

$$\partial_t p(U; t) = \frac{1}{2} D \hat{\Delta} p(U; t)$$

Laplace operator on the SU(2) group

- **Solution, with the initial condition:** $p(U; 0) = \delta(U - \mathbb{1})$

$$p(U; t) = \sum_{j=0}^{\infty} (2j + 1) \exp\left(-\frac{1}{2} j(j + 1) t\right) \sum_{m=-j}^j \mathcal{D}^j(U)_m^m$$

diffusion strength

rotation matrices

- **Conditional probability**

$$p(U_i | U_{i-1}) = p(U_i U_{i-1}^\dagger; t)$$

$t \rightarrow 0$ perfect noise correlation

$t \rightarrow \infty$ no correlation

Action of the channel

- **Probability distribution for unitaries**

$$p(U_1, \dots, U_N) = p(U_N | U_{N-1}) \cdots p(U_2 | U_1) = p(U_N U_{N-1}^\dagger; t) \cdots p(U_2^\dagger U_1; t)$$

- **The channel action**

$$\mathcal{E}(\rho) = \int dU_1 \cdots \int dU_N p(U_1, \dots, U_N) U_1 \otimes \cdots \otimes U_N \rho U_1^\dagger \otimes \cdots \otimes U_N^\dagger$$

$$\mathcal{E}(\rho) = \mathcal{I}_{N-1} (\mathcal{I}_{N-2} (\dots \mathcal{I}_1 (\mathcal{T}(\rho)) \dots))$$

$$\mathcal{T}(\rho) = \int dU U^{\otimes N} \rho U^{\dagger \otimes N}$$

$$\mathcal{I}_i(\rho) = \int dU p(U; t) \underbrace{\mathbb{1} \otimes \cdots \otimes \mathbb{1}}_i \otimes \underbrace{U \otimes \cdots \otimes U}_{N-i} \rho \underbrace{\mathbb{1} \otimes \cdots \otimes \mathbb{1}}_i \otimes \underbrace{U^\dagger \otimes \cdots \otimes U^\dagger}_{N-i}$$

Action of the channel

- **Output states have a twirled structure (T, \mathcal{I}_i commute)**

$$\mathcal{E}(\rho) = \mathcal{I}_{N-1} (\mathcal{I}_{N-2} (\dots \mathcal{I}_1 (T(\rho)) \dots))$$

- **Input states can be restricted to have the twirled structure, so the channel action can be described as**

$$\bigoplus_{j=0}^{N/2} \frac{p_j}{2j+1} \mathbb{1}_{\mathcal{H}_j} \otimes \rho_j \xrightarrow{\mathcal{E}} \bigoplus_{j=0}^{N/2} \frac{p'_j}{2j+1} \mathbb{1}_{\mathcal{H}_j} \otimes \rho'_j$$

- **Using properties of rotation matrices $\mathcal{D}_m^j(U)$, it is possible to derive analytical expression for the action of the channel**

$\mathcal{E}(\rho) =$ lengthy expression involving $e^{-\frac{1}{2}j(j+1)t}$ and Wigner $6j$ symbols

Example: Three qubit channel

- **Structure of a three qubit twirled state**

$$\rho = \frac{p}{2} (\mathbb{1}_{\mathcal{H}_{1/2}} \otimes \rho_{1/2}) \oplus \frac{1-p}{4} \mathbb{1}_{\mathcal{H}_{3/2}}$$

effectively two dimensional subspace

one dimensional subspace

$$|0\rangle \quad (j_{12} = 0, j_{123} = 1/2)$$

$$|1\rangle \quad (j_{12} = 1, j_{123} = 1/2)$$

$$|2\rangle \quad (j_{12} = 1, j_{123} = 3/2)$$

- **We have a qutrit channel, with no coherence between $|0\rangle, |1\rangle$ subspace and $|2\rangle$**

- **If correlations of noise were perfect (no diffusion), the channel would allow for $\log_2 3$ bits of classical communication and 1 qubit of quantum communication**

Fidelity of transmitting a qubit

- **Transmitting a qubit**

$$|\psi\rangle = \cos(\theta/2)|e_1\rangle + \sin(\theta/2)e^{i\phi}|e_2\rangle$$

$$|e_1\rangle = (|0\rangle + \sqrt{3}|1\rangle)/2$$

$$|e_2\rangle = (\sqrt{3}|0\rangle - |1\rangle)/2$$


substituting the output state $|2\rangle$, with a maximally mixed state of a qubit we can write the effective qubit channel in terms of evolution of the Bloch vector

$$\mathbf{r}_{\text{out}} = \begin{pmatrix} e^{-t} & 0 & 0 \\ 0 & e^{-2t} & 0 \\ 0 & 0 & \frac{2e^{-2t} + e^{-t}}{3} \end{pmatrix} \mathbf{r}_{\text{in}} + \begin{pmatrix} 0 \\ 0 \\ \frac{e^{-2t} - e^{-t}}{3} \end{pmatrix}$$

anisotropic shrinkning with displacement, states with $\phi = 0, \pi$ will tend to have high fidelity (weakest shrinking)

Fidelity of transmitting a qubit

for $t = 1$



$$\mathbf{r}_{\text{out}} = \begin{pmatrix} e^{-t} & 0 & 0 \\ 0 & e^{-2t} & 0 \\ 0 & 0 & \frac{2e^{-2t} + e^{-t}}{3} \end{pmatrix} \mathbf{r}_{\text{in}} + \begin{pmatrix} 0 \\ 0 \\ \frac{e^{-2t} - e^{-t}}{3} \end{pmatrix}$$

anisotropic shrinkning with displacement, states with $\phi = 0, \pi$ will tend to have high fidelity (weakest shrinking)

Summary

Phys. Rev. A 76, 022302 (2007)

- **Introduction of a natural model of an N qubit channel with imperfectly correlated random unitary rotations acting on consecutive qubits**
- **Derivation of an analytic formula for the action of the channel on an arbitrary input state**
- **Detailed analysis of the case $N=3$**
 - **fidelity of the channel**
 - **optimal classical capacity, and corresponding states**
 - **orthogonal states that provide almost optimal classical capacity even for non perfect noise correlations**
 - **threshold of diffusion strength above which coherent information vanishes**
- **Future work:**
 - **develop a perturbative approach for weak diffusion for large number of qubits, find optimal capacities and corresponding states**
 - **analyse within this framework 'estimate and correct' strategy for sending information**