

## Effects of Inorganic Mercury on Osteoclasts and Osteoblasts of the Goldfish Scales In Vitro

Suzuki, Nobuo

Noto Marine Laboratory, Institute of Nature and Environmental Technology, Kanazawa University

Yachiguchi, Koji

Noto Marine Center

Hayakawa, Kazuichi

Graduate School of Natural Science and Technology, Kanazawa University

Omori, Katsunori

Japan Aerospace Exploration Agency

他

<https://doi.org/10.5109/19635>

---

出版情報：九州大学大学院農学研究院紀要. 56 (1), pp.47-51, 2011-02. 九州大学大学院農学研究院  
バージョン：  
権利関係：



## Effects of Inorganic Mercury on Osteoclasts and Osteoblasts of the Goldfish Scales *In Vitro*

Nobuo SUZUKI<sup>1\*</sup>, Koji YACHIGUCHI<sup>2</sup>, Kazuichi HAYAKAWA<sup>3</sup>, Katsunori OMORI<sup>4</sup>, Koji TAKADA<sup>5</sup>, Makoto J. TABATA<sup>6</sup>, Kei-Ichiro KITAMURA<sup>7</sup>, Masato ENDO<sup>8</sup>, Shigehito WADA<sup>9</sup>, Ajai K. SRIVASTAV<sup>10</sup>, Vishwajit S. CHOWDHURY<sup>11</sup>, Yuji OSHIMA<sup>12</sup> and Atsuhiko HATTORI<sup>13</sup>

Department of Bioscience and Biotechnology, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University, Hakozaki 6–10–1, Higashi-ku, Fukuoka, 812–8581, Japan

(Received October 26, 2010 and accepted November 8, 2010)

The fish scales are the major source of internal calcium requirement due to having a higher internal calcium reservoir than the body skeleton during the periods of drastic calcium demand, such as sexual maturation. Therefore, we developed original *in vitro* assay system using goldfish scales that contain osteoclasts and osteoblasts, and examined the direct effect of inorganic mercury ( $\text{HgCl}_2$ ) on osteoclasts and osteoblasts. In this assay system, we measured the activities of tartrate-resistant acid phosphatase (TRAP) and alkaline phosphatase (ALP) as respective indicators of each activity in osteoclasts and osteoblasts. TRAP activity in the scales significantly decreased by the treatment of  $\text{HgCl}_2$  ( $10^{-6}$  to  $10^{-3}$  M) during 6 hrs of incubation. In addition, mRNA expressions of osteoclastic markers: TRAP and cathepsin K significantly decreased compared with control. In our knowledge, this is the first report of a direct effect of inorganic mercury on osteoclasts. On the other hand, ALP activity decreased after exposures of  $\text{HgCl}_2$  at a concentration of  $10^{-6}$ ,  $10^{-5}$  or  $10^{-4}$  M for 36 and 64 hrs, although its activity did not change after 6 and 18 hrs. The mRNA expression of metallothionein (MT) which is a metal-binding-protein that protects the organism from heavy metal, significantly increased by  $\text{HgCl}_2$  ( $10^{-4}$  M) although insulin-like growth factor-I (osteoblastic marker) was less than those of control scales by treatment with  $\text{HgCl}_2$  ( $10^{-4}$  M). These results suggests that osteoblasts may synthesize MT and protect from mercury until 18 hrs incubation. Thus, the scale *in vitro* assay system would be a useful means for analysis of heavy metal on bone metabolism.

**Keywords:** inorganic mercury, *in vitro* assay, osteoclasts, osteoblasts, scales

### INTRODUCTION

Heavy metals such as mercury, cadmium and copper are known to be extremely toxic to organisms. Mercury has been recognized as an environmental contaminant since the Minamata disaster in the late 1950s. Minamata disease which was caused by the consumption of marine fishes severely polluted with mercury from local industrial discharge due to this Minamata disaster (Takeuchi *et al.*, 1978; Takeuchi, 1982). This extremely adverse situation occurred because of mercury, a highly toxic compound, was severely bio-accumulated (in case of long-

fanned eels: approx. 1,000,000 times higher than environmental water) by fish (Redmayne *et al.*, 2000).

The effect of mercury on the central nervous system has widely studied and revealed that mercury is a neurotoxic material, and its poisoning effect is characterized by the damage in discrete portions of the brain, such as the visual cortex and the granule layer of the cerebellum (Castoldi *et al.*, 2001). As bio-accumulation of mercury in bone is lower than that in neural tissues (Boyer *et al.*, 1978; Doyle, 1979; Berglund *et al.*, 2000), much attention has not been given to bone in this area of research.

Recently, Lake *et al.* (2006) reported that the corre-

<sup>1</sup> Noto Marine Laboratory, Institute of Nature and Environmental Technology, Kanazawa University, Housu-gun, Ishikawa 927–0553, Japan

<sup>2</sup> Noto Marine Center, Housu-gun, Ishikawa 927–0552, Japan

<sup>3</sup> Graduate School of Natural Science and Technology, Kanazawa University, Kakuma, Ishikawa 920–1192, Japan

<sup>4</sup> Japan Aerospace Exploration Agency, Tsukuba, Ibaraki 305–8505, Japan

<sup>5</sup> Department of Biochemistry, Jikei University School of Medicine, Minato-ku, Tokyo 105–8461, Japan

<sup>6</sup> Section of Biostructural Science, Graduate School of Tokyo Medical and Dental University, Bunkyo-ku, Tokyo 113–8549, Japan

<sup>7</sup> Faculty of Health Sciences, Institute of Medical, Pharmaceutical and Health Sciences, Kanazawa University, Kanazawa, Ishikawa 920–0942, Japan

<sup>8</sup> Department of Marine Biosciences, Faculty of Marine Science, Tokyo University of Marine Science and Technology, Konan, Minato-ku, Tokyo 108–8477, Japan

<sup>9</sup> Department of Oral and Maxillofacial Surgery, Faculty of Medicine, University of Toyama, Sugitani, Toyama 930–0194, Japan

<sup>10</sup> Department of Zoology, University of Gorakhpur, Gorakhpur 273009, India

<sup>11</sup> International Education Center, Department of Bioresource Sciences, Faculty of Agriculture, Kyushu University, Fukuoka 812–8581, Japan

<sup>12</sup> Department of Bioscience and Biotechnology, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University, Hakozaki 6–10–1, Higashi-ku, Fukuoka, 812–8581, Japan

<sup>13</sup> Department of Biology, College of Liberal Arts and Sciences, Tokyo Medical and Dental University, Ichikawa, Chiba 272–0827, Japan

\* Corresponding author (E-mail: nobuo@kenroku.kanazawa-u.ac.jp)

lation between the total mercury concentration of the scales and that of the muscles was high ( $r=0.89$ ), and suggested the suitability for prediction of muscle tissue by the assessment of available mercury in the fish scales. It is known that the scales are calcified tissue which contains osteoclasts and osteoblasts (Bereiter-Hahn and Zylberberg, 1993; Suzuki *et al.*, 2000; Yoshikubo *et al.*, 2005; Suzuki *et al.*, 2007) and is reported that the scales are a better potential internal calcium reservoir than the body skeletons, jaws and otoliths, examined by the  $^{45}\text{Ca}$ -labelling study for the calcified tissues of goldfish and killifish (Mugiya and Watabe, 1977). In fishes, thus, the scale accumulates mercury and seems to be a sensitive tissue for mercury.

Recently, we have developed a novel *in vitro* assay system using goldfish scale (Suzuki *et al.*, 2000; Suzuki and Hattori, 2002) because the scale is a very active tissue of calcium regulation in fish described above. In the present study, therefore, we examined the effect of inorganic mercury ( $\text{HgCl}_2$ ) on the scale osteoclasts and osteoblasts. To confirm the effects of  $\text{HgCl}_2$  on osteoclasts and osteoblasts, the mRNA expressions of osteoclastic markers (tartrate-resistant acid phosphatase: TRAP and cathepsin K) and osteoblastic marker (insulin-like growth factor-I: IGF-I) were investigated using reverse-transcription (RT)-PCR. Furthermore, the mRNA expression of metallothionein (MT), which is a metal-binding-protein that protects the organism from heavy metal (Hamer, 1986; Klaassen *et al.*, 1999), was also examined using RT-PCR.

## MATERIALS AND METHODS

### Animals

Our previous study (Suzuki *et al.*, 2000) indicated that the sensitivity for calcemic hormone such as estrogen and calcitonin was higher in mature female than mature male in goldfish (*Carassius auratus*). Therefore, mature female goldfish ( $n=12$ ,  $35.50 \pm 1.30$  g) were purchased from commercial source (Higashikawa Fish Farm, Yamatokoriyama, Japan) and used in the scale *in vitro* assay. All experimental procedures were conducted in accordance with the Guide for the Care and Use of Laboratory Animals of Kanazawa University.

### Effect of $\text{HgCl}_2$ on TRAP and ALP activities in the cultured scales of goldfish

A 1% penicillin-streptomycin mixture (ICN Biomedicals Inc., OH, USA) was added to Eagle's minimum essential medium (MEM; ICN Biomedicals Inc.). HEPES (Research Organics Inc., OH, USA) (20 mM) was added into MEM and adjusted to pH 7.0. After filtration, MEM was used in this experiment for analyzing the effect of  $\text{HgCl}_2$  on TRAP and ALP activities in the cultured goldfish scales. Scales collected from goldfish under anesthesia with ethyl 3-aminobenzoate, methanesulfonic acid salt (MS-222, Sigma-Aldrich, Inc., MO, USA) and incubated for 6 hrs in MEM supplemented with  $10^{-8}$ – $10^{-3}$  M  $\text{HgCl}_2$  (Wako Pure Chemical Industries, Ltd., Osaka, Japan) and compared with control ( $\text{HgCl}_2$ -free medium).

To evaluate the effect of  $\text{HgCl}_2$  on osteoclasts and osteoblasts, furthermore, scales were incubated with  $\text{HgCl}_2$  ( $10^{-7}$ ,  $10^{-6}$ ,  $10^{-5}$ , and  $10^{-4}$  M) for comparatively longer exposure times, namely 18, 36, and 64 hrs. After incubation, scales were fixed in 10% formalin in a 0.05 M cacodylate buffer (pH 7.4) followed by a storage in a 0.05 M cacodylate buffer at 4 °C until analysis.

The measurement methods of TRAP and ALP activities have been described by Suzuki and Hattori (2002). We detected the respective enzyme activity from one scale by transferring each scale into a 96-well-microplate and directly incubating it with the substrate in each well. The procedure of TRAP measurement was as follows. Each scale was transferred to its own well in a 96-well microplate after measurement of the scale weight. An aliquot of 200  $\mu\text{l}$  of 10 mM para-nitrophenyl-phosphate and 20 mM tartrate in a 0.1 M sodium acetate buffer (pH 5.3) was added to each well. This plate was then incubated at 20 °C for 30 min while being shaken. After incubation, the reaction was stopped by adding 50  $\mu\text{l}$  of 2 N NaOH. One hundred and fifty  $\mu\text{l}$  of a colored solution was transferred to a new plate, and the absorbance was measured at 405 nm. The absorbance was converted into the amount of produced para-nitrophenol (pNP) using a standard curve for pNP. The results are shown as means  $\pm$  SEM of eight scales.

ALP activities were measured using an alkaline buffer (100 mM Tris-HCl, pH 9.5; 1 mM  $\text{MgCl}_2$ ; 0.1 mM  $\text{ZnCl}_2$ ). Other conditions were the same as those for the measurement of TRAP activity.

### Changes of TRAP, cathepsin K, IGF-I, and MT mRNA expression in $\text{HgCl}_2$ -treated scales for 18 hrs of culture

Scales were collected from goldfish under anesthesia with MS-222. To examine changes in TRAP, cathepsin K, IGF-I, and MT mRNAs that responded to  $\text{HgCl}_2$ , these scales were incubated for 18 hrs in MEM (containing antibiotic and 20 mM HEPES) supplemented with  $\text{HgCl}_2$  ( $10^{-4}$  M) and compared with the control (without metals). We previously reported that IGF-I mRNA expression decreased at 18 hrs of incubation (Suzuki and Hattori, 2003). Therefore, this incubation period was adopted. After incubation, the scales were frozen at  $-80$  °C for mRNA analysis.

Total RNAs were prepared from the goldfish scales using a total RNA isolation kit (Nippon Gene, Tokyo, Japan). RT-PCR was performed using Oligotex-dT 30 Super (Takara Bio Inc., Otsu, Japan) as an oligo dT primer to prevent genomic DNA contamination (Suzuki *et al.*, 1997). The gene-specific primers (TRAP 5': AACTTC-CGCATTCCTCGAACAG; TRAP 3': GGCCAGCCACCAGGAGATAA; cathepsin K 5': GCTAT-GGAGCCACACAAAAGG; cathepsin K 3': CTGCGCTTCCAGCTCTCACAT) reported by Azuma *et al.* (2007) were used. IGF-I and MT cDNAs were also amplified using gene specific primers (IGF-I 5': GGAGACGCTGTGCGGG; IGF-I 3': CCTCAGCTCACAGCTCTG; MT 5': ATGGATCCGTGCGAATGC; MT 3':

CTCCTCATTGACAGCAGCT). These were designed from the nucleotide sequences of respective cDNA (IGF-I: Kermouni *et al.*, 1998; MT: Chan, 1994).  $\beta$ -actin cDNA using a primer set (5':CACTGTGCCCATCTACGAG; 3':CCATCTCCTGCTCGAAGTC) (Chan *et al.*, 1998) were also amplified. The conditions for PCR amplification were denaturation for 0.5 min at 96 °C, annealing for 1 min at 55 °C, and extension for 2 min at 72 °C, followed by a single cycle at 72 °C for 30 min. The cycle numbers for the amplification in TRAP, cathepsin K, IGF-I, MT, and  $\beta$ -actin cDNAs were determined by ensuring that PCR amplification was at submaximum and the intensity of the band corresponded exactly to the amount of starting material. The PCR products were analyzed on a 2.5% NuSive GTG agarose gel (FMC BioProducts, ME, USA) and stained with ethidium bromide. The band densities were estimated using a computer program (NIH Image J). The mRNA levels of TRAP, cathepsin K, IGF-I and

MT were normalized to the mRNA level of  $\beta$ -actin.

### Statistical analysis

The statistical significance was assessed by one-way ANOVA followed by Dunnett test. The significance level chosen was as  $P < 0.05$ .

## RESULTS

### Effect of HgCl<sub>2</sub> on TRAP activity in the cultured scales of goldfish

HgCl<sub>2</sub> significantly decreased the TRAP activities of the scales by 6 hrs of incubation ( $P < 0.01$  for  $10^{-5}$  M;  $P < 0.001$  for  $10^{-4}$  and  $10^{-3}$  M) (Fig. 1). Thus, increased doses of HgCl<sub>2</sub> resulted in greater effects on decreasing TRAP activities dose-dependently.

By the long incubation time period (18 to 64 hrs), only at  $10^{-4}$  M, significant difference ( $P < 0.01$ ) between HgCl<sub>2</sub>-treated scales and control scales was obtained by




Fig. 1. Effect of HgCl<sub>2</sub> ( $10^{-8}$  to  $10^{-3}$  M) on TRAP activity in the cultured scales incubated for 6 hrs. \*\*, \*\*\* indicate statistically significant differences at  $P < 0.01$  and  $P < 0.001$ , respectively, from the value in the control scales.




Fig. 3. Effect of HgCl<sub>2</sub> ( $10^{-8}$  to  $10^{-3}$  M) on ALP activity in the cultured scales incubated for 6 hrs. There was no significant difference between HgCl<sub>2</sub>-treated scales and control scales.




Fig. 2. Effect of HgCl<sub>2</sub> ( $10^{-7}$  to  $10^{-4}$  M) on TRAP activity in the cultured scales incubated for 18, 36, and 64 hrs. \*\* indicates statistically significant difference at  $P < 0.01$  from the values in the control scales.




Fig. 4. Effect of HgCl<sub>2</sub> ( $10^{-7}$  to  $10^{-4}$  M) on ALP activity in the cultured scales incubated for 18, 36, and 64 hrs. \*, \*\*, \*\*\* indicate statistically significant differences at  $P < 0.05$ ,  $P < 0.01$  and  $P < 0.001$ , respectively, from the values in the control scales.

18 hrs of incubation (Fig. 2).


### Effect of HgCl<sub>2</sub> on ALP activity in the cultured scales of goldfish

The ALP activity of the control scales by 6 hrs of incubation was  $5.33 \pm 0.41$  (nmol produced pNP/mg scale/hr) which did not show any difference with HgCl<sub>2</sub>-treated groups ( $10^{-5}$  to  $10^{-3}$ M) (Fig. 3). Thus, the ALP activity did not change during 6 hrs of incubation with HgCl<sub>2</sub> compared to the control.

However, the ALP activity in the HgCl<sub>2</sub>-treated scales decreased significantly by 36 hrs ( $P < 0.05$  for  $10^{-6}$ M,  $10^{-5}$ M or  $10^{-4}$ M) and 64 hrs ( $P < 0.05$  for  $10^{-6}$ M;  $P < 0.01$  for  $10^{-5}$ M and  $P < 0.001$  for  $10^{-4}$ M) of incubation from the values of the control scales although it did not change at 18 hrs of incubation (Fig. 4).

### Changes of TRAP, cathepsin K, IGF-I, and MT mRNA expression in HgCl<sub>2</sub>-treated scales

After 18 hrs of incubation, the mRNA expressions of TRAP, cathepsin K and IGF-I in HgCl<sub>2</sub>-treated scales were significantly ( $P < 0.001$  for TRAP;  $P < 0.001$  for cathepsin K and  $P < 0.001$  for IGF-I) lower than those in the control scales (Fig. 5). Conversely, the mRNA expression was significantly ( $P < 0.001$ ) increased for MT when treated with HgCl<sub>2</sub> (Fig. 5).



**Fig. 5.** Changes in the mRNA expression of TRAP, cathepsin K, IGF-I, and MT in HgCl<sub>2</sub> ( $10^{-4}$ M)-treated scales of goldfish incubated for 18 hrs of culture. \*\*\* indicates statistically significant difference at  $P < 0.001$  from the values in the control scales.

## DISCUSSION

The present study demonstrated that fish scale sensitively responded to HgCl<sub>2</sub>. A high co-relation of mercury between scales and muscles was reported in largemouth bass (Lake *et al.*, 2006). This indicates that accumulation of mercury is occurred in the fish scale although mercury did not accumulate in the vertebral bone of fish (Camusso *et al.*, 1995). It is also well-known that the scale is a more active tissue in fish calcium regulation than vertebral bone (Mugiya and Watabe, 1977; Yamada,

1961; Berg, 1968; Bereiter-Hahn and Zylberberg, 1993). Therefore, we strongly believe that the fish scale is capable to accumulate mercury and respond to mercury similarly like calcium.

In mammals, the influence of mercury on bone metabolism has been studied only by *in vivo* experiments and investigated in bone formation or osteoblastic activity (Yonaga *et al.*, 1985; Jin *et al.*, 2002). Mercury inhibited the growth of tibia in rats (Yonaga *et al.*, 1985) and decreased serum levels of osteoblastic markers (ALP and osteocalcin) (Jin *et al.*, 2002). In our knowledge, our study is the first to indicate direct effect of inorganic mercury on osteoclasts. The inhibitory action of HgCl<sub>2</sub> on osteoclasts after 6 hrs incubation was stronger than that of 18 to 64 hrs incubation. As for organic mercury, similar results were obtained in our scale assay system (Suzuki *et al.*, 2004). Furthermore, we recently succeed to clone osteoclastic markers: TRAP and cathepsin K in fish for the first time (Azuma *et al.*, 2007) and examined mRNA expressions of these markers in the HgCl<sub>2</sub>-treated scales. In the present study, we confirmed that the both mRNA expressions of TRAP and cathepsin K decreased as TRAP enzyme activity did.

It was found that the mRNA expression of MT in HgCl<sub>2</sub>-treated scales increased in the present study. This result is similar to that in mammals because it has been demonstrated that MT plays a protective role in mercury-induced toxicity in bone (Jin *et al.*, 2002). Fish are aquatic animals with scales that are always exposed to environmental water. In an *in vitro* experiment for 6 and 18 hrs of incubation, therefore, osteoblasts may be resistant to mercury as a result of MT production. On the other hand, IGF-I mRNA expression decreased compared to the control. As IGF-I participates in osteoblastic growth and differentiation, we speculated that mercury has toxic effect on osteoblasts under long-term exposure.

We previously demonstrated that the osteogenesis of regenerating scale is very similar to that of mammalian membrane bone and a good model of osteogenesis (Yoshikubo *et al.*, 2005). Using this system, furthermore, we first demonstrated that calcitonin, a hypocalcemic hormone, suppressed osteoclastic activity in teleosts as well as in mammals (Suzuki *et al.*, 2000) and that melatonin, a major hormone secreted from the pineal gland, suppressed the functions in both osteoclasts and osteoblasts (Suzuki and Hattori, 2002). Osteoblasts in the scale responded to estrogen as they do in mammals (Yoshikubo *et al.*, 2005). In addition, the effects of endocrine disrupters, such as bisphenol-A (Suzuki and Hattori, 2003) and tributyltin (Suzuki *et al.*, 2006), and heavy metals, i.e., cadmium and organic mercury (Suzuki *et al.*, 2004), on osteoblasts and osteoclasts have been examined. Moreover, we indicated that cadmium (even at  $10^{-13}$ M) responded to TRAP activity in the scale (Suzuki *et al.*, 2004). Considering these results together with present data, our scale assay system will be useful for analysis of environmental contaminant on bone metabolism.



## ACKNOWLEDGMENTS

This study was supported in part by grants to N.S. (Grant-in-Aid for Scientific Research (C) No. 21500404), to A. H. (Grant-in-Aid for Scientific Research (C) No. 21570062) sponsored by the Japan Society for the Promotion of Science and to K. H. the Environment Research and Technology Development Fund (B-0905) sponsored by the Ministry of the Environment, Japan, and Health and Labour Sciences Research Grants of Ministry of Health, Labour and Welfare, Japan.

## REFERENCES

- Azuma, K., M. Kobayashi, M. Nakamura, N. Suzuki, S. Yashima, S. Iwamuro, M. Ikegame, T. Yamamoto and A. Hattori 2007 Two osteoclastic markers expressed in multinucleate osteoclasts of goldfish scales. *Biochem. Biophys. Res. Commun.*, **362**: 594–600
- Bereiter-Hahn, J. and L. Zylberberg 1993 Regeneration of teleost fish scale. *Comp. Biochem. Physiol.*, **105A**: 625–641
- Berg, A. 1968 Studies on the metabolism of calcium and strontium in freshwater fish. I. Relative contribution of direct and intestinal absorption. *Mem. Ist. Ital. Idrobiol.*, **23**: 161–196
- Berglund, M., A. Åkesson, P. Bjellerup and M. Vahter 2000 Metal–bone interactions. *Toxicol. Lett.*, **112–113**: 219–225
- Boyer, C. I. Jr., E. J. Andrews, A. deLahunta, C. A. Bache, W. H. Gutenmann and D. J. Lisk 1978 Accumulation of mercury and selenium in tissues of kittens fed commercial cat food. *Cornell. Vet.*, **68**: 365–374
- Camusso, M., L. Vigano and R. Balestrini 1995 Bioconcentration of trace metals in rainbow trout: A field study. *Ecotoxicol. Environ. Saf.*, **31**: 133–141
- Castoldi, A. F., T. Coccini, S. Ceccatelli and L. Manzo 2001 Neurotoxicity and molecular effects of methylmercury. *Brain Res. Bull.*, **55**: 197–203
- Chan, K.-M. 1994 PCR-cloning of goldfish and tilapia metallothionein complementary DNAs. *Biochem. Biophys. Res. Commun.*, **205**: 368–374
- Chan, K.-W., K.-L. Yu, J. Rivier and B. K.-C. Chow 1998 Identification and characterization of a receptor from goldfish specific for a teleost growth hormone-releasing hormone-like peptide. *Neuroendocrinology*, **68**: 44–56
- Doyle, J. J. 1979 Toxic and essential elements in bone: A review. *J. Anim. Sci.*, **49**: 482–497
- Hamer, D. H. 1986 Metallothionein. *Annu. Rev. Biochem.*, **55**: 913–951
- Jin, G.-B., S. Inoue, T. Urano, S. Cho, Y. Ouchi and J.-C. Cyong 2002 Induction of anti-metallothionein antibody and mercury treatment decreases bone mineral density in mice. *Toxicol. Appl. Pharmacol.*, **185**: 98–110
- Kermouni, A., S. S. Mahmoud, S. Wang, M. Moloney and H. R. Habibi 1998 Cloning of a full-length insulin-like growth factor-I complementary DNA in the goldfish liver and ovary and development of a quantitative PCR method for its measurement. *Gen. Comp. Endocrinol.*, **111**: 51–60
- Klaassen, C. D., J. Liu and S. Choudhuri 1999 Metallothionein: An intracellular protein to protect against cadmium toxicity. *Annu. Rev. Pharmacol. Toxicol.*, **39**: 267–294
- Lake, J. L., S. A. Ryba, J. R. Serbst and A. D. Libby 2006 Mercury in fish scales as an assessment method for predicting muscle tissue mercury concentrations in largemouth bass. *Arch. Environ. Contam. Toxicol.*, **50**: 539–544
- Mugiya, Y. and N. Watabe 1977 Studies on fish scale formation and resorption II: Effect of estradiol on calcium homeostasis and skeletal tissue resorption in the goldfish, *Carassius auratus*, and the killifish, *Fundulus heteroclitus*. *Comp. Biochem. Physiol.*, **57A**: 197–202
- Redmayne, A. C., J. P. Kim, G. P. Closs and K. A. Hunter 2000 Methyl mercury bioaccumulation in long-finned eels, *Anguilla dieffenbachii*, from three rivers in Otago, New Zealand. *Sci. Total Environ.*, **262**: 37–47
- Suzuki, N., C. Eguchi, T. Hirai and Y. Sasayama 1997 Nucleotide sequences of reptile calcitonins: Their high homology to chicken calcitonin. *Zool. Sci.*, **14**: 833–836
- Suzuki, N., T. Suzuki and T. Kurokawa 2000 Suppression of osteoclastic activities by calcitonin in the scales of goldfish (freshwater teleost) and nibbler fish (seawater teleost). *Peptides* **21**: 115–124
- Suzuki, N. and A. Hattori 2002 Melatonin suppresses osteoclastic and osteoblastic activities in the scales of goldfish. *J. Pineal Res.*, **33**: 253–258
- Suzuki, N. and A. Hattori 2003 Bisphenol A suppresses osteoclastic and osteoblastic activities in the cultured scales of goldfish. *Life Sci.*, **73**: 2237–2247
- Suzuki, N., M. Yamamoto, K. Watanabe, A. Kambegawa and A. Hattori 2004 Both mercury and cadmium directly influence calcium homeostasis resulting from the suppression of scale bone cells: The scale is a good model for the evaluation of heavy metals in bone metabolism. *J. Bone Miner. Metab.*, **22**: 439–446
- Suzuki, N., M. J. Tabata, A. Kambegawa, A. K. Srivastav, A. Shimada, H. Takeda, M. Kobayashi, S. Wada, T. Katsumata and A. Hattori 2006 Tributyltin inhibits osteoblastic activity and disrupts calcium metabolism through an increase in plasma calcium and calcitonin levels in teleosts. *Life Sci.*, **78**: 2533–2541
- Suzuki, N., K. Kitamura, T. Nemoto, N. Shimizu, S. Wada, T. Kondo, M. J. Tabata, F. Sodeyama, K. Ijiri and A. Hattori 2007 Effect of vibration on osteoblastic and osteoclastic activities: Analysis of bone metabolism using goldfish scale as a model for bone. *Adv. Space Res.*, **40**: 1711–1721
- Takeuchi, T., K. Eto, S. Oyanag and H. Miyajima 1978 Ultrastructural changes of human sural nerves in the neuropathy induced by intrauterine methylmercury poisoning (so-called fetal Minamata disease). *Virchows Arch. B Cell Path.*, **27**: 137–154
- Takeuchi, T. 1982 Pathology of Minamata disease: With special reference to its pathogenesis. *Acta Pathol. Jpn.*, **32 Suppl 1**: 73–99
- Yamada, J. 1961 Studies on the structure and growth of the scales in the goldfish. *Mem. Fac. Fish Hokkaido Univ.*, **9**: 181–226
- Yonaga, T., Y. Fujino, R. Tamura, K. Kurabayashi, T. Uraya, K. Aono and K. Yoshimura 1985 Effect of organic and inorganic mercury compounds on the growth of incisor and tibia in rats. *Anat. Anz.*, **159**: 373–383
- Yoshikubo, H., N. Suzuki, K. Takemura, M. Hoso, S. Yashima, S. Iwamuro, Y. Takagi, M. J. Tabata and A. Hattori 2005 Osteoblastic activity and estrogenic response in the regenerating scale of goldfish, a good model of osteogenesis. *Life Sci.*, **76**: 2699–2709