

Effects of Light Intensity on the Growth and Lipid Accumulation of Microalga *Scenedesmus* sp. 11-1 Under Nitrogen Limitation

Junhan Liu · Cheng Yuan · Guangrong Hu · Fuli Li

Received: 8 September 2011 / Accepted: 22 February 2012 /
Published online: 14 March 2012
© Springer Science+Business Media, LLC 2012

Abstract *Scenedesmus* spp. have been reported as potential microalgal species used for the lipid production. This study investigated the effects of light intensity (at three levels: 50, 250, and 400 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) on the growth and lipid production of *Scenedesmus* sp. 11-1 under N-limited condition. Carotenoid to chlorophyll ratio was higher when algae 11-1 grew under 250 and 400 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ than that under 50 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, while protein contents was lower. Highest biomass yield (3.88 g L^{-1}), lipid content (41.1 %), and neutral lipid content (32.9 %) were achieved when algae 11-1 grew at 400 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. Lipid production was slight lower at 250 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ level compared to 400 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. The major fatty acids in the neutral lipid of 11-1 were oleic acid (43–52 %), palmitic acid (24–27 %), and linoleic acid (7–11 %). In addition, polyunsaturated fatty acids had a positive correlation with total lipid production, and monounsaturated fatty acids had a negative one.

Keywords Light intensity · Microalgae · Biodiesel · Neutral lipid · Fatty acid composition

Introduction

Consumption of fossil fuels is one of the most concerned factors that contribute to the climate change. It is a pressing mission for humans to find an alternative fuel. The biodiesel based on soybean, *Trachycarpus fortunei*, rapeseed, and other oil plants has been reported to be suitable as a transport fuel. However, oil plants have low energy intensities per hectare due to the large amount of land use; it is still difficult for biodiesel from oil plants to compete with fossil fuel.

J. Liu · C. Yuan · G. Hu · F. Li (✉)
Key laboratory of Biofuels, Qingdao Institute of BioEnergy and Bioprocess Technology,
Chinese Academy of Sciences, Qingdao 266101, China
e-mail: lifl@qibebt.ac.cn

C. Yuan
Ocean College of Hebei Agricultural University, Qinhuangdao 066003, China

Present Address:

J. Liu
Sinopharm Chemical Reagent Co., Ltd, Shanghai 20002, China

Presently, microalgae are considered to be a potential feedstock for biodiesel production [1]. Compared to other oil plants, usage of microalgae would not be a direct threat of food supplies due to its low cultivation space requirement and high lipid productivity. It was reported that microalgae started to accumulate lipids when some nutrients are exhausted. Lipid contents of various microalgal species range from 20 to 50 % [1], and the lipid content in the same species changes in response to cultivation conditions. Former studies indicated that *Scenedesmus* spp. was a potential microalgal species for lipid production, and the production was relative to incubation conditions, including CO₂ supplement, nutrient condition, and temperature. [2–4].

The fatty acid composition of lipid is a key factor impacting the properties of biodiesel. It was reported that the fatty acid composition of *Scenedesmus* sp. responded to different conditions, including nitrogen concentration and temperature [4, 5]. In addition to nutrient levels and temperature, the light intensity is also an important factor affecting the growth and lipid accumulation, and it could also affect fatty acid composition in the microalgae *Parietochloris incise* and *Dunaliella viridis* [6, 7]. However, effect of the light intensity on lipid production and fatty acid composition in *Scenedesmus* spp. has not been reported.

Microalgae were frequently cultivated at nitrogen deficiency mediums that allowed a nitrogen-free period present during its cultivation to attain higher lipid productivity [5, 8, 9]. In this study, we investigated the effect of light intensity on the growth characteristics, total and neutral lipids accumulations, and fatty acid compositions of neutral lipids of *Scenedesmus* sp. 11-1 under nitrate deficiency condition.

Methods

Strain and Culturing Conditions

Microalga strain 11-1 was isolated from a wastewater pond in Qingdao, China, conserved in culture bank of Qingdao Institute of Bioenergy and Bioprocess Technology, and identified as *Scenedesmus* sp. by 18S rRNA gene sequencing and morphology characterization. The culture medium was modified BG-11, in which the concentration of NaNO₃ was decreased to 4.41 mmol L⁻¹, one fourth of normal BG-11 [10]. The strain was cultivated in a column with a diameter of 42 mm, effective volume of 0.6 L, and blowing air supplemented with CO₂ (2 %, v/v) at bottom at 25 °C. Trichromatic lamps (Y228-T5, NVC Lighting Technology Corporation, China) were used as the light source, and their spectrum range was 380–780 nm. All of the six cultures were cultivated at the light intensity of 50 μmol photons m⁻² s⁻¹ for the first 2 days. They were then divided into three groups grown under high light intensity (HL, 250 μmol photons m⁻² s⁻¹ on each side, 400 μmol photons m⁻² s⁻¹ in total), medium light intensity (ML, 250 μmol photons m⁻² s⁻¹ on one side) and low light intensity (LL, 50 μmol photons m⁻² s⁻¹ on one side), respectively. This experiment was conducted in duplicate.

Lipid Extraction

Microalgal cells were harvested by centrifugation at 3,000 rpm for 10 min. Cell pellets were lyophilized using a freeze drier (ALPHA1-2LD plus, Martin Christ, Germany). The microalgal cells were harvested by centrifugation at 8,000 rpm for 10 min and lyophilized using a freeze drier. The total lipids contained in the algal cells were extracted with a modified chloroform–methanol–water solvent system [11]. Dried algal powder (30–50 mg), 4 ml chloroform, and 2 mL methanol were added into 10-mL glass tube I and shaken for 10 s to disperse the powder. Tube I was incubated at 200 rpm for 12 h at 30 °C, centrifuged for 10 min at

3,500 rpm, and 6 mL of the supernatant was collected and transferred to tube II. Two milliliters methanol and 3.6 ml deionized water were added to reach a final chloroform/methanol/water ratio of 10:10:9. The chloroform layer in tube II was transferred into a preweighed tube III (w1) and dried for 30 min under a flow of N_2 at 60 °C after centrifuging at 3,500 rpm for 5 min. Tube III was dried using a vacuum-drying oven (Lantian DZF-6050, Hangzhou, China) at 0.09 MPa and 60 °C for 1.5 h and then weighed (w2). The lipid weight was calculated by subtracting w1 from w2. The extraction was performed in duplicates for each sample.

Separation of Neutral Lipid and Transesterification

The neutral lipid was separated from the total lipid by column chromatography [12] and weighted.

Methyl esters were extracted from the microalgae lipids by heating in 2 % H_2SO_4 –methanol solution at 85 °C for 2.5 h. After cooling to room temperature, 1 mL of high-performance liquid chromatography grade hexane and 1 mL of saturated NaCl solution were added. The mixture was centrifuged for 10 min at 3,000 rpm, and the upper layer containing fatty acid methyl esters (FAMES) was obtained for further analysis.

FAME samples were analyzed on a gas chromatography (7890A, Agilent, USA)–mass spectrometry (5975C, Agilent, USA) in electron impact mode with a HP-Innowax polyethylene glycol column (30 m × 250 μm × 0.25 μm). The carrier gas was helium. The injection temperature was 250 °C. The initial column temperature was 25 °C. It was elevated to 200 °C at a temperature gradient of 25 °C min^{-1} and was then elevated to 230 °C at a temperature gradient of 3 °C min^{-1} . The temperature was kept constant at 230 °C for 11 min. The fatty acid compositions were recorded as percentages of total fatty acids.

Determination of Nitrate Concentration

The nitrate concentrations were detected by a UV spectrophotometer (Cary 50, Varian, USA) at 220 and 275 nm as described before [13].

Determination of Pigment Concentration

Two milliliters microalgal cell suspension was centrifuged at 3,000 rpm for 5 min, and the cell sediments were resuspended in 5 ml 95 % (v/v) ethanol in dark for 12 h. Afterwards, the cell suspension was centrifuged for 10 min at 3,000 rpm. The absorbance (A) of the




Fig. 1 Growth of *Scenedesmus* sp. 11-1 and consumption of nitrate in BG-11 medium with an initial concentration of nitrate of 4.41 mM

supernatant was measured at 665, 649, and 470 nm using a spectrophotometer (Cary 50, Varian, USA). The concentrations of pigments, including chlorophyll *a*, chlorophyll *b*, chlorophyll, and carotenoid, were calculated by the following equations [14]:

$$C_a = 13.95 \times A_{665} - 6.88 \times A_{649}$$

$$C_b = 24.96 \times A_{649} - 7.32 \times A_{665}$$

$$C_{chl} = C_a + C_b$$

$$C_{car} = \frac{1,000 \times A_{470} - 2.05 \times C_a - 114.8 \times C_b}{245}$$

C_a , C_b , C_{chl} , and C_{car} represented the concentration (mg L^{-1}) of chlorophyll *a*, chlorophyll *b*, chlorophyll, and carotenoid, respectively.

Determination of Protein Yield and Elements Analysis

One milliliter microalgal cells were obtained by centrifugation at 3,000 rpm for 5 min. The cells were resuspended and incubated in a 0.5 mol L^{-1} NaOH solution for 10 min at 100°C , and total protein yield was then measured by BCA Protein Assay Kit (Bestbio, China).

The elements of algae powder were analyzed using an elemental analyzer (Vario EL cube, Elementer, Germany). The heating value Q (kJ g^{-1}) was calculated according to Dulong's formula [15]:

$$Q = 0.338 \times C + 1.442 \times \left(H - \frac{O}{8} \right)$$

where C , H , and O are weight percentages of carbon, hydrogen, and oxygen in algal dried cells, respectively.

Statistical Analysis

Two-tailed paired t tests were applied to ascertain significant differences using SPSS 10, and the level of statistical significance was $P < 0.05$.

Results and Discussion

The Effect of the Light Intensity on the Growth of *Scenedesmus* sp. 11-1

As shown in Fig. 1, it took 6, 8, and 10 days for strain 11-1 to reach stationary phase under the light intensities of 400, 250, and $50 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$, respectively. The total biomass yields were 3.88, 3.62, and 2.55 g L^{-1} under HL, ML, and LL, respectively (Fig. 1).

The difference of growth rates among the HL, ML, and LL cultures indicated that $400 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ was more suitable for the growth of strain 11-1 than other light intensities if nutrients were abundant in the medium. However, nitrogen deficiency became the more important factor to limit the growths of HL and ML cultures after day 4 than light intensity, which could explain why the two cultures had a similar level of final biomass yield. An increased growth rate corresponding to the light intensity was also noted in *Dunaliella tertiolecta* when it was cultivated at 100, 200, and $350 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ [16]. In the study about microalgae *D. viridis*, the growth rate rose with the increased light intensity below the level of $700 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ but decreased when the light intensity

changed from 700 to 1,500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ [17]. The typical midday outdoor light intensity is about 2,000 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ in equatorial regions. The light saturation constants of some algae are much lower than the value of sunlight. For example, the light saturations were 185 [18] and 200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ [19] for *Phaeodactylum tricornerutum* and *Porphyridium cruentum*, respectively. Thus, the high lipid productivity should be achieved in a photobioreactor reducing the light intensity below the light saturation constant when a microalga was cultivated outdoor.

Effect of Light Intensity on Pigment Content

The volumetric contents of carotenoid and chlorophyll both reached their highest level on day 4 in HL and ML cultures (Fig. 2a, b), while they reached the highest level on day 6 in LL




Fig. 2 The changes of contents of chlorophyll (a) and carotenoid (b) and ratios of carotenoid to chlorophyll (c)


culture. The contents of pigments (percentage of dry cell weight) especially chlorophyll decreased significantly from day 2 (Fig. 2a, b). The ratios of the chlorophyll content on day 12 were approximately 1:77, 1:26, and 1:4 of those on day 2 in HL, ML, and LL cultures, respectively. These reductions may be caused by the nitrate depletion since nitrogen is an essential element for the synthesis of chlorophyll. Meanwhile, the light available per cell was reduced due to the increased cells density.

The ratios of carotenoid to chlorophyll increased from 0.16 on day 2 to 1.01, 0.49, and 0.20 on day 12 under HL, ML, and LL, respectively (Fig. 2c), which indicated that the ratios had a positive relationship with light intensity. Given that the spectral absorption peak of carotenoid is in blue light band and that of chlorophyll in both of red and blue light band, the blue light would be more preferable for microalgal growth than the red light if that ratio increased. Previous reports reported that carotenoid contents would increase in the condition of nitrogen starvation [20] or high light intensity [21, 22]. Astaxanthin, as a main part of carotenoids, may play a role in protection against photo damages by reducing amount of light available to the light harvesting pigment–protein complex [23] and acts as an antioxidant-inhibiting lipid preoxidation [24]. The increase in the ratio of carotenoid to chlorophyll was even found to correlate closely with the volumetric content of total fatty acids [25].

Effect of Light Intensity on Protein Content

The percentages of nitrogen element content of dried algal cells were 1.37, 1.47, and 2.14 % in HL, ML, and LL cultures on day 12, respectively (Fig. 3a). There was a negative

Fig. 3 Effect of light intensity on nitrogen element contents on day 12 (a) and protein contents (b)



correlation between nitrogen and lipid content in algal cells, which confirmed the accelerate effect of nitrogen limitation on lipid accumulation.

The contents of protein were 6.1, 6.4, and 11.1 % in dried algal cells from HL, ML, and LL cultures on day 12, respectively (Fig. 3b). The microalgae in LL culture had higher protein content than those in ML and HL cultures after day 4. Previous studies presented similar results: When cells of *P. tricornutum* were grown at 18, 36, and 72 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, the lower light was responsible for a lower biomass yield and lower lipid content, but higher protein content [6].

Effect of Light Intensity on Total Lipids and Neutral Lipids Production

The highest lipid content (41.1 %) and yield (1.60 g L^{-1}) were both obtained from the HL culture on day 12 (Fig. 4a, b). A slightly decreased lipid content (39.2 %) and yield (1.42 g L^{-1}) were found in the cells of ML culture. However, the lipid content (26.2 %) and yield (0.67 g L^{-1}) were significantly lower in LL culture than those in HL and ML cultures. Thus, the biomass yield and lipid content could be promoted by increasing the light intensity from 50 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ to a level at the range of 250–400 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. After cultivation of 12 days, the lipid yields obtained on day 12 were 79, 127, and 245 % higher than those obtained on 6 days cultivation in HL, ML, and LL cultures, respectively (Fig. 4b). Thus, it was suggested to enhance cultivation time to achieve a higher lipid yield of microalgae.

The percentages of neutral lipids of total lipid were 80.0, 82.3, and 62.9 % in microalgal cells in HL, ML, and LL cultures on day 12, respectively (Fig. 5a), and they were corresponding to 32.9, 32.3, and 16.5 % as the percentages of dry cell weight for HL,

Fig. 4 Effect of light intensity on total lipid contents (a) and yields (b) of cells on days 6 and 12





Fig. 5 Effect of light intensity on neutral lipid contents in total lipid (a) and yields (b) on day 12



ML, and LL cultures, respectively. The neutral lipids were only 37.0 and 36.7 % of total lipid in HL and ML cultures, respectively, on day 6. These results indicated a significant accumulation of neutral lipid from days 6 to 12. The harvesting time should be determined not only by lipid productivity but also by lipid content in algal cells in order to cut down the cost in following processes of biodiesel production. Moreover, although there was no significant enhancement in biomass in stationary phase (Fig. 1), the heating value of biomass could be increased due to the accumulation of lipid, an organic with high heating value. The heating value increased from 23.9 kJ g^{-1} on days 6 to 27.6 kJ g^{-1} on day 12 in ML culture, while it increased from 24.6 to 28.2 kJ g^{-1} in HL culture. It is distinctly advantageous to use the biomass with high heating value in a pyrolysis process. Therefore, it could be acceptable to delay harvesting microalgae to a moment after the greatest lipid productivity appeared. The final neutral lipid yields on day 12 were 1.28, 1.16, and 0.38 g L^{-1} in HL, ML, and LL cultures, respectively (Fig. 5b). HL and ML cultures significantly enhanced neutral lipid yields than LL culture, which indicated that 250–400 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ would be a suitable range of light intensity for the accumulation of neutral lipid in column bioreactors used for the batch culturing system. Considering energy income, light intensity of 250 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ would be better.

Triacylglycerol, the main part of neutral lipids, serves as a sink of excessive energy absorbed by photosynthetic apparatus in microalgae [26]. Triacylglycerol synthesis requires large amounts of ATP and NAD(P)H produced by the photosynthesis. Therefore, it may be helpful in the dissipation of excess light energy and prevention of the photochemical damage of algal cells [27]. Thus, more carbon flux generated from the photosynthesis is channeled to

the lipid accumulation on a unit biomass basis when individual cell is exposed to a large quantity of light energy [7, 28]. Therefore, the high and medium lights could promote the final biomass yield, lipid content, and yield, especially neutral lipid content.

Fatty Acid Composition of Neutral Lipids

The major fatty acids of neutral lipid were oleic acid (43–52 %), palmitic acid (24–27 %), and linoleic acid (7–11 %) in strain 11-1 (Table 1). Eighty percent of the total fatty acids were composed of the three fatty acids. The fatty acids with carbon-16 and carbon-18 atoms were the main components in the neutral lipid, which was similar to traditional vegetable oils and adequate for biodiesel production [29–31].

The fatty acid profiles of neutral lipids were similar when strain 11-1 grew under HL and ML. Higher polyunsaturated fatty acid (PUFA) contents and lower monounsaturated fatty acid (MUFA) contents were noted in the cells when it grew under HL and ML compared to LL (Table 1). The increase in PUFA associated to light intensity was also noted in the microalgae *Pavlova lutheri* [28, 32]. By contrast, Liang et al. [33] reported that six strains of marine diatoms had higher contents of PUFA in total lipid when growing under a lower light intensity. They also found that the increase in PUFA and the decrease in MUFA were correlated with the increased neutral lipid content [33]. These results were similar to the study of Solovchenko et al. [7] but opposite to the study of Lin et al. [5]. Although the lipid containing a high content of PUFA may make the biodiesel susceptible to oxidation, it is easy to improve the saturation degree in certain chemical processes [1].

Table 1 Fatty acid composition of neutral lipid under different light intensities

	400 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (%, total fatty acids)	250 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (%, total fatty acids)	50 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (%, total fatty acids)
Saturated fatty acids (SFAs)			
C16:0	25.50±1.09 ^a	26.06±0.06	24.28±0.25
C18:0	4.98±0.03	5.28±0.24	5.35±0.15
C20:0	1.19±0.00	1.26±0.01	0.72±0.04
Sum	31.67±1.12	32.60±0.08	30.35±0.36
Monounsaturated fatty acids (MUFAs)			
C16:1	1.80±0.10	1.39±0.03	1.27±0.00
C18:1	43.75±0.09	44.13±0.37	51.96±1.31
C20:1	0.14±0.20	0.29±0.01	0.16±0.01
sum	45.70±0.21	45.81±0.34	53.40±1.32
Polyunsaturated fatty acids (PUFAs)			
C16:2	3.18±0.12	2.79±0.08	1.47±0.14
C16:3	0.94±0.12	0.66±0.16	0.32±0.01
C16:4	0.98±0.06	1.16±0.04	1.67±0.25
C18:2	10.51±0.16	10.34±0.01	7.40±0.20
C18:3	5.02±0.35	4.83±0.02	4.34±0.29
C18:4	1.46±0.01	1.35±0.03	1.07±0.05
sum	22.11±0.80	21.12±0.33	16.27±0.93

^a Results expressed as percentage of the total fatty acids. Data presented as mean values±standard deviation (SD) of duplicates (each treatment)

Although a higher lipid yield of 2.16 g L^{-1} ($0.12 \text{ g L}^{-1} \text{ day}^{-1}$ as the volumetric productivity) was recorded in *Scenedesmus obliquus*, it should be noted that a supplementation of 1.5 % glucose in the first 10 days' cultivation and an optimization of nutrient conditions in the next 8 days' cultivation were present [3]. Previous studies showed a volumetric lipid productivity of $0.065 \text{ g L}^{-1} \text{ day}^{-1}$ in *Neochloris oleoabundans* and $0.079 \text{ g L}^{-1} \text{ day}^{-1}$ in *S. obliquus*. They were both considered to be excellent lipid producers [2, 9]. The highest volumetric lipid productivity of 11-1 was $0.133 \text{ g L}^{-1} \text{ day}^{-1}$ in 12-day cultivation. Meanwhile, strain 11-1 showed higher lipid productivity than the other 20 strains isolated in the same period of our lab, whose lipid productivities ranged from 0.043 to $0.080 \text{ g L}^{-1} \text{ day}^{-1}$. According to the previous reports [2, 8], *Scenedesmus* sp. 11-1 could be an excellent candidate used for biodiesel production due to its high lipid (especially neutral lipid) productivity and high oleic acid content. Moreover, *Scenedesmus* sp. could be cultivated using the nutrients in wastewater [34, 35], which provided a way to realize the economic viability in industry [36, 37]. Meanwhile, the light intensities used in the studies of Mandal and Mallick [3] and Ho et al. [2] were 75 and $60 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$, respectively, which may be low to achieve the highest lipid productivity. A higher light intensity, like $250 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$, was suggested for evaluating lipid productivity of *Scenedesmus* sp.

Conclusion

The high light intensities at a range of $250\text{--}400 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ were found to be beneficial to the growth and lipid accumulation of *Scenedesmus* sp. 11-1 under nitrogen limited condition. The amounts of total and neutral lipids were significantly higher in the late growth stage (days 6–12) due to the nitrogen limitation. The increase in light intensity promoted the content of PUFA in neutral lipid and the ratio of carotenoid to chlorophyll. The highest total and neutral lipid productivities of strain 11-1 were 0.133 and $0.106 \text{ g L}^{-1} \text{ day}^{-1}$, respectively. Microalgae strain 11-1 has potential to be a promising candidate for biodiesel production.

Acknowledgment This study was supported by grant KGXC2-YW-374-3 from Chinese Academy of Sciences and grant 2008ZX07422-003-5-2 from the Ministry of Science and Technology of China.

References

1. Chisti, Y. (2007). *Biotechnology Advances*, 25, 294–306.
2. Ho, S., Chen, W., & Chang, J. (2010). *Bioresource Technology*, 101, 8725–8730.
3. Mandal, S., & Mallick, N. (2009). *Applied Microbiology and Biotechnology*, 84, 281–291.
4. Li, X., Hu, H., & Zhang, Y. (2010). *Bioresource Technology*, 102, 3098–3102.
5. Lin, Q., & Lin, J. (2010). *Bioresource Technology*, 102, 1615–1621.
6. Qian, K., & Michael, B. (1993). *Applied Biochemistry and Biotechnology*, 38, 93–103.
7. Solovchenko, A. E., Khozin-Goldberg, I., Didi-Cohen, S., Cohen, Z., & Merzlyak, M. N. (2008). *Journal of Applied Phycology*, 20, 245–251.
8. Griffiths, M., & Harrison, S. (2009). *Journal of Applied Phycology*, 21, 493–507.
9. Pruvost, J., Van Vooren, G., Cogne, G., & Legrand, J. (2009). *Bioresource Technology*, 100, 5988–5995.
10. Andersen, R. (2005). *Algal culturing techniques*. Burlington: Elsevier.
11. Bligh, E. G., & Dyer, W. J. (1959). *Canadian Journal of Biochemistry and Physiology*, 37, 911–917.
12. Borgström, B. (1952). *Acta Physiologica Scandinavica*, 25, 101–110.
13. Beschkov, V., Velizarov, S., Agathos, S. N., & Lukova, V. (2004). *Biochemical Engineering Journal*, 17, 141–145.
14. Lichtenthaler, H. K., & Wellburn, A. R. (1983). *Biochemical Society Transactions*, 603, 591–592.

15. Minowa, T., Yokoyama, S-y, Kishimoto, M., & Okakurat, T. (1995). *Fuel*, *74*, 1735–1738.
16. Tang, H., Abunasser, N., Garcia, M. E. D., Chen, M., Simon Ng, K. Y., & Salley, S. O. (2010). *Applied Energy*, *88*, 3324–3330.
17. Gordillo, F., Goutx, M., Figueroa, F., & Niell, F. (1998). *Journal of Applied Phycology*, *10*, 135–144.
18. Mann, J. E., & Myers, J. (1968). *Journal of Phycology*, *4*, 349–355.
19. Molina, E., Ación Fernández, F. G., García Camacho, F., Camacho Rubio, F., & Chisti, Y. (2000). *Journal of Applied Phycology*, *12*, 355–368.
20. Boussiba, S., & Vonshak, A. (1991). *Plant & Cell Physiology*, *32*, 1077–1082.
21. Solovchenko, A. E., Khozin-Goldberg, I., Didi-Cohen, S., Cohen, Z., & Merzlyak, M. N. (2008). *Russian Journal of Plant Physiology*, *55*, 455–462.
22. Zhekisheva, M., Boussiba, S., Khozin-Goldberg, I., Zarka, A., & Cohen, Z. (2002). *Journal of Phycology*, *38*, 325–331.
23. Bidigare, R. R., Ondrusek, M. E., Kennicutt, M. C., Iturriaga, R., Harvey, H. R., Hoham, R. W., et al. (1993). *Journal of Phycology*, *29*, 427–434.
24. Hagen, C., Braune, W., & Greulich, F. (1993). *Journal of Photochemistry and Photobiology B: Biology*, *20*, 153–160.
25. Solovchenko, A., Khozin-Goldberg, I., Cohen, Z., & Merzlyak, M. (2009). *Journal of Applied Phycology*, *21*, 361–366.
26. Hu, Q., Sommerfeld, M., Jarvis, E., Ghirardi, M., Posewitz, M., Seibert, M., et al. (2008). *The Plant Journal*, *54*, 621–639.
27. Roessler, P. G. (1990). *Journal of Phycology*, *26*, 393–399.
28. Courchesne, N. M. D., Parisien, A., Wang, B., & Lan, C. Q. (2009). *Journal of Biotechnology*, *141*, 31–41.
29. Sydney, E. B., da Silva, T. E., Tokarski, A., Novak, A. C., de Carvalho, J. C., Woiciechowski, A. L., et al. (2010). *Applied Energy*, *88*, 3291–3294.
30. Abou-Shanab, R. A. I., Hwang, J., Cho, Y., Min, B., & Jeon, B. (2011). *Applied Energy*, *88*, 3300–3306. doi:10.1016/j.apenergy.2011.01.060.
31. Harrington, K. J. (1986). *Biomass*, *9*, 1–17.
32. Guihéneuf, F., Mimouni, V., Ulmann, L., & Tremblin, G. (2009). *Journal of Experimental Marine Biology and Ecology*, *369*, 136–143.
33. Liang, Y., Mai, K., Sun, S., & Yu, D. (2001). *Chinese Journal of Oceanology and Limnology*, *19*, 249–254.
34. Kim, M. K., Park, J. W., Park, C. S., Kim, S. J., Jeune, K. H., Chang, M. U., et al. (2007). *Bioresource Technology*, *98*, 2220–2228.
35. Bhatnagar, A., Chinnasamy, S., Singh, M., & Das, K. C. (2010). *Applied Energy*, *88*, 1–7.
36. Pittman, J. K., Dean, A. P., & Osundeko, O. (2011). *Bioresource Technology*, *102*, 17–25.
37. Rawat, I., Ranjith Kumar, R., Mutanda, T., & Bux, F. (2010). *Applied Energy*, *88*, 3411–3424. doi:10.1016/j.apenergy.2010.11.025.