

Erratum to: Effects of redox-active interlayer anions on the oxygen evolution reactivity of NiFe-layered double hydroxide nanosheets

Daojin Zhou¹, Zhao Cai¹, Yongmin Bi¹, Weiliang Tian^{1,2}, Ma Luo¹, Qian Zhang¹, Qian Zhang¹, Qixian Xie¹, Jindi Wang¹, Yaping Li¹, Yun Kuang¹, Xue Duan¹, Michal Bajdich³, Samira Siahrostami⁴ (✉), and Xiaoming Sun¹ (✉)

¹ State Key Laboratory of Chemical Resource Engineering, College of Energy, Beijing Advanced Innovation Center for Soft Matter Science and Engineering, Beijing University of Chemical Technology, Beijing 100029, China

² Key Laboratory of Chemical Engineering in South Xinjiang, College of Life Science, Tarim University, Alar 843300, China

³ SUNCAT Center for Interface Science and Catalysis, SLAC National Accelerator Laboratory, Menlo Park, California 94025, USA

⁴ SUNCAT Center for Interface Science and Catalysis, Department of Chemical Engineering, Stanford University, Stanford, California 94305, USA

© Tsinghua University Press and Springer-Verlag GmbH Germany, part of Springer Nature 2019.

Erratum to

Nano Research 2018, 11(3): 1358–1368

https://doi.org/10.1007/s12274-017-1750-9

The labels in Fig. 8 in the original version of this article were unfortunately misplaced. The corrected figure is as follow.




Figure 8 Binding energy and Bader charge of (a) Ni and (b) Fe sites as a function of the experimentally measured onset potential in as-prepared LDHs (●: Binding energy; ▲: Bader charge).

The online version of the original article can be found at
https://doi.org/10.1007/s12274-017-1750-9