



**VICTORIA UNIVERSITY**  
MELBOURNE AUSTRALIA

*Effects of Resistance Training on Muscle Size and Strength in Very Elderly Adults: A Systematic Review and Meta-Analysis of Randomized Controlled Trials*

This is the Accepted version of the following publication

Grgic, Jozo, Garofolini, Alessandro, Orazem, John, Sabol, Filip, Schoenfeld, Brad J and Pedisic, Zeljko (2020) Effects of Resistance Training on Muscle Size and Strength in Very Elderly Adults: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Sports Medicine*, 50. pp. 1983-1999. ISSN 0112-1642

The publisher's official version can be found at  
<https://link.springer.com/article/10.1007%2Fs40279-020-01331-7>  
Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/41906/>

1 **Resistance training increases muscle size and strength in very elderly adults: a**  
2 **systematic review and meta-analysis of randomized controlled trials**

3 Jozo Grgic,<sup>1</sup> Alessandro Garofolini,<sup>1</sup> John Orazem,<sup>2</sup> Filip Sabol,<sup>3,4</sup> Brad J. Schoenfeld,<sup>5</sup> Zeljko  
4 Pedisic<sup>1</sup>

5

6 <sup>1</sup>Institute for Health and Sport (IHES), Victoria University, Melbourne, Australia

7 <sup>2</sup>School of Health Sciences, Human Services and Nursing, Lehman College, Bronx, USA

8 <sup>3</sup>Fitness Academy, Zagreb, Croatia

9 <sup>4</sup>Faculty of Kinesiology, University of Zagreb, Zagreb, Croatia

10 <sup>5</sup>Department of Health Sciences, Lehman College, Bronx, USA

11 **Short title:** Resistance training and the very elderly

12 **Corresponding author:**

13 Jozo Grgic

14 Institute for Health and Sport (IHES), Victoria University, Melbourne, Australia

15 [jozo.grgic@live.vu.edu.au](mailto:jozo.grgic@live.vu.edu.au)

## 16 **Abstract**

17 **Background:** Effects of resistance training on muscle strength and hypertrophy are well-  
18 established in adults and younger elderly. However, less is currently known about these  
19 effects in the very elderly (i.e., 75 years of age and older).

20 **Objective:** To examine the effects of resistance training on muscle size and strength in very  
21 elderly individuals.

22 **Methods:** Randomized controlled studies that explored the effects of resistance training in  
23 very elderly on muscle strength, handgrip strength, whole-muscle hypertrophy, and/or muscle  
24 fiber hypertrophy were included in the review. Meta-analyses of effect sizes (ESs) were used  
25 to analyze the data.

26 **Results:** Twenty-two studies were included in the review. The meta-analysis found a  
27 significant effect of resistance training on muscle strength in the very elderly (difference in  
28 ES = 0.97; 95% confidence interval [CI]: 0.50, 1.44;  $p = 0.001$ ). In a subgroup analysis that  
29 included only the oldest-old participants (80+ years of age), there was a significant effect of  
30 resistance training on muscle strength (difference in ES = 1.28; 95% CI: 0.28, 2.29;  $p =$   
31 0.020). For handgrip strength, we found no significant difference between resistance training  
32 and control groups (difference in ES = 0.26; 95% CI: -0.02, 0.54;  $p = 0.064$ ). For whole-  
33 muscle hypertrophy, there was a significant effect of resistance training in the very elderly  
34 (difference in ES = 0.30; 95% CI: 0.10, 0.50;  $p = 0.013$ ). We found no significant difference  
35 in muscle fiber hypertrophy between resistance training and control groups (difference in ES  
36 = 0.33; 95% CI: -0.67, 1.33;  $p = 0.266$ ). There were minimal reports of adverse events  
37 associated with the training programs in the included studies.

38 **Conclusions:** We found that very elderly can increase muscle strength and muscle size by  
39 participating in resistance training programs. Resistance training was found to be an effective  
40 way to improve muscle strength even among the oldest-old.

41 **Key points:**

42 ► We found that very elderly adults can increase their muscle strength and size by  
43 participating in resistance training programs.

44 ► These effects were observed with resistance training interventions that generally included  
45 low weekly training volumes and frequencies.

46 ► There were minimal reports of adverse events associated with the training programs.

47

48

## 49 **1 Introduction**

50 Dynapenia is the age-associated loss of muscle strength [1]. Low muscle strength increases  
51 the risk of mobility limitations and mortality in older adults [1-4]. Sarcopenia is a progressive  
52 skeletal muscle characterized by a degenerative loss of muscle mass and function [5]. It is  
53 associated with an increased likelihood of physical disability, falls, fractures, and mortality  
54 [5]. Resistance training is the most widely recognized mode of exercise for increasing muscle  
55 strength and muscle size. The effectiveness of resistance training in achieving these outcomes  
56 among youth, adults, and older adults is well established [6-8]. The effects of resistance  
57 training on older adults have been recently reviewed by Fragala et al. [9]. However, this  
58 review considered studies conducted among adults aged 50 years and older, with less focus  
59 placed on the effects of resistance training on muscle strength and hypertrophy in the very  
60 elderly (i.e., 75 years of age and older) [10, 11].

61

62 Muscle hypertrophy occurs when muscle protein synthesis exceeds muscle protein  
63 degradation over time [12]. Research has established that, compared to their younger  
64 counterparts, older adults experience a reduced muscle protein synthetic response to protein  
65 intake, a physiological adaptation termed "anabolic resistance" [13]. Muscle hypertrophy in  
66 response to resistance training is associated with myonuclear addition via satellite cell  
67 recruitment [14]. In this context, data suggest that resistance training induces significant  
68 addition of myonuclei per muscle fiber in young adults [15]. However, no significant satellite  
69 cell or myonuclear addition was found in older adults that performed 12-16 weeks of  
70 resistance training [15, 16]. Therefore, some researchers speculate that there might be an age-  
71 related ceiling above which an individual cannot further increase muscle size with resistance  
72 training [17]. Additionally, there are estimates that older individuals have up to a 47%

73 reduction in the number of motor units, and this reduction might be associated with  
74 compromised gains in muscle strength with resistance training in this population [18, 19].

75

76 The seminal work by Fiatarone et al. [20] suggested that participation in resistance training  
77 increases muscle strength and muscle size, even at the advanced stages of aging. In this  
78 single-arm study, ten participants with an average age of 90 years (range: 86 to 96 years)  
79 performed eight weeks of resistance training. After the intervention, knee extension one-  
80 repetition maximum (1RM) strength improved by 15 kg, accompanied by an increase in  
81 quadriceps muscle size of 9%. However, in a more recent randomized controlled study [16],  
82 12-weeks of resistance training in a group of participants aged 83 to 94 years did not  
83 significantly increase their muscle size.

84

85 In 2013, a systematic review by Stewart et al. [11] provided a summary of studies that  
86 explored the effects of different modes of physical training (including resistance training) on  
87 muscle size and strength in adults aged 75 years or older. Even though this review concluded  
88 that resistance training is an effective exercise intervention for increasing muscle size and  
89 strength in this age group, the conclusions were based only on two included studies. It is  
90 important to note that several studies that satisfied the inclusion criteria of Stewart et al. [10]  
91 were not identified and included in the review [21-29]. Furthermore, since 2013, new original  
92 studies have been published on this topic, adding new relevant data to further our  
93 understanding of muscular adaptations to resistance training in very elderly adults [16, 30-34].

94

95 The aim of this systematic review and meta-analysis was, therefore, to examine the effects of  
96 resistance training on strength and muscle size in very elderly individuals. A systematic

97 review on this topic is needed, given that: (a) the evidence presented in studies examining the  
98 effects of resistance training in this age group is conflicting; and (b) there are no recent  
99 systematic reviews on this topic. Findings on this topic could have a substantial public health  
100 impact because the very elderly represent one of the fastest-growing age groups in the  
101 population, and it is estimated that only 8.7% of adults aged 75 years or older participate in  
102 muscle-strengthening activities [35, 36].

103

## 104 **2 Methods**

### 105 **2.1 Search strategy**

106 For this systematic review, we followed the Preferred Reporting Items for Systematic  
107 Reviews and Meta-Analyses guidelines [37]. In total, we searched through nine databases:  
108 Academic Search Elite, CINAHL, ERIC, Open Access Theses and Dissertations, Open  
109 Dissertations, PsycINFO, PubMed/MEDLINE, Scopus, and SPORTDiscus. In all of these  
110 databases, we used the following search syntax (or equivalent) to search through titles,  
111 abstracts, and keywords of indexed documents: ("very elderly" OR "oldest old" OR "oldest-  
112 old" OR "very old" OR "advancing age" OR "advancing years" OR "old-old" OR "old old"  
113 OR septuagenarian\* OR nonagenarian\* OR octogenarian\* OR centenarian\* OR "75 and  
114 older" OR "80 and older" OR "85 and older" OR "90 and older" OR "95 and older" OR "75  
115 years" OR "80 years" OR "85 years" OR "90 years" OR "95 years") AND ("resistance  
116 training" OR "resistance exercise" OR "weight lifting" OR "weightlifting" OR "strength  
117 exercise" OR "strength training" OR "strengthening" OR "resistive exercise" OR "resistive  
118 training") AND ("muscle hypertrophy" OR "muscular hypertrophy" OR "muscle mass" OR  
119 "lean body mass" OR "fat-free mass" OR "fat free mass" OR "muscle fiber" OR "muscle size"  
120 OR "muscle fibre" OR "muscle thickness" OR "cross-sectional area" OR "cross sectional

121 area" OR "computed tomography" OR "magnetic resonance imaging" OR "muscle power"  
122 OR "strength" OR "1RM" OR "isokinetic" OR "isometric"). We also performed secondary  
123 searches that consisted of: (a) screening the reference lists of studies that were included in the  
124 review; and (b) examining the reference lists of previous related reviews [7, 11, 38-43]. To  
125 reduce the probability of study selection bias, two authors of the review (JG and AG)  
126 conducted the study selection independently. After both authors completed their searches, the  
127 lists of included and excluded studies were compared between them. Any discrepancies  
128 between the two authors in the included and excluded studies were resolved through  
129 discussion and agreement. The databases were searched on January 20<sup>th</sup>, 2020.

130

## 131 **2.2 Inclusion criteria**

132 Studies that satisfied the following criteria were included in the review: (a) the participants  
133 were aged 75 years or older; (b) the participants were randomized into the intervention and  
134 control group(s); (c) the exercise intervention was comprised of resistance training while the  
135 control group did not exercise; (d) the study assessed muscle strength and/or muscle size pre-  
136 and post-intervention; and (e) the training protocol lasted for a minimum of six weeks. All  
137 forms of strength tests, including isotonic, isometric, isokinetic, and handgrip tests were  
138 deemed relevant. For muscle hypertrophy, we considered studies that assessed changes at the  
139 whole-muscle (macroscopic methods) and/or muscle fiber level (microscopic methods).

140

## 141 **2.3 Data extraction**

142 In each of the included studies, we extracted the following data: (a) author names and year of  
143 publication; (b) characteristics of the sample size, including their age and sex; (c) specifics of  
144 the resistance training intervention (e.g., the number of performed sets, exercise selection); (d)



145 adverse events reported during the intervention (if any); (e) exercise used for the muscle  
146 strength test and/or body site and tool used for the muscle hypertrophy assessment; and (f) pre  
147 and post-intervention mean  $\pm$  standard deviation (SD) of the strength and/or hypertrophy  
148 outcomes. For the studies that reported standard errors, we converted them to SDs. Two  
149 authors of the review (JG and FS) performed the data extraction independently. After both  
150 authors completed the data extraction from all studies, the coding sheets were compared  
151 between the authors. In case of any discrepancies in the data extraction files, the data was re-  
152 checked from the studies.

153

#### 154 **2.4 Methodological quality**

155 The methodological quality of the included studies was assessed using the 27-item Downs and  
156 Black checklist [44]. This checklist evaluates different aspects of the study design, with items  
157 1–10 referring to reporting, items 11–13 referring to external validity, items 14–26 referring  
158 to internal validity, and item 27 referring to statistical power. Given that the included studies  
159 explored the effects of a resistance training intervention, the standard 27-item checklist was  
160 modified by adding two items, item 28 and item 29. Item 28 was on the reporting of  
161 adherence to the training program, while item 29 was related to training supervision. For each  
162 item—including items 28 and 29—one point was allocated to the study if the criterion was  
163 satisfied; no points were allocated if the criterion was not satisfied. The maximum possible  
164 score on the modified version of the Downs and Black checklist was, therefore, 29 points.  
165 Based on the summary score, studies that had 21–29 points were classified as being of ‘good  
166 quality’, studies with 11–20 points were classified as being of ‘moderate quality’, while  
167 studies that scored less than 11 points were considered to be of ‘poor quality’ [45, 46] The  
168 methodological quality assessment was performed independently by two authors (JG and  
169 AG), with discussions and agreement for any observed differences in the initial scoring.

170

171 **2.5 Statistical analysis**

172 The meta-analyses for strength and hypertrophy outcomes were performed on the training  
173 intervention minus control difference in relative effect sizes (ESs). The data for strength and  
174 hypertrophy were converted to relative ES, calculated as the posttest-pretest mean change in  
175 each group, divided by the pooled pretest SD, with an adjustment for small sample bias [47].  
176 The variance of the ESs depends on the within-subject posttest-pretest correlation. Given that  
177 this correlation was not reported in any of the included studies, when possible it was  
178 estimated by back-solving from paired t-test *p*-values or SDs of posttest-pretest change scores.  
179 Among studies for which the correlation could be derived from the available data, the median  
180 value was 0.85. A more conservative value of 0.75 was used for all studies. Sensitivity  
181 analyses (not presented) were performed using correlations ranging from 0.25 to 0.85, and  
182 their results were consistent with those using 0.75. In order to account for correlated ESs  
183 within studies, we used a robust variance meta-analysis model, with an adjustment for small  
184 samples [48]. In the main meta-analysis for muscle strength, we included all available studies.  
185 A sensitivity analysis was performed by excluding the two studies [26, 29] that used upper-  
186 body exercises for the strength test. In a subgroup analysis, we explored the effects of  
187 resistance training on muscle strength only among the “oldest-old” (i.e., 80+ years). Handgrip  
188 strength was analyzed separately from other strength tests as this test is commonly used alone  
189 in predicting mortality and functional declines in the very elderly [49]. For hypertrophy, the  
190 following meta-analyses were performed: (a) for whole-muscle hypertrophy outcomes; and  
191 (b) for muscle fiber cross-sectional area (CSA). All differences in ESs were presented with  
192 their 95% confidence intervals (95% CIs). These differences were interpreted as: “trivial”  
193 ( $\leq 0.20$ ); “small” (0.21–0.50); “medium” (0.51–0.80); and “large” ( $> 0.80$ ). The potential  
194 presence publication bias was checked by examining funnel plot asymmetry and calculating

195 trim-and-fill estimates. The trim-and-fill estimates (not presented) were similar to the main  
196 results. Heterogeneity was explored using the  $I^2$  statistic, with values of  $\leq 50\%$ , 50–75%, and  
197  $>75\%$  indicating low, moderate, and high levels of heterogeneity, respectively. All meta-  
198 analyses were performed using the robumeta package within R version 3.6.1 and the trim-and-  
199 fill analyses were calculated using the metafor package [50, 51]. Group differences were  
200 considered statistically significant at  $p < 0.05$ .

201

## 202 **3 Results**

### 203 **3.1 Study selection**

204 The total number of search results in the nine databases was 2076. After excluding 2016  
205 search results based on title or abstract, 60 full-text papers were read. Of the 60 full-text  
206 papers, 17 studies were included. Secondary searches resulted in another 1559 search results  
207 and with the inclusion of five additional papers (Figure 1). Therefore, the final number of  
208 included studies was 22 [16, 21-34, 52-58]. Of note, in two cases, the strength and whole-  
209 muscle hypertrophy data were published separately from muscle fiber CSA data, even though  
210 the data collection was carried out in the same cohort [16, 30, 52, 53]. Additionally, one group  
211 of authors published the data on strength, whole-muscle CSA, and muscle fiber CSA in three  
212 separate papers, even though the data was collected in a single study [54-56].

213

### 214 **3.2 Study characteristics**

#### 215 **3.2.1 Muscle strength outcomes**

216 In the seventeen studies that explored muscle strength outcomes and met the inclusion  
217 criteria, the pooled number of participants was 880 (84% females; Table 1). The median  
218 sample size per study was 38 (range: 14 to 144 participants). The interventions lasted from 8

219 to 18 weeks. Training frequency was from 1 to 3 days per week. Eleven studies used  
220 isometric strength tests, four used isotonic strength tests, and three used isokinetic tests (one  
221 used both isometric and isokinetic tests). Two studies employed tests on upper-body  
222 exercises, while the remaining studies used lower body exercises (Table 2). Eight studies  
223 assessed handgrip strength (Table 2).

224

### 225 **3.2.2 Hypertrophy outcomes**

226 In the nine studies that explored hypertrophy outcomes and met the inclusion criteria, the total  
227 sample size was 204 participants (67% females; Table 1). The median sample size per study  
228 was 26 participants (range: 23 to 49 participants). The interventions lasted from 10 to 18  
229 weeks, with a training frequency of 2 to 3 days per week. Six studies reported data on whole-  
230 muscle hypertrophy. For this outcome, studies used computed tomography (three studies), B-  
231 mode ultrasound (two studies), and magnetic resonance imaging (one study). Three studies  
232 explored changes at the muscle fiber level. All studies assessed lower-body hypertrophy. The  
233 training programs used in the studies are summarized in Table 2.

234

### 235 **3.3 Methodological quality**

236 The average score on the modified 29-item Downs and Black checklist was 25 (range: 21 to  
237 28 points). All studies were classified as being of good methodological quality. Scores on all  
238 items of the checklist are reported in Table 3.

239

### 240 **3.4 Meta-analysis results for muscle and handgrip strength**

241 The meta-analysis found a significant effect of resistance training on muscle strength in the  
242 very elderly (difference in ES = 0.97; 95% CI: 0.50, 1.44;  $p = 0.001$ ;  $I^2 = 87\%$ ; Figure 2). In  
243 the sensitivity analysis, there was a significant effect of resistance training on lower-body

244 muscle strength in the very elderly (difference in ES = 0.96; 95% CI: 0.48, 1.45;  $I^2 = 87%$ ;  $p$   
245 = 0.001). In a subgroup analysis that included only the oldest-old participants (80+ years of  
246 age), there was a significant effect of resistance training on muscle strength (difference in ES  
247 = 1.28; 95% CI: 0.28, 2.29;  $p = 0.020$ ;  $I^2 = 86%$ ; Figure 3). For handgrip strength, we found  
248 no significant difference between resistance training and control groups (difference in ES =  
249 0.26; 95% CI: -0.02, 0.54;  $p = 0.064$ ;  $I^2 = 51%$ ; Figure 4).

250

### 251 **3.5 Meta-analysis results for whole-muscle and muscle fiber hypertrophy**

252 For whole-muscle hypertrophy, there was a significant effect of resistance training in the very  
253 elderly (difference in ES = 0.30; 95% CI: 0.10, 0.50;  $p = 0.013$ ;  $I^2 = 0%$ ; Figure 5). We found  
254 no significant difference in muscle fiber hypertrophy between resistance training and control  
255 groups (difference in ES = 0.33; 95% CI: -0.67, 1.33;  $p = 0.266$ ;  $I^2 = 7%$ ; Figure 6).

256

## 257 **4 Discussion**

258 The main finding of this systematic review and meta-analysis is that resistance training  
259 increases muscle strength in very elderly people, even among the oldest-old. We also found  
260 that resistance training results in muscle hypertrophy at the whole-muscle level in very  
261 elderly. The ES for strength and whole-muscle hypertrophy was large and small, respectively.  
262 Even though the pooled ES favored resistance training for muscle fiber hypertrophy and  
263 handgrip strength, these effects were not statistically significant.

264

### 265 **4.1 Muscle strength**

266 We found that resistance training produced substantial increases in muscle strength in the very  
267 elderly. Increases in muscle strength were also observed in a subgroup analysis of studies that  
268 included the oldest-old suggesting that resistance training enhances muscle strength even at an

269 advanced stage of aging. Xue et al. [59] reported that dynapenia is associated with increased  
270 mortality risk. Findings from the “Health, Aging and Body Composition Study” further  
271 indicated that knee extension strength—as measured by isokinetic dynamometry—is  
272 associated with a reduced risk of mortality [3]. Dynapenia also increases the risk of physical  
273 disability and reduces physical performance [1]. Therefore, muscle strength is identified as  
274 one of the key muscle qualities for physical independence in the very elderly [1, 4]. After the  
275 age of 75 years, muscle strength annually declines by about 2% to 4% (ES: 0.17 to 0.24) for  
276 those who do not perform regular resistance exercise [60-62]. Our findings suggest that  
277 participation in resistance training over 8 to 18 weeks, with a frequency of 1 to 3 days per  
278 week, can restore strength that has been potentially lost over several years of inactivity.  
279 Research has also established that lower limb muscle weakness is an important risk factor for  
280 falls in the older population [63]. When considering only the studies that used lower-body  
281 exercise for the strength test, an ES of 0.96 (95% CI: 0.48, 1.45) was found. These data  
282 highlight that increasing muscle strength through resistance training participation could be of  
283 great health benefit for the very elderly. Our findings are, therefore, highly relevant from a  
284 public health perspective. Moreover, data suggests that only 8.7% of adults aged 75 years and  
285 older participate in muscle-strengthening activities [36]. Thus, it is clear that finding ways to  
286 further promote participation and adherence to muscle-strengthening activities in this age  
287 group is of considerable public health interest.

288

#### 289 **4.2 Handgrip strength**

290 The handgrip strength test is widely used to evaluate muscle strength as it is noninvasive and  
291 inexpensive [64]. Given its simplicity, this test is often utilized in epidemiological studies  
292 [49]. In the sample of included studies, the pooled ES favored resistance training condition,  
293 but the effect was not statistically significant ( $p = 0.064$ ). In one of the included studies,

294 resistance training focused exclusively on the lower body, but strength was evaluated using  
295 the handgrip test [31]. This might not be entirely appropriate, given that the largest increases  
296 in strength are expected for the muscle groups that were covered in the training program [65,  
297 66]. Indeed, one study reported that 24 weeks of whole-body resistance training produced a  
298 substantial increase in 1RM knee extension and leg press strength (on average by 21 and 45  
299 kg, respectively), that were not accompanied by any significant changes in handgrip strength  
300 [67]. In line with this finding, some authors have speculated that there is only a limited ability  
301 to increase handgrip strength in adulthood [68]. While handgrip strength testing can certainly  
302 provide valuable information about physical functioning, the use of this test may, in some  
303 cases, provide limited insights into the efficacy of a given resistance training program.

304

### 305 **4.3 Whole-muscle hypertrophy**

306 We found that very elderly individuals can increase muscle size despite their advancing age,  
307 although the expected improvements may be small to modest (ES = 0.30; 95% CI: 0.10, 0.50).  
308 Nonetheless, the finding that the very elderly can increase their muscle size is highly relevant,  
309 given that sarcopenia may increase the risk of falls and fractures, increase frailty, decrease  
310 functional independence and quality of life as well as increase the risk of chronic disease and  
311 all-cause mortality [4]. There are estimates that in the very elderly muscle size is reduced at a  
312 rate of 0.64% to 0.98% per year (ES: 0.14 to 0.23) [60, 62]. Our results suggest that resistance  
313 training interventions lasting from 10 to 18 weeks with a training frequency of 2 to 3 days per  
314 week can increase muscle size that was potentially lost over multiple years of aging. This  
315 finding is of public great health importance, if we consider estimates that the prevalence of  
316 sarcopenia in adults older than 75 years ranges from 27% to 60% [69].

317

### 318 **4.4 Muscle fiber hypertrophy**

319 Despite the findings observed for whole-muscle hypertrophy, we did not find significant  
320 increases in muscle fiber CSA, even though in the sample of included studies the pooled ES  
321 of 0.33 favored resistance training. The lack of a significant finding in this analysis could be  
322 attributed to the small pooled sample size. Specifically, only three studies with a combined  
323 sample of 53 participants were included in this analysis. The small sample sizes in individual  
324 studies for this outcome were probably due to the difficulties in collecting muscle biopsy  
325 samples in this age group. In a group of 87 older adults that were considered for a Bergstrom  
326 needle muscle biopsy, only 19% to 59% of participants had adequate levels of muscle mass  
327 needed for biopsy sampling (depending on factors such as sex, age, and frailty) [70].  
328 Furthermore, some participants had suboptimal muscle thickness, suggesting that multiple  
329 samples might be required to obtain an adequate amount of muscle for the analysis. While  
330 future studies are needed to elucidate possible effects of resistance training on muscle fiber  
331 hypertrophy in the very elderly, there may be challenges in collecting the necessary data.

332

#### 333 **4.5 Adverse events**

334 A recent systematic review reported that fear of a heart attack, stroke, or even death, is one of  
335 the most common barriers to participation in resistance exercise for older adults [71].

336 Therefore, when conducting exercise intervention studies among older adults, the reporting of  
337 adverse events associated with the training intervention is essential. The included studies  
338 reported minimal adverse events (Table 2). Specifically, in some studies, there were reports of  
339 muscle soreness following the exercise sessions, and in one study there was an exacerbation  
340 of preexisting osteoarthritis in one participant (Table 2). There were no reported serious  
341 events directly related to exercise interventions. These results suggest that resistance training  
342 can be safe, even for the very elderly.

343



#### 344 **4.6 Methodological quality**

345 All included studies were of good methodological quality. Therefore, the results presented  
346 herein were not confounded by studies with poor methodological quality. Nonetheless, it is  
347 worth noting that four included studies did not report participants' adherence to the training  
348 program [22, 33, 34, 58]. Adherence to a given training program is one of the key variables  
349 that influence its overall efficacy [72]. Therefore, future studies should ensure that adherence  
350 data are reported.

351

#### 352 **4.7 Strengths and limitations of the review**

353 The strengths of this review are that: (a) the search for studies was conducted through nine  
354 databases using a search syntax with a broad range of relevant search terms; and (b) 17  
355 studies with over 800 participants were included in the analysis for muscle strength, which  
356 allowed for an additional subgroup analysis including only the oldest-old. This review's main  
357 limitation is that the meta-analysis on muscle fiber hypertrophy included only three studies  
358 with a combined sample of 53 participants. Besides, there was high heterogeneity in the  
359 analysis for muscle strength. However, it should be considered here that the effects from all  
360 studies in this analysis were in the same direction (i.e., favoring of resistance training), but  
361 their overall effectiveness varied. The variation in ESs could be associated with the  
362 differences between studies in duration, training programs, and strength tests.

363

#### 364 **4.8 Suggestions for future research**

365 The included studies generally utilized only one type of strength test. Given that the studies  
366 used isotonic training programs, it might be expected that resistance training would have the  
367 greatest effect on isotonic strength [73, 74]. However, the majority of studies used isometric  
368 tests to evaluate changes in muscle strength. Ultimately, the small number of studies

369 employing isotonic and isokinetic strength assessments limits the ability to further subanalyze  
370 the effects of resistance training on strength in different tests. Isotonic and isokinetic strength  
371 tests were used only in four and three studies, respectively (Table 2). Therefore, future studies  
372 on the topic may consider utilizing isotonic, isometric, and isokinetic strength measures in the  
373 same group of participants to directly explore if the effects of resistance training in the very  
374 elderly vary between different strength tests.

375

## 376 **5 Conclusion**

377 This systematic review and meta-analysis found that the very elderly can increase their  
378 muscle strength and size by participating in resistance training programs. Moreover,  
379 resistance training was found to be an effective way to improve muscle strength even among  
380 the oldest-old. Importantly, the resistance training interventions generally included low  
381 weekly training volumes and frequencies, suggesting that a relatively low time commitment is  
382 needed to reap these benefits. There were minimal reports of adverse events associated with  
383 the training programs in the included studies, thus suggesting that resistance training can be a  
384 safe mode of exercise for the very elderly. More research is needed on the effects of resistance  
385 training on handgrip strength and muscle fiber hypertrophy.

386 **Data Availability Statement**

387 The datasets generated and analyzed during the current systematic review and meta-analysis  
388 are available from the corresponding author on reasonable request.

389 **Contributors**

390 JG conceived the idea for the review. JG and AG conducted the study selection quality  
391 assessment. JG and FS conducted the data extraction. JO performed the statistical analysis. JG  
392 drafted the initial manuscript. All authors contributed to data interpretation, writing of the  
393 manuscript, and its revisions.

394 **Funding**

395 No sources of funding were used to assist in the preparation of this article.

396 **Conflict of Interest**

397 Jozo Grgic, Alessandro Garofolini, John Orazem, Filip Sabol, Brad J. Schoenfeld and Zeljko  
398 Pedisic have no conflicts of interest that are directly relevant to the content of this article.

399 **References**

- 400 1. Clark BC, Manini TM. What is dynapenia? *Nutrition*. 2012;28:495–503.
- 401 2. Visser M, Deeg DJ, Lips P, et al. Skeletal muscle mass and muscle strength in relation  
402 to lower-extremity performance in older men and women. *J Am Geriatr Soc*.  
403 2000;48:381–6.
- 404 3. Newman AB, Kupelian V, Visser M, et al. Strength, but not muscle mass, is associated  
405 with mortality in the health, aging and body composition study cohort. *J Gerontol A*  
406 *Biol Sci Med Sci*. 2006;61:72–7.
- 407 4. Clark BC, Manini TM. Functional consequences of sarcopenia and dynapenia in the  
408 elderly. *Curr Opin Clin Nutr Metab Care*. 2010;13:271–6.
- 409 5. Cruz-Jentoft AJ, Bahat G, Bauer J, et al. Sarcopenia: revised European consensus on  
410 definition and diagnosis. *Age Ageing*. 2019;48:16–31.
- 411 6. Faigenbaum AD, Kraemer WJ, Blimkie CJ, et al. Youth resistance training: updated  
412 position statement paper from the national strength and conditioning association. *J*  
413 *Strength Cond Res*. 2009;23:S60–79.
- 414 7. Liu CJ, Latham NK. Progressive resistance strength training for improving physical  
415 function in older adults. *Cochrane Database Syst Rev*. 2009;CD002759.
- 416 8. American College of Sports Medicine. American College of Sports Medicine position  
417 stand. Progression models in resistance training for healthy adults. *Med Sci Sports*  
418 *Exerc*. 2009;41:687–708.
- 419 9. Fragala MS, Cadore EL, Dorgo S, et al. Resistance training for older adults: position  
420 statement from the National Strength and Conditioning Association. *J Strength Cond*  
421 *Res*. 2019;33:2019–52.

- 422 10. Ouchi Y, Rakugi H, Arai H, et al. Redefining the elderly as aged 75 years and older:  
423 proposal from the Joint Committee of Japan Gerontological Society and the Japan  
424 Geriatrics Society. *Geriatr Gerontol Int.* 2017;17:1045–7.
- 425 11. Stewart VH, Saunders DH, Greig CA. Responsiveness of muscle size and strength to  
426 physical training in very elderly people: a systematic review. *Scand J Med Sci Sports.*  
427 2014;24:e1–10.
- 428 12. Phillips SM. A brief review of critical processes in exercise-induced muscular  
429 hypertrophy. *Sports Med.* 2014;44:71–7.
- 430 13. Burd NA, Gorissen SH, van Loon LJ. Anabolic resistance of muscle protein synthesis  
431 with aging. *Exerc Sport Sci Rev.* 2013;41:169–73.
- 432 14. Petrella JK, Kim JS, Mayhew DL, et al. Potent myofiber hypertrophy during resistance  
433 training in humans is associated with satellite cell-mediated myonuclear addition: a  
434 cluster analysis. *J Appl Physiol.* 2008;104:1736–42.
- 435 15. Petrella JK, Kim JS, Cross JM, et al. Efficacy of myonuclear addition may explain  
436 differential myofiber growth among resistance-trained young and older men and  
437 women. *Am J Physiol Endocrinol Metab.* 2006;291:E937–46.
- 438 16. Karlsen A, Bechshøft RL, Malmgaard-Clausen NM, et al. Lack of muscle fibre  
439 hypertrophy, myonuclear addition, and satellite cell pool expansion with resistance  
440 training in 83-94-year-old men and women. *Acta Physiol (Oxf).* 2019;227:e13271.
- 441 17. Lundberg TR, Gustafsson T. Fibre hypertrophy, satellite cell and myonuclear  
442 adaptations to resistance training: have very old individuals reached the ceiling for  
443 muscle fibre plasticity? *Acta Physiol.* 2019;227:e13287.
- 444 18. Doherty TJ, Vandervoort AA, Brown WF. Effects of ageing on the motor unit: a brief  
445 review. *Can J Appl Physiol.* 1993;18:331–58.

- 446 19. Doherty TJ, Vandervoort AA, Taylor AW, et al. Effects of motor unit losses on  
447 strength in older men and women. *J Appl Physiol.* 1993;74:868–74.
- 448 20. Fiatarone MA, Marks EC, Ryan ND, et al. High-intensity strength training in  
449 nonagenarians. Effects on skeletal muscle. *JAMA.* 1990;263:3029–34.
- 450 21. Bruunsgaard H, Bjerregaard E, Schroll M, et al. Muscle strength after resistance  
451 training is inversely correlated with baseline levels of soluble tumor necrosis factor  
452 receptors in the oldest old. *J Am Geriatr Soc.* 2004;52:237–41.
- 453 22. Caserotti P, Aagaard P, Larsen JB, et al. Explosive heavy-resistance training in old  
454 and very old adults: changes in rapid muscle force, strength and power. *Scand J Med  
455 Sci Sports.* 2008;18:773–82.
- 456 23. Giné-Garriga M, Guerra M, Pagès E, et al. The effect of functional circuit training on  
457 physical frailty in frail older adults: a randomized controlled trial. *J Aging Phys Act.*  
458 2010;18:401–24.
- 459 24. Hruda KV, Hicks AL, McCartney N. Training for muscle power in older adults:  
460 effects on functional abilities. *Can J Appl Physiol.* 2003;28:178–89.
- 461 25. Judge JO, Whipple RH, Wolfson LI. Effects of resistive and balance exercises on  
462 isokinetic strength in older persons. *J Am Geriatr Soc.* 1994;42:937–46.
- 463 26. Kalapotharakos VI, Diamantopoulos K, Tokmakidis SP. Effects of resistance training  
464 and detraining on muscle strength and functional performance of older adults aged 80  
465 to 88 years. *Aging Clin Exp Res.* 2010;22:134–40.
- 466 27. Kim HK, Suzuki T, Saito K, et al. Effects of exercise and amino acid supplementation  
467 on body composition and physical function in community-dwelling elderly Japanese  
468 sarcopenic women: a randomized controlled trial. *J Am Geriatr Soc.* 2012;60:16–23.

- 469 28. Serra-Rexach JA, Bustamante-Ara N, Hierro Villarán M, et al. Short-term, light- to  
470 moderate-intensity exercise training improves leg muscle strength in the oldest old: a  
471 randomized controlled trial. *J Am Geriatr Soc.* 2011;59:594–602.
- 472 29. Skelton DA, Young A, Greig CA, et al. Effects of resistance training on strength,  
473 power, and selected functional abilities of women aged 75 and older. *J Am Geriatr*  
474 *Soc.* 1995;43:1081–7.
- 475 30. Bechshøft RL, Malmgaard-Clausen NM, Gliese B, et al. Improved skeletal muscle  
476 mass and strength after heavy strength training in very old individuals. *Exp Gerontol.*  
477 2017;92:96–105.
- 478 31. Benavent-Caballer V, Rosado-Calatayud P, Segura-Ortí E, et al. Effects of three  
479 different low-intensity exercise interventions on physical performance, muscle CSA  
480 and activities of daily living: a randomized controlled trial. *Exp Gerontol.*  
481 2014;58:159–65.
- 482 32. Cadore EL, Casas-Herrero A, Zambom-Ferraresi F, et al. Multicomponent exercises  
483 including muscle power training enhance muscle mass, power output, and functional  
484 outcomes in institutionalized frail nonagenarians. *Age (Dordr).* 2014;36:773–85.
- 485 33. Kim H, Suzuki T, Kim M, et al. Effects of exercise and milk fat globule membrane  
486 (MFGM) supplementation on body composition, physical function, and hematological  
487 parameters in community-dwelling frail Japanese women: a randomized double blind,  
488 placebo-controlled, follow-up trial. *PLoS One.* 2015;10:e0116256.
- 489 34. Sahin UK, Kirdi N, Bozoglu E, et al. Effect of low-intensity versus high-intensity  
490 resistance training on the functioning of the institutionalized frail elderly. *Int J Rehabil*  
491 *Res.* 2018;41:211–7.
- 492 35. Christensen K, Doblhammer G, Rau R, et al. Ageing populations: the challenges  
493 ahead. *Lancet.* 2009;374:1196–208.

- 494 36. National Center for Health Statistics. Survey Description, National Health Interview  
495 Survey, 2015. Hyattsville, Maryland: National Center for Health Statistics, 2016.
- 496 37. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews  
497 and meta-analyses: the PRISMA statement. *Ann Intern Med.* 2009;151:264–9.
- 498 38. Fielding RA. The role of progressive resistance training and nutrition in the  
499 preservation of lean body mass in the elderly. *J Am Coll Nutr.* 1995;14:587–94.
- 500 39. Guizelini PC, de Aguiar RA, Denadai BS, et al. Effect of resistance training on muscle  
501 strength and rate of force development in healthy older adults: A systematic review  
502 and meta-analysis. *Exp Gerontol.* 2018;102:51–8.
- 503 40. Liberman K, Forti LN, Beyer I, et al. The effects of exercise on muscle strength, body  
504 composition, physical functioning and the inflammatory profile of older adults: a  
505 systematic review. *Curr Opin Clin Nutr Metab Care.* 2017;20:30–53.
- 506 41. Porter MM. High-intensity strength training for the older adult—a review. *Top Geriatr*  
507 *Rehabil.* 1995;10:61–74.
- 508 42. Steib S, Schoene D, Pfeifer K. Dose-response relationship of resistance training in  
509 older adults: a meta-analysis. *Med Sci Sports Exerc.* 2010;42:902–14.
- 510 43. Straight CR, Lindheimer JB, Brady AO, et al. Effects of resistance training on lower-  
511 extremity muscle power in middle-aged and older adults: a systematic review and  
512 meta-analysis of randomized controlled trials. *Sports Med.* 2016;46:353–64.
- 513 44. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the  
514 methodological quality both of randomised and non-randomised studies of health care  
515 interventions. *J Epidemiol Community Health.* 1998;52:377–84.
- 516 45. Grgic J, Schoenfeld BJ, Davies TB, et al. Effect of resistance training frequency on  
517 gains in muscular strength: a systematic review and meta-analysis. *Sports Med.*  
518 2018;48:1207–20.



- 519 46. Grgic J, Schoenfeld BJ, Skrepnik M, et al. Effects of rest interval duration in  
520 resistance training on measures of muscular strength: a systematic review. *Sports Med.*  
521 2018;48:137–51.
- 522 47. Morris B. Estimating effect sizes from pretest-posttest-control group designs. *Organ*  
523 *Res Methods.* 2008;11:364–86.
- 524 48. Tanner-Smith EE, Tipton E, Polanin JR. Handling complex meta-analytic data  
525 structures using robust variance estimates: a tutorial in R. *J Dev Life Course*  
526 *Criminology.* 2016;2:85–112.
- 527 49. Sasaki H, Kasagi F, Yamada M, et al. Grip strength predicts cause-specific mortality  
528 in middle-aged and elderly persons. *Am J Med.* 2007;120:337–42.
- 529 50. Fisher Z, Tipton E, Zhipeng H. robumeta: robust variance meta-regression. R package  
530 version 2.0. 2017. <https://CRAN.R-project.org/package=robumeta>.
- 531 51. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat*  
532 *Softw.* 2010;36:1–48.
- 533 52. Fiatarone MA, O'Neill EF, Ryan ND, et al. Exercise training and nutritional  
534 supplementation for physical frailty in very elderly people. *N Engl J Med.*  
535 1994;330:1769–75.
- 536 53. Fiatarone Singh MA, Ding W, Manfredi TJ, et al. Insulin-like growth factor I in  
537 skeletal muscle after weight-lifting exercise in frail elders. *Am J Physiol.*  
538 1999;277:E135–43.
- 539 54. Sipilä S, Suominen H. Effects of strength and endurance training on thigh and leg  
540 muscle mass and composition in elderly women. *J Appl Physiol.* 1995;78:334–40.
- 541 55. Sipilä S, Multanen J, Kallinen M, et al. Effects of strength and endurance training on  
542 isometric muscle strength and walking speed in elderly women. *Acta Physiol Scand.*  
543 1996;156:457–64.

- 544 56. Sipilä S, Elorinne M, Alen M, et al. Effects of strength and endurance training on  
545 muscle fibre characteristics in elderly women. *Clin Physiol*. 1997;17:459–74.
- 546 57. Hvid LG, Strotmeyer ES, Skjødt M, et al. Voluntary muscle activation improves with  
547 power training and is associated with changes in gait speed in mobility-limited older  
548 adults - a randomized controlled trial. *Exp Gerontol*. 2016;80:51–6.
- 549 58. Kim H, Suzuki T, Saito K, et al. Effects of exercise and tea catechins on muscle mass,  
550 strength and walking ability in community-dwelling elderly Japanese sarcopenic  
551 women: a randomized controlled trial. *Geriatr Gerontol Int*. 2013;13:458–65.
- 552 59. Xue QL, Beamer BA, Chaves PH, et al. Heterogeneity in rate of decline in grip, hip,  
553 and knee strength and the risk of all-cause mortality: the women's health and aging  
554 study II. *J Am Geriatr Soc*. 2010;58:2076–84.
- 555 60. Mitchell WK, Williams J, Atherton P, et al. Sarcopenia, dynapenia, and the impact of  
556 advancing age on human skeletal muscle size and strength; a quantitative review.  
557 *Front Physiol*. 2012;3:260.
- 558 61. Goodpaster BH, Park SW, Harris TB, et al. The loss of skeletal muscle strength, mass,  
559 and quality in older adults: the health, aging and body composition study. *J Gerontol A*  
560 *Biol Sci Med Sci*. 2006;61:1059–64.
- 561 62. Delmonico MJ, Harris TB, Visser M, et al. Longitudinal study of muscle strength,  
562 quality, and adipose tissue infiltration. *Am J Clin Nutr*. 2009;90:1579–85.
- 563 63. Moreland JD, Richardson JA, Goldsmith CH, et al. Muscle weakness and falls in older  
564 adults: a systematic review and meta-analysis. *J Am Geriatr Soc*. 2004;52:1121–9.
- 565 64. Cronin J, Lawton T, Harris N, et al. A brief review of handgrip strength and sport  
566 performance. *J Strength Cond Res*. 2017;31:3187–217.

- 567 65. Saric J, Lisica D, Orlic I, et al. Resistance training frequencies of 3 and 6 times per  
568 week produce similar muscular adaptations in resistance-trained men. *J Strength Cond*  
569 *Res.* 2019;33:S122–9.
- 570 66. Sale D, MacDougall D. Specificity in strength training: a review for the coach and  
571 athlete. *Can J Appl Sport Sci.* 1981;6:87–92.
- 572 67. Tieland M, Verdijk LB, de Groot LC, et al. Handgrip strength does not represent an  
573 appropriate measure to evaluate changes in muscle strength during an exercise  
574 intervention program in frail older people. *Int J Sport Nutr Exerc Metab.* 2015;25:27–  
575 36.
- 576 68. Buckner SL, Dankel SJ, Bell ZW, Abe T, Loenneke JP. The association of handgrip  
577 strength and mortality: what does it tell us and what can we do with it? *Rejuvenation*  
578 *Res.* 2019;22:230–4.
- 579 69. Baumgartner RN, Koehler KM, Gallagher D, et al. Epidemiology of sarcopenia among  
580 the elderly in New Mexico. *Am J Epidemiol.* 1998;147:755–63.
- 581 70. Wilson W, Breen L, Lord JM, et al. The challenges of muscle biopsy in a community  
582 based geriatric population. *BMC Res Notes.* 2018;11:830.
- 583 71. Burton E, Farrier K, Lewin G, et al. Motivators and barriers for older people  
584 participating in resistance training: a systematic review. *J Aging Phys Act.*  
585 2017;25:311–24.
- 586 72. Gentil P, Bottaro M. Effects of training attendance on muscle strength of young men  
587 after 11 weeks of resistance training. *Asian J Sports Med.* 2013;4:101–6.
- 588 73. Buckner SL, Jessee MB, Mattocks KT, et al. Determining strength: a case for multiple  
589 methods of measurement. *Sports Med.* 2017;47:193–5.

- 590 74. Schoenfeld BJ, Grgic J, Ogborn D, et al. Strength and hypertrophy adaptations  
591 between low- vs. high-load resistance training: a systematic review and meta-analysis.  
592 J Strength Cond Res. 2017;31:3508–23.