

Universitat de Lleida

Document downloaded from:

<http://hdl.handle.net/10459.1/71160>

The final publication is available at:

<https://doi.org/10.1007/s10964-020-01289-9>

Copyright

(c) Springer Verlag, 2020

Journal of Youth and Adolescence

Effects of School-Based Multicomponent Positive Psychology Interventions on Well-Being and Distress in Adolescents: A Systematic Review and Meta-Analysis --Manuscript Draft--

Manuscript Number:	JOYO-D-20-00326R3
Full Title:	Effects of School-Based Multicomponent Positive Psychology Interventions on Well-Being and Distress in Adolescents: A Systematic Review and Meta-Analysis
Article Type:	Empirical Research
Keywords:	Positive psychology, Well-being, Intervention, Multicomponent, School, Meta-analysis
Corresponding Author:	Claudia Tejada-Gallardo Universitat de Lleida Lleida, Spain, SPAIN
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Universitat de Lleida
Corresponding Author's Secondary Institution:	
First Author:	Claudia Tejada-Gallardo
First Author Secondary Information:	
Order of Authors:	Claudia Tejada-Gallardo Ana Blasco-Belled Cristina Torrelles-Nadal Carles Alsinet
Order of Authors Secondary Information:	
Funding Information:	

Effects of School-Based Multicomponent Positive Psychology Interventions on Well-Being and Distress in Adolescents: A Systematic Review and Meta-Analysis

Claudia Tejada-Gallardo*

Universitat de Lleida

Avinguda de l'estudi general, nº 4, 25001, Lleida

+34 973 70 66 16

claudia.tejada@udl.cat

ORCID: 0000-0002-2867-7561

Ana Blasco-Belled

Universitat de Lleida

Avinguda de l'estudi general, nº 4, 25001, Lleida

+34 973 70 66 16

ana.blasco@udl.cat

ORCID: 000-0003-0641-0169

Cristina Torrelles-Nadal

Universitat de Lleida

Avinguda de l'estudi general, nº 4, 25001, Lleida

+34 973 70 66 16

cristina.torrelles@udl.cat

ORCID: 0000-0002-8669-110X

Carles Alsinet

Universitat de Lleida

Avinguda de l'estudi general, nº 4, 25001, Lleida

+34 973 70 66 16

carles.alsinet@udl.cat

ORCID: 0000-0001-9353-3591

*Corresponding author

Claudia Tejada-Gallardo is a PhD student at the University of Lleida, Spain. She has a master of science in psychology from the University of Maastricht, Holland. She is also a professor of social psychology at the University of Lleida, Spain. Her major research interests are adolescent's well-being, youth at risk, prevention and intervention approaches.

Ana Blasco-Belled is a post-doctoral researcher of the University of Lleida, Spain. She obtained her PhD in social psychology with international mention from the University of Lleida, Spain. Her research interests concern the study of well-being of adults and adolescents and the social dynamics of happiness.

Cristina Torrelles-Nadal is Lecturer in the Department of Psychology, University of Lleida, Spain. She received her PhD in Psychopedagogy, with international mention from the University of Lleida, Spain. Her research interests include well-being of adults and adolescents, gratitude, teamwork, happiness, optimism, emotional intelligence and prevention and intervention approaches.

Carles Alsinet is a social psychology professor and director of the Social Innovation Chair (SIC) at the University of Lleida, Spain. He received his PhD in psychology from the University of Lleida, Spain. The topics of his research are well-being, quality of life and social health of adults and adolescents. As a social consultant, he's currently developing intervention programs based on self-centered approaches to increase people's well-being.

**Effects of School-Based Multicomponent Positive Psychology Interventions on Well-Being and
Distress in Adolescents: A Systematic Review and Meta-analysis**

Abstract

Multicomponent positive psychology interventions are increasing in the general population but the study of its effectiveness in adolescents is still scarce, especially in the school context. Previous meta-analyses have reported that multicomponent positive psychology interventions increase well-being and reduce distress outcomes. However, the results on these outcomes limit their samples to adult populations. The aim of the current systematic review and meta-analysis is to evaluate and compare the immediate but also long-lasting effects of school-based multicomponent positive psychology interventions aimed at increasing well-being indicators of mental health (i.e., subjective and psychological well-being) and reducing the most common psychological distress indicators (i.e., depression, anxiety, and stress) in adolescents. A total of 9 randomized and non-randomized controlled trials from the searched literature met inclusion criteria for the meta-analysis. The results showed small effects for subjective well-being ($g = 0.24$), psychological well-being ($g = 0.25$), and depression symptoms ($g = 0.28$). Removing low-quality studies led to a slight decrease in the effect sizes for subjective well-being and a considerable increase for psychological well-being and depression symptoms. The relevant moderation analyses had an effect on subjective well-being and depression symptoms. The present systematic review and meta-analysis found evidence for the efficacy of school-based multicomponent positive psychology interventions in improving mental health in the short and long-term. Small effects for subjective well-being, psychological well-being, and depression symptoms were identified. Effects for psychological well-being and depression symptoms remained significant over time. In light of our results, education policy-makers and practitioners are encouraged to include positive practices within the schools' curriculum as effective and easily implemented tools that help to enhance adolescents' mental health. Further research is needed in order to strengthen the findings about school-based multicomponent positive psychology interventions in adolescents.

Keywords: Positive psychology, Well-being, Intervention, Multicomponent, School, Meta-analysis

Introduction

1 Psychological research and practice have traditionally focused on the treatment of mental illness, ignoring
2 aspects of well-being and positive functioning promotion (Chakhssi, Kraiss, Sommers-Spijkerman, &
3 Bohlmeijer, 2018). Adolescence is a crucial transitional stage in biological and psychological
4 development (Burger & Samuel, 2017), wherein optimal mental health is conceived as the combination of
5 high levels of well-being and few symptoms of mental illness (Keyes, 2009; World Health Organization,
6 2004). In contrast to the growing number of evidence-based interventions, which mainly focus on the
7 treatment of mental disorders and the reduction of psychological distress, few studies have examined the
8 contribution that interventions aiming at increasing well-being have in adolescents (Suldo, Savage, &
9 Mercer, 2014). Given the need for prevention and intervention during the period of adolescence, school-
10 based programs are an increasing and effective method to enhance positive functioning (Freire, Lima,
11 Teixeira, Araújo, & Machado, 2018). In this effort, positive psychology interventions originated as
12 scientifically-based interventions that focus on strengthening positive emotions, thoughts, and behaviors
13 through activities that can be easily implemented in daily routines (Schotanus-Dijkstra, Drossaert,
14 Pieterse, Walburg, & Bohlmeijer, 2015), and have proven their usefulness in promoting well-being and
15 reducing distress in the school setting (Water, 2011). Thus, the present study seeks to examine the effects
16 of school-based multicomponent positive psychology interventions on well-being indicators of mental
17 health (i.e., subjective and psychological well-being) and psychological distress indicators (i.e.,
18 depression, anxiety, and stress) in adolescents through a systematic review of the existing literature and a
19 meta-analysis.

School-Based Positive Psychology Interventions in Adolescents

20 Adolescents spend a significant part of their lifetime in the school and therefore this environment plays an
21 important role in their social, emotional, and psychological development (Waters, 2011). Psychological
22 distress can appear during adolescence, being the onset of mental health problems such as anxiety and
23 depression symptoms (Paus, Keshavan, & Giedd, 2008), which are the later predictors of stress' levels
24 (Shapero, Hankin, & Barrocas, 2013). Taking into account these considerations, schools could serve as a
25 vehicle for the promotion of positive youth development and effective learning (Clonan, Chafouleas,
26 McDougal, & Riley-Tillman, 2004). Positive education pursues the promotion of positive emotions,
27 positive relationships, and character strengths and, simultaneously, it attempts to foster skills for
28 happiness and well-being in the school context (Bernard & Walton, 2011). Positive education comprises
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 educational institutions that teach both the skills of well-being and the skills of achievement, mainly
2 throughout the application of school-based positive psychology interventions (Seligman et al., 2009). To
3 date, these interventions have been implemented in isolated classrooms and not as part of the school
4 curriculum. However, to reach a sustained effect, adolescents have to be exposed to positive psychology
5 interventions more constantly and through different high-school years (Waters, 2011). Given the need for
6 a positive youth development in schools, it is imperative to foster the introduction of these interventions
7 under the approach of positive education practices (Noble & McGrath, 2008). This approach argues that
8 there are five key foundations of well-being: (i.e., social and emotional competency, positive emotions,
9 positive relationships, engagement, and sense of meaning and purpose), which are the enabling factors
10 that should be built into positive psychology interventions.
11
12
13
14
15
16
17
18
19

20 **Prior Research on Positive Psychology Interventions and Multicomponent Positive Psychology** 21 **Interventions**

22 In contrast to positive psychology interventions, which are single-component interventions that include
23 one or more individual exercises targeting one component of well-being (e.g., gratitude, hope or character
24 strengths) (Froh, Kashdan, Ozimkowski, & Miller, 2009; Marques, Lopez, & Pais-Ribeiro, 2011; Proctor
25 et al., 2011), multicomponent positive psychology interventions are based on a variety of individual
26 exercises targeting two or more theoretically relevant well-being components that resemble the key
27 elements of positive education and are conducted within an integral program (Seligman, 2011, 2018).
28 According to the Synergistic Change Model (Rusk et al., 2018), which poses that targeting multiple
29 domains of positive functioning decreases the risk of relapse and increases the probability of spill-over
30 effects and synergy between activities, multicomponent positive psychology interventions are more likely
31 to provide long-term effects over positive psychology interventions (Rusk et al., 2018).
32
33
34
35
36
37
38
39
40
41
42
43

44 To understand the scope of multicomponent positive psychology interventions it is necessary to
45 acknowledge the two main theoretical approaches that define well-being: the hedonic and eudaimonic
46 perspectives. The hedonic perspective describes well-being in terms of pleasure maximization and pain
47 avoidance (Ryan & Deci, 2001), and subjective well-being has been commonly referred to as the
48 psychological construct affiliated to this conceptualization (Diener, 1984). By contrast, the eudaimonic
49 perspective claims the realization of one's true inner potential and virtue as a pathway to experiencing a
50 meaningful and fulfilling life (Vittersø, 2016; Waterman, 1993). Under this premise, well-being is a
51 relatively sustained experience that requires personal effort to be achieved (Waterman, 1993) and
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 involves how one is functioning in response to life's demands (psychological well-being; Ryff, 1989,
2 1995) and to larger society (social well-being; Keyes, 1998). Multicomponent positive psychology
3 interventions are composed of at least two components that target both perspectives of well-being:
4 hedonia and eudaimonia. Nevertheless, studies examining the effectiveness of these interventions applied
5 in the school context in adolescents are limited (Roth, Suldo, & Ferron, 2017), possibly due to the
6 difficulties of introducing school-based well-being initiatives in the school curriculum.
7
8
9

10
11 In a prior literature review of school-based single component positive psychology interventions,
12 adolescents' well-being and academic performance was fostered while distress was mitigated through the
13 application of positive interventions (Waters, 2011). Despite the increasing popularity of multicomponent
14 positive psychology interventions in the general population and the emergence of positive education as a
15 mechanism to introduce positive interventions in the school context, it comes as surprise that the
16 application and examination of multicomponent positive psychology interventions' effectiveness has not
17 been previously reviewed in adolescents, but only in adults (Hendriks, Schotanus-Dijkstra, Hassankhan,
18 de Jong, and Bohlmeijer, 2019; Koydemir, Sökmez, and Schütz, 2020). As a result, the present study
19 focused on a review of the existing literature on the topic and the subsequent effects of school-based
20 multicomponent positive psychology interventions in adolescents.
21
22
23
24
25
26
27
28
29
30
31

32 **Current study**

33
34 In recent years, there has been an increased popularity of studies on positive psychology interventions.
35 While some meta-analyses have demonstrated the efficacy of multicomponent positive psychology
36 interventions in adult samples, no single meta-analysis has investigated the effects of these interventions
37 in adolescents in the school context. To address this gap, the purpose of the present meta-analysis was to
38 examine the immediate but also long-lasting effects of school-based multicomponent positive psychology
39 interventions aimed at increasing well-being and reducing psychological distress symptoms of
40 adolescents aged between 10 and 18 years old.
41
42
43
44
45
46
47

48 The first aim was to identify and quantify the efficacy of well-being indicators of mental health
49 (subjective and psychological well-being) as primary outcomes, and psychological distress indicators
50 (depression, anxiety, and stress) as secondary outcomes, since it is necessary to the continued success of
51 the interventions. To note, not all studies included in the current meta-analysis measured aspects of
52 psychological distress, therefore all selected studies included (a) well-being mental health outcomes, but
53 not necessarily (b) psychological distress outcomes. As a secondary aim, moderation analyses were also
54
55
56
57
58
59
60
61
62
63
64
65

1 applied to explore how intervention effects may vary across different groups (i.e., publication year, study
2 design, type of intervention, control group, number of sessions, duration of intervention, quality of the
3 studies, and follow-up measures). Such analyses were used to investigate sources of heterogeneity, which
4 cannot be drawn from the pooled treatment effect estimate. Taken together, the present study attempted to
5 evaluate the effects of school-based multicomponent positive psychology interventions on well-being and
6 distress symptomatology and the features that improve their efficacy in order to establish
7
8
9
10
11
12 recommendations for future interventions to implement in school settings.

13 **Methods**

14
15
16 The preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) for the
17 conducting of systematic reviews and meta-analyses (Moher et al., 2015) were followed in the planning,
18 implementation and reporting of the present meta-analysis. This study was registered in PROSPERO
19 (#CRD42019139586), an international prospective register for systematic reviews.
20
21
22

23 **Search Strategy**

24
25
26 A systematic literature search was performed in the PsycINFO, PubMed, and Scopus databases. Another
27 procedure to identify eligible studies was the reference list screening of previous reviews and meta-
28 analyses for additional potentially eligible studies (Ciocanel, Power, Eriksen, & Gillings, 2017; Dunning
29 et al., 2018; Taylor, Oberle, Durlak, & Weissberg, 2017; Waters, 2011; Weiss, Westerhof, & Bohlmeijer,
30 2016). The last run was conducted in July 2019 and it was done by the first and second authors. The
31 search terms used were: positive psychology, well-being, happy, happiness, gratitude, optimism,
32 kindness, strengths, compassion, forgiveness, intervention, prevention, program, promotion, adolescent,
33 teenager, and youth. The search strings were combined according to the databases (see Appendix Table
34
35
36
37
38
39
40
41
42
43 1).

44 **Selection of Studies**

45
46 The inclusion criteria for the selected studies were formulated in accordance with the PICOS approach
47 and the studies were included, based on the following criteria: (1) the focus of the intervention was
48 adolescents (10–18 years old) from the general population, (2) interventions should target at least two
49 components of well-being as part of a program, (3) measures of the intervention should include outcomes
50 of subjective or psychological well-being and could also include secondary outcomes of depression,
51 anxiety, and stress symptoms, (4) school programs were the focus of the interventions, which included all
52 activities delivered in the school setting during school hours, (5) studies were randomized controlled trials
53
54
55
56
57
58
59
60
61
62
63
64
65

1 and non-randomized controlled trials that used a control condition, and (6) studies were published in peer-
2 reviewed journals without time restrictions. The exclusion criteria were the following: (1) young people
3 from the clinical population, (2) interventions that only targeted one component of well-being, such as
4 gratitude intervention, hope therapy, strength-based approaches and mindfulness-based interventions, (3)
5 studies not providing sufficient data for the analysis of the effect sizes at post-treatment and with
6 comparison groups, (4) studies published in book chapters, theoretical or narrative review papers and
7 dissertations, and (5) articles published in languages other than English or Spanish.

14 **Data Extraction**

15 Two data extractors (CTG and ABB) independently assessed the eligibility of the studies following a
16 standardized procedure. After removal of duplicates, the retrieved articles from the search were screened.
17 First, the title and abstract were screened by two independent reviewers (CTG and ABB). Secondly, the
18 full text was assessed for all eligible studies. In cases where there was a need to discuss further the
19 disagreements over abstracts or the full text, a third researcher (CTN) also took part until a consensus was
20 reached in all the processes. Data extraction templates were used to extract all the data from the included
21 studies. The information extracted was comprised of the following aspects: the source of the study
22 (author, publication, and date), participants' details (demographic characteristics and sample size of the
23 experimental and control groups), study design (randomized and non-randomized controlled trials), type
24 of intervention (exercises, number of sessions, duration of the intervention, and delivery) and outcomes at
25 pre, post, and follow-up measures (primary and secondary outcome measures—subjective and
26 psychological well-being and depression, anxiety, and stress symptoms respectively). When data on study
27 methods or results were missing, the authors were contacted to request the necessary information to
28 proceed with the analysis.

44 **Quality Assessment**

45 All the selected studies were assessed for their methodological quality using the newest Cochrane
46 collaboration tool for assessing risk of bias (RoB 2) in randomized controlled trials (Sterne et al., 2019)
47 by the first (CTG) and second author (ABB) independently. The Cochrane collaboration tool was
48 assessed with respect to five domains: (1) bias arising from the randomization process, (2) bias due to
49 deviation from intended interventions, (3) bias due to missing outcome data, (4) bias in measurement of
50 the outcome, and (5) bias in selection of the reported result. The rating consisted of answering (yes,
51 probably yes, probably no, no, and no information) signaling questions that elicited relevant information
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

for the assessment of risk of bias. The responses to the questions guided the final judgment; the study was judged to be at “low risk of bias” for all domains, if all domains were considered to be at low risk, as having “some concerns” if at least one domain was considered to raise some concerns, or at “high risk of bias” if at least one domain was considered to be at high risk of bias (Sterne et al., 2019). When discrepancies between authors' ratings regarding quality scores occurred, the authors discussed these differences about the article until they agreed on a common score. The same method was used to assess the risk of bias for randomized and non-randomized controlled trials to standardize the tool for both designs as the majority of studies were randomized controlled trials (and only two were non-randomized controlled trials). An inter-rater reliability of 0.98 was found, which represents an almost perfect level of agreement between authors (McHugh, 2012).

20 **Statistical Analyses**

21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Data were analyzed with the program Comprehensive Meta-Analysis (CMA V.3; Borenstein, Hedges, Higgins, & Rothstein, 2013). For each study, means, standard deviations, and sample sizes were extracted to calculate the effect size. Hedges' g effect size was used to calculate the differences between groups (i.e., multicomponent positive psychology intervention condition and control condition). Standardized effect sizes were calculated from the average score at pre-intervention and the average score at post-intervention from both groups separately and divided by the pooled standard deviations. Hedge's g measure was used instead of Cohens' d to calculate the effect sizes because the former is a more accurate measure when sample sizes are small, and when sample sizes are bigger the results for both statistics are roughly equivalent (Cuijpers, 2016). In the present study, small and big sample sizes were present, therefore in the calculation of the effect sizes of the different outcomes the instruments that explicitly measured the outcome were used. For subjective well-being and psychological well-being, a pooled effect size was calculated for all the instruments that measure each outcome. In this sense, one effect size was provided for each outcome in the study. The same procedure was applied to the outcomes of depression and anxiety symptoms. Concerning the stress outcome, it was not possible to calculate its effect size due to the lack of studies measuring it. Follow-up effect sizes were calculated, between-group, if there was a minimum of three studies per outcome.

Due to the diversity of the population (i.e., adolescents from different schools and countries), considerable heterogeneity across studies was expected. For this reason, a random-effects model was used (Borenstein, Hedges, Higgins, & Rothstein, 2010), which assumes that effect sizes are sampled from a

1 population of universal effect sizes. Separated meta-analyses were performed for (1) subjective well-
2 being, (2) psychological well-being, (3) depression symptoms, and (4) anxiety symptoms with a
3 confidence interval of 95% and using a two-tailed test. General effect sizes are considered small between
4 0–0.32, moderate between 0.33–0.55, and large between 0.56–1.2 (Lipsey & Wilson, 1993). Forest plots
5 for between-group effect sizes at post-intervention were created for each outcome variable. Heterogeneity
6 of effect sizes was also examined using the Q test and I^2 statistics. The Q test indicates whether there is
7 heterogeneity in the meta-analysis or not—significant values indicate heterogeneity, suggesting that one
8 or more variables moderated the effect size. The I^2 statistics captures the percentage of the variance
9 between the included studies that can be attributed to heterogeneity—a value of 0% indicates
10 homogeneity, while values of 25%, 50%, and 75 % indicate low, moderate, and high levels of
11 heterogeneity, respectively (Higgins & Thompson, 2002).

12 Different subgroup analyses were examined to test the possible moderating effects of the
13 following variables: (1) year of publication: the last five years and older, (2) study design: randomized
14 and non-randomized controlled trials, (3) intervention: multicomponent positive psychology intervention
15 and multicomponent positive psychology intervention combined with another form of intervention, (4)
16 control group: placebo or waitlist, (5) sessions: six sessions or less and more than six sessions, (6)
17 duration: six weeks or less and more than six weeks, (7) quality of studies: high quality, some concerns
18 and low quality, and (8) presence of follow-up: measurement at follow-up and no measurements at
19 follow-up.

20 Publication bias was assessed through funnel plots, the Egger's test, Duval and Tweedie's trim-
21 and-fill procedure, and the fail-safe N. First, funnel plots were created by plotting the overall mean effect
22 size against study size. A symmetric distribution of studies around the effect size indicates the absence of
23 publication bias, whereas an accumulation of studies on one side or another indicates publication bias
24 (Sterne, Egger, & Moher, 2008). Second, Egger's test was used to examine the symmetric distribution of
25 studies around the effect size with a quantitative test. Funnel plot asymmetry was considered if $p < 0.05$
26 (Egger, Smith, Schneider, & Minder, 1997). Third, Duval and Tweedie's trim-and-fill procedure was
27 applied. This procedure imputes the effect sizes of missing studies and produces an adjusted effect size
28 accounting for these missing studies (Duval & Tweedie, 2000). Finally, the fail-safe N indicates the
29 number of unpublished non-significant studies that would be required to lower the overall effect size
30 below significance (Egger et al., 1997). Findings were considered robust if the fail-safe $N \geq 5n + 10$,

where n is the number of studies (Rosenberg, 2005).

Results

Study Selection

A total of 11,585 studies were found in the electronic databases: 3,275 from PubMed, 3,862 from PsycInfo, 4,440 from Scopus and 8 from different reference lists. After removal of duplicates, 9,735 articles remained for title and abstract screening. In the next phase, 9,661 articles were discarded because they did not meet the inclusion criteria and 74 articles remained to be full-text reviewed. Of these, a total of 9 articles were included in the meta-analysis. The PRISMA analysis represents the process of study selection (Fig. 1).

Figure_1

Description of Studies

The studies included a total of 4,898 participants, 2,537 in the intervention condition and 2,357 in the control condition. The age of participants ranged from 10 to 18 ($M = 13.27$, $SD = 1.85$; 54% female) years old. The 9 studies included in the meta-analysis were targeting multicomponent positive psychology interventions and 2 of them were combined with another technique (acceptance and commitment therapy and positive youth development). Well-being therapy was a technique used in 2 studies and it was identify as a multicomponent positive psychology intervention because it was based on Ryff's model of psychological well-being (Schotanus-Dijkstra et al., 2015). Seven studies were randomized controlled trials and 2 were non-randomized controlled trials. The delivery mode was group intervention for all the studies. Five control conditions were active control groups (placebo, $n = 4$; anxiety-management school protocol, $n = 1$) and 4 were non-active control groups (waitlist). The number of sessions varied between 6 and 18 ($M = 11.56$, $SD = 4.53$). The duration of the program ranged from 4 to 30 weeks ($M = 13.44$; $SD = 9.58$). Six studies reported follow-up effects and the average follow-up time was 6.25 months ($SD = 2.75$) at post-intervention. The main characteristics of the studies are presented in Table 1.

Table_1

Study Measures

1 The study's primary outcomes were both subjective well-being and psychological well-being. The
2
3 outcomes classified as subjective well-being were the following: satisfaction with life, positive affect, and
4
5 student's life satisfaction. The outcomes classified as psychological well-being were the following:
6
7 flourishing and psychological well-being domains (i.e., self-efficacy, autonomy, environmental well-
8
9 being, personal growth, positive relationships, purpose in life, and self-acceptance). All studies included
10
11 at least one measure of each outcome. Concerning secondary outcomes of the meta-analysis, the measures
12
13 included were depression, anxiety, and stress symptomatology. However, only one study reported stress
14
15 outcomes and thus analyses were not performed for this outcome. In total, six studies measured subjective
16
17 well-being, five studies measured psychological well-being, four studies measured depression, and four
18
19 studies measured anxiety symptoms. Five studies included two measures of subjective well-being
20
21 (Boniwell et al., 2015; Roth et al., 2017; Shoshani & Steinmetz, 2014; Shoshani et al., 2016; Suldo,
22
23 Savage, & Mercer, 2014b), which were pooled by the authors of the meta-analysis. One study included
24
25 more than two measures of psychological well-being that were pooled (Tomba et al., 2010), and two
26
27 studies included two measures of anxiety symptoms that were also pooled (Ruini et al., 2009; Tomba et
28
29 al., 2010). In these cases, the effect sizes were calculated using the average effect size of various
30
31 outcomes of each study (Malle, 2006; Turner, Herbert, & Bernard, 2006). See Appendix Table 2 for
32
33 detailed information on the used instruments per outcome.
34
35

Quality Assessment

36
37
38 When a minimum of one domain did not meet the criteria, the study was labeled as being at high risk of
39
40 bias. When at least one domain was judged to raise some concerns, the study was labeled as having some
41
42 concerns and when all domains were judged to be at low risk of bias, the study was labeled as being at
43
44 low risk. Three studies were rated as being at high risk of bias (i.e., low quality), four studies were rated
45
46 as having some concerns and two studies did meet all the criteria for being rated as low risk of bias (i.e.,
47
48 high quality). The randomization process domain was the most poorly rated due to the non-randomized
49
50 controlled trials included in the study that were also assessed with the Cochrane collaboration tool for
51
52 assessing risk of bias (RoB 2). The quality assessment is displayed in Table 2.
53
54
55
56

57 Table_2
58
59
60
61
62
63
64
65

Post-Intervention Effects of Multicomponent Positive Psychology Interventions

Post-intervention effects were calculated for the main outcomes (subjective well-being and psychological well-being) and for the secondary outcomes (depression and anxiety symptoms). The analysis of effect sizes was done for all the studies and also for all the outcomes excluding low quality studies (i.e., studies rated as being at high risk of bias). Follow-up effects were also calculated for the studies that included a follow-up analysis and for outcomes excluding low quality studies. Results for studies excluding outliers were not calculate since neither of the studies included in the meta-analysis was considered an outlier. The main results are presented in Table 3.

Table_3

Post-intervention effects on subjective well-being

For subjective well-being ($n = 6$) a significant small effect size was observed ($g = 0.24$, 95% CI 0.11–0.38, $p = 0.000$) at post-intervention. The effect sizes of the studies ranged from 0.08 to 0.69. Heterogeneity analysis revealed a moderate level of heterogeneity ($I^2 = 49.09$, $Q = 9.82$, $p < 0.05$). When low quality studies were excluded, the effect size remained small ($g = 0.21$, 95% CI 0.05–0.37, $p < 0.01$) and the heterogeneity increased to a higher level ($I^2 = 62.91$, $Q = 8.09$, $p < 0.04$). The forest plot in Figure 2 shows the post-intervention effect on subjective well-being.

Post-intervention effects on psychological well-being

For psychological well-being ($n = 5$) a significant small effect size was observed ($g = 0.25$, 95% CI 0.01–0.51, $p < 0.05$). The effect sizes of the studies ranged from 0.02 to 0.60. Heterogeneity analysis revealed a high level of heterogeneity between studies ($I^2 = 82.29$, $Q = 22.58$, $p = 0.000$). When low quality studies were excluded, the effect size remained small but was non-significant ($g = 0.31$, 95% CI 0.03–0.67, $p < 0.12$) and heterogeneity remained high ($I^2 = 87.45$, $Q = 15.93$, $p = 0.000$). The forest plot in Figure 3 shows the post-intervention effect on psychological well-being.

Post-intervention effects on depression symptoms

For depression symptoms ($n = 4$) a significant small effect size was observed ($g = 0.28$, 95% CI 0.13–0.43, $p = 0.000$). The effect sizes of the studies ranged from 0.03 to 0.44. Heterogeneity analysis revealed that the level of heterogeneity was non-significant. When low quality studies were excluded, the effect size increased to moderate ($g = 0.34$, 95% CI 0.24–0.44, $p = 0.000$) and the heterogeneity remained non-

significant. The forest plot in Figure 4 shows the post-intervention effect on depression symptoms.

Post-intervention effects on anxiety symptoms

For anxiety symptoms ($n = 4$) the effect size at post-intervention was non-significant. When low quality studies were excluded, the effect size remained non-significant. Heterogeneity analysis revealed that no heterogeneity was found and hence did not require exploration in a subgroup analysis. The forest plot in Figure 5 shows the post-intervention effect on anxiety symptoms.

Figure_2, 3, 4 and 5

Subgroup Analysis

A total of eight moderators were defined in the meta-analysis: year of publication (last five years vs. older), design of the study (randomized vs. non-randomized controlled trials), type of intervention (multicomponent positive psychology intervention vs. multicomponent positive psychology intervention combined with another positive intervention), control group (placebo vs. waitlist), sessions (six or less vs. more than six), duration (six weeks or less vs. more than six weeks), quality of the studies (high, some concerns, and low), and follow-up measures (presence of follow-up vs. no follow-up). These categorical moderators were used in a subgroup analysis to examine the likely contributions in the overall effect sizes at post-intervention.

Regarding subjective well-being, significant moderating analyses indicated that non-randomized controlled trials showed a significant larger effect size ($n = 2$; $g = 0.38$, 95% CI 0.12–0.64, $p < 0.003$) compared to randomized controlled trials ($n = 4$; $g = 0.21$, 95% CI 0.05–0.37, $p < 0.009$). Depending on the type of intervention, moderating effects were also observed: the only study that employed a multicomponent positive psychology intervention combined with another type of positive intervention showed a significant larger effect size ($n = 1$; $g = 0.45$, 95% CI 0.00–0.90, $p < 0.05$) compared to studies with a multicomponent positive psychology intervention exclusively ($n = 5$; $g = 0.23$, 95% CI 0.09–0.37, $p < 0.001$). The type of control group also had a significant moderating effect: studies where the control group was a placebo showed larger effects ($n = 3$; $g = 0.38$, 95% CI 0.12–0.64, $p < 0.003$) compared to control group studies with waitlist ($n = 2$; $g = 0.15$, 95% CI 0.05–0.37, $p < 0.009$). The quality of the study was also a significant moderator for subjective well-being: studies with some concerns regarding their quality had a larger effect size ($n = 2$; $g = 0.48$, 95% CI 0.05–0.93, $p < 0.02$) compared to low

quality studies ($n = 2$; $g = 0.38$, 95% CI 0.12–0.64, $p < 0.003$). Finally, the fact of including a follow-up in the study was also considered a moderator: studies with no follow-up showed a significant larger effect size ($n = 2$; $g = 0.38$, 95% CI 0.12–0.64, $p < 0.003$) compared to the studies that included follow-up ($n = 4$; $g = 0.21$, 95% CI 0.05–0.37, $p < 0.009$).

For depression symptoms significant moderating analyses were found. The study that was published in the last five years had a significant larger effect size ($n = 1$; $g = 0.44$, 95% CI 0.20–0.69, $p = 0.000$) compared to the older published studies ($n = 3$; $g = 0.23$, 95% CI 0.05–0.41, $p < 0.01$). Depending on the type of intervention, moderating effects were also observed: the study that employed a multicomponent positive psychology intervention combined with another type of positive intervention showed a significant larger effect size ($n = 1$; $g = 0.44$, 95% CI 0.20–0.69, $p = 0.000$) compared to studies with a multicomponent positive psychology intervention exclusively ($n = 3$; $g = 0.23$, 95% CI 0.05–0.41, $p < 0.001$). The quality of the study was also a significant moderator for subjective well-being. Studies with some concerns regarding quality had larger effects ($n = 2$; $g = 0.37$, 95% CI 0.18–0.56, $p = 0.000$) compared to high quality studies ($n = 1$; $g = 0.33$, 95% CI 0.21–0.45, $p = 0.000$). Finally, the fact of including a follow-up in the study was also considered a moderator: the study with no follow-up analysis showed a significant larger effect size ($n = 1$; $g = 0.44$, 95% CI 0.20–0.69, $p = 0.000$) compared to the studies that included follow-up ($n = 3$; $g = 0.23$, 95% CI 0.05–0.41, $p < 0.01$). For psychological well-being and anxiety symptoms, any significant results were found. Table 4 shows the results of the subgroup analysis.

Table_4

Follow-Up Effects

Studies reporting follow-up data of periods ranging from six to twelve months were analyzed. In the present meta-analysis, when multiple follow-ups were present in the same study, only the longest interval was used. Analysis of follow-ups did not show any effect size on subjective well-being and anxiety symptoms. The follow-up effect size for psychological well-being was moderate ($g = 0.44$, 95% -0.45–1.31, $p < 0.04$) and for depression symptoms was small ($g = 0.31$, 95% 0.20–0.41, $p = 0.000$). When low quality studies were excluded, the results for psychological well-being did not show any significant effect. By contrast, the effect size remained small for depression symptoms ($g = 0.29$, 95% 0.11–0.47, $p < 0.001$) and turned out to be significantly small for anxiety symptoms ($g = 0.21$, 95% 0.10–0.33, $p =$

0.000).

Publication Bias

The funnel plot is an approach used to assess publication bias. Symmetry in the funnel plot indicates that the publications are representative, whereas asymmetry indicates that publication bias is likely to be present in the meta-analysis. In the current study, the funnel plot for each analyzed outcome (subjective well-being, psychological well-being, depression, and anxiety) was somewhat asymmetrical. However, the funnel plot is a subjective method and therefore cannot be relied on exclusively in the case of this result. Thus, the fail-safe N test was also conducted, which revealed that 53 studies for subjective well-being, 41 studies for psychological well-being, 27 studies for depression symptoms, and 3 studies for anxiety symptoms were needed to nullify the significant effect at $p > 0.05$. The Egger's regression intercept test was also used and the results were only significant for psychological well-being (-4.76 , $t = 4.74$, $df = 3$, $p < 0.01$), reporting non-significant results for subjective well-being (0.84 , $t = 0.94$, $df = 4$, $p < 0.40$), depression symptoms (-1.18 , $t = 0.56$, $df = 2$, $p < 0.62$), and anxiety symptoms (-1.18 , $t = 1.07$, $df = 2$, $p < 0.39$). Finally, Duval and Tweedie's trim-and-fill method was also tested and possible missing studies were imputed to observe potential changes on the effect sizes. For psychological well-being ($g = 0.29$, $95\% \text{ CI} = 0.05\text{--}0.52$) and anxiety symptoms ($g = 0.16$, $95\% \text{ CI} = 0.07\text{--}0.26$) the effect sizes increased, for subjective well-being decreased ($g = 0.18$, $95\% \text{ CI} = 0.05\text{--}0.31$) and for depression symptoms remained the same ($g = 0.28$, $95\% \text{ CI} = 0.13\text{--}0.43$). Importantly, the studies included in the meta-analysis, and specifically on each outcome, are not many in order to consider the results potentially significant. Thus, it is likely that missing publications might have affected the results of the present meta-analysis.

Discussion

Despite the increasing popularity of multicomponent positive psychology interventions in the general population and the emergence of positive education as a mechanism to introduce positive interventions in the schools, the effectiveness of multicomponent positive psychology interventions has not been previously examined in adolescents. To address this gap, the present study investigated the effects of school-based multicomponent positive psychology interventions on well-being (subjective and psychological) and psychological distress symptoms (depression, anxiety and stress) in adolescents through a meta-analysis. Results indicated that school-based multicomponent positive psychology interventions enhanced subjective and psychological well-being and reduced depression symptoms but

1 not anxiety. Multicomponent positive psychology interventions can increase adolescent's mental health in
2 the short and long term, hence, education policy-makers and practitioners are encouraged to implement
3 these interventions within the schools' curriculum, especially in combination with other evidence-based
4 positive interventions, in efforts to provide adolescent with effective tools that ensure their mental health
5 in school.
6
7
8
9

10 **Efficacy of Multicomponent Positive Psychology Interventions on Mental Health in Adolescents**

11 The current meta-analysis showed that school-based multicomponent positive psychology interventions
12 in adolescents yielded to increases in subjective well-being ($g = 0.24$) and psychological well-being ($g =$
13 0.25) and reductions in symptoms of depression ($g = 0.28$), in accordance with research on adults
14 (Hendriks et al., 2019; Koydemir et al., 2020). Although the effects were small, this suggests that
15 multicomponent positive psychology interventions are an effective strategy to boost well-being in the
16 school context. A possible explanation for the small effects would be that adolescents' well-being tend to
17 decrease from early to late adolescence (González-Carrasco, Casas, Malo, Viñas, & Dinisman, 2017). A
18 remarkable finding about the efficacy of multicomponent positive psychology interventions is related to
19 the long-term effects found for psychological well-being and depression symptoms: on the one hand, the
20 effects on these outcomes increased from the short to the long term, which contrasts with previous meta-
21 analyses about positive interventions in adults (Bolier et al., 2013; Chakhssi et al., 2018). On the other
22 hand, the effects on subjective well-being were larger in the short-term (just after the intervention) but
23 decreased in the long run, similar to research on adults (Bolier et al., 2013; Hendriks et al., 2019).
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 According to the hedonic and eudaimonic traditions of well-being, multicomponent positive
39 psychology interventions may have an immediate impact during and right after the intervention is
40 conducted, raising the levels of emotional well-being. Using positive practices that target the key
41 elements of positive education (social and emotional competency, positive emotions, positive
42 relationships, engagement, and sense of meaning and purpose) can entail a pleasant experience for
43 adolescents. While this sense of subjective well-being dissipates over time, a deeper and more sustained
44 sense of meaning and fulfillment is reinforced at follow-up. The fact that adolescents tend to seek many
45 small and momentary pleasures but also have life objectives to accomplish (González-Carrasco et al.,
46 2019) may contribute to explain why multicomponent positive psychology interventions provide both
47 short-term and long-term gains.
48
49
50
51
52
53
54
55
56
57

58 The promotion of mental health (and also education) is captured by the 2030 Agenda for
59
60
61
62
63
64
65

1 Sustainable Development Goals (UN General Assembly, 2015), an international agreement to ensure 17
2 global challenges for sustainable development. Depression is considered as a common indicator of mental
3 illness (Keyes, 2002) characterized by anhedonia (low subjective well-being) and aspects of individual
4 and social malfunctioning (low psychological well-being) (American Psychiatric Association [APA],
5 2013), and it has become a paramount concern in mental health and societal progress given the high rates
6 of people affected in recent years (World Health Organization, 2017). More specifically, depression
7 symptoms affect negatively the daily functioning of adolescents (Derdikman-Eiron et al., 2011) and they
8 have been associated with poorer social relationships and academic outcomes, substance abuse, and
9 increased risk of suicide (Beesdo, Knappe, & Pine, 2009; Birmaher et al., 1996). In light of this,
10 researchers and practitioners are called to provide accessible tools to mitigate this issue that might thus be
11 of societal advantage, and the present study found empirical evidence to support the notion that
12 multicomponent positive psychology interventions are an effective long-term strategy to reduce
13 depressive symptomatology among adolescents.

26 **Which Characteristics Can Make Multicomponent Positive Psychology Interventions More** 27 **Beneficial?**

28 Subgroup analysis revealed that the characteristics of the intervention could lead to effective
29 contributions of the multicomponent positive psychology interventions on subjective well-being and
30 depression symptoms. The main features of these interventions included year of publication, study
31 design, type of intervention, control group, quality of studies, and measurement of follow-up. For
32 depression symptoms, the fact that studies published over the last five years were more effective suggests
33 that interventions might evolve and adapt over time to be more effective. The study design was
34 considered as another significant moderator. The non-randomization of participants for the intervention or
35 control group was observed as a more effective process than the randomization of participants for
36 subjective well-being outcomes. This can be explained by the fact that in non-randomized controlled
37 trials it is easier to control study characteristics (e.g., allocation sequence generation or allocation
38 sequence concealment [Sterne et al., 2019]), which may lead to more (pre)fabricated or beneficial results
39 than randomized controlled trials. Also, in many cases, peer journals preferred the publication of studies
40 with favorable (i.e., significant) results rather than unfavorable (i.e., non-significant) results—this is
41 known as the "file drawer problem" (Dalton, Aguinis, Dalton, Bosco, & Pierce, 2012).

42 The type of intervention was a significant moderator for subjective well-being and depression
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 symptoms. When multicomponent positive psychology interventions were combined with another type of
2 positive intervention, the results turned out to be more effective. This combination increases the
3 probability of merging complementary approaches (Burckhardt, Manicavasagar, Batterham, & Hadzi-
4 Pavlovic, 2016) and therefore providing and covering a wider range of intervention goals. The control
5 group used in the studies turned out to be a significant moderator for subjective well-being and depression
6 symptoms. In studies where the control group was a placebo, the effect was larger than studies where the
7 control group was in the waitlist. The fact that in psychosocial interventions the control group engaged in
8 other activities, despite being unrelated to the intervention, is considered more valid than relying on the
9 classical design of the waitlist control group (Popp & Schneider, 2015). In addition, the quality of studies
10 also had a significant moderating effect on subjective well-being and depression symptoms. In line with a
11 previous meta-analysis (Bolier et al., 2013), for subjective well-being studies with some concerns
12 regarding quality showed larger effects compared to studies with low and high quality. For depression
13 symptoms, it was found the same pattern; however, previous meta-analysis with adults (Chakhssi et al.,
14 2018; Hendriks et al., 2019) showed contradictory results, which might have been caused as a result of
15 the different tools used to assess risk of bias in the different meta-analytic reviews. Finally, significant
16 differences in the effects of the follow-up moderator for subjective well-being and depression symptoms
17 were found. Studies in which follow-up was not examined had larger effects than studies including
18 follow-up measures. As stated previously, this may be related to the assumption that studies with non-
19 significant results are less likely to be published (Sterne et al., 2019). When follow-up was tested, the
20 effectiveness of interventions might be lowered since the peak of effectiveness normally happens right
21 after the intervention. The intervention effects from previous meta-analyses in adults (Bolier et al., 2013;
22 Hendriks et al., 2019) on subjective well-being are likely to be reduced over time, thus the results
23 unsurprisingly suggest that positive interventions targeting subjective well-being may only increase
24 momentary well-being in adolescents.

48 **Limitations**

49 Although the current systematic review and meta-analysis addressed some gaps in the literature, the
50 findings must be understood within the confines of its limitations. First, caution is warranted when
51 interpreting the results of the study due to the small number of studies included in the analysis of each
52 outcome and subgroups. For this reason, future studies of school-based multicomponent positive
53 psychology interventions in adolescents are needed to draw firmer conclusions on the efficacy of these
54
55
56
57
58
59
60
61
62
63
64
65

1 interventions. Second, regarding the quality of the studies included in the meta-analysis, it creates a
2 sample with only two studies being rated as low risk of bias. This aspect highlights the importance of
3 considering the assessment of the quality of studies in meta-analytic reviews to be aware of possible
4 limitations of the findings. Third, one of the outcomes of the study (stress) could not be analyzed since
5 there was a lack of studies reporting the stress' effects. Fourth, the inclusion criteria (e.g., adolescent
6 population, intervention in the school setting, articles from peer-reviewed journals, or articles in English
7 or Spanish) were very restrictive and results should be interpreted in light of the chosen features. Lastly,
8 the study only included studies published in peer-reviewed journals and the possibility to omit
9 unpublished results or studies with null findings is present.

10
11
12
13
14
15
16
17
18 The benefits of school-based multicomponent positive psychology interventions in adolescents
19 are evident; however, the studies presented few limitations to discuss. Schools programs are implemented
20 with time restrictions because they are not commonly integrated in the school curriculum but rather
21 handled and implemented by external professionals. Future programs should be implemented by school
22 teachers or school psychologists previously trained, which might enable to introduce the program in
23 schools and through different years. Furthermore, the combination of a multicomponent positive
24 psychology intervention with another evidence-based positive intervention resulted more effective than
25 the single application of a multicomponent positive psychology intervention. Among the reviewed
26 studies, only 2 used the combination of both interventions. Additionally, the number of participants in
27 some studies was weakly powered (i.e., less than 50 participants) and therefore lacked predictive value,
28 which might be translated into exaggerated effects (Button et al., 2013). It is important to highlight that an
29 optimal sample size is required to yield representative results. Finally, all studies included some risk of
30 bias, suggesting that a rigorous methodology and reporting should be followed to increase the quality of
31 studies. The main issue is the lack of information provided by the studies – for instance, many studies
32 lacked information regarding the allocation sequence of participants, session attendance, and the blinding
33 of the assessor to intervention status. This issue should be addressed in the context of journal publication
34 and advocates for more high-quality studies that provide all information needed to allow a doable
35 assessment of research quality.

36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 **Implications**

55 Overall, school-based multicomponent positive psychology interventions aiming at increasing well-being
56 and reducing depression symptoms in adolescents were effective. Ensuring mental health and education
57
58
59
60
61
62
63
64
65

1 are paramount to the Sustainable Development Goals, since mental illness can indeed be a risk factor of
2 education attainment. Providing adolescents with low-cost, easily-delivered tools that promote their
3 mental health and optimal development are key to ensure health and education goals. This meta-analysis
4 showed that multicomponent positive psychology interventions can offer plausible resources to meet
5 these goals: in the short-run, these interventions can help to maximize pleasant experiences in the school
6 context that may also benefit adolescents' academic performance. In the long-run, they can strengthen the
7 psychological functioning of adolescents and provide them with a sense of meaning and fulfillment. The
8 present findings encourage to introduce multicomponent positive psychology interventions in the school
9 curriculum for building up psychological resources and mitigating the most common societal burden
10 identified worldwide – depression symptomatology.
11
12
13
14
15
16
17
18
19

20 Based on moderator analysis, we would recommend professionals to implement multicomponent
21 positive psychology interventions in conjunction with other evidence-based positive interventions that
22 enhance well-being, and researchers to preferentially include a placebo control group. Education policy-
23 makers and practitioners are potentially called to exert a relevant role in the application and guidance of
24 positive practices that, in the end, might facilitate the learning process and academic performance of
25 adolescents. As the continued willingness from professionals working with adolescents for the
26 incorporation of the positive education premises in the schools to develop both positive youth functioning
27 and effective learning, education policies should consider the introduction of positive practices like
28 multicomponent positive psychology interventions in the school's curriculum to enhance the well-being
29 of adolescents and reduce their psychological distress. In line with the above-mentioned considerations,
30 teaching both the skills of well-being and the skills of achievement will be associated with greater
31 academic achievement and engagement of adolescents in the school context (Datu, 2018). Still, it is
32 necessary to establish rigorous methodological approaches to lead to higher-quality studies in the field of
33 positive psychology interventions, such as using randomized controlled trials and follow-up
34 measurements to increase the quality of studies and reduce publication bias. The present study presents
35 evidence on the efficacy of multicomponent positive psychology interventions, proposes following
36 rigorous methodological approaches, and exposes new gaps in the conceptualization of these
37 interventions in the school context.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56

57 **Conclusion**

58
59
60
61
62
63
64
65

1 The effectiveness of multicomponent positive psychology interventions in adolescents' school-based
2 settings was, to date, not systematically evaluated. This meta-analysis addressed this gap by synthesizing
3 the efficacy of these interventions on subjective and psychological well-being, and symptoms of
4 depression, anxiety, and stress. Results demonstrated that multicomponent positive psychology
5 interventions can be effective in improving subjective and psychological well-being and reducing
6 depression symptoms in adolescents. No effects were found for symptoms of anxiety, while effects on
7 stress could not be analyzed due to lack of studies. The effects on psychological well-being and
8 depression symptoms remained significant in the long-term, posing that multicomponent positive
9 psychology interventions offer an opportunity to ensure mental health during adolescents' development in
10 schools. Academic policies and education practitioners should consider the inclusion of these
11 interventions within the school curriculum to promote adolescents' mental health and optimal
12 development.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- 1 American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders*.
 2 Washington, D.C.: American Psychiatric Association.
 3
 4
 5
 6 Beesdo, K., Knappe, S., & Pine, D. S. (2009). Anxiety and anxiety disorders in children and adolescents:
 7 Developmental issues and implications for DSM-V. *Psychiatric Clinics of North America*, 32(3),
 8 483–524. <https://doi.org/10.1016/j.psc.2009.06.002>
 9
 10
 11
 12 Bernard, M. E., & Walton, K. (2011). The effect of you can do it! Education in six schools on student
 13 perceptions of well-being, teaching-learning and relationships. *The Journal of Student Wellbeing*,
 14 5(1), 22. <https://doi.org/10.21913/jsw.v5i1.679>
 15
 16
 17
 18 Birmaher, B., Ryan, N. D., Williamson, D. E., Brent, D. A., Kaufman, J., Dahl, R. E., ... Nelson, B.
 19 (1996). Childhood and adolescent depression: A review of the past 10 years. Part I. *Journal of the*
 20 *American Academy of Child and Adolescent Psychiatry*, 35(11), 1427–1439.
 21 <https://doi.org/10.1097/00004583-199611000-00011>
 22
 23
 24
 25
 26 Bolier, L., Haverman, M., Westerhof, G. J., Riper, H., Smit, F., & Bohlmeijer, E. (2013). Positive
 27 psychology interventions: A meta-analysis of randomized controlled studies. *BMC Public Health*,
 28 13(1). <https://doi.org/10.1186/1471-2458-13-119>
 29
 30
 31
 32 *Boniwell, I., Osin, E. N., & Martinez, C. (2015). Teaching happiness at school: Non-randomised
 33 controlled mixed-methods feasibility study on the effectiveness of Personal well-being lessons.
 34 *Journal of Positive Psychology*, 11(1), 85–98. <https://doi.org/10.1080/17439760.2015.1025422>
 35
 36
 37
 38 Borenstein, M., Hedges, L., Higgins, J., & Rothstein, H. (2013). *Comprehensive meta-analysis version*
 39 *3.0*. Englewood, NJ: Biostat.
 40
 41
 42 Borenstein, M., Hedges, L. V, Higgins, J. P. T., & Rothstein, H. R. (2010). A basic introduction to fixed-
 43 effect and random-effects models for meta-analysis. *Research Synthesis Methods*, 1(2), 97–111.
 44 <https://doi.org/10.1002/jrsm.12>
 45
 46
 47
 48 *Burckhardt, R., Manicavasagar, V., Batterham, P. J., & Hadzi-Pavlovic, D. (2016). A randomized
 49 controlled trial of strong minds: A school-based mental health program combining acceptance and
 50 commitment therapy and positive psychology. *Journal of School Psychology*, 57, 41–52.
 51 <https://doi.org/10.1016/j.jsp.2016.05.008>
 52
 53
 54
 55
 56 Burger, K., & Samuel, R. (2017). The role of perceived stress and self-efficacy in young people's life
 57 satisfaction: A longitudinal study. *Journal of Youth and Adolescence*, 46(1), 78–90.
 58
 59
 60
 61
 62
 63
 64
 65

<https://doi.org/10.1007/s10964-016-0608-x>

- 1
2 Button, K. S., Ioannidis, J. P. A., Mokrysz, C., Nosek, B. A., Flint, J., Robinson, E. S. J., & Munafò, M.
3
4 R. (2013). Power failure: Why small sample size undermines the reliability of neuroscience. *Nature*
5
6 *Reviews Neuroscience*, *14*(5), 365–376. <https://doi.org/10.1038/nrn3475>
- 7
8 Chakhssi, F., Kraiss, J. T., Sommers-Spijkerman, M., & Bohlmeijer, E. T. (2018). The effect of positive
9
10 psychology interventions on well-being and distress in clinical samples with psychiatric or somatic
11
12 disorders: A systematic review and meta-analysis. *BMC Psychiatry*, *18*(1), 1–17.
13
14 <https://doi.org/10.1186/s12888-018-1739-2>
- 15
16 Ciocanel, O., Power, K., Eriksen, A., & Gillings, K. (2017). Effectiveness of positive youth development
17
18 interventions: A meta-analysis of randomized controlled trials. *Journal of Youth and Adolescence*,
19
20 *46*(3), 483–504. <https://doi.org/10.1007/s10964-016-0555-6>
- 21
22 Clonan, S. M., Chafouleas, S. M., McDougal, J. L., & Riley-Tillman, T. C. (2004). Positive psychology
23
24 goes to school: Are we there yet? *Psychology in the Schools*, *41*(1), 101–110.
25
26 <https://doi.org/10.1002/pits.10142>
- 27
28 Cuijpers, P. (2016). *Meta-analysis in mental health: A practical guide*. Amsterdam, The Netherland: Pim
29
30 Cuijpers Uitgeverij.
- 31
32 Dalton, D. R., Aguinis, H., Dalton, C. M., Bosco, F. A., & Pierce, C. A. (2012). Revisiting the file drawer
33
34 problem in meta-analysis: An assessment of published and nonpublished correlation matrices.
35
36 *Personnel Psychology*, *65*(2), 221–249. <https://doi.org/10.1111/j.1744-6570.2012.01243.x>
- 37
38 Datu, J. A. D. (2018). Flourishing is associated with higher academic achievement and engagement in
39
40 filipino undergraduate and high school students. *Journal of Happiness Studies*, *19*(1), 27–39.
41
42 <https://doi.org/10.1007/s10902-016-9805-2>
- 43
44 Derdikman-Eiron, R., Indredavik, M. S., Bratberg, G. H., Taraldsen, G., Bakken, I. J., & Colton, M.
45
46 (2011). Gender differences in subjective well-being, self-esteem and psychosocial functioning in
47
48 adolescents with symptoms of anxiety and depression: Findings from the Nord-Trøndelag health
49
50 study. *Scandinavian Journal of Psychology*, *52*(3), 261–267. [https://doi.org/10.1111/j.1467-](https://doi.org/10.1111/j.1467-9450.2010.00859.x)
51
52 [9450.2010.00859.x](https://doi.org/10.1111/j.1467-9450.2010.00859.x)
- 53
54 Diener, E. (1984). Subjective well-being. *Psychological Bulletin*, *95*(3), 542–575.
55
56 <https://doi.org/10.1037/0033-2909.95.3.542>
- 57
58 Dunning, D. L., Griffiths, K., Kuyken, W., Crane, C., Foulkes, L., Parker, J., & Dalgleish, T. (2018).
59
60
61
62
63
64
65

Research review: The effects of mindfulness-based interventions on cognition and mental health in children and adolescents – a meta-analysis of randomized controlled trials. *Journal of Child Psychology and Psychiatry and Allied Disciplines*. <https://doi.org/10.1111/jcpp.12980>

Duval, S., & Tweedie, R. (2000). Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*, *56*(2), 455–463. <https://doi.org/10.1111/j.0006-341X.2000.00455.x>

Egger, M., Smith, G. D., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test measures of funnel plot asymmetry. *BMJ*, *315*(7109), 629–634. <https://doi.org/10.1136/bmj.315.7109.629>

*Freire, T., Lima, I., Teixeira, A., Araújo, M. R., & Machado, A. (2018). Challenge to be+. A group intervention program to promote the positive development of adolescents. *Children and Youth Services Review*, *87*, 173–185. <https://doi.org/10.1016/j.childyouth.2018.02.035>

Froh, J. J., Kashdan, T. B., Ozimkowski, K. M., & Miller, N. (2009). Who benefits the most from a gratitude intervention in children and adolescents? Examining positive affect as a moderator. *The Journal of Positive Psychology*, *4*(5), 408–422. <https://doi.org/10.1080/17439760902992464>

Froh, J. J., Sefick, W. J., & Emmons, R. A. (2008). Counting blessings in early adolescents: An experimental study of gratitude and subjective well-being. *Journal of School Psychology*, *46*(2), 213–233. <https://doi.org/10.1016/j.jsp.2007.03.005>

González-Carrasco, M., Casas, F., Viñas, F., Malo, S., Gras, M. E., & Bedin, L. (2017). What leads subjective well-being to change throughout adolescence? An exploration of potential factors. *Child Indicators Research*, *10*(1), 33–53. <https://doi.org/10.1007/s12187-015-9359-6>

González-Carrasco, M., Vaqué, C., Malo, S., Crous, G., Casas, F., & Figuer, C. (2019). A Qualitative longitudinal study on the well-being of children and adolescents. *Child Indicators Research*, *12*(2), 479–499. <https://doi.org/10.1007/s12187-018-9534-7>

Hendriks, T., Schotanus-Dijkstra, M., Hassankhan, A., de Jong, J., & Bohlmeijer, E. (2019). The efficacy of multi-component positive psychology interventions: A Systematic review and meta-analysis of randomized controlled trials. *Journal of Happiness Studies*. <https://doi.org/10.1007/s10902-019-00082-1>

Higgins, J. P. T., & Thompson, S. G. (2002). Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine*, *21*(11), 1539–1558. <https://doi.org/10.1002/sim.1186>

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Keyes, L. M. (1998). Social well-being. *Social Psychology Quarterly*, *61*(2), 121–137.
<https://doi.org/10.2307/2787065>
- Keyes, L. M. (2002). The mental health continuum: From languishing to flourishing in life. *Journal of Health and Social Behavior*, *43*(2), 207–222. <https://doi.org/10.2307/3090197>
- Keyes, L. M. (2009). The nature and importance of positive mental health in America's adolescents. In *Handbook of Positive Psychology in Schools* (pp. 9–23). <https://doi.org/10.4324/9780203884089-10>
- Koydemir, S., Sökmez, A. B., & Schütz, A. (2020). A meta-analysis of the effectiveness of randomized controlled positive psychological interventions on subjective and psychological well-Being. *Applied Research in Quality of Life*. <https://doi.org/10.1007/s11482-019-09788-z>
- Lipsey, M. W., & Wilson, D. B. (1993). The efficacy of psychological, educational, and behavioral Treatment: Confirmation from meta-analysis. *American Psychologist*, *48*(12), 1181–1209.
<https://doi.org/10.1037/0003-066X.48.12.1181>
- Malle, B. F. (2006). Supplemental material for the actor–observer asymmetry in attribution: A (surprising) meta-analysis. *Psychological Bulletin*, *132*(6), 895–919. <https://doi.org/10.1037/0033-2909.132.6.895.supp>
- McHugh, M. L. (2012). Interrater reliability: The kappa statistic. *Biochemia Medica*, *22*(3), 276–282.
<https://doi.org/10.11613/bm.2012.031>
- Moher, D., Shamseer, L., Clarke, M., Ghersi, D., Liberati, A., Petticrew, M., ... Whitlock, E. (2015). Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews*, *4*(1), 1–9. <https://doi.org/10.1186/2046-4053-4-1>
- Noble, T., & McGrath, H. (2008). The positive educational practices framework: A tool for facilitating the work of educational psychologists in promoting pupil wellbeing. *Educational and Child Psychology*, *25*(2), 119–134.
- Paus, T., Keshavan, M., & Giedd, J. N. (2008). Why do many psychiatric disorders emerge during adolescence? *Nature Reviews Neuroscience*, *9*(12), 947–957. <https://doi.org/10.1038/nrn2513>
- Popp, L., & Schneider, S. (2015). Attention placebo control in randomized controlled trials of psychosocial interventions: Theory and practice. *Trials*, *16*(1), 1–3. <https://doi.org/10.1186/s13063-015-0679-0>
- Rosenberg, M. S. (2005). The file-drawer problem revisited: A general weighted method for calculating

fail-safe numbers in meta-analysis. *Evolution*, 59(2), 464–468. <https://doi.org/10.1111/j.0014-3820.2005.tb01004.x>

*Roth, R. A., Suldo, S. M., & Ferron, J. M. (2017). Improving middle school students' subjective well-being: Efficacy of a multicomponent positive psychology intervention targeting small groups of youth. *School Psychology Review*, 46(1), 21–41. <https://doi.org/10.17105/10.17105/spr46-1.21-41>

*Ruini, C., Ottolini, F., Tomba, E., Belaise, C., Albieri, E., Visani, D., ... Fava, G. A. (2009). School intervention for promoting psychological well-being in adolescence. *Journal of Behavior Therapy and Experimental Psychiatry*, 40(4), 522–532. <https://doi.org/10.1016/j.jbtep.2009.07.002>

Ryan, R. M., & Deci, E. L. (2001). On happiness and human potentials: A review of research on hedonic and eudaimonic well-being. *Annual Review of Psychology*, 52(1), 141–166. <https://doi.org/10.1146/annurev.psych.52.1.141>

Ryff, C. D. (1989). Happiness is everything, or is it? Explorations on the meaning of psychological well-being. *Journal of Personality and Social Psychology*, 57(6), 1069–1081. <https://doi.org/10.1037/0022-3514.57.6.1069>

Ryff, C. D. (1995). Psychological Well-Being in Adult Life. *Current Directions in Psychological Science*, 4(4), 99–104.

Schotanus-Dijkstra, M., Drossaert, C. H., Pieterse, M. E., Walburg, J. A., & Bohlmeijer, E. T. (2015). Efficacy of a multicomponent positive psychology self-help intervention: Study protocol of a randomized controlled trial. *JMIR Research Protocols*, 4(3), 1–16. <https://doi.org/10.2196/resprot.4162>

Seligman, M. (2011). *Flourish: A visionary new understanding of happiness and well-being*. New York, NY: Simon & Schuster.

Seligman, M. (2018). PERMA and the building blocks of well-being. *Journal of Positive Psychology*, 13(4), 333–335. <https://doi.org/10.1080/17439760.2018.1437466>

Seligman, M., Ernst, R. M., Gillham, J., Reivich, K., & Linkins, M. (2009). Positive education: Positive psychology and classroom interventions. *Oxford Review of Education*, 35(3), 293–311. <https://doi.org/10.1080/03054980902934563>

Shapero, B. G., Hankin, B. L., & Barrocas, A. L. (2013). Stress generation and exposure in a multi-wave study of adolescents: Transactional processes and sex differences. *Journal of Social and Clinical Psychology*, 32(9), 989–1012. <https://doi.org/10.1521/jscp.2013.32.9.989>

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- *Shoshani, A., & Steinmetz, S. (2014). Positive psychology at school: A School-based intervention to promote adolescents' mental health and well-being. *Journal of Happiness Studies*, *15*(6), 1289–1311. <https://doi.org/10.1007/s10902-013-9476-1>
- *Shoshani, A., Steinmetz, S., & Kanat-Maymon, Y. (2016). Effects of the Maytiv positive psychology school program on early adolescents' well-being, engagement, and achievement. *Journal of School Psychology*, *57*, 73–92. <https://doi.org/10.1016/j.jsp.2016.05.003>
- Sterne, J. A. C., Savović, J., Page, M. J., Elbers, R. G., Blencowe, N. S., Boutron, I., ... Higgins, J. P. T. (2019). RoB 2: A revised tool for assessing risk of bias in randomised trials. *The BMJ*, *366*. <https://doi.org/10.1136/bmj.l4898>
- Sterne, J. A., Egger, M., & Moher, D. (2008). Addressing reporting biases. In *Cochrane Handbook for Systematic Reviews of Interventions* (pp. 297–333). Chichester, UK: John Wiley & Sons, Ltd. <https://doi.org/10.1002/9780470712184.ch10>
- *Suldo, S. M., Savage, J. A., & Mercer, S. H. (2014). Increasing middle school students' life satisfaction: Efficacy of a positive psychology group intervention. *Journal of Happiness Studies*, *15*(1), 19–42. <https://doi.org/10.1007/s10902-013-9414-2>
- Taylor, R. D., Oberle, E., Durlak, J. A., & Weissberg, R. P. (2017). Promoting positive youth development through school- based social and emotional learning interventions: A meta- analysis of follow- up effects. *Child Development*, *88*(4), 1156–1171. <https://doi.org/10.1111/cdev.12864>
- *Tomba, E., Belaise, C., Ottolini, F., Ruini, C., Bravi, A., Albieri, E., ... Fava, G. A. (2010). Differential effects of well-being promoting and anxiety-management strategies in a non-clinical school setting. *Journal of Anxiety Disorders*, *24*(3), 326–333. <https://doi.org/10.1016/j.janxdis.2010.01.005>
- Turner, Herbert M., I., & Bernard, R. M. (2006). Calculating and synthesizing effect sizes. *Contemporary Issues in Communication Science and Disorders*, *33*, 42–55. https://doi.org/10.1044/cicsd_33_s_42
- UN General Assembly. (2015). *Transforming our world: The 2030 agenda for sustainable development (A/RES/70/1)*. United Nations.
- Vittersø, J. (2016). The most important idea in the world: An introduction. In *Handbook of Eudaimonic Well-Being* (pp. 1–24). Cham: Springer International Publishing. https://doi.org/10.1007/978-3-319-42445-3_1
- Waterman, A. S. (1993). Two conceptions of happiness: Contrasts of personal expressiveness (eudaimonia) and hedonic enjoyment. *Journal of Personality and Social Psychology*, *64*(4), 678–

691. <https://doi.org/10.1037/0022-3514.64.4.678>

1 Waters, L. (2011). A review of school-based positive psychology interventions. *Australian Educational*
2 *and Developmental Psychologist*, 28(2), 75–90. <https://doi.org/10.1375/aedp.28.2.75>
3

4 Weiss, L. A., Westerhof, G. J., & Bohlmeijer, E. T. (2016). Can we increase psychological well-being?
5 The effects of interventions on psychological well-being: A meta-analysis of randomized controlled
6 trials. *PLoS ONE*, 11(6). <https://doi.org/10.1371/journal.pone.0158092>
7
8
9

10 World Health Organization (2004). *Promoting Mental Health: Concepts, emerging evidence and*
11 *practice. Summary report Geneva*. Geneva.
12
13

14 World Health Organization (2017). *Depression and other common mental disorders global health*
15 *etimates*. Geneva.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 1. Study selection flow diagram




Figure 2. Forest plot of post-intervention effect sizes for subjective well-being




Figure 3. Forest plot of post-intervention effect sizes for psychological well-being




Figure 4. Forest plot of post-intervention effect sizes for depression symptoms




Figure 5. Forest plot of post-intervention effect sizes for anxiety symptoms

Table 1

Main characteristics of the included studies (N = 9)

First author (year)	Location	Sample size (age), % female	Design	Intervention type	Delivery	Duration in weeks (n sessions/hours)	Control group	Follow-up	Outcome measures
Boniwell (2015)	England, UK	164 (12 – 13 years), 50%	non-RCT	MPPI	Group	9 weeks (18 sessions)	Placebo	–	SWB: SLSS, MSLSS, PANAS-C
Burckhardt (2016)	Australia	267 (15 – 18 years), 50%	RCT	MPPI + ACT	Group	12 weeks (16 sessions)	Placebo	–	PWB: FS Dep/Anx/Str: DASS-21
Freire (2018)	Portugal	99 (13 – 17 years), 56%	non-RCT	MPPI + PYD	Group	8 weeks (8 sessions)	Placebo	–	SWB: SWLS PWB: PWBS-A
Roth (2017)	USA	42 (11 – 13 years), 50%	RCT	MPPI	Group	10 weeks (10 sessions)	Waitlist	5 and 7 weeks	SWB: BMSLSS, SLSS, PANAS-C
Ruini (2009)	Italy	227 (14 – 16 years), 61%	RCT	MPPI (WBT)	Group	6 weeks (6 sessions)	Placebo	6 months	PWB: PWB Anx: RCMAS, SQ; Dep: SQ
Shoshani (2014)	Israel	1038 (11 – 14 years), 50,5%	RCT	MPPI	Group	30 weeks (15 sessions)	Waitlist	6 and 12 months	SWB: SWLS Anx/Dep: BSI
Shoshani (2016)	Israel	2517 (12 – 15 years) 50%	RCT	MPPI	Group	30 weeks (15 sessions)	Waitlist	8 and 12 months	SWB: SWLS, PANAS

Table 1

Continued

First author (year)	Location	Sample size (age), % female	Design	Intervention type	Delivery	Duration in days or weeks (n sessions/hours)	Control group	Follow-up	Outcome measures
Suldo (2014)	USA	40 (10 – 12 years old), 62%	RCT	MPPI	Group	10 weeks (10 sessions)	Waitlist	6 months	SWB: BMSLSS, SLSS, PANAS-C
Tomba (2010)	Italy	162 (mean = 11.41 years) 57%	RCT	MPPI (WBT)	Group	6 weeks (6 sessions)	AM	6 months	PWB: PWB Anx: RCMAS, SQ; Dep: SQ

ACT: acceptance and commitment therapy; AM: anxiety-management school protocol; Anx: anxiety; Dep: depression; MMPI: multicomponent positive psychology intervention; non-RCT: non-randomized controlled-trial; PYD: positive youth development; PWB: psychological well-being; RCT: randomized controlled-trial; Str: stress; SWB: subjective well-being; WBT: well-being therapy. Abbreviations of questionnaires are listed in Appendix Table 2.

Table 2

Quality assessment of the studies included in the meta-analysis

Study	Domain 1	Domain 2	Domain 3	Domain 4	Domain 5	Overall score
Boniwell (2015)	HR	SC	LR	SC	LR	HR
Burckhardt (2016)	LR	LR	LR	SC	LR	SC
Freire (2016)	HR	LR	LR	SC	SC	HR
Roth (2017)	LR	LR	LR	SC	LR	SC
Ruini (2009)	LR	SC	LR	LR	HR	HR
Shoshani (2014)	LR	LR	LR	LR	LR	LR
Shoshani (2016)	LR	LR	LR	LR	LR	LR
Suldo (2014)	SC	LR	LR	SC	SC	SC
Tomba (2010)	LR	LR	LR	SC	LR	SC

Domain 1: risk of bias arising from the randomization process; Domain 2: risk of bias due to deviations from the intended interventions (effect of assignment to intervention); Domain 3: risk of bias due to missing outcome data; Domain 4: risk of bias in measurement of the outcome; Domain 5: risk of bias in selection of the reported results. LR: low risk; SC: some concerns; HR: high risk

Table 3
Between-group effects

Outcome measures	Studies	Hedges' g	95% CI	Z	Heterogeneity		Fail-safe N
					Q-value	I ²	
<i>Studies post-intervention</i>							
SWB	6	0.24	(0.11-0.37)	3.65 ^{***}	9.82 [*]	49.09	53
PWB	5	0.25	(-0.01-0.51)	1.86 [*]	22.58 ^{***}	82.29	41
Depression	4	0.28	(0.13-0.43)	3.69 ^{***}	5.64	46.84	27
Anxiety	4	0.14	(0.04-0.24)	2.96 ^{**}	2.17	0.00	3
<i>Studies post-intervention excluding low quality studies</i>							
SWB	4	0.21	(0.05-0.37)	2.59 [*]	8.09 [*]	62.91	
PWB	4	0.31	(-0.03-0.67)	1.74	15.93 ^{***}	87.45	
Depression	3	0.34	(0.24-0.44)	6.56 ^{***}	1.01	0.00	
Anxiety	3	0.15	(0.05-0.25)	2.99 ^{**}	1.75	0.00	
<i>Follow-up studies</i>							
SWB	4	0.13	(0.03-0.23)	2.54 [*]	4.10	30.34	
PWB	3	0.44	(-0.45-1.31)	1.80 [*]	87.65 ^{***}	97.71	
Depression	3	0.31	(0.20-0.41)	5.80 ^{***}	1.78	0.00	
Anxiety	3	0.15	(0.05-0.26)	2.99 ^{**}	1.74	0.00	
<i>Follow-up studies excluding low quality studies</i>							
PWB	2	0.66	(-0.39-1.72)	1.22	40.16 ^{***}	97.51	
Depression	2	0.29	(0.11-0.47)	3.24 ^{**}	1.51	34.05	
Anxiety	2	0.21	(0.10-0.33)	3.75 ^{***}	0.54	0.00	

PWB: psychological well-being; SWB: subjective well-being

*p < 0.05; **p < 0.01; ***p < 0.001

Table 4

Subgroup analysis

Outcome	Criteria	Value	N	Hedges' <i>g</i>	95% CI	I ²	Z
SWB	Year	Last 5 years	4	0.27	(0.20-0.34) ^{***}	0.00	7.21
		Older	2	0.08	(-0.03-0.20)	0.00	1.46
	Design	RCT	4	0.21	(0.05-0.37) ^{**}	62.73	2.06
		non-RCT	2	0.38	(0.12-0.64) ^{**}	0.00	2.94
	Intervention	MPPI	5	0.23	(0.09-0.37) ^{**}	54.63	3.24
		MPPI combined	1	0.45	(-0.00-0.90) [*]	0.00	1.95
	Control	Placebo	2	0.38	(0.12-0.64) ^{**}	0.00	2.94
		Waitlist	4	0.21	(0.05-0.37) ^{**}	62.73	2.60
	Sessions	≤ 6 sessions	–	–	–	–	–
		> 6 sessions	6	0.24	(0.11-0.37) ^{***}	49.09	3.65
	Duration	≤ 6 weeks	–	–	–	–	–
		> 6 weeks	6	0.24	(0.11-0.37) ^{***}	49.09	3.65
	Quality	High	2	0.17	(0.00-0.34) [*]	0.00	2.19
		Concerns	2	0.48	(0.05-0.93) [*]	82.20	2.19
		Low	2	0.38	(0.12-0.64) ^{**}	0.00	2.94
	Follow-up	Yes	4	0.21	(0.05-0.37) ^{**}	62.73	2.60
		No	2	0.38	(0.12-0.64) ^{**}	0.00	2.94
	PWB	Year	Last 5 years	2	0.12	(-0.08-0.34)	0.00
Older			3	0.33	(-0.00-0.67) [*]	85.29	1.91
Design		RCT	4	0.29	(0.00-0.57) [*]	84.62	2.00
		non-RCT	1	0.02	(-0.42-0.47)	0.00	0.09
Intervention		MPPI	3	0.33	(-0.00-0.67) [*]	85.21	1.91
		MPPI combined	2	0.12	(0.08-0.34)	0.00	1.87
Control		Placebo	3	0.15	(-0.00-0.32)	0.00	1.86
		Waitlist	2	0.38	(-0.08-0.85)	87.66	1.62
Sessions		≤ 6 sessions	2	0.16	(-0.03-0.36)	0.00	1.66
		> 6 sessions	3	0.29	(-0.08-1.50)	86.66	1.50
Duration		≤ 6 weeks	2	0.16	(-0.03-0.36)	0.00	1.66
		> 6 weeks	3	0.29	(-0.08-0.68)	86.66	1.50
Quality		High	2	0.60	(0.48-0.73) ^{***}	0.00	9.58
		Concerns	2	0.14	(-0.04-0.33)	0.00	1.52
		Low	1	0.15	(-0.07-0.38)	0.00	1.34
Follow-up		Yes	3	0.33	(-0.00-0.67) [*]	85.21	1.91
		No	2	0.12	(-0.08-0.34)	0.00	1.18
Depression		Year	Last 5 years	1	0.44	(0.20-0.69) ^{***}	0.00
	Older		3	0.23	(0.05-0.41) [*]	50.42	2.53
	Design	RCT	4	0.28	(0.13-0.43) ^{***}	46.68	3.69

		non-RCT	–	–	–	–	–
Anxiety	Intervention	MPPI	3	0.23	(0.05-0.41)*	50.42	2.53
		MPPI combined	1	0.44	(0.20-0.69)***	0.00	3.61
	Control	Placebo	2	0.24	(-0.15-0.64)	80.34	1.19
		Waitlist	2	0.32	(0.20-0.43)***	0.00	5.55
	Sessions	≤ 6 sessions	2	0.13	(-0.08-0.35)	15.21	1.19
		> 6 sessions	2	0.35	(0.24-0.46)***	0.00	6.38
	Duration	≤ 6 weeks	2	0.13	(-0.08-0.35)	15.21	1.19
		> 6 weeks	2	0.35	(0.24-0.46)***	0.00	6.38
	Quality	High	1	0.33	(0.21-0.45)***	0.00	5.32
		Concerns	2	0.37	(0.18-0.56)***	0.00	3.86
		Low	1	0.03	(-0.22-0.29)	0.00	0.27
	Follow-up	Yes	3	0.23	(0.05-0.41)*	50.42	2.53
		No	1	0.44	(0.20-0.69)***	0.00	3.61
	Year	Last 5 years	1	0.02	(-0.22-0.25)	0.00	0.08
		Older	3	0.17	(0.06-0.27)**	0.00	3.19
	Design	RCT	4	0.14	(0.04-0.24)**	0.00	2.96
		non-RCT	–	–	–	–	–
	Intervention	MPPI	3	0.17	(0.06-0.27)**	0.00	3.19
		MPPI combined	1	0.01	(-0.22-0.25)	0.00	0.08
	Control	Placebo	2	0.03	(-0.14-0.21)	0.00	0.38
Waitlist		2	0.18	(0.07-0.30)**	0.00	3.27	
Sessions	≤ 6 sessions	2	0.11	(-0.08-0.31)	0.00	1.17	
	> 6 sessions	2	0.13	(-0.03-0.29)	40.61	1.56	
Duration	≤ 6 weeks	2	0.11	(-0.08-0.31)	0.00	1.17	
	> 6 weeks	2	0.13	(-0.03-0.29)	40.61	1.56	
Quality	High	3	0.15	(0.05-0.25)**	0.00	2.99	
	Concerns	1	0.06	(-0.19-0.32)	0.00	0.47	
	Low	–	–	–	–	–	
Follow-up	Yes	3	0.17	(0.06-0.27)**	0.00	3.19	
	No	1	0.01	(-0.22-0.25)	0.00	0.08	

MPPI: multicomponent positive psychology intervention; PWB: psychological well-being; RCT: randomized controlled trial; non-RCT: non-randomized controlled trial; SWB: subjective well-being
 * p < 0.05; ** p < 0.01; *** p < 0.001

Appendix Table 1

Strings of the search

PUBMED: ((*"positive psych*"*[Title/Abstract] OR *"well-being"*[Title/Abstract]
OR *wellbeing*[Title/Abstract] OR *happy*[Title/Abstract] OR *happiness*[Title/Abstract]
OR *gratitude*[Title/Abstract] OR *optimism*[Title/Abstract] OR *kindness*[Title/Abstract]
OR *strengths*[Title/Abstract] OR *compassion*[Title/Abstract] OR *forgiveness*[Title/Abstract])) AND
(*intervention**[Title/Abstract] OR *prevention**[Title/Abstract] OR *program**[Title/Abstract]
OR *promotion*[Title/Abstract])) AND (*adolescent**[Title/Abstract] OR *youth*[Title/Abstract]
OR *teen**[Title/Abstract]) (filter: English, Spanish)

PSYCINFO: AB (*"positive psych*"* OR *"well-being"* OR *wellbeing* OR *happy* OR *happiness* OR
gratitude OR *optimism* OR *kindness* OR *strengths* OR *compassion* OR *forgiveness*) AND AB (*intervention** OR *prevention** OR *program** OR *promotion*) AND AB (*adolescent** OR *teenager** OR
youth OR *teen**) and TI (*"positive psych*"* OR *"well-being"* OR *wellbeing* OR *happy* OR *happiness*
OR *gratitude* OR *optimism* OR *kindness* OR *strengths* OR *compassion* OR *forgiveness*) AND TI (*intervention** OR *prevention** OR *program** OR *promotion*) AND TI (*adolescent** OR *teenager** OR
youth OR *teen**) (filter: academic journals, English, Spanish)

SCOPUS: (TITLE-ABS-KEY (*"positive psych*"* OR *"well-*
being" OR *wellbeing* OR *happy* OR *happiness* OR *gratitude* OR *optimism* OR *kindness* OR *str*
engths OR *compassion* OR *forgiveness*) AND TITLE-ABS
KEY (*intervention** OR *prevention** OR *program** OR *promotion*) AND TITLE-ABS-
KEY (*adolescent** OR *youth* OR *teen**)) AND DOCTYPE (ar) AND (LIMIT-
TO (SUBJAREA , "PSYC") OR LIMIT-TO (SUBJAREA , "SOCI")) AND (LIMIT-
TO (LANGUAGE , "English") OR LIMIT-TO (LANGUAGE , "Spanish"))

Appendix Table 2

Abbreviations of questionnaires

Subjective well-being: BMSLSS: The Brief Multidimensional Student's Life Satisfaction Scale;

MSLSS: Multidimensional Students' Life Satisfaction Scale; PANAS-C: Positive and Negative Affect

Schedule for Children; SLSS: Students' Life Satisfaction Scale; SWLS: Satisfaction with Life Scale

Psychological well-being: FS: Flourishing Scale; PWB: Psychological Well-Being; PWBS-A:

Psychological Well-Being Scale for Adolescents

Depression: BSI: Brief Symptoms Inventory; DASS-21: Depression, Anxiety and Stress Scale; SQ:

Kellner's Symptom Questionnaire

Anxiety: BSI: Brief Symptoms Inventory; DASS-21: Depression, Anxiety and Stress Scale; RCMAS:

Revised Children Manifest Anxiety Scale; SQ: Kellner's Symptom Questionnaire

Stress: DASS-21: Depression, Anxiety and Stress Scale

Authors' Contributions CTG conceived of the study, participated in the initial literature search, the evaluation of quality of studies, conducted the preliminary statistical analysis and wrote the first draft of the manuscript; ABB advised the design, participating in the identification of relevant studies, the evaluation of quality of studies, interpretation of the study findings and helped to draft the manuscript; CTN participated in the literature search, data extraction and statistical analysis; CA contributed to the conceptualization of the study, interpretation of the findings and oversaw study execution. All authors read and approved the final manuscript.

Data Sharing Declaration The datasets generated and/or analyzed during the current study are not publicly available but are available from the corresponding author on reasonable request.

Compliance with Ethical Standards

Conflicts of Interest The authors declare that they have no conflict of interest.

Preregistration The present manuscript was preregistered (study design, data extraction, quality assessment and data synthesis) in PROSPERO, an international prospective register for systematic reviews, with the following registration number: #CRD42019139586.

Ethical Approval The current manuscript used published studies as the sole data source and did not involve any other data collection or direct interactions with human participants. As such, seeking approval from a human subjects' review committee is irrelevant.

Informed Consent This research did not involve any data collection or direct interaction with participants and therefore informed consent was not obtained.