

鳥取大学研究成果リポジトリ

Tottori University research result repository

タイトル Title	Effects of substituting alfalfa hay for concentrate on energy utilization and feeding cost of crossbred Simmental male calves in Gansu Province, China
著者 Author(s)	Kobayashi, Nobuyuki; Hou, Fujiang; Tsunekawa, Atsushi; Chen, Xianjiang; Yan, Tianhai; Ichinohe, Toshiyoshi
掲載誌・巻号・ページ Citation	Grassland science , 63 (4) : 245 - 254
刊行日 Issue Date	2017-08-01
資源タイプ Resource Type	学術雑誌論文 / Journal Article
版区分 Resource Version	著者版 / Author
権利 Rights	© 2017 Japanese Society of Grassland Science
DOI	10.1111/grs.12169
URL	http://repository.lib.tottori-u.ac.jp/5712

1 Short running title: Energy use of Chinese alfalfa by calf

2

3 Effects of substituting alfalfa hay for concentrate on energy utilization and feeding cost of
4 crossbred Simmental male calves in Gansu Province, China

5

6 Nobuyuki Kobayashi^{1,2}, Fujiang Hou³, Atsushi Tsunekawa², Xianjiang Chen³, Tianhai Yan⁴
7 and Toshiyoshi Ichinohe⁵

8

9

10 1 The United Graduate School of Agricultural Sciences, Tottori University, Tottori, Japan

11 2 Arid Land Research Center, Tottori University, Tottori, Japan

12 3 College of Pastoral Agriculture Science and Technology, Lanzhou University, Lanzhou,
13 China

14 4 Agri-Food and Biosciences Institute, Belfast, Northern Ireland, UK

15 5 Faculty of Life and Environmental Science, Shimane University, Matsue, Japan

16

17

18

19

20

21 Correspondence

22 Toshiyoshi Ichinohe, Faculty of Life and Environmental Science, Shimane University,
23 Matsue 690-8504, Japan.

24 Email: toshi@life.shimane-u.ac.jp

25

1 **Abstract**

2

3 From August to September 2015 (trial 1 (T1)) and September to November 2015 (trial 2
4 (T2)), the effects of replacing concentrate feed (C) with alfalfa hay (AH) on the daily gain
5 (DG), dietary energy utilization status and the economic advantage of AH feeding for
6 growing beef cattle were studied in crossbred male Simmental calves (n = 18) in Gansu
7 Province, China. The target DG was set as 1 kg for the both trials. Animals in each trial
8 were allocated to a conventional feeding group (CTRL), a low-level AH feeding group
9 (LA), and a high-level AH feeding group (HA). In a one-way-layout design, they were fed
10 iso-energetic experimental diets comprising harvested corn stover (CS) and C (T1-CTRL,
11 T2-CTRL), diets replacing 22% (T1-LA) or 44% (T1-HA) of the quantity of C for
12 T1-CTRL with AH, and diets replacing 13% (T2-LA) or 25% (T2-HA) of the quantity of C
13 for T2-CTRL with AH. Measurements of feed intake and DG, respiration and metabolism
14 trials were performed for 49 and 41 days in T1 and T2, respectively. Average DG did not
15 reach the target value for HA in T1 and CTRL in T2. Energy metabolizability was slightly
16 greater for CTRL than for LA and HA in T1 and significantly greater for CTRL than for the
17 other groups in T2. There was no marked difference in energy metabolizability between LA
18 and HA in the both trials. Dietary substituting AH for C did not impair the feed intake of the
19 animals, but it did not improve feed efficiency. In terms of economic feasibility, low-level
20 AH inclusion in the diets of growing beef cattle appeared more profitable at the 1-kg DG
21 level as compared with CTRL and high-level AH inclusion, and should be practiced in the
22 drylands of Gansu Province, China.

23

24 **Keywords**

1 Alfalfa hay; beef cattle; China; dryland; energy metabolism.

2

1 **Introduction**

2

3 The rates of beef cattle production and beef consumption are increasing rapidly in China
4 (FAO 2015) because of the dramatic shift in dietary preferences that have coincided with
5 the rapid economic growth and the improvement of living standards. The increase in animal
6 feed demand has resulted in yearly increases in the quantities of imported concentrate feed
7 ingredients and forages in China (MAFF 2013).

8 Gansu Province is one of the major beef cattle production areas of China (National
9 Bureau of Statistics of China 2014) and is being prioritized as a beef production area by the
10 Chinese Government (Chinese State Committee for Development and Reform 2013). In
11 many provinces in China, including Gansu Province, livestock grazing has been restrained
12 to prevent the desertification of natural pastures in accordance with the directive “restore
13 agricultural land to forest and pasture” which had issued in 2003 (Han *et al.* 2008).
14 Therefore, foraging systems for confined beef cattle with both maximum utilization of
15 roughage produced on farm and high-level production performance are needed to be
16 established in Gansu Province. The effect of stall-feeding regimens conventionally used in
17 Gansu Province should be evaluated to develop beef cattle production models based on
18 harvested roughage diets in the huge dryland area of western China. Moreover, feeding
19 studies based on locally available feeds would help to upgrade the current Chinese Feeding
20 Standard for Beef Cattle (CFSBC) (Ministry of Agriculture of the People’s Republic of
21 China 2004).

22 Rong *et al.* (2004) reported that alfalfa has high drought tolerance, and there is strong
23 potential to increase the area under alfalfa cultivation because of its high feeding values and
24 economic return to farmers (Guo *et al.* 2007). In Huanxian County in Gansu Province, the

1 area under alfalfa cultivation has expanded and it accounted for 28% of all irrigated
2 farmland in the Yellow River Basin of Gansu Province in 2006 (Hou *et al.* 2008). Since
3 alfalfa grown in Gansu Province can be considered as a self-sufficient substitute for
4 expensive commercial concentrates, utilization of the local alfalfa might reduce the beef
5 cattle feeding cost in Gansu Province.

6 Simmental crossbred is the major types of beef cattle kept by farmers in Gansu
7 Province. Currently, there is an absence of data from both *in vivo* and *in situ* studies on the
8 energy utilization status of conventional beef cattle diets in Gansu Province and the effects
9 of alfalfa hay (AH) feeding on energy metabolizability. Hence, by performing two seasonal
10 feeding trials (in the warm and cool seasons) and energy balance trials, we aimed to
11 evaluate the effect of AH feeding level on the body weight (BW) gain, efficiency of dietary
12 energy utilization, and feeding cost of growing Simmental cattle in Gansu Province, China.

13

14 **Materials and methods**

15

16 **Study site**

17 The experiment was conducted at the Linze Research Station of the College of Pastoral
18 Agriculture Science and Technology, Lanzhou University, located in Linze County, Gansu
19 Province, China (Figure 1). The research station is located at latitude 39.24°N and
20 longitude 100.06°E, and its elevation is 1390 m above sea level (Zhu *et al.* 1997). The total
21 precipitation in 2013 was 89 mm, all of which fell from May to September. The annual
22 average temperature in the same year was 8.3 °C. The average temperature in the warm
23 season (from August to September) and the cool season (from September to November) in
24 2013 was 18.1 and 7.7 °C, respectively (data supplied from the Linze Research Station).

1 The study site is categorized as a typical arid zone (United Nations Environment
2 Programme 1997).

3

4 **Cattle, diets, and feeding management**

5 Two feeding trials were conducted, from 6 August to 16 September 2015 (trial 1 (T1) in the
6 warm season) and from 24 September to 27 October 2015 (trial 2 (T2) in the cool season),
7 with the aim of achieving a 1-kg daily gain (DG). Eighteen male crossbred Simmental
8 calves with a mean body weight (BW) of 175.8 ± 23.8 kg (7 months of age) and with a mean
9 BW of 218.8 ± 27.4 kg (8 months of age) were purchased from a local calf market and used
10 for T1 and T2, respectively. We have conducted an interview to beef cattle farmers on their
11 feeding management, and concluded that the amount of concentrate feeding were markedly
12 higher in the cool season than in the warm season in Gansu province. In designing
13 experimental feeding regimens, we considered the conventional feeding practices of beef
14 cattle farmers in Gansu Province.

15 Based on the initial BW, the 18 calves in each trial were allocated to one of the
16 following three dietary treatments, namely a conventional feeding group (control (CTRL),
17 $n = 6$), a low-level AH feeding group (LA, $n = 6$) and a high-level AH feeding group (HA,
18 $n = 6$). In the CTRL groups of T1 and T2 (T1-CTRL, T2-CTRL), allocated calves were
19 offered corn stover (CS) as a sole forage diet together with concentrate feed (C). The diet
20 fed to the CTRL group in T1 and T2 was designed in reference to the conventional diets
21 used by beef cattle farmers in Gansu Province. The calves in the LA group were offered CS
22 with a low level of AH and C; those in HA group were offered CS with a high level of AH
23 and C. The experimental diet in each treatment was designed to suffice metabolizable
24 energy (ME) and metabolizable protein requirements for a 1-kg DG of a bull calf on the

1 basis of the estimation equation of the AFRC (1993) and the tabular values listed in the
2 CFSBC (2004) using weekly measured BW of the calves. Tabular values listed in the
3 *Standard Tables of Feed Composition in Japan (2009)* (NARO 2010) were used when
4 convert the digestible energy values of CS, AH and feed ingredients of C listed in CFSBC
5 (2004) into the ME concentration. Assured nutritive values of commercial concentrate
6 comprising C was also used to calculate the ME concentration of C. Since the ME
7 requirement for 1-kg DG of $\text{kg}^{-0.75}$ BW of calf was considered to be greater for T2 than that
8 for T1 (CFSBC 2004), concentrate-based experimental diets were designed in T2 to suffice
9 the ME requirement by taking account of the estimated DM intake (DMI) calculated by the
10 equation of AFRC (1993). In T1-LA, 22% (on a dry matter (DM) weight basis) of the daily
11 allowance of C in T1-CTRL was replaced with AH; in T1-HA, 44% of the daily C in
12 T1-CTRL was replaced with AH. In T2-LA, 13% of the daily C in T2-CTRL was replaced
13 with AH; and in T2-HA, 25% of the daily C in T2-CTRL was replaced with AH.

14 The CS and AH used in T1 and T2 were produced at the Linze Research Station of the
15 College of Pastoral Agriculture Science and Technology. The CS used in the both trials was
16 harvested in September 2014. The AH used in this experiment was harvested in May 2015
17 (first-cut hay, harvested before the bloom stage of growth in 2015) for T1, and in July
18 (second-cut hay) and September (third-cut hay) 2015 for T2. The AH used for T1 and T2
19 was coarsely chopped, mixed and stored for use as the experimental feed. The C consisted
20 of commercial concentrate (31%), wheat bran (10%) and corn grain (59%). The corn grain
21 was produced at the Linze Research Station in September 2014. Throughout the feeding
22 trials, the calves were housed in individual pens and had free access to fresh water and
23 mineral blocks. Coarsely chopped CS and AH (5- to 10-cm lengths) and C were offered via
24 a separate trough for each animal as separate meals each day. The chopped roughage was

1 offered twice a day (07.30 and 19.30 hours), and C was offered once a day at 14.30 hour.

2

3 **Measurements and sample collection**

4 In both T1 and T2, the amount of feed offered and theorts were weighed and recorded daily
5 throughout the experimental period to calculate daily feed intake. Representative samples
6 of CS, AH, and C were collected several times during the feeding study and composited for
7 analyses of their chemical composition. The calves were weighed at the beginning and the
8 end of the feeding study, as well as weekly. On the basis of the BW of the animals, the feed
9 provision level for each animal allocated to the CTRL, LA, and HA groups was calculated
10 weekly.

11 At the conclusion of the feeding trials, respiratory measurements were conducted for 2
12 days, following 5-days adaptation, at the Linze research station for 15 calves in T1 and 12
13 calves in T2 (selected from the three dietary groups, i.e. five calves in T1 and four calves in
14 T2 per group) by using ventilated open-circuit respiration chambers. Oxygen consumption
15 and CO₂ and CH₄ production were measured by both infrared absorption based gas
16 analyzer (CO₂ and CH₄) and paramagnetic-based O₂ gas analyzer (VA-3000, Horiba Ltd.,
17 Kyoto, Japan). Daily heat production (HP) by each animal was calculated by using the
18 equation of Brower (1965). The ME for maintenance (ME_m) of male calves was estimated
19 by using a linear regression equation between net energy (NE) intake (obtained by
20 subtracting HP from ME intake (MEI)) and MEI, with both expressed on the basis of
21 metabolic body size (kg^{0.75} BW), as presented by Freetly *et al.* (2006). During the
22 respiration trial, experimental diets were offered to the animals on the same schedule as in
23 the feeding trials. During a 5-days period of adaptation of the animals to the respiration
24 chamber and 2 days of respiratory measurement, total urine collection and spot feces

1 collection were conducted to determine the daily urinary and fecal gross energy (GE)
2 excretion of calves allocated to the CTRL, LA and HA groups. Excreted urine was
3 collected into a container containing 50 mL of 10% (v/v) H₂SO₄ to keep urine pH below 3.
4 Urine samples (about 100 mL) were taken from each animal and stored at -15 °C for
5 further analysis. Daily fecal excretion was estimated by using acid detergent lignin (ADL)
6 as an internal marker to determine digestibility.

7

8 **Chemical analysis**

9 Collected feed and fecal samples were dried at 60 °C in a forced-air oven. The dried
10 samples were ground and sieved to pass through a 1-mm screen. The concentrations of GE,
11 DM, crude protein (CP), ash-free neutral detergent fiber (NDFom) and ADL in the dried
12 feed samples and fecal samples were determined. Urine samples were determined for GE
13 concentrations. The concentrations of DM and CP were determined by using the method of
14 the Association of Official Analytical Chemists (AOAC 1984). The concentrations of
15 NDFom and ADL were determined by the procedure of Van Soest *et al.* (1991). The GE
16 concentration of the samples was determined with a bomb calorimeter (CA-4AJ, Shimadzu
17 Corp., Kyoto, Japan). Urine samples were freeze-dried for 48 hours in a freeze dryer
18 (VD550R, Taitec Corp., Koshigaya, Japan), and the GE concentration was determined with
19 the bomb calorimeter (CA-4AJ, Shimadzu Corp.).

20

21 **Economic analysis**

22 To examine the effect of substituting AH for C in terms of economic feasibility, the
23 differences in feeding cost among the dietary treatment groups were calculated in both T1
24 and T2. Feed cost was calculated as the sum of expenses for purchase of alfalfa hay and

1 commercial concentrate by using their market price (1.7 yuan kg⁻¹ for alfalfa hay, 2.9 yuan
2 kg⁻¹ for concentrate) and the daily intake of AH and C for each cattle obtained in T1 and T2.
3 In addition, the economic benefit of the calves' DG was estimated by subtracting the feed
4 cost from the expected income (profit) from DG calculated with the market price of calves
5 (22 yuan kg⁻¹ BW). The estimates were converted at the rate of 1 US\$ = 6.36 yuan (based
6 on the averages between 6 August and 1 November 2015).

7

8 **Statistical analysis**

9 Differences in means among the three groups in each trial were tested by using one-way
10 ANOVA and Tukey's test. Possible seasonal differences in the efficiency of energy
11 utilization were not considered because of the difference in feeding regimen in T1 and T2.
12 All statistical analyses were performed with statistical software R (Version 3.1.1, The R
13 Foundation for Statistical Computing, Vienna, Austria). Dietary effect was considered
14 significant at $P \leq 0.05$.

15

16 **Results and discussion**

17

18 **Chemical composition of diet**

19 The GE concentration and chemical composition of CS did not differ between T1 and T2
20 (Table 1). Minson (1980) reported that at least a CP concentration of 8% (on a DM basis) is
21 needed if forage is given as a sole diet to ruminants. The concentrations of CP in CS ranged
22 from approximately 4 to 5% on a DM basis and that of NDFom was about 70% of DM,
23 indicating that CS could not be used as the basal diet for beef cattle to achieve high levels
24 of production; thus a concentrate or a leguminous forage supplement should be needed to

1 achieve 1-kg DG in crossbred Simmental male calves when fed CS as the basal forage.
2 Reflecting the differences in harvesting stage of alfalfa, the concentrations of CP, NDFom
3 and ADL of AH differed slightly between T1 and T2. The CP concentrations of AH in T1
4 and T2 (15.8 and 12.0% on a DM basis, respectively) were lower than the tabular values
5 listed by NARO (2010) (ranging from 19.1 to 21.8% on a DM basis) but were close to the
6 value listed in CFSBC (2004) (13.1% on a DM basis). Our finding, the CP concentrations
7 of AH in the both trials were close to that of C, suggests that it may be possible to replace C
8 with AH to provide CP in CS based diet feeding for beef calves. The NDFom and ADL
9 concentrations of the second- and third-cut AH in T2 were slightly higher than those of the
10 first-cut AH in T1. The NDFom concentrations in AH harvested before the bloom stage
11 were shown to vary from 36.0 to 39.3% on a DM basis (NARO 2010), the higher NDFom
12 concentrations of AH that we obtained might have been caused by unavoidable
13 contamination at harvest with other poaceous forage crops grown at the research station.
14 The difference in the NDFom of C between T1 and T2 might be likely a result of variations
15 in the nutrients concentration of the wheat bran and corn grain added to the commercial
16 concentrate.

17 The daily feed allowance for calves at the start of T1 and T2 are shown in Table 2. As
18 described above, the feed allowance for each animal was calculated weekly on the basis of
19 the BW with keeping the feed proportion of the feeding treatments in each trial. The values
20 shown in Table 2 were those fed at 175.8 kg and 218.8 kg BW in T1 and T2, respectively.
21 The estimated CP concentration of the diets fed to the calves ranged from 8.9 to 9.7% on a
22 DM basis in T1 and from 11.7 to 11.9% on a DM basis in T2 (Table 2). The estimated
23 NDFom concentration of the diets ranged from 45.1 to 52.5% on a DM basis in T1 and
24 from 29.1 to 37.5% on a DM basis in T2. The estimated NDFom concentration tended to

1 increase as AH feeding level increased. In contrast, the variation in estimated CP
2 concentration among the feeding groups in each of T1 and T2 was less than that of the
3 estimated NDFom concentration. The ADL concentrations were more than 2% on a DM
4 basis for all of the diets, and were sufficiently high eligible to use as an internal marker to
5 estimate fecal DM excretion (Munitifering 1982).

6

7 **Feed intake, metabolizable energy intake, daily gain, feed efficiency, and digestibility**

8 The average feed intakes of the calves over the 1-month period of each feeding trial are
9 summarized for each feeding group in Table 3. Feed refusal of C was observed for the
10 CTRL group in both T1 and T2. The refusal of forage (CS and AH) for T2-HA (0.13 kg
11 DM day⁻¹) appeared to be greater than that for T2-CTRL (0.03 kg DM day⁻¹) and T2-LA
12 (0.07 kg DM day⁻¹). There were no significant differences in total DMI, expressed as kg
13 day⁻¹ and % BW, among the T1 feeding groups. The NDFom concentration of daily
14 ingested feed (on a DM basis) for T1-CTRL, T1-LA and T1-HA was calculated as 39.9,
15 44.3 and 49.5%, respectively, and it appeared to increase with increasing AH feeding level.
16 Forage fiber concentration clearly affects voluntary feed intake through the physical
17 regulation caused by the rumen fill (Ichinohe *et al.* 1994). In contrast, Hales *et al.* (2014)
18 has reported no difference in DMI when alfalfa hay is substituted at various mixing ratios
19 for concentrate-based diets. Our results in T1 also revealed that dietary substitution with
20 AH, regardless of the consequent increase in NDFom intake, did not reduce the total DMI.
21 In T2, however, total DMI (kg day⁻¹ and % BW) differed significantly ($P < 0.05$) among
22 the three feeding groups; that for T2-CTRL had the lowest value, and the total DMI for
23 T2-HA was significantly greater ($P < 0.05$) than that for T2-LA. The NDFom concentration
24 of the ingested feed (on a DM basis) for T2-CTRL, T2-LA and T2-HA was calculated to be

1 33.3, 37.5 and 40.7%, respectively. As observed in T1, they increased with increasing AH
2 feeding level. Opposite to the result observed in T1, the increase of NDFom intake with
3 increasing AH feeding level resulted in the significant increase in total DMI in T2. Our
4 results for total DMI in T2 supported the findings of a study of growing ruminants by
5 Defoor *et al.* (2002), in which they suggested that growing heifers ate more with increasing
6 NDFom concentration of the diet to compensate for the reduction in dietary energy intake.
7 Furthermore, the result of feed intake in T2 appeared to indicate that differences in the
8 dietary forage to concentrate ratio affected feed intake as suggested by McCarthy *et al.*
9 (1989), Krause *et al.* (2002) and Oba and Allen (2003). Kennedy and Murphy (1988)
10 reviewed digesta particle outflow from the rumen, and summarized that additionally to the
11 physiological rumen fill caused by forage feeding, the cold temperature also enhanced
12 rumen digesta outflow rate. Hence, it was thought that both of cold temperature in October
13 2015 at Linze research station and differences in the ruminal NDFom contents caused by
14 the differed NDFom intake might have resulted in the significant differences in total DMI
15 among the treatments in T2. Our results suggested that a dietary AH allowance up to 50%
16 of C in warm season to 80% of C in cool season (Table 3) did not have any detrimental
17 effects on feed intake in the both trials.

18 We calculated the average daily MEI and DG of the calves (Figure 2). The MEI did not
19 differ significantly among the feeding groups in T1, although T1-HA group showed slightly
20 lower value than the other feeding groups ($P > 0.05$). In contrast, the MEI was significantly
21 higher ($P < 0.05$) for T2-HA than for T2-CTRL, and it did not differ between T2-LA and
22 T2-HA. Over the 1-month feeding period, the values of MEI expressed as a percentage of
23 the ME required for 1-kg DG in male calves were calculated as the following: 97% for
24 T1-CTRL, 90% for T1-LA, 78% for T1-HA, 98% for T2-CTRL, 108% for T2-LA and

1 119% for T2-HA. No significant difference in MEI between T2-CTRL and T2-LA or
2 between T2-LA and T2-HA was detected ($P > 0.05$; Figure 2). The average DG values (kg
3 day⁻¹) throughout the observation period were calculated for T1-CTRL, 0.9; T1-LA, 1.0;
4 T1-HA, 0.6; T2-CTRL, 0.7; T2-LA, 1.2 and T2-HA, 1.2. The lower average DG for T1-HA
5 than those for the other groups in T1 ($P < 0.05$) might be caused by the numerically lower
6 MEI for T1-HA. The value of average DG for T2-CTRL was significantly lower ($P < 0.05$)
7 than those for the other groups in T2, whereas those for T2-LA and T2-HA did not differ
8 significantly ($P > 0.05$). The average DG achieved the target growth performance for
9 T2-LA and T2-HA in T2 as designed. Hales *et al.* (2014) reported that the retained energy
10 (RE) for BW gain in finishing beef cattle decreased as dietary concentration of AH
11 increased. The numerically lower MEI for T1-HA than for T1-CTRL and T1-LA might
12 have reduced the RE in T1-HA as compared with those in the other two groups, and thus
13 resulted in the significantly lower DG value ($P < 0.05$). However, the lack of DG difference
14 between T1-CTRL and T1-LA indicated that the AH feeding level in T1-LA treatment did
15 not cause the RE depression that might have been thought to be occurred in the T1-HA
16 group. The lower DG value observed in T2-CTRL as compared with those in the other two
17 groups in T2 may have been attributable to the lower intake of C in T2-CTRL. As indicated
18 in Table 2 and Table 3, daily consumption of C accounted for 65% of allowance (4.8 kg
19 DM day⁻¹) in T2-CTRL group. Since T2-CTRL treatment was not able to suffice the ME
20 requirement to achieve the target 1-kg DG, concentrate allowance of T2-CTRL appeared to
21 be too much when fed CS based diet to growing calf in the cool season in Gansu Province.
22 Our results for average MEI and DG suggested that AH provision at the LA-group in both
23 T1 and T2 had no negative effect on energy retention for BW gain of growing male calves.

24 The trend of the average feed conversion ratio (FCR) among the three feeding groups

1 differed between T1 and T2 (Table 3). The value of FCR for T1-CTRL and T1-LA did not
2 differ from each other but was significantly higher ($P < 0.05$) than that for T1-HA. The
3 FCR values in T1 paralleled the average DG (Figure 2), indicating that the T1-LA diet was
4 comparable to a conventional concentrate-based diet in Gansu Province in the warm season.
5 The FCR did not differ significantly between the feeding groups in T2, but it was
6 numerically higher for T2-LA than for the other feeding groups. Although there was no
7 clear relationship between DG and FCR in T2, the merit of low-level substitution of AH for
8 C, as compared with high-level AH feeding, was thus emphasized in both T1 and T2.

9 The digestibility of DM, CP and NDFom in the three feeding groups did not differ
10 significantly in T1 ($P > 0.05$, Table 3). Conversely, the digestibility of DM, CP and NDFom
11 for T2-CTRL was significantly higher than for T2-LA and T2-HA ($P < 0.05$). Afore
12 discussed, the depression of NDFom digestibility for AH feeding groups in T2 might have,
13 partly, been explained by a higher outflow rate of rumen digesta in the cool season
14 (Kennedy and Murphy 1988). The difference in AH feeding level between LA and HA did
15 not affect nutrient digestibility in the both trials.

16

17 **Energy utilization efficiency**

18 Gross energy intake did not differ significantly ($P > 0.05$) among the feeding groups in T1,
19 whereas it differed significantly ($P < 0.05$) among the feeding groups in T2 with showing
20 the highest value for T2-HA, followed by T2-LA and then T2-CTRL (Table 4). In T1, the
21 energy digestibility of the feeding groups did not differ significantly ($P > 0.05$), although
22 the increase in fecal energy excretion (from 528.2 for T1-CTRL to 633.5 $\text{kJ kg}^{-0.75} \text{BW day}^{-1}$
23 ¹ for T1-HA; data not shown) with increasing AH feeding level resulted in the slight
24 decrease in energy digestibility. In contrast, energy digestibility was significantly ($P < 0.05$)

1 greater for T2-CTRL than for the other two groups; in T2, the values were in the reverse
2 order of ranking of total DMI (Table 3). As was the trends observed for digestion
3 coefficients of gross energy in the two trials, energy metabolizability did not differ among
4 groups in T1 ($P > 0.05$), but it was significantly greater ($P < 0.05$) for T2-CTRL than for
5 the other T2 groups. Plausibly, the lack of difference in MEI and energy metabolizability (P
6 > 0.05) between T2-LA and T2-HA might explain the lack of difference in DG between
7 these two groups (Figure 2). Increasing the dietary substitution rate of AH did not markedly
8 improve energy metabolizability in T1 and T2. In both T1 and T2, CH₄ production was
9 significantly greater for the HA feeding group than for the other two groups ($P < 0.05$).
10 Hales *et al.* (2014), in a study using dry-rolled corn-based diets supplemented with AH,
11 reported that fecal energy loss and methane production increased with decreasing energy
12 digestibility as the AH feeding level increased owing to an increase in NDF intake and
13 coinciding decrease in NDF digestibility. The numerical and significant differences in
14 NDFom digestibility were observed in T1 and T2, respectively, and the relationship across
15 CH₄ production, NDFom digestibility and NDFom intake observed in the current study
16 appeared to be consistent with those of Hales *et al.* (2014).

17 The values for energy digestibility and metabolizability (Table 4) agreed with the results
18 of Liu *et al.* (2013), who calculated 66% energy digestibility and 58% energy
19 metabolizability in Xiangzhong Black bulls. There was no large difference between
20 crossbred Simmental calves and other breeds in terms of energy metabolizability. On the
21 other hand, MEm, which was calculated as 652 kJ kg^{-0.75} BW day⁻¹ in T1 and 600 kJ kg^{-0.75}
22 BW day⁻¹ in T2 (Figure 3), seemed to be higher than the previous reported values of 469 kJ
23 kg^{-0.75} BW day⁻¹ (Henrique *et al.* 2005), 460 kJ kg^{-0.75} BW day⁻¹ for Nelore bulls, 485 kJ
24 kg^{-0.75} BW day⁻¹ for Nelore steers (Tedeschi *et al.* 2002), 506 kJ kg^{-0.75} BW day⁻¹ for

1 Chinese water buffalo (Qin *et al.* 2011) and $506 \text{ kJ kg}^{-0.75} \text{ BW day}^{-1}$ for Xiangzhong Black
2 bulls (Liu *et al.* 2013). This suggests that the relatively larger MEm requirement of
3 crossbred Simmental bull calves might have resulted in a large ME requirement for BW
4 gain than that of other species.

5 In this study, T1-LA, T2-LA and T2-HA treatment exceeded the targeted DG value, and
6 the ratio of DG to MEI (DG ME^{-1} , in $\text{g MJ}^{-1} \text{ day}^{-1}$) was calculated as 23.2, 20.1 and 19.3
7 for T1-LA, T2-LA and T2-HA, respectively. These values were higher than those reported
8 previously as $19.2 \text{ g MJ}^{-1} \text{ day}^{-1}$ for Zebu cows and $15.7 \text{ g MJ}^{-1} \text{ day}^{-1}$ for Brown Swiss
9 cows (Jose *et al.* 2010) and were lower than that of $31.1 \text{ g MJ}^{-1} \text{ day}^{-1}$ in Xiangzhong Black
10 bulls (Liu *et al.* 2013). Thus the ME utilization efficiency for BW gain for crossbred
11 Simmental bull calves appeared to be lower than that for Chinese indigenous Xiangzhong
12 Black cattle. The less ME utilization efficiency of crossbred Simmental male calves
13 suggested the need for greater energy allowance to ensure a target production performance
14 as compared with Xiangzhong Black cattle.

15 The experimental group that achieved 1-kg DG, T1-LA, T2-LA and T2-HA group was
16 estimated to suffice 113, 125 and 136% of the NE requirement for maintenance and
17 fattening (NEmf) (CFSBC 2004), respectively. The NEmf provision estimates by CFSBC
18 (2004) appeared to be lower than the actual NE intakes of the calves in each feeding group.
19 Liu *et al.* (2013) reported that 13-months-old Xiangzhong Black cattle achieved BW gain
20 of 1 kg day^{-1} and the animal growth performance was well mirrored by the NEmf of
21 CFSBC (2004). This suggests that there is a difference among breeds in the extent to which
22 they meet the NE requirement for BW gain based on the estimation equation of CFSBC
23 (2004). These differences among cattle breeds should be incorporated into the current
24 CFSBC (2004) to predict NE requirements of beef cattle accurately at a given production

1 level.

2

3 **Economic evaluation of feeding AH to calves**

4 We found a significant decrease from T1-LA to T1-HA and a significant increase from
5 T2-CTRL to T2-HA in feed costs ($P < 0.05$) (Table 5), due to a decrease in DMI of C
6 relatively more than an increase of DMI of AH from T1-LA to T1-HA, and an increase in
7 DMI of AH without a decrease in DMI of C from T2-CTRL to T2-LA and T2-HA (Table 3).
8 The feed costs of LA groups in both trails were numerically less than T1-CTRL for T1 and
9 significantly less than T2-HA for T2, respectively. The groups that achieved a 1-kg DG
10 were T1-LA, T2-LA and T1-HA (Figure 2). The feed cost was lower ($P < 0.05$) and the
11 economic benefit was numerically higher for T2-LA than for T2-HA. In order to achieve a
12 1-kg DG, feeding AH at the LA level seemed preferable in terms of economic feasibility in
13 Gansu Province.

14

15 **Conclusions**

16

17 Replacement of C with AH at differing levels did not cause DMI depression associated
18 with increased NDFom intake, but high level AH feeding caused an increase in methane
19 emission. The AH supplementation at a low level increased the DG of growing beef calves.
20 Energy metabolizability did not differ significantly between the low-level AH diet and the
21 high-level AH diet. Considering the importance of economic feasibility, the low-level AH
22 diet gave a relatively high economic benefit. Low-level AH inclusion in the diets of
23 growing beef cattle is preferably practiced in the drylands of Gansu province, China. As the
24 DMI and DG lowered for T2-CTRL than for the other groups in T2, the feeding practice

1 allowing diet with being relatively high proportion of concentrate in the cool season may
2 have to be carefully considered. The MEm requirements of crossbred Simmental calves
3 were calculated to be higher than for other breeds, suggesting that the energy allowance
4 needs to be carefully calculated to ensure the energy requirement of crossbred Simmental.
5 To our knowledge, this is the first report of the determination of energy metabolism using a
6 respiration trial in beef cattle fed domestic basal forages produced in Gansu Province,
7 China. In conjunction with an energy metabolism study, further evaluation of the nitrogen
8 utilization status of growing Chinese Simmental beef calves will be needed to establish a
9 feeding regimen appropriate to the drylands of western China.

10

11 **Acknowledgments**

12

13 We thank the students of the College of Grassland Science, Lanzhou University, China, for
14 supporting the animal experiments and for analyzing feed and fecal samples. This study
15 was supported by the Marginal Region Agriculture Project of Tottori University, the
16 National Key Project of Scientific and Technical Supporting Programs of China
17 (2014CB138706), the National Natural Science Foundation of China (No. 31172249), and
18 the Program for Changjiang Scholars and Innovative Research Team in the University of
19 China (IRT13019).

20

21 **References**

22

23 AFRC (1993) *Energy and Protein requirements of Ruminants. An advisory manual*
24 *prepared by the AFRC Technical Committee on Responses to Nutrients.* CAB

1 International, Wallingford, UK.

2 AOAC (Association of Official Analytical Chemists) (1984) *Official Method of Analysis*,
3 *14th edn.* AOAC, Arlington, VA.

4 Brower E (1965) Report of sub-committee on constants and factors. In: *Energy Metabolism*
5 *of Farm Animals* (Ed. Blaxter KL), EAAP Publ. No. 11, Academic Press, London, 441–
6 443.

7 Chinese State Committee for Development and Reform (2013) Chinese National Plan for
8 developing production of beef and sheep meat 2013–2020. Chinese State Committee for
9 Development and Reform, Beijing, China. (In Chinese.)

10 Defoor PJ, Galyean ML, Salyer GB, Nunnery GA, Parsons CH (2002) Effects of roughage
11 source and concentration on intake and performance by finishing heifers. *J Anim Sci* 80:
12 1395–1404.

13 FAO (2015) FAOSTAT. Statistics Division, FAO, Rome, available from URL:
14 <http://faostat3.fao.org/download/Q/QL/E> [cited 30 November 2015].

15 Freetly HC, Nienaber JA, Brown-Brandl T (2006) Partitioning of energy during lactation of
16 primiparous beef cows. *J Anim Sci* 84: 2157–2162.

17 Guo H, Zhao J, Yand X, Huang B, Zhang M, Wand N (2007) Moisture use efficiency and
18 economic benefit of some crops in the east of Gansu. *Chinese J Soil Sci* 38: 709–712. (In
19 Chinese.)

20 Hales KE, Brown-Brandl TM, Freely HC (2014) Effects of decreased dietary roughage
21 concentration on energy metabolism and nutrient balance in finishing beef cattle. *J Anim*
22 *Sci* 92: 264–271.

23 Han Z, Zheng Q, Abe J, Zhou Z (2008) Pasturage prohibition and pasturage pause and
24 stock raising management in Inner Mongolia China: transaction zone between cropping

- 1 area and nomadic area. *Agric Marketing J of Japan* 17: 80–85. (In Japanese.)
- 2 Henrique DS, Vieira RAM, Malafaia PAM, Mancini MC, Goncalves AL (2005) Estimation
3 of the total efficiency of metabolizable energy utilization for maintenance and growth by
4 cattle in tropical conditions. *R Bras Zootec* 34: 1006–1016.
- 5 Hou FJ, Nan ZB, Xie YZ, Li XL, Lin HL, Ren JZ (2008) Integrated crop-livestock
6 production systems in China. *Rangeland J* 30: 221–231.
- 7 Kennedy PM, Murphy MR (1988) The nutritional implications of differential passage of
8 particles through the ruminant alimentary tract. *Nutr Res Rev* 1: 189–208.
- 9 Ichinohe T, Tamura T, Ueda K, Okubo M, Asahida Y (1994) The particle size distribution
10 of ingested boli, rumen digesta and feces in sheep fed orchardgrass hay harvested at
11 different stage of maturity. *Anim Sci Technol* 65: 701–708.
- 12 Jose VC, Juan CK, Juan GM (2010) Estimation of Metabolizable Energy Requirements for
13 Maintenance and Efficiency of Weight Gain in Bos Taurus (Brown Swiss) and Bos
14 Indicus (Zebu) Cows in Tropical Mexico. *J Animal Vet Adv* 9: 522–528.
- 15 Krause KM, Combs DK, Beauchemin KA (2002) Effects of forage particle size and grain
16 fermentability in mid lactation cows. 1. Milk production and diet digestibility. *J Daily*
17 *Sci* 85: 1936–1946.
- 18 Liu D, Fu D, Qu M, Zhu Y, Li F, Deng X (2013) Energy metabolism and requirement of 12
19 to 13 months old Xiangzhong Black cattle. *Chinese J Anim Nutr* 25: 3013–3019. (In
20 Chinese.)
- 21 MAFF (2013) *Special Study for Governmental Administration: Country Report of 2012 -*
22 *China and Thailand*. Policy Research Institute, Ministry of Agriculture, Forestry and
23 Fisheries, Tokyo, Japan. (In Japanese.)
- 24 McCarthy RD, Klusmeyer TH, Vicini JL, Clark JH, Nelson DR (1989) Effects of source of

1 protein and carbohydrate on ruminal fermentation and passage of nutrients to the small
2 intestine of lactating cows. *J Dairy Sci* 72: 2002–2016.

3 Minson DJ (1980) Nutritional differences between tropical and temperate pastures. In:
4 *Grazing Animals* (Ed Morley FHW), Commonwealth Agricultural Bureaux, Farnham
5 Royal, Slough, 167–182.

6 Ministry of Agriculture of the People's Republic of China (2004) Feeding Standard for
7 Beef Cattle. Ministry of Agriculture of the People's Republic of China, Beijing,
8 available from URL:
9 <http://wenku.baidu.com/view/a112b1a1c77da26925c5b0f1.html> [cited 18 July 2015].
10 (In Chinese.)

11 Munitifering RB (1982) Evaluation of various lignin assays for determining ruminal
12 digestion of roughages by lambs. *J Anim Sci* 55: 432–438.

13 National Bureau of Statistics of China (2014) *China Statistics Yearbook 2013*. China
14 Statistics Press, Beijing, China. (In Chinese.)

15 NARO (National Agricultural and Food Research Organization) (2010) *Standard Tables of*
16 *Feed Composition in Japan (2009)*. Japan Livestock Industry Association, Tokyo. (In
17 Japanese.)

18 Oba M, Allen MS (2003) Effects of corn grain conservation method on feeding behavior
19 and productivity of lactating dairy cows at two dietary starch concentrations. *J Dairy Sci*
20 86: 174–183.

21 Qin G, Zou C, Pang C, Yang B, Liang X, Liu J, Xia Z, Wen Q, Yan T (2011) Evaluation of
22 fasting metabolism of growing water buffalo (*Bubalus, Bubalis*). *Anim Sci J* 82: 735–
23 740.

24 Rong Y, Han J, Hu Y, Li G (2004) The review of grazing alfalfa varieties. *Grassl China* 26:

- 1 49-56. (In Chinese.)
- 2 Tedeschi LO, Boin C, Fox DG, Leme PR, Alleoni GF, Lanna DPD (2002) Energy
3 requirement for maintenance and growth of Nellore bulls and steers fed high-forage
4 diets. *J Anim Sci* 80: 1671–1682.
- 5 United Nations Environment Programme (1997) *World Atlas of Desertification*. UNEP,
6 Nairobi.
- 7 Van Soest PJ, Robertson JB, Lewis BA (1991) Methods for dietary fiber, neutral detergent
8 fiber, and nonstarch polysaccharides in relation to animal nutrition. *J Dairy Sci* 74:
9 3583–3597.
- 10 Zhu X, Shen Y, Yan S, Zhao Y (1997) The base research of Linze grassland ecological test
11 station. *Pratacultural Sci* 1997 (2): 14–35. (In Chinese.)

Table 1 Gross energy concentration and chemical composition of feed ingredients of experimental diets fed to Simmental beef calves

Item	GE, kJ g ⁻¹ DM	Chemical composition, % DM		
		CP	NDFom	ADL
Trial 1				
Corn stover	16.3	5.2	70.3	5.7
Alfalfa hay	16.7	15.8	52.2	7.5
Concentrate [‡]	17.6	13.1	17.1	2.2
Trial 2				
Corn stover	16.2	4.3	72.0	5.1
Alfalfa hay [†]	16.5	12.0	52.8	8.7
Concentrate [‡]	17.7	12.8	23.4	2.4

ADL, acid detergent lignin; CP, crude protein; DM, dry matter; GE, gross energy; NDFom, ash-free neutral detergent fiber.

Trial 1 was conducted in the warm season (from 6 August to 23 September 2015).

Trial 2 was conducted in the cool season (from 24 September to 3 November 2015).

[†]Mixture of the offered alfalfa hay harvested in July and September 2015 (mixed ratio =50:50) were analyzed.

[‡]Concentrate comprised of 59.0% of corn grain, 31.1% of commercial concentrate (comprising soybean meal, sunflower meal, rape seed meal and cotton seed meal, urea, sodium chloride, vitamin and mineral premix; composition not clear), and 9.9% of wheat bran as fed basis.

Table 2 Feed allowance and chemical composition of experimental diets fed to Simmental beef calves

Item	Trial 1			Trial 2		
	T1-CTRL	T1-LA	T1-HA	T2-CTRL	T2-LA	T2-HA
Feed ingredient [†] , kg DM day ⁻¹						
Corn stover	3.0	3.0	3.0	0.6	0.6	0.6
Alfalfa hay	0.0	0.7	1.4	0.0	1.0	1.9
Concentrate	2.7	2.1	1.5	4.8	4.2	3.6
Chemical composition [‡] , % DM						
CP	8.9	9.3	9.7	11.9	11.8	11.7
NDFom	45.1	48.9	52.5	29.1	33.7	37.5
ADL	3.4	4.0	4.7	2.9	4.3	5.2

ADL, acid detergent lignin; CP, crude protein; DM, dry matter; NDFom, ash-free neutral detergent fiber

Trial 1 and Trial 2: for details, see footnote to Table 1.

T1-CTRL and T2-CTRL, no alfalfa hay feeding (control); T1-LA and T2-LA, low level of alfalfa hay feeding; T1-HA and T2-HA, high level of alfalfa hay feeding.

[†]Calculated by using the equation of AFRC (1993) on the basis of the initial body weight of male calves for Trial 1 (175.8 kg on average) and Trial 2 (218.8 kg on average) to suffice the metabolizable energy requirement for a 1-kg daily gain.

[‡]Values were estimated on the basis of the chemical composition of feed ingredients (Table 1) and the ingredient compositions of the experimental diets.

Table 3 Feed intake, daily gain, feed efficiency, dietary energy intake, and digestibility in Simmental beef calves									
Item	Trial 1				Trial 2				
	T1-CTRL	T1-LA	T1-HA	SEM	T2-CTRL	T2-LA	T2-HA	SEM	
Feed intake									
Corn stover, kg DM day ⁻¹	1.5	1.8	2.0	0.02	0.8	0.8	0.8	0.00	
Alfalfa hay, kg DM day ⁻¹	0.0	0.4	0.9	0.01	0.0	1.2	2.4	0.01	
Concentrate, kg DM day ⁻¹	3.1	2.7	1.8	0.10	3.1	3.3	3.1	0.12	
Total DMI, kg day ⁻¹	4.6	4.9	4.7	0.11	3.8 ^c	5.3 ^b	6.3 ^a	0.12	
Total DMI, % BW	2.4	2.5	2.6	0.10	1.7 ^c	2.2 ^b	2.6 ^a	0.08	
Feed conversion ratio, kg DG day ⁻¹ kg ⁻¹ DMI	0.20 ^a	0.21 ^a	0.13 ^b	0.02	0.18	0.23	0.18	0.02	
Digestibility, %									
DM	64.6	60.8	59.3	5.44	76.7 ^a	66.0 ^b	62.5 ^b	1.53	
CP	58.6	55.5	59.7	7.04	72.9 ^a	62.9 ^b	61.3 ^b	2.36	
NDFom	52.6	50.7	49.2	6.28	60.1 ^a	49.4 ^b	48.2 ^b	2.40	
<p>Trial 1, Trial 2, T1-CTRL, T1-LA, T1-HA, T2-CTRL, T2-LA, and T2-HA: for details, see Tables 1 and 2 or the Materials and methods section of the text.</p> <p>BW, body weight (kg); CP, crude protein; DG, daily gain; DM, dry matter; DMI, dry matter intake; NDFom, ash-free neutral detergent fiber; SEM, standard error of the mean.</p> <p>^{a, b, c} Means with different superscripts within a row for each of Trial 1 and Trial 2 differ significantly ($P \leq 0.05$).</p>									

Table 4 Energy utilization efficiencies and methane emission in Simmental beef calves								
Items	Trial 1				Trial 2			
	T1-CTRL	T1-LA	T1-HA	SEM	T2-CTRL	T2-LA	T2-HA	SEM
Gross energy intake, $\text{kJ kg}^{-0.75} \text{ BW day}^{-1}$	1543.0	1612.4	1592.7	42.1	1157.7 ^c	1482.6 ^b	1767.1 ^a	40.4
Energy digestibility (DE GE^{-1}), %	65.2	60.8	59.2	5.4	77.1 ^a	66.7 ^b	63.5 ^b	1.7
Energy metabolizability (ME GE^{-1}), %	58.6	54.4	51.7	5.5	67.2 ^a	57.9 ^b	55.6 ^b	1.6
Methane production, $\text{L kg}^{-0.75} \text{ BW day}^{-1}$	1.6 ^b	1.8 ^b	2.2 ^a	0.1	1.6 ^b	2.0 ^b	2.5 ^a	0.1

Trial 1, Trial 2, T1-CTRL, T1-LA, T1-HA, T2-CTRL, T2-LA, and T2-HA: for details, see Tables 1 and 2 or the Materials and methods section of the text.

BW, body weight; DE, digestible energy; GE, gross energy; ME, metabolizable energy; SEM, standard error of the mean.

^{a, b} Means with different superscripts within a row for each of Trial 1 and Trial 2 differ significantly ($P \leq 0.05$).

Table 5 Economic evaluation of alfalfa introduction for feeding growing Simmental bull calves in Gansu Province

Item	Trial 1				Trial 2			
	T1-CTRL	T1-LA	T1-HA	SEM	T2-CTRL	T2-LA	T2-HA	SEM
Feed cost, US\$ day ⁻¹ head ⁻¹	1.64 ^a	1.55 ^a	1.28 ^b	0.06	1.70 ^c	2.20 ^b	2.47 ^a	0.07
Profit [†] , US\$ day ⁻¹ head ⁻¹	3.24 ^a	3.57 ^a	2.19 ^b	0.26	2.38 ^b	4.16 ^a	3.97 ^a	0.42
Economic benefit [‡] , US\$ day ⁻¹ head ⁻¹	1.60 ^{ab}	2.03 ^a	0.91 ^b	0.27	0.68	1.96	1.50	0.44

Trial 1, Trial 2, T1-CTRL, T1-LA, T1-HA, T2-CTRL, T2-LA, and T2-HA: for details, see Tables 1 and 2 or the Materials and methods section of the text.

SEM, standard error of the mean.

Values for feed cost were calculated on the basis of the results for feed intake (Table 3) as obtained in the feeding trials.

[†]Calculated by multiplying DG (Figure 2) by the market price of a growing calf.

[‡]Calculated by subtracting the feed cost from the profit[†].

^{a, b} Means with different superscripts within a row for each of Trial 1 and Trial 2 differ significantly ($P \leq 0.05$).

Figure legends

Figure 1 Locations of Gansu Province and Linze Research Station of Lanzhou University, and the eight western provinces prioritized for beef production in China.

Figure 2 Metabolizable energy intake and daily gain of calves. CTRL, control group; LA, low-level alfalfa hay feeding group; HA, high-level alfalfa hay feeding group. ^{a, b}Means of metabolizable energy intake and daily gain with different superscripts within each of Trial 1 and Trial 2 differ significantly among the three feeding groups ($P \leq 0.05$). Empty bars, Metabolizable energy intake ($\text{kJ kg}^{-0.75} \text{ BW day}^{-1}$); solid bars, Daily gain (g day^{-1}).

Figure 3 Linear regression of net energy intake (NEI, $f(x)$) and metabolizable energy intake (MEI, x) of calves in Trial 1 (warm season) and Trial 2 (cool season). The NEI was estimated as $\text{MEI} - \text{HP}$ (heat production). Units are expressed on a metabolic body size basis ($\text{kJ kg}^{-0.75} \text{ BW day}^{-1}$). Trial 1: $f(x) = 1.02 \pm 0.06 x - 663 \pm 60.4$, $R^2 = 0.90$, $n = 30$. Trial 2: $f(x) = 0.77 \pm 0.10 x - 462 \pm 120.3$, $R^2 = 0.71$, $n = 24$. Metabolizable energy for maintenance (ME_m) is the interpolant of x at the point where $f(x)$ is zero.



Figure 1 Locations of Gansu Province and Linze Research Station of Lanzhou University in the eight western provinces prioritized for beef production in China.




Figure 2 Relationship between metabolizable energy intake and daily gain among groups in trial 1 and trial 2.





Figure 3 Relationship between differences of metabolizable energy intake and heat production and metabolic energy intake of each cattle, per metabolic body size.