

Full Length Research Paper

Effects of various medium compositions and wounding treatments on *in vitro* growth and regeneration of bird of paradise (*Strelitzia reginae*)

J. J. North¹, P. A. Ndakidemi^{2*} and C. P. Laubscher

¹Faculty of Applied Sciences, Cape Peninsula University of Technology, P.O. Box 652, Cape Town 8000, South Africa.

²The Nelson Mandela African Institute of Science and Technology, P. O. Box 447-Arusha-Tanzania.

Accepted 16 February, 2012

The aim of this study was to investigate the use of antioxidants, wounding treatments and hormone concentrations in efforts to overcome phenolic oxidation and stimulate axillary bud proliferation. Significant results were achieved with 1-naphthalene acetic acid (NAA) and 6-benzylaminopurine (BAP) concentrations on explants discoloration and callus formation. The antioxidant treatments, activated charcoal (AC) and ascorbic acid (AA) significantly affected explant discoloration, the induction of callus and the length of roots developed. Wounding treatments resulted in a reduction of plant height, an increase in both explants discoloration and callus formation. The most effective treatment in reducing explants discoloration at the media contact point was achieved in interactive effects of higher NAA and BAP concentrations (0.1 mg.l⁻¹ NAA; 3 mg.l⁻¹ BAP and 0.5 mg.l⁻¹ NAA; 5 mg.l⁻¹ BAP) without wounding. Interactions between antioxidants and wounding treatments resulted in the absolute absence of callus induction in all treatments involving ascorbic acid.

Key words: Ascorbic acid, activated charcoal, 6-benzylaminopurine (BAP), callus formation, discoloration, 1-naphthalene acetic acid (NAA).

INTRODUCTION

The bird of paradise (*Strelitzia reginae*) is a plant of significant commercial value (Paiva et al., 2004). However, its commercial exploitation and success is limited by its naturally low rate of multiplication (Ziv and Halevy, 1983). Micropropagation as an advanced propagation and cloning method could overcome the constraints posed by the slow conventional propagation methods, thus, allowing for the large scale production which is needed to exploit its horticultural potential. In most reported investigations (Promtep, 1981; Ziv and Halevy, 1983; Paiva et al., 2004; Kantharaju et al., 2008),

only partial success and a low rate of multiplication were obtained, indicating major problems with growing and multiplying this plant *in vitro*. Furthermore, the successful regeneration from zygotic embryo explants has not been reported. There are no reports on success or attempts made in the stimulation of axillary bud proliferation from embryo-derived plantlets.

Axillary bud proliferation exploits the normal ontogenetic route for branch development by lateral meristems (Gamborg and Phillips, 2002). In *Strelitzia*, there is an absolute absence of branching from axillary buds *in vivo*. This may be a result of a strong apical dominance effect (van de Pol and van Hell, 1988). A method of eradicating apical dominance *in vitro* is required to promote branching and increase the multiplication rate of *Strelitzia*.

Since apical dominance has been proved to be under the control of various growth regulators (Wickson and Thimann, 1958; Woolley and Wareing, 1972; Cline, 1994),

*Corresponding author. E-mail: ndakidemipa@gmail.com.

Abbreviations: MS, Murashige and Skoog (1962); AC, activated charcoal; AA, ascorbic acid; BAP, 6-benzylaminopurine; NAA, 1-naphthalene acetic acid.

the proportions of these substances in the media can be manipulated to break dormancy and produce shoots (Razdan, 1993). The concentration and combination of auxins and cytokinins is a key factor which determines successful plant regeneration (Razdan, 1993). In order to increase axillary bud development in *S. reginae*, the optimal balance between these two groups of growth regulators needs to be determined.

Axillary meristems are generally the source of buds that form when leaders are damaged (Burrows, 1989), thus, indicating the positive effects of meristem wounding and even apical bud removal on stimulating the growth of suppressed axillary buds. This method of eliminating apical dominance introduces branching to increase the multiplication rate. Similar to *in vivo* methods, an *in vitro* wounding method is needed to reduce apical dominance and promote axillary bud development. Significant effects of *in vitro* wounding on shoot and root induction have been reported for various species such as *Pyrus malus* (Korban and Skirvin, 1985; Browning et al., 1987; Welander, 1988) and *Yucca elephantipes* (Mauseth and Halperin, 1975; Bentz et al., 1988).

The failure of tissue culture attempts in the propagation of *S. reginae* is largely due to the oxidative browning of explants (Ziv and Halevy, 1983; Paiva et al., 2004, Kantharaju et al., 2008). The excessive production of polyphenols leads to the browning and eventual death of explants (Ziv and Halevy, 1983; Pan and van Staden, 1998; Zeweldu and Ludder, 1998; Birmeta and Welander, 2004; Diro and van Staden, 2004). Tissue injury stimulates the production of phenols (Dodds and Roberts, 1995). Thus, polyphenolic exudation will be exaggerated in response to the wounding techniques employed in this experiment, thus, making the need for antioxidants in the culture media even more evident. The addition of the antioxidant, activated charcoal (AC), to culture media to adsorb toxic substances is widely reported (Horner et al., 1977; Fridborg et al., 1978; Weatherhead et al., 1979; Theander and Nelson, 1988). However, the adsorption properties of AC are non-selective and capable of adsorbing high concentrations of various growth regulators (Pan and van Staden, 1998). As mentioned, the ratio and concentration of auxins and cytokinins in the media is a key factor in determining successful plant regeneration (Razdan, 1993). Thus, the addition of AC to shoot proliferation media may have adverse effects and inhibit growth and regeneration *in vitro* (Pan and van Staden, 1998); thus, highlighting the need to introduce another antioxidant to promote growth and regeneration. Ascorbic acid (AA) is an antioxidant used to control the oxidation of phenols (Chawla, 2002). A comparative study of these two antioxidants would gain further insight into the adsorption of the growth regulators and identify the most successful antioxidant for use in this stage of culture. The objective of this study was to assess the effects various auxin and cytokinin concentrations, antioxidant and wounding treatments have on shoot

formation, explants height, explants discoloration (the entire explants and at the medium contact point only), callus formation and root length.

MATERIALS AND METHODS

Plant material

Embryo-derived *in vitro* seedlings of *S. reginae* were used in this experiment. Germinated plantlets were subjected to 2 wounding treatments; unwounded explants (control) and explants longitudinally sectioned through the apical meristem.

Culture conditions and media

Explants were transferred to different regeneration media. The basal medium comprised the Murashige and Skoog (1962) (MS) salts supplemented with 100 mg.l⁻¹ myo-inositol, 0.1 mg.l⁻¹ thiamine-HCl, 0.1 mg.l⁻¹ pyridoxine, 2 mg.l⁻¹ glycine and 30 g.l⁻¹ sucrose. Various concentrations of 6-benzylaminopurine (BAP) 0, 2, 3, 5 and 6 mg.l⁻¹ and 1-naphthalene acetic acid (NAA) 0, 0.1 and 0.5 mg.l⁻¹ were added to the media. The antioxidants of 2.5 g.l⁻¹ AC and 0.05 g.l⁻¹ AA were each separately added to the various media. The experiment consisted of ten medium types (Table 1). Ten replicates were used for each treatment. The media was solidified with 7 g.l⁻¹ agar. The pH of the media was adjusted to 5.95 prior to autoclaving at 121°C for 20 min. The unwounded cultures were incubated in a growth room with a 16 h light and 8 h dark cycle at 25±2°C.

Data collection and analysis

Data on number of shoots developed per explant, shoot length, root number and length, degree of plantlet discoloration and callus formation were collected at weekly intervals. Based on visual observations, the degree of plantlet discoloration (entire explant and at the media contact point only) was rated on a scale of 1 to 5 (1 = No discoloration and 5 = Extreme discoloration), modified from the rating scale given by Ziv and Halevy (1983). The degree of callus formation was rated as: 1 = none, 2 = low, 3 = medium and 4 = high. This rating scale was modified from that given by Ziv and Halevy (1983). Data collected were analyzed for statistical significance using unbalanced factorial analysis of variance (ANOVA) where 3 levels of NAA (0, 0.1 and 0.5 mg.l⁻¹) were split in 5 levels of BAP (0, 2, 3, 5, 6 mg.l⁻¹) and two antioxidants (2.5 g.l⁻¹ AC and 0.05 g.l⁻¹ AA) were added to each level of BAP making a total of ten treatments (Table 1). These computations were done with the STATISTICA Software Programme version 2010 (StatSoft Inc., Tulsa, OK, USA). The Fisher least significance difference was used to compare treatment means at P = 0.05 level of significance (Steel and Torrie, 1980).

RESULTS AND DISCUSSION

Effect of various NAA and BAP concentrations, antioxidants and wounding treatments on explant height

The height of explants was not affected by variations of NAA and BAP concentrations. Similarly, the antioxidant treatments did not produce significant results. Although a

Table 1. Concentrations and combinations of auxin and cytokinin supplements and antioxidants tested in axillary bud proliferation of regenerated embryos.

Treatment	Auxin and cytokinin concentration (mg.l ⁻¹)		Antioxidant (g.l ⁻¹)
	NAA	BAP	
1	0	0	2.5 activated charcoal
2	0	0	0.05 ascorbic acid
3	0.1	2	2.5 activated charcoal
4	0.1	2	0.05 ascorbic acid
5	0.1	3	2.5 activated charcoal
6	0.1	3	0.05 ascorbic acid
7	0.5	5	2.5 activated charcoal
8	0.5	5	0.05 ascorbic acid
9	0.5	6	2.5 activated charcoal
10	0.5	6	0.05 ascorbic acid

slight increase in explants height was observed in the presence of AC. Wounding had a significant influence on explants height, with maximum height observed in entire, unwounded explants (Table 2). Wounded explants displayed a 28% reduction in height, compared with the unwounded. These results are in line with those reported by Bhatia et al. (2005), who found that shoot height in tomato was much lower in shoots regenerated from wounded explants compared with those that originated from intact cotyledons. Wounding induces stress in plant tissues and suppresses plant growth (Zhang and Turner, 2008).

Effect of various NAA and BAP concentrations, antioxidants and wounding treatments on discoloration of the explants

The variations in NAA and BAP concentrations did not significantly affect explants discoloration in this experiment. Although a higher degree of discoloration was observed in the control (the treatment free of plant hormones). Antioxidant treatments had a significant ($P \leq 0.05$) effect on reducing explants discoloration (Table 3). The antioxidant, AA, was 20, 18 and 19% more effective in reducing entire explant discoloration than AC in weeks 6, 7 and 8, respectively. AA treatments have been widely reported to have positive effects on reducing the oxidative browning of explants (Wu and du Toit, 2004; Abeyaratne and Lathiff, 2002). A study showing parallel results with this study was achieved in Cavendish banana cv. Formosa (Ko et al., 2009) in which AC was not as effective as AA in reducing the incidence of lethal browning. Ko et al. (2009) suggested that AA may have been absorbed by the plantlets, translocated to leaves and prevented the oxidation of phenolic compounds on the target site. It is conceivable that this may also be the case in this study.

Throughout the experiment, a significantly higher ($P \leq 0.001$) degree of explants discoloration was observed in wounded explants (Table 3). A 37% higher level of discoloration was observed in wounded explants in week 2, which increased to 55% over the duration of the experiment, compared with the unwounded explants. Tissue injury stimulates the production of phenols (Dodds and Roberts, 1995), a defensive mechanism common in plants in response to any type of tissue damage (Pan and van Staden, 1998; Ndakidemi and Dakora, 2003). Thus, the production of polyphenolic compounds is exaggerated in response to wounding (George, 1993; Zeweldu and Ludders, 1998; Strosse et al., 2009). The excessive production of polyphenols results in browning and eventual death of tissues (Pan and van Staden, 1998).

Effect of various NAA and BAP concentrations, antioxidants and wounding treatments on discoloration of explants at the media contact point

In week 2, the various NAA and BAP concentrations significantly ($P \leq 0.01$) affected explant discoloration at the medium contact point (Table 4). The treatment containing the highest level of NAA (0.5 mg.l⁻¹) and BAP (6 mg.l⁻¹) was the most effective in reducing discoloration. Whereas, the highest level of discoloration was observed in the control (the treatment without NAA and BAP). These results are in line with those of Xu et al. (2009) who reported increased levels of NAA and BAP to effectively reduce the discoloration of *Dioscorea opposita* explants.

Results revealed that AC was significantly ($P \leq 0.001$) more effective than AA in reducing explant discoloration at the medium contact point (Table 4). In weeks 2, 3 and 4, the respective reductions of 33, 34 and 31% were observed in AC treatments relative to AA treatments. It is

Table 2. Effect of various NAA and BAP concentrations, antioxidants and wounding treatments on explant height (mm).

Treatment	Time (Weeks)								
	1	2	3	4	5	6	7	8	9
Concentration									
0 mg.l ⁻¹ NAA; 0 mg.l ⁻¹ BAP	13.45±1.26 ^a	16.60±1.69 ^a	18.50±1.85 ^a	18.80±1.87 ^a	19.00±1.91 ^a	19.55±1.99 ^a	19.90±2.09 ^a	20.30±2.24 ^a	20.65±2.35 ^a
0.1 mg.l ⁻¹ NAA; 2 mg.l ⁻¹ BAP	13.70±1.29 ^a	16.55±1.89 ^a	20.40±1.94 ^a	20.95±1.96 ^a	21.20±2.17 ^a	21.80±2.21 ^a	22.25±2.28 ^a	22.95±2.32 ^a	23.25±2.39 ^a
0.1 mg.l ⁻¹ NAA; 3 mg.l ⁻¹ BAP	14.50±1.10 ^a	19.35±1.90 ^a	22.35±2.54 ^a	22.95±2.61 ^a	24.60±2.77 ^a	25.40±2.90 ^a	26.20±2.98 ^a	26.65±2.94 ^a	26.80±2.94 ^a
0.5 mg.l ⁻¹ NAA; 5 mg.l ⁻¹ BAP	16.20±1.26 ^a	19.25±1.40 ^a	21.20±1.74 ^a	21.75±1.59 ^a	21.70±1.46 ^a	22.30±1.49 ^a	22.90±1.64 ^a	23.50±1.78 ^a	23.60±1.83 ^a
0.5 mg.l ⁻¹ NAA; 6 mg.l ⁻¹ BAP	13.65±1.04 ^a	17.60±1.55 ^a	19.65±1.98 ^a	21.25±2.26 ^a	22.25±2.34 ^a	22.80±2.40 ^a	23.05±2.42 ^a	23.40±2.50 ^a	23.45±2.50 ^a
Antioxidants									
AA	14.32±0.73 ^a	16.68±0.93 ^a	18.90±1.06 ^a	19.62±1.14 ^a	20.24±1.19 ^a	20.94±1.23 ^a	21.46±1.27 ^a	22.32±1.34 ^a	22.50±1.38 ^a
AC	14.28±0.78 ^a	19.06±1.17 ^a	21.94±1.43 ^a	22.66±1.44 ^a	23.26±1.51 ^a	23.80±1.57 ^a	24.26±1.63 ^a	24.40±1.65 ^a	24.60±1.67 ^a
Wounding									
Wounded	13.56±0.58 ^a	15.92±0.92 ^b	18.06±1.18 ^b	18.34±1.16 ^b	18.66±1.22 ^b	19.12±1.31 ^b	19.24±1.35 ^b	19.60±1.43 ^b	19.72±1.47 ^b
Unwounded	15.04±0.89 ^a	19.82±1.14 ^a	22.78±1.28 ^a	23.94±1.34 ^a	24.84±1.39 ^a	25.62±1.38 ^a	26.48±1.41 ^a	27.12±1.40 ^a	27.38±1.40 ^a
3-way ANOVA (F-statistic)									
Main effects									
Concentration	0.8 ^{ns}	0.7 ^{ns}	0.6 ^{ns}	0.6 ^{ns}	0.9 ^{ns}	1.0 ^{ns}	1.1 ^{ns}	1.0 ^{ns}	0.9 ^{ns}
Antioxidant	0.0 ^{ns}	2.6 ^{ns}	3.0 ^{ns}	2.9 ^{ns}	2.7 ^{ns}	2.3 ^{ns}	2.0 ^{ns}	1.1 ^{ns}	1.1 ^{ns}
Wounding	1.7 ^{ns}	7.1 ^{**}	7.2 ^{**}	9.8 ^{**}	11.2 ^{**}	11.7 ^{***}	13.6 ^{***}	14.0 ^{***}	14.1 ^{***}
Interactions									
Concentration*Antioxidant	0.4 ^{ns}	1.1 ^{ns}	1.2 ^{ns}	1.3 ^{ns}	1.5 ^{ns}	1.5 ^{ns}	1.2 ^{ns}	1.2 ^{ns}	1.1 ^{ns}
Concentration*Wounding	0.4 ^{ns}	1.1 ^{ns}	0.8 ^{ns}	0.9 ^{ns}	0.7 ^{ns}	0.8 ^{ns}	0.7 ^{ns}	0.7 ^{ns}	0.8 ^{ns}
Antioxidant*Wounding	0.0 ^{ns}	0.2 ^{ns}	0.0 ^{ns}	0.0 ^{ns}	0.0 ^{ns}	0.2 ^{ns}	0.1 ^{ns}	0.2 ^{ns}	0.2 ^{ns}
Conc*Anto*Wounding	0.3 ^{ns}	0.8 ^{ns}	0.7 ^{ns}	0.5 ^{ns}	0.5 ^{ns}	0.8 ^{ns}	0.8 ^{ns}	1.1 ^{ns}	1.1 ^{ns}

** , P≤0.01; *** , P≤0.001; ns, not significantly different. Values (Mean ± MSE, n = 10) followed by dissimilar letters in a column are significantly different by least significant difference test at P=0.05.

at the point of contact between the plantlet and the culture medium that oxidative browning is exaggerated due to oxygen coming into contact with the tissue and the required nutrients (North et al., 2010). Results in this study indicate that AC played a key role, and was more effective than

AA, in reducing plantlet discoloration at this point of contact. The browning and subsequent death of cultured explants is a major problem that usually occurs as a result of phenolic compounds (Ozyigit, 2008). Wounded tissues release these polyphenolic compounds, which diffuse into the

medium (Strosse et al., 2009) and are detrimental to the further development of explants which become necrotic and die (Ziv and Halevy, 1983). The addition of AC to culture media has been widely reported to reduce tissue browning (Chang et al., 2001; Wang et al., 2005; Guo et al., 2007;

Table 3. Effect of various NAA and BAP concentrations, antioxidants and wounding treatments on discoloration of the explants.

Treatment	Time (Weeks)								
	1	2	3	4	5	6	7	8	9
Concentration									
0 mg.l ⁻¹ NAA; 0 mg.l ⁻¹ BAP	1.10±0.10 ^a	1.75±0.22 ^a	1.80±0.24 ^a	1.85±0.24 ^a	2.00±0.26 ^a	2.30±0.30 ^a	2.45±0.35 ^a	2.45±0.35 ^a	2.55±0.37 ^a
0.1 mg.l ⁻¹ NAA; 2 mg.l ⁻¹ BAP	1.00±0.00 ^a	1.45±0.15 ^a	1.60±0.18 ^a	1.60±0.18 ^a	1.75±0.23 ^a	1.85±0.24 ^a	2.10±0.26 ^a	2.25±0.26 ^a	2.30±0.27 ^a
0.1 mg.l ⁻¹ NAA; 3 mg.l ⁻¹ BAP	1.00±0.00 ^a	1.35±0.11 ^a	1.50±0.17 ^a	1.70±0.19 ^a	1.85±0.21 ^a	1.90±0.20 ^a	2.10±0.25 ^a	2.20±0.26 ^a	2.35±0.26 ^a
0.5 mg.l ⁻¹ NAA; 5 mg.l ⁻¹ BAP	1.00±0.00 ^a	1.40±0.11 ^a	1.50±0.15 ^a	1.55±0.18 ^a	1.75±0.25 ^a	1.95±0.32 ^a	1.95±0.32 ^a	2.00±0.32 ^a	2.10±0.33 ^a
0.5 mg.l ⁻¹ NAA; 6 mg.l ⁻¹ BAP	1.00±0.00 ^a	1.45±0.11 ^a	1.65±0.17 ^a	1.65±0.17 ^a	1.65±0.17 ^a	1.80±0.20 ^a	2.15±0.24 ^a	2.75±0.29 ^a	3.00±0.35 ^a
Antioxidants									
AA	1.04±0.04 ^a	1.54±0.08 ^a	1.60±0.10 ^a	1.70±0.11 ^a	1.70±0.11 ^a	1.74±0.12 ^b	1.94±0.14 ^b	2.08±0.15 ^b	2.26±0.18 ^a
AC	1.00±0.00 ^a	1.42±0.10 ^a	1.62±0.13 ^a	1.64±0.14 ^a	1.90±0.17 ^a	2.18±0.19 ^a	2.36±0.21 ^a	2.58±0.22 ^a	2.66±0.22 ^a
Wounding									
Wounded	1.00±0.00 ^a	1.82±0.07 ^a	2.06±0.09 ^a	2.12±0.10 ^a	2.28±0.12 ^a	2.56±0.15 ^a	2.90±0.15 ^a	3.14±0.15 ^a	3.40±0.16 ^a
Unwounded	1.04±0.04 ^a	1.14±0.09 ^b	1.16±0.10 ^b	1.22±0.11 ^b	1.32±0.12 ^b	1.36±0.13 ^b	1.40±0.14 ^b	1.52±0.15 ^b	1.52±0.15 ^b
3-way ANOVA (F-statistic)									
Main effects									
Concentration	1.0 ^{ns}	1.6 ^{ns}	0.7 ^{ns}	0.5 ^{ns}	0.5 ^{ns}	0.9 ^{ns}	0.6 ^{ns}	1.5 ^{ns}	2.0 ^{ns}
Antioxidant	1.0 ^{ns}	1.1 ^{ns}	0.0 ^{ns}	0.2 ^{ns}	1.3 ^{ns}	5.7*	4.0*	5.7*	3.3
Wounding	1.0 ^{ns}	36.7***	44.5***	37.5***	30.7***	42.1***	51.1***	59.4***	73.9***
Interactions									
Concentration*Antioxidant	1.0 ^{ns}	0.3 ^{ns}	0.8 ^{ns}	1.8 ^{ns}	1.1 ^{ns}	1.5 ^{ns}	0.4 ^{ns}	0.6 ^{ns}	0.2 ^{ns}
Concentration*Wounding	1.0 ^{ns}	1.1 ^{ns}	2.1 ^{ns}	1.7 ^{ns}	1.7 ^{ns}	1.7 ^{ns}	0.3 ^{ns}	0.3 ^{ns}	0.3 ^{ns}
Antioxidant*Wounding	1.0 ^{ns}	4.6*	0.2 ^{ns}	0.0 ^{ns}	0.1 ^{ns}	1.2 ^{ns}	4.0*	1.5 ^{ns}	0.5 ^{ns}
Conc*Anto*Wounding	1.0 ^{ns}	0.4 ^{ns}	1.4 ^{ns}	0.9 ^{ns}	1.4 ^{ns}	1.7 ^{ns}	0.4 ^{ns}	0.8 ^{ns}	0.9 ^{ns}

*, P≤0.05; ***, P≤0.001; ns, not significantly different. Values (Mean ± MSE, n = 10) followed by dissimilar letters in a column are significantly different by Least significant difference test at P=0.05. Rating scale used is 1 to 5 (1, No discoloration; 5, extreme discoloration).

North et al., 2010). The promotary effects of AC may be attributed mainly to its irreversible adsorption of inhibitory compounds and decrease phenolic oxidation (Thomas, 2008). Phenols leached from the tissues may readily be absorbed

by the AC, reducing the discoloration more effectively in those tissues coming into direct contact with the AC supplemented media.

Wounded explants displayed a significantly higher level of discoloration at the medium contact

point throughout the duration of the experiment. In week 2, a 15% discoloration was observed which progressively increased to 57% in the final week. As mentioned, wounding stimulates the production of phenols (George, 1993; Dodds and

Table 4. Effect of various NAA and BAP concentrations, antioxidants and wounding treatments on discoloration of explants at the media contact point.

Treatment	Time (Weeks)								
	1	2	3	4	5	6	7	8	9
Concentration									
0 mg.l ⁻¹ NAA; 0 mg l ⁻¹ BAP	1.30±0.22 ^a	1.75±0.22 ^a	1.75±0.27 ^a	1.75±0.27 ^a	1.75±0.27 ^a	1.95±0.28 ^a	2.35±0.36 ^a	2.40±0.36 ^a	2.45±0.37 ^a
0.1 mg.l ⁻¹ NAA; 2 mg.l ⁻¹ BAP	1.00±0.00 ^a	1.20±0.09 ^b	1.30±0.15 ^a	1.35±0.15 ^a	1.70±0.23 ^a	1.90±0.24 ^a	1.95±0.23 ^a	2.05±0.23 ^a	2.25±0.28 ^a
0.1 mg.l ⁻¹ NAA; 3 mg.l ⁻¹ BAP	1.00±0.00 ^a	1.40±0.15 ^b	1.40±0.15 ^a	1.45±0.17 ^a	1.50±0.18 ^a	1.70±0.21 ^a	2.05±0.25 ^a	2.05±0.25 ^a	2.30±0.32 ^a
0.5 mg.l ⁻¹ NAA; 5 mg.l ⁻¹ BAP	1.00±0.00 ^a	1.25±0.12 ^b	1.55±0.18 ^a	1.55±0.17 ^a	1.95±0.29 ^a	2.15±0.32 ^a	2.20±0.32 ^a	2.30±0.32 ^a	2.30±0.32 ^a
0.5 mg.l ⁻¹ NAA; 6 mg.l ⁻¹ BAP	1.00±0.00 ^a	1.15±0.08 ^b	1.30±0.13 ^a	1.40±0.13 ^a	1.55±0.17 ^a	1.80±0.25 ^a	2.05±0.29 ^a	2.35±0.30 ^a	2.70±0.37 ^a
Antioxidants									
AA	1.04±0.04 ^a	1.62±0.09 ^a	1.76±0.13 ^a	1.78±0.13 ^a	1.84±0.13 ^a	1.96±0.13 ^a	2.10±0.14 ^a	2.22±0.14 ^a	2.44±0.19 ^a
AC	1.08±0.08 ^a	1.08±0.08 ^b	1.16±0.09 ^b	1.22±0.09 ^b	1.54±0.16 ^a	1.84±0.19 ^a	2.14±0.22 ^a	2.24±0.22 ^a	2.36±0.23 ^a
Wounding									
Wounded	1.00±0.00 ^a	1.46±0.09 ^a	1.70±0.11 ^a	1.76±0.11 ^a	2.12±0.15 ^a	2.48±0.16 ^a	2.84±0.17 ^a	3.02±0.17 ^a	3.36±0.18 ^a
Unwounded	1.12±0.09 ^a	1.24±0.10 ^b	1.22±0.12 ^b	1.24±0.12 ^b	1.26±0.12 ^b	1.32±0.12 ^b	1.40±0.12 ^b	1.44±0.13 ^b	1.44±0.13 ^b
3-way ANOVA (F-statistic)									
Main effects									
Concentration	1.8 ^{ns}	4.5 ^{**}	1.5 ^{ns}	1.0 ^{ns}	0.8 ^{ns}	0.6 ^{ns}	0.4 ^{ns}	0.5 ^{ns}	0.5 ^{ns}
Antioxidant	0.2 ^{ns}	28.0 ^{***}	18.6 ^{***}	15.7 ^{***}	2.7 ^{ns}	0.4 ^{ns}	0.0 ^{ns}	0.0 ^{ns}	0.1 ^{ns}
Wounding	1.8 ^{ns}	4.7 [*]	11.9 ^{***}	13.5 ^{***}	22.4 ^{***}	33.6 ^{***}	40.8 ^{***}	53.1 ^{***}	69.0 ^{***}
Interactions									
Concentration*Antioxidant	0.2 ^{ns}	0.8 ^{ns}	0.7 ^{ns}	0.6 ^{ns}	1.3 ^{ns}	1.1 ^{ns}	0.7 ^{ns}	0.6 ^{ns}	0.9 ^{ns}
Concentration*Wounding	1.8 ^{ns}	3.6 ^{**}	1.4 ^{ns}	1.3 ^{ns}	2.1 ^{ns}	1.7 ^{ns}	0.5 ^{ns}	0.5 ^{ns}	0.9 ^{ns}
Antioxidant*Wounding	0.2 ^{ns}	13.9 ^{***}	11.9 ^{***}	11.5 ^{**}	1.0 ^{ns}	0.0 ^{ns}	1.5 ^{ns}	2.5 ^{ns}	1.1 ^{ns}
Conc*Anto*Wounding	0.2 ^{ns}	0.5 ^{ns}	0.3 ^{ns}	0.4 ^{ns}	1.2 ^{ns}	0.8 ^{ns}	0.2 ^{ns}	0.1 ^{ns}	0.3 ^{ns}

P*≤0.01; *P*≤0.001; ns: not significantly different. Values (Mean ± MSE, n = 10) followed by dissimilar letters in a column are significantly different by least significant difference test at *P*=0.05. Rating scale used is 1 to 5 (1, No discoloration; 5, extreme discoloration).

Roberts, 1995; Zeweldu and Ludders, 1998; Strosse et al., 2009). Phenolic interactions expressed as oxidative browning of the explants

can lead to the death of the plant material (Taji and Williams, 1996). Oxygen free radicals, generated by wounding (Salin and Bridges, 1981;

Thompson et al., 1987), can also lead to the oxidative browning of explants. Reducing contact with oxygen reduces the rate of oxidation of

Table 5. Effect of various NAA and BAP concentrations, antioxidants and wounding treatments on callus development.

Treatment	Time (Weeks)						
	3	4	5	6	7	8	9
Concentration							
0 mg.l ⁻¹ NAA; 0 mg.l ⁻¹ BAP	0.00±0.00 ^b	0.00±0.00 ^b	0.00±0.00 ^b	0.05±0.05 ^a	0.10±0.10 ^a	0.15±0.15 ^a	0.15±0.15 ^a
0.1 mg.l ⁻¹ NAA; 2 mg.l ⁻¹ BAP	0.05±0.05 ^b	0.05±0.05 ^b	0.05±0.05 ^b	0.15±0.11 ^a	0.15±0.11 ^a	0.10±0.10 ^a	0.10±0.10 ^a
0.1 mg.l ⁻¹ NAA; 3 mg.l ⁻¹ BAP	0.10±0.07 ^b	0.10±0.07 ^b	0.15±0.11 ^{ab}	0.25±0.18 ^a	0.30±0.18 ^a	0.30±0.18 ^a	0.35±0.21 ^a
0.5 mg.l ⁻¹ NAA; 5 mg.l ⁻¹ BAP	0.30±0.11 ^a	0.30±0.11 ^a	0.30±0.11 ^a	0.40±0.15 ^a	0.60±0.23 ^a	0.60±0.23 ^a	0.60±0.23 ^a
0.5 mg.l ⁻¹ NAA; 6 mg.l ⁻¹ BAP	0.15±0.08 ^{ab}	0.15±0.08 ^{ab}	0.15±0.08 ^{ab}	0.20±0.12 ^a	0.30±0.18 ^a	0.30±0.18 ^a	0.30±0.18 ^a
Antioxidants							
AA	0.18±0.05 ^a	0.18±0.05 ^a	0.20±0.06 ^a	0.34±0.10 ^a	0.46±0.13 ^a	0.46±0.13 ^a	0.48±0.14 ^a
AC	0.06±0.03 ^b	0.06±0.03 ^b	0.06±0.03 ^b	0.08±0.05 ^b	0.12±0.07 ^b	0.12±0.07 ^b	0.12±0.07 ^b
Wounding							
Wounded	0.24±0.06 ^a	0.24±0.06 ^a	0.26±0.07 ^a	0.42±0.11 ^a	0.58±0.14 ^a	0.58±0.15 ^a	0.60±0.15 ^a
Unwounded	0.00±0.00 ^b	0.00±0.00 ^b	0.00±0.00 ^b	0.00±0.00 ^b	0.00±0.00 ^b	0.00±0.00 ^b	0.00±0.00 ^b
3-way ANOVA (F-statistic)							
Main effects							
Concentration	3.79**	3.8**	2.8*	1.3 ^{ns}	1.8 ^{ns}	1.6 ^{ns}	1.6 ^{ns}
Antioxidant	5.14*	5.1*	5.2*	6.6*	6.9*	6.1*	6.5*
Wounding	20.57***	20.6***	17.8***	17.3***	20.0***	17.9***	18.0***
Interactions							
Concentration*Antioxidant	1.21 ^{ns}	1.2 ^{ns}	1.2 ^{ns}	0.8 ^{ns}	0.8 ^{ns}	0.7 ^{ns}	0.8 ^{ns}
Concentration*Wounding	3.79 ^{ns}	3.8 ^{ns}	2.8 ^{ns}	1.3 ^{ns}	1.8 ^{ns}	1.6 ^{ns}	1.5 ^{ns}
Antioxidant*Wounding	5.14*	5.1*	5.2*	6.6*	6.9*	6.1*	6.5*
Conc*Anto*Wounding	1.21 ^{ns}	1.2 ^{ns}	1.2 ^{ns}	0.8 ^{ns}	0.8 ^{ns}	0.7 ^{ns}	0.8 ^{ns}

*, P≤0.05; **, P≤0.01; ***, P≤0.001; ns, not significantly different. Values (Mean ± MSE, n = 10) followed by dissimilar letters in a column are significantly different by least significant difference test at P=0.05. The degree of callus formation is rated as: 1, none; 2, low; 3, medium; 4, high.

phenols at the wounded site (Elmore et al., 1990). It is at the media contact point that an adequate supply of oxygen comes into contact with the plant material, resulting in an increased level of oxidative browning of these tissues.

Effect of various NAA and BAP concentrations, antioxidants and wounding treatments on callus development

Callus formation from the explants was induced in

NAA and BAP treatments. The highest level of callus was formed in 0.5 mg.l⁻¹ NAA and 5 mg.l⁻¹ BAP (Table 5). This significant increase was opposed to the control (the treatment without NAA and BAP), in which no callus was formed. Reports

Table 6. Effect of various NAA and BAP concentrations, antioxidants and wounding treatments on the length of roots developed (mm).

Treatment	Root length (mm)
Concentration	
0 mg.l ⁻¹ NAA; 0 mg.l ⁻¹ BAP	4.90±3.35 ^a
0.1 mg.l ⁻¹ NAA; 2 mg.l ⁻¹ BAP	6.55±3.52 ^a
0.1 mg.l ⁻¹ NAA; 3 mg.l ⁻¹ BAP	2.95±2.39 ^a
0.5 mg.l ⁻¹ NAA; 5 mg.l ⁻¹ BAP	9.10±5.15 ^a
0.5 mg.l ⁻¹ NAA; 6 mg.l ⁻¹ BAP	2.80±1.99 ^a
Antioxidants	
AA	1.98±1.15 ^b
AC	8.54±2.78 ^a
Wounding	
Wounded	3.48±1.59 ^a
Unwounded	7.04±2.62 ^a
3-way ANOVA (F-statistic)	
Main effects	0.59 ^{ns}
Concentration	4.54 [*]
Antioxidant	1.34 ^{ns}
Wounding	
Interactions	0.23 ^{ns}
Concentration*Antioxidant	0.46 ^{ns}
Concentration*Wounding	1.19 ^{ns}
Antioxidant*Wounding	1.55 ^{ns}
Conc*Anto*Wounding	

*, P≤0.05; ns, not significantly different. Values (Mean ± MSE, n = 10) followed by dissimilar letters in a column are significantly different by least significant difference test at P = 0.05.

of NAA and BAP combinations supporting the development of callus have been well documented for several plant species (Koroch et al., 2002; Ray et al., 2011; Nurazah et al., 2009). Results presented here are in agreement with these studies, where it was observed that low concentrations of NAA in combination with increased rates of BAP resulted in the induction of callus. Similar to our study, Ray et al. (2011) reported that explants cultured on medium without hormones did not produce callus.

A significantly (P≤0.05) higher level of callus formation was observed in AA treatments than in AC treatments (Table 5). In week 1, callus formation was 66% greater in AA treatments, which gradually increased to 75% in the final week of the experiment, as compared with AC treatments. Several studies report the presence of AC to significantly reduce the formation of callus. The addition of AC prevented callus induction in sorghum and cotton (Zhang et al., 2000; Nguyen, et al., 2007) and reduced callus induction in black wattle and *Oxalis triangularis* (Teng and Ngai, 1999; Quoirin et al., 2001). There was

no callus formation in unwounded explants. Whereas, wounding increased callus formation in wounded treatments increased from 0.24 in week 3 to 0.60 in week 9, resulting in a 60% increase (Table 5). The injury that explants experience in response to these wounding techniques may also influence the morphogenesis response in a way similar to that of plants in the natural environment, where wounding often stimulates callus formation (George, 1993).

Effect of various NAA and BAP concentrations, antioxidants and wounding treatments on the length of roots developed

Root length was not significantly affected by NAA and BAP concentrations. Similarly, wounding treatments did not significantly affect root length, although a slight increase in length was observed in unwounded explants than in the wounded. In AC treatments, roots developed were 77% longer than roots developed in AA treatments (Table 6). The positive influence of AC on rooting is




Figure 1. Interactive effects of antioxidants and wounding on entire explant discoloration in (A) week 2 and (B) week 7. Rating scale used is 1 to 5 (1, No discoloration; 5, extreme discoloration). Wounding 1= Wounded; Wounding 2= unwounded.

widely reported (Makunga et al., 2006; Mulwa and Bhalla, 2006; Yan et al., 2006; Agarwal and Kanwar, 2007; Xiao et al. 2007; Makunga and van Staden, 2008; Firoozabady et al., 2006; Feyissa et al., 2005; Loc et al., 2005). Enhanced root growth may be due to the ability of AC to adsorb the polyphenols produced through chemical processes within the media, which may act as growth inhibitors (Madhusudhanan and Rahiman, 2000). AC may also enhance rooting by eliminating light, providing a favourable physical environment to the rhizosphere (Gantait et al., 2009).

Interactive effects of various NAA and BAP concentrations, antioxidants and wounding treatments

The results represent the significant ($P \leq 0.05$) interactive effects of antioxidant and wounding treatments on discoloration of the entire explants in week 2 (Figure 1A) and week 7 (Figure 1B). AC effectively reduced the level of discoloration in both wounded and unwounded treatments in both weeks. The highest level of explant discoloration was observed in AA treatments, either with




Figure 2. Interactive effects of wounding and concentration on explant discoloration at the medium contact point (EDMCP) in week 2, Rating scale used is 1 to 5 (1, No discoloration; 5, extreme discoloration). Wounding 1= Wounded; Wounding 2= unwounded.

wounding (week 2) or without wounding (week 7). In a study on *Ensete ventricosum*, which is related to *S. reginae* (Strosse et al., 2009), similar results were found. The addition of AA did not effectively reduce polyphenol exudation. However, the most effective treatment was AC, which prevented polyphenol exudation in wounded explants (Birmeta and Welander, 2004). The addition of AC to culture media is a recognized practice and its influence may be attributed to the adsorption of inhibitory substances in the medium (Horner et al., 1977; Fridborg et al., 1978; Weatherhead et al., 1979; Theander and Nelson, 1988) and a drastic decrease in the phenolic oxidation of tissues (Carlberg et al., 1983; Liu, 1993; Teixeira et al., 1994). AC has a very fine network of pores with large inner surface area on which many substances can be adsorbed (Thomas, 2008).

The interactive effects of wounding treatments and NAA and BAP concentrations on explants discoloration at the medium contact point produced significant results in weeks 2, (Figure 2). In this week, the highest level of

discoloration was observed in 0.1 mg.l⁻¹ NAA and 2 mg.l⁻¹ BAP, in wounded and unwounded treatments. Whereas the lowest levels were observed in higher NAA and BAP concentrations (0.1 mg.l⁻¹ NAA; 3 mg.l⁻¹ BAP and 0.5 mg.l⁻¹ NAA; 5 mg.l⁻¹ BAP), in both unwounded and at a concentration of 0.5mg.l⁻¹ NAA and 5mg.l⁻¹ BAP in wounded explants. Similarly, in a study on *D. opposita* increased NAA and BAP concentrations effectively and reduced the discoloration of explants (Xu et al., 2009). In *Sorghum bicolor*, brown pigments completely inhibited shoot growth (Baskaran and Jayabalan, 2005). The addition of BAP to these cultures enabled the further growth of shoots. In a study on *Gossypium arboreum*, Smith et al. (1977) reported that both NAA and BAP alone did not support good growth or survival of explants. Various reports have revealed a strong synergistic effect in BAP and NAA interactions in banana (Novak et al., 1989; Okole and Schulz, 1996; Cote et al., 2000; Khalil et al., 2002; Srangsam and Kanchanapoom, 2007). The addition of these two plant growth regulators at an




Figure 3. Interactive effects of antioxidants and wounding on explant discoloration at the medium contact point (EDMCP) in week 2. Rating scale used is 1 to 5 (1, No discoloration; 5, extreme discoloration). Wounding 1= Wounded; Wounding 2= unwounded.

increased concentration may reduce discoloration by enabling the establishment and further growth of plantlets

The interactive effects of antioxidants and wounding treatments gave rise to significant results on explants discoloration at the medium contact point in week 2 (Figure 3).

Both the lowest and highest levels of discoloration were observed in AA treatments. In wounded explants, AA was did not effectively reduce discoloration and resulted in the highest level. Whereas in unwounded explants, AA was the most effective antioxidant, resulting in the lowest level of discoloration, in wounding treatments AC significantly reduced explants discoloration at the medium contact point. As mentioned previously, wounded tissues produce an excess of polyphenolic compounds (George, 1993; Zeweldu and Ludders, 1998; Strosse et al., 2009), which result in an increased level of discoloration due to oxidative browning of explants. As mentioned previously, AC is most effective in reducing discoloration in wounded explants, by adsorbing phenols exuded. In complete, intact plantlets it is possible that AA may reduce discoloration by absorbing AA and prevent the oxidation of phenols on the target site (Ko et al., 2009).

The results in Figures 4 and 5 show the significant ($P \leq 0.05$) interactive effects of antioxidants and wounding on the formation of callus. The presence of AC completely inhibited the development of callus in both wounded and unwounded explants. An increased callus

formation was observed in AA treatments, the highest level formed in wounded explants. This trend was observed throughout the duration of the experiment.

Although majority of reports confirm the positive role of AC in promoting the growth and development of plant tissues, AC inducing negative results is also reported in some cases (Thomas, 2008).

The difficulty in using AC is that its characteristic of high adsorptive power is non-selective. In addition to adsorbing inhibitory phenols, it is able to absorb high concentrations of growth regulators required by plant tissues (Fridborg et al., 1978; Ebert and Taylor, 1990; Nissen and Sutter, 1990; Ebert et al., 1993; Pan and van Staden, 1998; Thomas, 2008). As reported earlier, callus formation was induced by increased NAA and BAP concentrations, as opposed to no callus formation in the control (the treatment without NAA and BAP). Thus, it could be presumed that AC adsorbed the growth regulators required to induce callus formation.

In conclusion, although plant regeneration via axillary bud proliferation was not significantly increased, insight into the effects of NAA and BAP concentrations, antioxidants and wounding techniques on plant height, reducing explants discoloration, callus formation and root length was achieved. Future studies would be aimed at increasing the rate of axillary bud development and shoot regeneration from callus in efforts to improve the clone propagation in *S. reginae*.




Figure 4. Interactive effects of antioxidants and wounding on callus formation in (A) WK3, (B) week 4, (C) week 5, (D) week 6. The degree of callus formation is rated as: 1, None; 2, low; 3, medium; 4, high. Wounding 1= Wounded; Wounding 2= unwounded.




Figure 4. Contd.

ACKNOWLEDGEMENTS

The financial assistance of the National Research

Foundation (NRF) towards this research is hereby acknowledged. Opinions expressed and conclusions arrived at, are those of the author and are not necessarily




Figure 5, Interactive effects of antioxidants and wounding on callus formation in (A) WK7, (B) week 8 and (C) week 9. The degree of callus formation is rated as: 1, None; 2, low; 3, medium; 4, high. Wounding 1= Wounded; Wounding 2= unwoun

to be attributed to the NRF. A further thank you to Dr. Joachim Makoi for his technical assistance.

REFERENCES

- Abeyaratne WM, Lathiff MA (2002). In vitro propagation of Rathambala (*Musa AAA*) and the occurrence of phenotypic variations in the pseudostem. *Ann. Sri Lanka Department Agric.*, 4: 191-197.
- Agarwal S, Kanwar K (2007). Comparison of genetic transformation in *Morus alba* L. via different regeneration systems. *Plant Cell Rep.*, 26: 177-185.
- Bentz SE, Parliman BJ, Talbott HJ, Ackerman WL (1988). Factors affecting in vitro propagation of *Yucca glauca*. *Plant Cell Tissue Org. Cult.*, 14: 111-120.
- Bhatia P, Ashwath N, Midmore DJ (2005). Effects of genotype, explant orientation and wounding on shoot regeneration in tomato. *In Vitro Cell Dev. Biol.-Plant*, 41: 457-464.
- Birmeta G, Welander M (2004). Efficient micropropagation of *Ensete ventricosum* applying meristem wounding: a three-step protocol. *Plant Cell Rep.*, 23: 277-283.
- Browning G, Ognjanov V, Passey AJ, James DJ (1987). Multiple shoot and root regeneration from pear embryo cotyledon explant in vitro. *J. Hort. Sci.*, 62: 305-311.
- Burrows GE (1989). Developmental anatomy of axillary meristems of *Araucaria cunninghamii* released from apical dominance following shoot apex decapitation in vitro and in vivo. *Botanical Gazette*, 150(4): 369-377.
- Carlberg I, Glimelius K, Eriksson T (1983). Improved culture ability of potato protoplasts by use of activated charcoal. *Plant Cell Rep.*, 2: 223-225.
- Chang SH, Ho CK, Chen ZZ, Tsay JY (2001). Micropropagation of *Taxus mairei* from mature trees. *Plant Cell Rep.*, 20: 496-502.
- Chawla HS (2002). Introduction to Plant Biotechnology 2nd rev. ed. Science Publishers Inc., Enfield, New Hampshire.
- Cline MG (1994). The role of hormones in apical dominance. New approaches to an old problem in plant development. *Physiol. Plant*, 90: 230-237.
- Cote FX, Folliot M, Domergue R, Dubois C (2000). Field performance of embryogenic suspension-derived banana plants (*Musa AAA*, cv. Grande naine). *Euphytica*, 11: 245-251.
- Diro M, van Staden J (2004). Germination of zygotic embryos and in vitro growth of seedlings of wild types of *Ensete ventricosum*. *S. Afr. J. Bot.*, 70(4): 635-639.
- Dodds JH, Roberts LW (1995). *Experiments in Plant Tissue Culture*. 3rd rev ed. Cambridge: Cambridge University Press.
- Ebert A, Taylor F, Blake J (1993). Changes of 6-benzylaminopurine and 2,4 dichlorophenoxyacetic acid concentrations in plant tissue culture media in the presence of activated charcoal. *Plant Cell Tissue Org. Cult.*, 33: 157-163.
- Ebert A, Taylor HF (1990). Assessment of the changes of 2,4 dichlorophenoxyacetic acid concentrations in plant tissue culture media in presence of activated charcoal. *Plant Cell Tissue Org. Cult.*, 20: 165-172.
- Elmore HW, Samples B, Sharma S, Harrison M (1990). Influence of cultural and physiochemical factors on ascorbate stability in plant tissue culture media. *Plant Cell Tissue Org. Cult.*, 20: 131-135.
- Feyissa T, Welander M, Negash L (2005). Micropropagation of *Hagenia abyssinica*: a multipurpose tree. *Plant Cell Tissue Org. Cult.*, 80: 119-127.
- Firoozabady E, Heckert M, Gutterson N (2006). Transformation and regeneration of pineapple. *Plant Cell Tissue Org. Cult.*, 84: 1-16.
- Fridborg G, Pedersen M, Landstorm LE, Eriksson T (1978). The effect of activated charcoal on tissue cultures adsorption of metabolites inhibiting morphogenesis. *Physiol. Plant*, 43: 104-106.
- Gamborg OL, Phillips GC (1995). *Plant Cell, Tissue and Organ Culture: Fundamental Methods*. Springer, Verlag Berlin Heidelberg, p. 22.
- Gantait S, Mandal N, Das P (2009). Impact of auxins and activated charcoal on in vitro rooting of *Dendrobium chrysotoxum* Lindl. cv. Golden Boy. *J. Trop. Agric.*, 47(1-2): 84-86.
- George EF (1993). Factors affecting growth and morphogenesis. In: George EF (ed). *Plant propagation by tissue culture*. Exegetics Ltd., London, 231-271.
- Guo Y, Bai J, Zhang Z (2007). Plant regeneration from embryogenic suspension-derived protoplasts of ginger (*Zingiber officinale* Rosc.). *Plant Cell Tissue Org. Cult.*, 89: 151-157.
- Horner M, McComb JA, McComb AJ, Street HE (1977). Ethylene production and plantlet formation by *Nicotiana* anthers cultured in the presence and absence of charcoal. *J. Exp. Bot.*, 28: 1366-1372.
- Kantharaju M, Kulkarni BS, Reddy BS, Hedge RV, Patil BC, Adiga JD (2008). Effect of antioxidant on in vitro establishment of *Strelitzia reginae* through shoot tip explants. *J. Agric. Sci.*, 21(2): 324-325.
- Khalil SM, Cheah KT, Perez EA, Gaskill DA, Hu JS (2002). Regeneration of banana (*Musa spp.* AAB cv. Dwarf Brazilian) via secondary somatic embryogenesis. *Plant Cell Rep.*, 20: 1128-1134.
- Ko WH, Su CC, Chen CL, Chao CP (2009). Control of lethal browning of tissue culture plantlets of Cavendish banana cv. Formosa with ascorbic acid. *Plant Cell Tissue Org. Cult.*, 96: 137-141.
- Korban SS, Skirvin RM (1985). In vitro shoot regeneration from an intact and a sectioned embryo-axis of apple seeds. *Plant Sci.*, 39: 61-66.
- Koroch A, Juliani HR, Kapteyn J, Simon JE (2002). In vitro regeneration of *Echinacea purpurea* from leaf explants. *Plant Cell Tissue Org. Cult.*, 69: 79-83.
- Liu MSC (1993). Plant regeneration in cell suspension culture of sugarcane as affected by activated charcoal, medium composition and tissue culture. *Taiwan Sugar*. May-June, p. 18-25.
- Loc NH, Duc DT, Kwon TH, Yang MS (2005). Micropropagation of zedoary (*Curcuma zedoaria* Roscoe)-a valuable medicinal plant. *Plant Cell Tissue Org. Cult.*, 81: 119-122.
- Madhusudhanan K, Rahiman BA (2000). The effect of activated charcoal supplemented media to browning of in vitro cultures of *Piper* species. *Biol. Plant*, 43(2): 297-299.
- Makunga NP, Jager AK, van Staden J (2006). Improved in vitro rooting and hyperhydricity in regenerating tissues of *Thapsia garganica* L. *Plant Cell Tissue Org. Cult.*, 86: 77-86.
- Makunga NP, van Staden J (2008). An efficient system for the production of clonal plantlets of the medicinally important aromatic plant: *Salvia Africana-lutea* L. *Plant Cell Tissue Org. Cult.*, 92: 63-72.
- Mauseth JD, Halperin W (1975). Hormonal control of organogenesis in *Opuntia polycantha* (Cactaceae). *Amer. J. Bot.*, 62: 869-877.
- Mulwa RMS, Bhalla PL (2006). In vitro plant regeneration from immature cotyledon explants of macadamia (*Macadamia tetraphylla* L. Johnson). *Plant Cell Rep.*, 15: 473-497.
- Ndakidemi PA, Dakora FD (2003). Legume seed flavonoids and nitrogenous metabolites as signals and protectants in early seedling development. *Functional Plant Biol.*, 30: 729-745.
- Nguyen TV, Thu TT, Claeys M, Angenon G (2007). Agrobacterium mediated transformation of sorghum (*Sorghum bicolor* (L.) Moench) using an improved in vitro regeneration system. *Plant Cell Tissue Org. Cult.*, 91(2): 155-164.
- Nissen SJ, Sutter EG (1990). Stability of IAA and IBA in nutrient medium to several tissue culture procedures. *Hort. Sci.*, 25: 800-802.
- North JJ, Ndakidemi PA, Laubscher CP (2010). The potential of developing an in vitro method for propagating Strelitziaceae. *Afr. J. Biotechnol.*, 9(45): 7583-7588.
- Novak FJ, Afza R, Van Duren M, Perea-Dallos M, Cronger BV, Xiaolang T (1989). Somatic embryogenesis and plant regeneration in suspension cultures of dessert (AA and AAA) and cooking (ABB) bananas (*Musa spp.*) *Biotechnol.*, 7: 154-159.
- Nurazah Z, Radzali M, Syahida A, Maziah M (2009). Effects of plant growth regulators on callus induction from *Cananga odorata* flower petal explant. *Afr. J. Biotechnol.*, 8(12): 2740-2743.
- Okole BN, Schulz FA (1996). Micro-cross sections of banana and plantains (*Musa spp.*): morphogenesis and regeneration of callus and shoot buds. *Plant Sci.*, 116: 185-195.
- Ozyigit II, Kahraman MV, Ercan O (2007). Relation between explants age, total phenols and regeneration response in tissue cultured cotton (*Gossypium hirsutum* L.). *Afr. J. Biotechnol.*, 6: 3-8.
- Paiva PDO, Paiva R, Pasqual M, Paiva LV (2004). In vitro establishment of *Strelitzia reginae* Banks.). *Ciência Agrotecnol.*, 28(5): 1031-1037.
- Pan MJ, van Staden J (1998). The use of charcoal in in vitro culture – A review. *Plant Growth Regul.* 26: 155-163.

- Promtep K (1981). Embryo and tissue culture of *Strelitzia reginae* ait. Agris record: XB8110894.
- Quoirin M, Da Silva MC, Martins KG, De Oliveira DE. (2001). Multiplication of juvenile black wattle by microcuttings. Plant Cell Tissue Org. Cult., 66: 199-205.
- Ray BP, Hassan L, Sarker SK (2011). In vitro cultivation and regeneration of *Solanum melongena* (L.) using stem, root and leaf explants. Nepal J. Biotechnol., 1(1): 49-54.
- Razdan MK (1993). Introduction to Plant Tissue Culture. Andover, Hampshire: Intercept Limited.
- Salin ML, Bridges SM (1981). Chemiluminescence in wounded root tissue. Plant Physiol., 67: 43-46.
- Smith RH, James Price H, Thaxton JB (1977). Defined conditions for the initiation and growth of cotton callus in vitro I. *Gossypium arboreum*. Soc. Vitro Biol., 13(5): 329-334.
- Srangsam A, Kanchanapoom K (2007). Establishment of in vitro Culture of Musa AA Group 'Kluai Sa' and Musa AA Group 'Kluai Leb Mue Nang' and the Analysis of Ploidy Stability. Sci. Asia, 33: 437-442.
- Steel RGD, Torrie JH (1980). Principles and procedures of statistics: A biometrical approach (2nd Ed.). McGraw Hill, New York.
- Strosse H, Van den Houwe I, Panis B (2009). Banana Improvement. FAO Corporate Document Repository. mhtml:file://E:\Zingiberaceae\Banana Improvement.mht
- Taji A, Williams R (1996). Tissue culture of Australian plants. University of New England, Armidale NSW, Australia.
- Teixeira JB, Söndahl MR, Kirby EG (1994). Somatic embryogenesis from immature inflorescences of oil palm. Plant Cell Rep., 13: 247-250.
- Teng WL, Ngai YW (1999). Regeneration of *Oxalis triangularis* spp. triangularis from suspension cells cultured in three different systems (solid, liquid-flask and bioreactor cultures). Plant Cell Rep., 18: 701-706.
- Theander O, Nelson DA (1988). Aqueous, high temperature transformation of carbohydrates relative to utilization of biomass. Adv. Carbohydr. Chem. Biochem., 46: 273-326.
- Thomas TD (2008). The role of activated charcoal in plant tissue culture. Biotechnol. Adv., 26: 618-631.
- Thompson JE, Legge RL, Barber RF (1987). The role of free-radicals in senescence and wounding. New Phytol., 105: 317-344.
- van de Pol PA, van Hell TF (1988). Vegetative Propagation of *Strelitzia reginae*. Acta Hort. 226: 581-586.
- Wang Q, Jaana Laamanen J, Uosukainen M, Valkonen JPT (2005). Cryopreservation of in vitro grown shoot tips of raspberry (*Rubus idaeus* L.) by encapsulation-vitrification and encapsulation-dehydration. Plant Cell Rep., 24: 280-288.
- Weatherhead MA, Burdon J, Henshaw GG (1979). Effects of activated charcoal as an additive to plant tissue culture media. Z. Pflanzenphysiol., 94: 399-405.
- Welander MJ (1988). Plant regeneration from leaf and stem segments of shoots raised in vitro from mature apple trees. J. Plant Physiol., 132: 738-744.
- Wickson M, Thimann KV (1958). The antagonism of auxin and kinetin in apical dominance. Physiol. Plant. 11: 62-74.
- Woolley DJ, Wareing PF (1972). The interaction between growth promoters in apical dominance. New Phytol., 71: 781-793.
- Wu HC, du Toit ES (2004). Reducing oxidative browning during in vitro establishment of *Protea cynaroides*. Sci. Hort., 100: 355-358.
- Xiao W, Huang XL, Huang X, Chen YP, Dai XM, Zhao JT (2007). Plant regeneration from protoplasts of *Musa acuminata* cv. Mas (AA) via somatic embryogenesis. Plant Cell Tissue Org. Cult., 90: 191-200.
- Xu J, Yin H, Wang W, Mi Q, Liu X (2009). Effects of sodium nitroprusside on callus induction and shoot regeneration in micropropagated *Dioscorea opposita*. Plant Growth Regul., 59: 279-285.
- Yan N, Hu H, Huang J, Xu K, Wang H, Zhou Z (2006). Micropropagation of *Cypripedium flavum* through multiple shoots of seedlings derived from mature seeds. Plant Cell Tissue Org. Cult., 84: 113-117.
- Zeweldu T, Ludders P (1998). Preliminary investigation in Ensete (*Ensete spp.*). J. Appl. Bot., 72(1-2): 25-27.
- Zhang BH, Liu F, Yao CB (2000). Plant regeneration via somatic embryogenesis in cotton. Plant Cell Tissue Org. Cult., 60: 89-94.
- Zhang Y, Turner JG (2008) Wound-Induced Endogenous Jasmonates Stunt Plant Growth by Inhibiting Mitosis. PLoS ONE, 3(11): e3699.
- Ziv M, Halevy AH (1983). Control of Oxidative Browning and in vitro Propagation of *Strelitzia reginae*. Hort. Sci., 18(4): 434-436.