

Supplementary information

Effects of ZnO Film Growth Route and Nanostructure on Electron Transport and Recombination in Dye-Sensitized Solar Cells.

Constance Magne^{a,b}, Thomas Moehl^c, Mathieu Urien^b, Michael Grätzel^c and Thierry Pauporté^{a*}

^a Laboratoire d'Electrochimie, chimie des Interfaces et Modélisation pour l'Energie, UMR-CNRS 7575, Chimie-Paristech, 11 rue P. et M. Curie, 75231 Paris cedex 05, France. *Author for correspondence : Fax: (33)1 44 27 67 50; Tel: (33)1 55 62 63 83; E-mail: thierry-pauporte@chimie-paristech.fr

^b Saint-Gobain Recherche, 39 quai Lucien Lefranc, 93303 Aubervilliers Cedex, France,

^c Laboratoire Photonique et Interfaces, Ecole Polytechnique fédérale de Lausanne, CH-1015 Lausanne, Switzerland.




Fig. S1 : Nyquist plots of the cell voltage effects on the impedance spectra recorded in the light for the three investigated ZnO solar cells (a) ED1; (b) ED2 and (c) NP. The full lines are the fits of the spectra.




Fig. S2 : (a-c) Nyquist plot of the impedance spectrum of ED2 cell measured under illumination at an applied voltage of 0.55V. (b) is a high frequency zoom view of (a), showing the counter-electrode contribution.




Fig. S3 : R_{ct} and R_{tr} of D149/ZnO cells measured under light (100 mW.cm^{-2} , AM1.5G) versus the corrected applied potential.




Fig. S4: Electron lifetimes measured by EIS in the dark for the NP-ZnO cell versus applied voltage. The blue line is the raw data and the red line is obtained after ohmic drop correction.




Fig.S5 : Lifetimes (τ_n) (full line) and transport times (τ_d) (dashed line) in the dark as a function of the conductivity.




Fig. S6: Diffusion length of the electron (L_n) in the ZnO photoelectrodes measured in the dark as a function of the applied voltage corrected of the ohmic drop.