

Efficacy of Progressive Resistance Training on Balance Performance in Older Adults

A Systematic Review of Randomized Controlled Trials

Rhonda Orr,¹ Jacqui Raymond¹ and Maria Fiatarone Singh^{1,2,3}

- 1 Discipline of Exercise and Sport Science, Faculty of Health Sciences, University of Sydney, Sydney, New South Wales, Australia
- 2 Faculty of Medicine, University of Sydney, Sydney, New South Wales, Australia
- 3 Hebrew SeniorLife and Jean Mayer USDA Human Nutrition Center on Aging, Tufts University, Boston, Massachusetts, USA

Contents

Abstract	317
1. Methods	319
1.1 Search Strategy	319
1.2 Inclusion and Exclusion Criteria	319
1.3 Selection	320
1.4 Data Abstraction	320
1.5 Quantitative Data Synthesis and Validity Assessment	320
2. Results	323
2.1 Study Characteristics	323
2.1.1 Study Design	323
2.1.2 Quality Assessment and Validity	323
2.1.3 Participant Characteristics	324
2.1.4 Intervention Characteristics	324
2.1.5 Control Activities	328
2.1.6 Withdrawals	328
2.1.7 Balance Outcome Measures	328
2.2 Balance Outcomes	328
2.2.1 Static Balance	334
2.2.2 Dynamic Balance	335
2.2.3 Functional Balance	335
2.2.4 Computerized Dynamic Posturography	336
2.2.5 Subgroup Analysis	336
3. Discussion	336
3.1 Implications for Research	340
4. Conclusion	340

Abstract

The serious health, social and economic consequences of falls are well documented. Lower extremity muscle weakness and power as well as balance impairment are major independent intrinsic contributors to falls and amenable to intervention. Progressive resistance training (PRT) is widely accepted as an appropriate modality for treating sarcopenia and has been reported to improve balance. However, other studies affirm no significant effect of PRT on balance.

To date, there is no clear, definitive statement or synthesis of studies that has examined the effect of PRT on balance. Therefore, our objective was to systematically review the literature to probe the merit of PRT as a single intervention on balance performance in older adults. We conducted a comprehensive search of major electronic databases to October 2006, with citation searches and bibliographic searches of journal articles and literature/systematic reviews. Two independent reviewers screened for eligibility and assessed the quality of the studies using the Physiotherapy Evidence Database scale for validity assessment. Randomized controlled trials of PRT only, with any balance outcome in participants with a mean age of ≥ 60 years (individual minimum age > 50 years) were included. Trials that contained more than one intervention, providing the PRT and control groups matched the inclusion criteria, were also included. Because of the heterogeneity of interventions and balance outcomes, a meta-analysis was not performed. However, corrected effect sizes with confidence intervals were determined for each study outcome. Twenty-nine studies were compatible with the inclusion/exclusion criteria and were eligible for review. Participants ($n = 2174$) included healthy, community-dwelling, mobility-limited, frail cohorts and those with chronic comorbidities. Balance outcomes conducted were extensive and were broadly categorized by the authors as: static, dynamic, functional and computerized dynamic posturography. Some studies used more than one balance outcome. The number of balance tests in all totalled 68. Fourteen studies (15 tests representing 22% of all balance tests) reported improvements, significantly greater than controls, in balance performance following PRT. Improvements were not linked to a particular type of balance performance. The inconsistent effect of PRT on balance may be explained by heterogeneity of cohort and balance tests, variability in methodology of the balance test and sample size, inadequate dose of PRT and/or compliance to training, or lack of statistical power. Standardization of balance testing methodology and better reporting of procedures may ensure greater comparability of results in future studies. It is also possible that PRT alone is not a robust intervention for balance control. This is the first systematic synthesis of the literature to examine the effectiveness of PRT alone on balance performance in older adults. The limited evidence presented in currently published data has not consistently shown that the use of PRT in isolation improves balance in this population. However, further research should explore optimal resistance training regimens that: focus on the muscles most pertinent to balance control, best target neuromuscular adaptations that protect against postural challenges and elucidate mechanism(s) by which PRT may affect balance control.

Functional decline and frailty, serious clinical sequelae of a fall, make falls a major public health concern. Falls are multifactorial in origin. Although dizziness, postural hypotension, depression, cognition, visual disturbance and slow reaction time are associated with falls,^[1] lower extremity muscle weakness and balance impairment (balance dysfunction or postural instability) have been identified as major independent contributors to falls.^[1-4] Fallers have lower quadriceps,^[4] ankle dorsiflexor^[5-7]

and ankle plantarflexor^[5,7] muscle strength than non-fallers. Similarly, poor balance was strongly associated with falls risk in community dwelling^[6,8] and institutionalized older adults^[9] and stroke patients.^[10]

Loss of strength has been observed to be associated with a decline in functional performance,^[11,12] thus prompting the investigation of muscle weakness as a contributor to balance itself. Increased ankle plantarflexor,^[13] knee and hip strength^[14] was

associated with better standing balance in healthy^[13] and disabled^[14] older adults. Furthermore, a strong relationship between decline in strength and poorer balance performance has been identified in community-dwelling older adults and nursing home residents.^[15]

More recently, lower leg power has been observed to contribute to poorer functional performance^[16] and is suggested to be a better indicator of falls risk than strength.^[17] The age-related reduction in muscle power appears to be determined by impaired contraction velocity, as well as muscle mass/force-generating capacity. This attenuated ability to contract postural muscles rapidly may contribute to an increased risk of falls.^[18] Elderly fallers demonstrated lower leg power than non-fallers^[17] and reduced leg power has been shown to be an early indicator of balance deficits.^[19]

Because sarcopenia and muscle weakness are factors amenable to improvement, progressive resistance training (PRT) [strength training exercise with progressive overload where muscles exert a force against an external load or contract isometrically] has been widely accepted as an appropriate modality of treatment. Gains in strength following PRT may be accompanied by improved body composition, metabolic health, blood pressure, cognition, depressive symptomatology, sleep and reaction time, making PRT an efficacious intervention in terms of time and effort for older adults. Power training, performed with a fast-velocity concentric action and slow eccentric action, is considered as a specific form of resistance training. PRT and power training have been reported to improve balance performance.^[20-33] In contrast, a recent systematic review and meta-analysis of the effect of PRT on physical function could find no apparent merit of PRT on standing balance in 12 studies.^[34] Moreover, the FICSIT (Frailty and Injuries: Cooperative Studies of Intervention Techniques)^[35] and Moreland et al.^[3] meta-analyses lacked supporting evidence for a significant effect of PRT on falls events.

The literature examining PRT and balance performance is characterized by an extreme heterogeneity in populations, training regimen (dose, mode, progression) and balance outcome variables making clear comparisons between studies difficult. There has been no clear definitive statement or

synthesis of these studies. Few studies have closely examined whether the dose of resistance exercise determines a positive balance outcome, an aspect critical to the design of optimal exercise programmes to enhance balance. Furthermore, no 'gold standard' of clinical balance exists, and numerous and varied tests have been devised to assess and quantify different components of balance/postural stability, each having their own merits and limitations. Therefore, our objective was to present the first systematic synthesis of evidence from randomized controlled trials (RCTs) in order to determine the efficacy of PRT as a singular intervention on balance performance in older adults. Due to the heterogeneity of key variables, a meta-analysis was not performed.

1. Methods

1.1 Search Strategy

Electronic database searches were performed in: CINAHL, Cochrane Central Register of Controlled Trials, Cochrane Musculoskeletal Injuries Group, Cochrane Reviews, MEDLINE, SportDiscus, EMBASE, Science Direct, Current Contents, Web of Science; PEDro and PubMed, from earliest record to October 2006. Permutations of keyword combinations for the following categories were used:

1. Intervention: resistance/strength/power/weight training/exercise, randomized trials, and clinical trials.
2. Participants: aged, elderly, geriatric, older adults, or senior.
3. Outcome: balance, postural stability, postural control, body sway, or neuromuscular performance.

Bibliographies of all eligible papers and systematic reviews identified from electronic database searches were manually searched for any papers missed by the database searches. Non-English papers were included, but excluded if translation was unsuccessful.

1.2 Inclusion and Exclusion Criteria

Potentially relevant papers describing RCTs that investigated PRT as an exercise intervention and balance performance as an outcome in older adults, were examined.

Studies of older, men or women, mean age ≥ 60 years (minimum age 50 years) were considered. Participants could include healthy, community-dwelling cohorts, nursing home residents, frail, mobility- or functionally-limited adults or persons with pathology.

We examined RCTs in which one group performed PRT or power training. Studies that included balance training or multimodal training (i.e. training additional to resistance exercise) were excluded. PRT could be conducted by a number of modalities, for example, weights machines, free weights, isometric exercise, elastic tubing (Therabands™)¹ or other equipment (weighted vests/belts/balls, soft weights, rice bags, filled water bottles).

RCTs in which participants assigned to the control group participated in usual daily activity/usual care, or activities that enhanced the blinding of the intervention, e.g. sham exercise, flexibility training, educational seminars, were considered.

All modes of static, dynamic and functional balance performance and postural challenge assessment (computerized dynamic posturography) were included in this review. Descriptions of the types of tests used to assess balance performance in this review are presented in table I. Static balance is defined as the ability to maintain the centre of gravity over a narrow base of support in an upright position.^[36,37] Dynamic balance is the ability to maintain equilibrium whilst the body's centre of gravity is in motion (i.e. the centre of gravity does not stay within the base of support). Functional balance assesses whether a balance problem exists, predicts falls risk or assesses the ability to carry out everyday tasks or activities.^[37] Computerized dynamic posturography assesses postural stability on a computer-interfaced dynamic force platform. This test allows the quantitative examination of balance under static and dynamic conditions and can test sensory organization and reactions to surface displacement.

1.3 Selection

All papers identified by the search strategy were screened independently by two researchers (RO and JR), first by title and then by abstract using the

eligibility criteria. Any differences were resolved by discussion and mutual consent, or by a third assessor (MFS). We did not include abstracts and conference papers from annual meetings because of the paucity of data. Remaining papers were retrieved for further scrutiny to determine eligibility. If data provided were insufficient, we attempted to contact authors for further information.

1.4 Data Abstraction

Data were extracted on to pretested standardized forms using the following headings: Study Characteristics: *Study Design* (including *Statistical Analysis*), *Participant Characteristics*, *Intervention Characteristics*, *Control Activities* and *Balance Outcome Measures* and *Balance Outcomes*.

1.5 Quantitative Data Synthesis and Validity Assessment

The primary outcome was change in balance performance. Because of the heterogeneity of balance tests used and outcomes measured, we used the standardized difference method as the effect size (ES) measure. ES was determined by subtracting the mean change in balance outcome in the PRT group from the mean change in balance outcome in the control group, and dividing the difference by the pooled baseline standard deviation of the PRT and control groups.^[48] The ES was then corrected for small-sample bias.^[49] A weighted mean difference was calculated as the difference in means of the change scores in the intervention and control groups, using the method of the Cochrane Collaboration group.^[50] For studies that included multiple outcomes (e.g. more than one balance test performed, or one group training at high intensity and one training at low intensity, or at differing weekly frequency), each balance test was treated independently.

Some studies performed *a priori* or *post hoc* power testing based on primary outcomes. Because power analysis may have been performed on outcome measures other than balance performance in these studies, *post hoc* power testing was completed on balance data using the *G*Power* power analysis

1 The use of trade names is for product identification purposes only and does not imply endorsement.

Table 1. Assessment of balance performance by balance type

Balance test	Description of balance test	Balance measure	Reliability/validity of measure
Static, eyes open and/or eyes closed			
Single-leg stance	Stand on one leg with other leg flexed, allowing foot to clear the floor and balance as long as possible. No assistive devices are permitted ^[36,38]	Time up to 5–60 sec or as long as possible without grasping for support or moving feet (most studies use 30 sec)	Validity demonstrated by relationship with gait performance, falls status, IADL, frailty ^[39]
Parallel stance, double-leg stance, Classic Romberg	Feet placed together, side by side ^[36,38]	Time up to 10–30 sec, without grasping for support or moving feet	Reproducibility coefficients 0.7–0.99 ^[38]
Semi-tandem stance	Toe of one foot placed adjacent to heel of other foot ^[36,38]	Time up to 10–30 sec	
Tandem stance, Sharpened Romberg	Heel of one foot placed directly in front of toe of other foot ^[36,38]	Time up to 10–30 sec	
FICST-3 (or -4)	Combined parallel, semi-tandem, tandem stance (plus single-leg stance)	Score 0–4 (0–5), ^a determined by the time each stance could be held for, up to 10 sec	$r = 0.66$ over 3–4 mo test-retest interval ^[36]
Postural sway on compliant surface ^[6]	Tandem stance on a 15-cm thick medium-density foam rubber mat, wearing swaymeter (40-cm rod attached to waist by a belt and extends posteriorly. A pen mounted at the end of the rod records sway excursions on graph paper)	Sway path (mm) traced in 30 sec	Test-retest reliability coefficients ($n = 37$). Eyes open: $r = 0.81$; $p < 0.01$ (95% CI 0.66, 0.90). Eyes closed: $r = 0.73$; $p < 0.01$ (95% CI 0.53, 0.85) ^[6]
Balance ability on a tilt board ^[38]	Stand with feet perpendicular to tilt board's rotational axis to measure anterior-posterior or omni-directional instability	Time up to 30 sec	
Dynamic, eyes open and/or eyes closed			
Tandem walk – forward or backward	Walk as quickly and carefully as possible placing one foot directly in front of (or behind) the toe (heel) of the other foot without making errors (i.e. when one foot is not placed in contact with the other, taking a step to the side)	Time (average of 2 trials), number of errors	Test-retest (1 wk) $r = 0.94$ ^[22]
Functional reach ^[40]	Maximum distance that can be reached forward beyond arm's length while maintaining a standing fixed base of support. Predictive of recurrent falls risk	Mean distance achieved over 3 trials or score 0–3 ^b determined by distance reached. Two practice trials allowed	CV = 2.5%, ICC = 0.92, ICC (intertester) = 0.98
Balance beam walk ^[38]	Walk as quickly as possible along a narrow (8.5 cm) or wide (17 cm) beam over a distance of 6–9 m	Time on beam(s), distance walked before stepping off (m)	

Continued next page

Table 1. Contd

Balance test	Description of balance test	Balance measure	Reliability/validity of measure
Functional			
Berg Balance Test ^(10,41)	Level of balance quantified by 14 tasks: sit to stand, stand to sit, transfers, standing with feet together, with eyes closed, with one foot in front, on one foot, reaching forward, retrieving an object from the floor, turning to look behind, turning 360°, placing alternate foot on stool	Score 0–56° determined from an ordinal scale of 0 (cannot perform) to 4 (can perform) for each of the 14 activities	Inter-rater reliability coefficient = 0.98, intra-rater reliability coefficient = 0.98, internal consistency Cronbach's α = 0.96. Correlations with laboratory measures of sway (–0.55), global ratings of therapists (0.81) and care-givers (0.47–0.61) ⁽¹⁰⁾
CS-PFP – balance and coordination domain ⁽⁴²⁾	Functional tasks included transferring laundry, donning and removing a jacket and seatbelt, sweeping, vacuuming, bed making, stair climbing, getting in and out of a bathtub	Score 0–12 (average of 5 standardized and summed domain scores). ^d Separate domain scores given. The tasks are quantified by time, distance or weight. CS-PFP balance and coordination is derived from tasks quantified by time	Inter-rater reliability coefficient = 0.99, test-retest coefficient = 0.96, Cronbach's α = 0.91. CS-PFP-B&C correlated with the 5 physical performance capacity measures it reflected (r = –0.21 to –0.91) ⁽⁴²⁾
ADAP – balance and coordination domain ⁽²⁴⁾	16 common functional tasks, such as transferring laundry, boarding a bus performed at peak effort. The test is modelled on CS-PFP	Score 0–100 for each task. The tasks are quantified by time (speed), distance or weight. ^e ADAP – balance and coordination is derived from tasks quantified by time	ICC total score = 0.96, ICC domain scores = 0.75–0.95 ⁽²⁴⁾
Computerized dynamic posturography, computerized balance platforms or force plates¹			
Chattecx Dynamic Balance System ⁽⁴³⁾	Single- or double-leg stance, static platform, double-leg stance on platform moving forward and backward, double-leg stance on platform tilting up and down (4° amplitude, frequency 1 Hz)	Time up to 30 sec. Sway amplitude (cm). Loss of balance	ICC = 0.41–0.90 ICC = 0.06 (double-leg stance, static platform, eyes closed) ⁽⁴⁴⁾
Sensory Organisation Test ⁹ using EquiTest ⁽⁴⁵⁾	The platform's support surface and visual surround are fixed or can sway with the participant (sway referenced). Three consecutive 20-sec trials are given in each of 3 visual (eyes open, eyes closed, sway-referenced) and 2 support (fixed, sway-referenced) conditions. Timing and magnitude of surface forces are measured in response to these perturbations	Functional base of support. Equilibrium quotient (sway within and exceeding limits of stability). Balance strategy score (Newtons shear force/kg body mass). Loss of balance when sway exceeds limits of stability (8.5° anteriorly, 4° posteriorly)	ICC (1 week apart) = 0.15–0.7 for first trial and 0.26–0.68 for average of 3 trials ⁽⁴⁶⁾
Motor Coordination Test using EquiTest ⁽⁴⁵⁾	Double-leg stance on platform that provides a series of sudden anterior and posterior transitions (250, 300, 400 msec duration) and toes-up, toes-down rotations (8° amplitude, 400 msec duration)	Latency – time (msec) from perturbation to actively resisting sway. Sway momentum	
Balance Master ⁽⁴⁷⁾	Move centre of gravity as quickly as possible to attain 8 targets set at 75% of their limits of stability. Visual conditions: eyes open, eyes closed or with visual feedback	% postural sway with eyes open, eyes closed, and with video feedback. Total number of 75% limits of stability targets attained. Time taken to attain each target	
a	FICSIT-3 score = 0–4, FICSIT-4 score = 0–5; 0 = unable to perform parallel stance; 0.5 = parallel stance held <10 sec; 1 = parallel stance held for 10 sec, semi-tandem stance <10 sec; 2.0 = parallel and semi-tandem stance held for 10 sec, unable to perform tandem stance; 3.0 = parallel and semi-tandem stance held for 10 sec, tandem stance <10 sec; 4 = all 3 stances held for 10 sec; 5 = all 4 stances held for 10 sec. ⁽³⁸⁾		

Continued next page

Table I. Contd

b	Functional Reach score 0-3: 0 = unable or unwilling to reach; 1 = reach ≤ 6 inches (15.2 cm); 2 = 6 inches < reach < 10 inches (25.4 cm); 3 = reach ≥ 10 inches. Relative risk for falls: 0 = 28 times more likely to fall; 1 = 4 times more likely to fall; 2 = 2 times more likely to fall; 3 = not likely to fall. ^[40]
c	Berg Balance Test score 0-56. The Berg Balance Test rates the ability to maintain balance when performing 14 movements used in everyday tasks. For scores of 54-56, each 1 point drop is associated with a 3-4% increase in fall risk. For scores of 46-55, each 1 point drop is associated with a 6-8% increase in fall risk. A score <45 reflects those unsafe in independent ambulation. ^[10] A score <36 indicates close to 100% risk of falls.
d	The CS-PFP is a comprehensive measure of physical functional performance. Fifteen everyday tasks of independent living are summarized in 5 physical domains that reflect upper- and lower-body strength, upper-body flexibility, endurance and balance and coordination. ^[42] Measures of time, weight and distance are transformed to standard scores and scaled 0-12. The lowest 10% of scores are designated as 1 and highest 10% scored as 12. Scores between the 10th and 90th percentile are grouped into 8 percentile bands and scored as 2-11. Inability or refusal to perform task scores zero. The total CS-PFP is the sum of the standard scores, scaled 0-12.
e	The ADAP is a comprehensive measure of physical functional performance comprising 16 common tasks that are summarized in 5 physical domains that reflect upper- and lower-body strength, upper-body flexibility, endurance and balance and coordination. ADAP provides a total score and 5 physical domain scores. Each task was scaled from 0 to 100. ^[24]
f	Assess postural stability is represented by movements of the centre of mass, on a stable and/or moving, destabilizing/perturbing surface, with eyes open and/or closed.
g	The Sensory Organisation Test evaluates which of the senses (visual, vestibular or somatosensory) [or combinations] and motor strategies contribute to balance deficits. The 6 conditions are: (1) eyes open/firm surface; (2) eyes closed/firm surface; (3) sway referenced vision/firm surface; (4) eyes open/sway-referenced surface; (5) eyes closed/sway-referenced surface; and (6) sway referenced vision/sway-referenced surface.
	ADAP = Assessment of Daily Activity Performance; CS-PFP = Continuous-Scale Physical Performance Score; CS-PFP-B&C = CS-PFP Balance and Coordination domain; CV = coefficient of variation; FICSIT = Frailty and Injuries: Cooperative Studies of Intervention Techniques; IADL = Instrumental Activities of Daily Living; ICC = intra-class correlation coefficient.

program.^[51] As a consequence of the heterogeneity of interventions, balance outcomes and cohorts, a meta-analysis was not performed.

Quality assessment of eligible papers was undertaken independently by RO and JR using the Physiotherapy Evidence Database (PEDro) scale,^[52] an 11-item scale that includes the 3-item Jadad scale^[53] and 9-item Delphi list.^[54] The PEDro scale rates RCTs from 0 to 10. One question is used to establish external validity and is not included in the score. The PEDro scale reports an inter-rater reliability intraclass correlation coefficient (ICC) = 0.68 (95% CI 0.57, 0.76). Any differences were resolved by discussion and mutual consent or rating by a third assessor (MFS).

2. Results

The flow of papers, from potentially relevant to selection, from January through October 2006 is displayed in figure 1. A total of 29 studies were eligible for review according to the inclusion/exclusion criteria. Thirty-seven of the 66 papers retrieved were excluded for the following reasons: utilized combined training programmes, not isolated PRT^[55-70] balance training,^[71,72] not PRT,^[73-75] no progression of resistance training,^[76] not a RCT,^[77-85] balance outcome not included,^[86,87] mean age <60 years,^[88] or the paper contained the same cohort or was a preliminary paper to one included in this review.^[89-91]

2.1 Study Characteristics

2.1.1 Study Design

Of the 29 single-blind RCTs, one study was a multicentred study^[92] and two were conducted at two sites.^[26,93] Most RCTs consisted of a PRT group and a control group; four studies comprised four groups;^[32,33,94,95] and eight studies included three groups.^[23-28,31,93]

2.1.2 Quality Assessment and Validity

Quality assessment and internal validity of the included trials are summarized in table II and table III. The inter-rater correlation of PEDro scores between the researchers was 0.9, greater than reported by PEDro raters.^[52] Two studies randomized more participants to PRT because of greater anticipated




Fig. 1. Flow of papers from potentially relevant to selected for review.

withdrawals from the control group.^[93,96] Twelve papers (43% of studies) comprising 39% of all balance tests scored ≥ 6 on the PEDro scale, signifying better quality studies.

2.1.3 Participant Characteristics

A total of 2174 participants were randomized (1020 to PRT, 866 to control, 268 to other modes of training, 20 not specified^[29]). Most participants were healthy and community-dwelling ($n = 1090$).^[21,22,24,27,29,32,33,47,94,95,99,101,103-105] Six RCTs^[11,28,31,96,100,102] comprised functionally impaired or mobility-impaired participants ($n = 432$). Frail ($n = 383$)^[30,92,93] and institutionalized participants ($n = 20$)^[98] and individuals with coronary heart disease (CHD) [$n = 42$],^[20] osteoarthritis ($n = 103$)^[26] and low bone mass ($n = 243$)^[25] were also eligible for inclusion in this review. The mean age of all participants ranged from 60–88 years and 68.3% were women. Not all studies reported ages for the intervention groups, but the average age ranged from 61 to 88 years where reported, and was comprised of 70% women. The average age of control group participants ranged from 57 to 88 years where reported, and was comprised of 70.7% women. One study contained only men^[29] and seven studies examined women only.^[22,24-26,28,30,104] Major co-

morbidities included CHD, osteoarthritis, osteoporosis, hypertension, type 2 diabetes mellitus, respiratory and kidney disorders and cancer. Studies were conducted in Australia,^[21,32,101] Canada,^[25] Finland,^[30] the Netherlands,^[24,93,96] New Zealand,^[92] Portugal,^[29] the UK^[47,104] and the US.^[11,20,22,23,26,27,31,33,47,94,95,98-103,105] Ethnicity, although not well reported, was principally Caucasian (76.5–100%), with African American (7–34%) and Asian (7%) cohorts also participating. The groups were generally comparable at baseline. Some studies indicated some significant baseline differences in age, weight, body composition, ankle and hand-grip strength or function.

2.1.4 Intervention Characteristics

The intervention and control group characteristics are shown in the supplementary material ['ArticlePlus'] at <http://sportsmedicine.adisonline.com>. A total of 852 (from 1020) completed PRT and a total of 126 participants (range 0–18 per study) were reported as withdrawals. Attrition rates averaged 13%, and varied greatly between the studies from 0%^[11] to 39%,^[31] with three RCTs not specifying intervention withdrawals.^[26,29,102]

Setting and Training Equipment

Interventions were mainly conducted in a gym/community setting;^[11,21,22,24-27,30,32,33,94,95,98,101] combined gym and home-based location^[28,47,93,96,104,105] or were solely home-based programmes.^[92,100,102] Some studies did not report a training venue, but a gym/community setting was likely from the description of training equipment.^[20,29,31,99,101,103]

A diversity of training equipment was used: pneumatic^[22,25,31,32] and weight-stack^[20,23,29,30,33,95,101,103] resistance training machines; free weights;^[20,21,24,26,27,92,98] elastic bands/tubing;^[24,28,47,93,96,98,100,102,104,105] body weight;^[27,28,93,96,104] and weighted equipment such as balls, vests, rice bags, sandbags^[11,27,28,33,98,104] solely, or in combination. Twenty studies used a single type of training equipment (12 used resistance training machines,^[22,23,25,29-32,94,95,99,101,103] three used free weights,^[21,26,92] four used elastic bands/tubing,^[92,100,102,105] one used weighted vests^[11]) and nine studies used a combination of equipment.^[20,24,27,28,33,93,96,98,104]

Table II. Quality assessment and internal validity of studies

Randomization process	Allocation concealment	Baseline comparability	Assessor blinding	ITT analysis	Attrition rate (% with- draws)	A priori power testing	Post hoc calculation of power of balance tests ^a	Methodological quality (0–10) ^b	Reference
Stratified by physical function domain of SF-36	NR	Yes	NR	No	21	NR	0.71	4	20
Computer-generated random numbers list	Stated, but no specific information given	Yes	Yes	No	9	No	0.59	8	21
Computer-generated algorithm (permutated blocks) stratified by residence	Sealed envelopes	Yes	Yes	Yes	10	Yes	0.10	8	98
NR	NR	Yes	Yes	NR	0	No	0.15	6	11
NR but more randomized to RT because of greater anticipated withdrawals from control group	NR	Yes	Yes, but not the same assessor for pre- and post-measures	No	32	Yes, but proved to be insufficient power	0.06–0.22	4	93
NR	NR	Yes	NR	No	20	Yes	0.05	5	99
Randomly permutated blocks	NR	Yes	Yes	No	12	Post hoc	0.05–0.15	7	94
Block randomized and stratified by level of function	NR	Yes	Yes	No	12	No	0.06	6	100
Computer-generated random numbers table	NR	Yes	Yes verified blinding status, 37% correct	NR	17.6	Yes	0.32	6	24
NR	NR	Yes	NR	NR	10.4	NR	0.05–0.12	5	101
Randomly permutated blocks of 4 by staff not involved in data collection	NR	Yes	Yes	No	7	No	0.05–0.39	6	102
Computer-generated randomization sequence in blocks of 6	No	Yes	Yes verified blinding status, <38% correct	Yes	6.7	Yes	0.30	8	92
Stratified by baseline postural sway	NR	Yes	Yes	Yes	6	Yes	0.80	7	25
Stratified variable, block randomized to give even numbers in each group	NR	Yes	NR	No	NR	NR	0.09–0.88	4	26
Stratified by sex	NR	Yes	No	NR	31.4	Post hoc	0.05–0.42	4	31
NR	NR	No	NR	Yes	4.8	NR	0.46	5	22

Continued next page

Table II. Contd

Randomization process	Allocation concealment	Baseline comparability	Assessor blinding	ITT analysis	Attrition rate (% withdrawals)	A priori power testing	Post hoc calculation of power of balance tests ^a	Methodological quality (0–10) ^b	Reference
NR	NR	NR	NR	No	5	NR	0.46	5	23
Computer-generated random numbers list, blocks of 4, stratified by sex	Serially numbered opaque sealed envelopes	Yes	No	No	8.9	Yes	0.10–0.39	6	32
NR	NR	Yes	No	Yes	19	No	0.26–0.94	5	27
NR	NR	NR	NR	No	8.3	No	0.05	5	103
Random numbers table after matching for age and habitual physical activity	NR	Yes	No	No	16.7	Yes	0.21	5	104
20 matched for age then randomized, but one participant 'chose' training group	NR	Yes	NR	No	10	Yes	Cannot calculate	4	28
NR	NR	NR	NR	NR	NR	NR	0.74	2	29
NR	NR	Yes	NR	No	14.7	Yes	0.24–0.75	5	95
Randomized in blocks of 4	Closed envelopes	Yes	NR	Yes	23.5	No	0	6	30
NR	NR	Yes	No	No	19	Post hoc	0.05–0.21	5	105
NR, but more randomized to RT because of greater anticipated withdrawals from control group	NR	No	NR	No	34	NR	0.05–0.73	3	47
NR, but more randomized to RT because of greater anticipated withdrawals from control group	NR	Yes	Yes	No	8.3	NR	0.05–0.06	5	96
Blocked allocation schedule generated by algorithm and stratified by sex	NR	Yes	Yes	No	3.8	Post hoc	0.06–0.23	6	33

^a Post hoc calculations were performed by the authors using G*Power.^[51]

^b Methodological rating based on Physiotherapy Evidence Database (PEDro) score.^[52]

ITT = intention-to-treat; NR = not reported; RT = resistance training; SF-36 = Medical Outcomes Study Short Form-36.^[97]

Table III. Summary of the numbers of trials reporting the characteristics of study design described in table II

Characteristic of study design	Yes		No		Not reported	
	no. of trials	references	no. of trials	references	no. of trials	references
Method of randomization	16	20,21,24-26,28,30-33,92,94,98,100,102,104			13	11,22,23,27,29,93,95,47,96,99,101,103,105
Allocation concealment	4	21,30,32,98	1	92	24	11,20,22-29,31,33,47,93-96,99-105
Baseline comparability	24	11,20,21,24-28,30-33,92-95,96,98-102,104,105	2	22,47	3	23,29,103
Blinding of assessors	12	11,21,24,25,33,92-94,96,98,100,102	5	27,31,32,104,105	12	20,22,23,26,28-30,95,47,99,101,103
Power testing	10	24,25,28,32,92,93,95,98,99,104	6	11,21,30,100,102,103	9	20,22,23,26,27,29,47,96,101
	4 ^a	31,33,94,105				
ITT analysis	6	22,25,27,30,92,98	18	20,21,23,26,28,32,33,93-95,47,96,99,100,102-105	5	11,24,29,31,101
Attrition	26	11,20,22-25,27-29,32,33,47,92-96,98,99,101-105			3	21,31,100

a *Post hoc.*

ITT = intention-to-treat.

Training Prescription

The prescription of PRT was generally well described. The study duration averaged 22.7 weeks (range 8–104 weeks). The mean training session duration was 58.8 minutes (range 35–90 minutes) and frequency of training was 2–3 days/week. Two to three sets per session were prescribed in all studies.

Studies using resistance machines quantified the training intensity most objectively using percentage of one repetition maximum (1RM).^[20,22,23,25,29-33,92,94,95,99,101,103] some also used subjective assessment, i.e. ratings of perceived exertion (RPE).^[20,22,32,33] Several studies prescribed variable intensity in each session. For example, each of the three sets was performed at successively increasing intensities of 50%, 60% and 70% 1RM,^[99] or intensity varied according to the exercise: shoulder press was performed at 45% 1RM, chest press at 50% 1RM, and fly at 60% 1RM.^[23] Four studies used RPE alone,^[11,21,24,102] but Jette et al.^[102] did not specify the value. Skelton et al.^[104] prescribed intensity at <70% heart rate reserve. Five studies did not report intensity,^[26,28,98,100] but the methodology suggested training was performed at relatively low intensity. Four studies used power training; one at 40% 1RM,^[31] one at 20%, 50% and 80% 1RM,^[32] one at increasing loads with each set (45%, 60% and 75% 1RM)^[101] and the other at moderate intensity

using weighted vests.^[11] We categorized studies according to training intensity: high ($\geq 70\%$ 1RM),^[20-22,25,30-33,94,95,103] moderate (41–69% 1RM),^[11,23,24,29,32,47,92,99,104,105] and low ($\leq 40\%$ 1RM).^[26,27,31,32,93,96,98,100,102,104]

Progression

Although described in all studies, the protocol for progression was somewhat diverse. Training load was increased: following 1RM testing 1-,^[32] 2-,^[38,92,103] 4-^[20,22,29,31,95,99] and 6-weekly;^[23] by maintaining RPE;^[21,24] when a number of repetitions or sets and repetitions were achieved;^[25,26,28,93,96,100,101,104] when fatigued;^[47] when possible;^[27] or with no details other than 'as muscle strength increased'.^[30]

Supervision and Compliance

Training was fully supervised in 18 of 29 cases,^[11,20-22,24,25,27,29,30,32,33,93,95,96,98-101] with five studies explicitly reporting supervision by trained staff;^[20,21,24,32,96] whereas 11 studies were unsupervised,^[92] monitored intermittently^[28,47,102,104,105] or supervision not reported.^[23,26,31,94,103] Twenty-five of 29 studies reported compliance rates;^[11,20,22-29,32,33,47,92-96,98,99,101-105] three studies did not.^[21,31,100] One study described compliance to training as 'satisfactory'.^[30] Overall compliance to training at gym sessions ranged from 29% to 100% attendance. Compliance to training

ranged from 70% to 100% attendance at gym sessions on average; however, two studies reported individual participation rates in PRT as low as 29%^[93] and 47%.^[27] Overall compliance to home-based training was monitored by training diaries (supported by a booklet^[93,96]) and reported to be 69–88%. Individual home-training adherence rates were reported as low as 10%^[93] and one study noted that participant reporting in the training logs was not good.^[96] Research staff paid home visits and gave regular phone calls to maximize adherence.^[34,102] Three studies did not describe participant adherence.^[21,31,100]

Adverse Events

Adverse events were reported in 17 of 29 trials,^[11,20-22,24,25,27,28,31-33,94,96,98,101,104,105] with 7 of 17 trials reporting no adverse events.^[20,27,28,98,101,104,105] Adverse events attributed to training were mainly musculoskeletal in nature and none were reported as serious. These included exacerbation of osteoarthritic pain,^[96] trochanteric bursitis, muscle, joint and lower-back pain.

2.1.5 Control Activities

Control group activities varied across the studies and included: no treatment (usual daily activity with no increase in physical activity levels^[23,24,95,96,99]), wait-list for an exercise programme on completion of the study,^[22,31,32,94,100] attention-control (health educational talks,^[26,31,33] group driver education classes,^[47,105] phone calls and home visits^[92]). Some studies offered alternative exercise, including slow-velocity chair-based exercises with low/no resistance,^[11] functional exercises,^[30] flexibility/stretching^[20,21] and/or deep breathing classes to subjects blinded as to which intervention was the preferred treatment arm.^[25] Only Liu-Ambrose and colleagues^[25] described the exercises as 'sham'. Some wait-list controls later joined the study as intervention participants^[27,28,98,102] or became a combined training group.^[101] No control activity details were reported in three studies.^[29,93,103]

2.1.6 Withdrawals

In all trials, 1795 participants were reported to have completed the studies; 256 (average 13.2%, range 2.5–31.9%) dropped out. No completion or withdrawal details were reported for a further 123 participants randomized.^[26,29,102] A total of 753

(from 1020) participants completed PRT intervention and 126 participants (range 0–18 per study) dropped out from PRT. A total of 644 (from 866) participants completed control activities and 80 participants (range 0–13 per study) dropped out from control activities. Three RCTs did not specify intervention or control withdrawals,^[26,29,102] accounting for a further 141 and 142 participants, respectively. PRT attrition rates averaged 13%, and varied greatly between the studies from 0%^[11] to 39%.^[31] Compliance to control activities also varied between the studies, with withdrawals averaging 11.4% and ranged from 0%^[21,22,31] to 23%.^[93]

2.1.7 Balance Outcome Measures

In all of the studies reviewed, balance was only one of a number of functional performance outcomes measured. Muscle strength, muscle power, mobility, functional capacity, physical health and cognitive function were also evaluated. Balance was a secondary outcome measure in two studies.^[25,92] Balance tests were categorized as static, dynamic, functional or using computerized dynamic posturography (table I). We found that there was substantial variability in the methodological conduct of static and dynamic tests as well as a paucity of specific description, such as which leg was tested (self-selected, dominant, non-dominant, both), whether the test was conducted with eyes open or closed, shoes on or off, the number of practice trials permitted, the number of trials allowed, the distance walked, the period for which the test was conducted, (set duration or for as long as possible) and/or what times were reported (best, mean or total of trials). Some studies permitted one trial only while others permitted up to five trials. Some studies permitted one or two practice trials then recorded the best of three trials.

2.2 Balance Outcomes

Details of balance tests and balance outcomes following PRT are presented in table IV. In studies with two or more groups undertaking PRT under several conditions, e.g. high intensity versus low intensity, or frequency of 3 versus 1 day/wk, or strength training versus power training, each group was treated as having independent data. Furthermore, some studies included more than one balance

Table IV. Balance tests and balance outcomes following progressive resistance training (PRT)

Subjects	PRT intensity and duration	Balance outcome characteristics			Balance results			Reference		
		balance type	balance test	measure (max time, distance, or score)	trials allowed	% change \pm SD in PRT group ^a	significant between-group effect		WMD (95% CI)	effect size (95% CI)
Healthy, CD	MI, 12 wk	Static	SLS EO	NR	Mean of 2			0.00 (-10.65, 10.65)	0.00 (-0.53, 0.53)	105
Healthy, CD	HI, 52 wk	Dynamic	SLS EC	NR				-0.25 (-1.22, 0.72) ^b	-0.14 (0.67, 0.39) ^b	22
		Dynamic	BTW	No. errors over 2.4 m (8 ft)	Mean of 2	-14.3	p = 0.005	0.50 (-0.37, -1.37)	0.31 (-0.23, 0.84)	
Healthy, CD	MI, 12 wk	Dynamic	BTW	Time over 6.1m (20 ft)	Mean of 2			-5.9 (-10.45, -1.35) ^b	-0.82 (-1.48, -0.17) ^b	23
		Static	SLS EC	Time up to 45 sec (combined right and left leg results)	2 practice then best and mean of 3		p < 0.05	1.62 (-0.08, 3.32)	0.61 (-0.04, 1.25)	
Healthy, CD	MI, 12 wk	Dynamic	Functional reach	Distance (cm)	NR	1		1.90 (-1.32, 5.12)	0.37 (-0.25, 1.0)	104
Functionally limited, CD	LI, 8 wk	Static	SLS EO	As long as possible	NR, but practice trial given	42		Cannot calculate	Cannot calculate	28
		Dynamic	SLS EC	As long as possible		54	Yes, p NR			
Healthy, CD	MI, 14 wk	Dynamic	BTW	NR		8.9				47
			Functional reach	Distance (cm)	Best of 3	1		0.04 (-0.02, 0.10)	0.39 (-0.22, 1.00)	
		PG (Balance Master)	% Postural sway EO	1					-0.07 (-0.16, 0.02) ^b	-0.49 (-1.10, 0.12) ^b
			% Postural sway/visual FB	1					-0.01 (-0.09, 0.07) ^b	-0.08 (-0.68, 0.53) ^b
	No. 75% LOS targets attained	1					1.52 (0.38, 2.66)	0.81 (0.19, 1.44)		

Continued next page

Table IV. Contd

Subjects	PRT intensity and duration	Balance outcome characteristics				Balance results		Reference			
		balance type	balance test	measure (max time, distance, or score)	trials allowed	% change \pm SD in PRT group ^a	significant between-group effect		WMD (95% CI)	effect size (95% CI)	
Healthy, CD	HI, 12 wk	PG (Equest)	LOB during SOT	No. LOB	1			-1.00 (-2.57, 0.57) ^b	-0.39 (-1.01, 0.22) ^b	33	
			Functional base of support	Time up to 30 sec	Best of 2			-0.01 (-0.11, 0.09) ^b	-0.08 (-0.83, 0.68) ^b		
	HI, 24 wk	Static	SLS EO				p \leq 0.005	1.20 (-6.2, 8.6)	0.12 (-0.60, -0.83)		
		Static	SLS EO	Time up to 10 sec	Best of 2			-1.00 (-5.31, 3.31) ^b	-0.10 (-0.66, 0.45) ^b	94	
Healthy, CD		Dynamic	Tandem stand EO	10 sec	1			0.0 (-1.56, 1.56)	0.00 (-0.055, 0.55)		
			Standing on tilt board (AP direction)	Time up to 30 sec	1			-3.0 (-6.74, -0.74) ^b	-0.45 (-1.01, 0.11) ^b		
		Standing on tilt board (omni direction)	Time up to 30 sec	1				-2.0 (-6.68, 2.68)	-0.24 (-0.80, 0.32)		
		Wide-beam balance walk	Time over 6 m	1				-0.10 (-0.62, 0.42) ^b	-0.11 (-0.66, 0.45) ^b		
		Narrow-beam balance walk	Time over 6 m	1				0.50 (-0.52, 1.52)	0.27 (-0.28, 0.83)		
		Static	Tandem stand EO	Time up to 60 sec	1 practice, 3 trials (best or average NR)	16		p < 0.05	14.00 (5.13, 22.87)	0.69 (0.24, 1.14)	27
		Dynamic	SLS EO			98			14.8 (6.64, 22.96)	0.80 (0.34, 1.25)	
			SLS EC			9			1.50 (0.22, 2.78)	0.51 (0.07, 0.96)	
			Tandem walk	Time over 3 m (10 ft)		26			-3.50 (-8.68, 1.68) ^b	-0.30 (-0.74, 0.14) ^b	
		Tandem walk	No. of missteps over 3 m		37			-1.30 (-2.59, -0.01) ^b	-0.44 (-0.88, 0.00) ^b		

Continued next page

Table IV. Contd

Subjects	PRT intensity and duration	Balance outcome characteristics			Balance results		Reference		
		balance type	balance test	measure (max time, distance, or score)	trials allowed	% change ± SD in PRT group ^a		significant between-group effect	WMD (95% CI)
Frail, moderately limited, CD	LI, 10 wk	Dynamic	Functional reach	NR	2 practice, then mean of 3 trials		0.20 (-1.10, 1.50)	0.06 (-0.36, 0.49)	100
	LI, 24 wk	Static	Single leg-stand EO	Time up to 30 sec	1 practice, then mean of 2 trials	20	-0.15 (-2.28, 1.98) ^b	-0.02 (-0.29, 0.25) ^b	102
Healthy, CD	HI, 24 wk	Dynamic	Forward tandem walk	No. of successful steps up to 10			0.81 (-0.13, 1.75)	0.23 (-0.04, 0.50)	
		Dynamic	Functional Reach	Distance (cm)			-0.33 (-1.20, 0.54)	-0.10 (-0.37, 0.17)	
CD with knee osteoarthritis	LI, 78 wk	Dynamic	BTW 3 d/wk	Time over 6 m	Best of 3		-7.70 (-13.27, -2.13) ^b	-1.16 (-2.04, -0.27)	95
		Static on force platform	DLS EO	Postural sway distance (in 10 sec)	4 first trial was practice, but not disclosed		-3.3 (-11.8, -5.2)	-0.55 (-1.39, 0.28) ^b	
Functionally limited, CD	LI, 10 wk	Static	FICSIT 3	Time up to 10 sec, score 1-6	NR	26 (CI -17, 69)	-8.70 (-17.30, -0.10) ^b	-0.87 (-1.75, 0.01) ^b	96
		Dynamic	SLS EO	Time up to 10 sec			-0.05 (-0.13, 0.03) ^b	-0.29 (-0.76, 0.18) ^b	
CD with some physical disability	LI, 24 wk	Dynamic	BTW 1 d/wk				-0.18 (-0.29, -0.07) ^b	-0.76 (-1.24, -0.27) ^b	
		Static	SLS EO	Time up to 10 sec			1.00 (0.07, 1.93)	0.51 (0.03, 0.98)	
Healthy, CD	HI, 24 wk	Dynamic	BTW 3 d/wk	Time over 6 m	Best of 3		-7.70 (-13.27, -2.13) ^b	-1.16 (-2.04, -0.27)	95
		Dynamic	BTW 1 d/wk				-3.3 (-11.8, -5.2)	-0.55 (-1.39, 0.28) ^b	
CD with knee osteoarthritis	LI, 78 wk	Static on force platform	DLS EO	Postural sway distance (in 10 sec)	4 first trial was practice, but not disclosed		-8.70 (-17.30, -0.10) ^b	-0.87 (-1.75, 0.01) ^b	
		Dynamic	SLS EO	Time up to 10 sec			-0.05 (-0.13, 0.03) ^b	-0.29 (-0.76, 0.18) ^b	
Functionally limited, CD	LI, 10 wk	Static	FICSIT 3	Time up to 10 sec, score 1-6	NR	26 (CI -17, 69)	-0.18 (-0.29, -0.07) ^b	-0.76 (-1.24, -0.27) ^b	
		Dynamic	SLS EO	Time up to 10 sec			1.00 (0.07, 1.93)	0.51 (0.03, 0.98)	
Healthy, CD	HI, 24 wk	Dynamic	BTW 3 d/wk	Time over 6 m	Best of 3		-7.70 (-13.27, -2.13) ^b	-1.16 (-2.04, -0.27)	95
		Dynamic	BTW 1 d/wk				-3.3 (-11.8, -5.2)	-0.55 (-1.39, 0.28) ^b	
CD with knee osteoarthritis	LI, 78 wk	Static on force platform	DLS EO	Postural sway distance (in 10 sec)	4 first trial was practice, but not disclosed		-8.70 (-17.30, -0.10) ^b	-0.87 (-1.75, 0.01) ^b	
		Dynamic	SLS EO	Time up to 10 sec			-0.05 (-0.13, 0.03) ^b	-0.29 (-0.76, 0.18) ^b	
Functionally limited, CD	LI, 10 wk	Static	FICSIT 3	Time up to 10 sec, score 1-6	NR	26 (CI -17, 69)	-0.18 (-0.29, -0.07) ^b	-0.76 (-1.24, -0.27) ^b	
		Dynamic	SLS EO	Time up to 10 sec			1.00 (0.07, 1.93)	0.51 (0.03, 0.98)	
Healthy, CD	HI, 24 wk	Dynamic	BTW 3 d/wk	Time over 6 m	Best of 3		-7.70 (-13.27, -2.13) ^b	-1.16 (-2.04, -0.27)	95
		Dynamic	BTW 1 d/wk				-3.3 (-11.8, -5.2)	-0.55 (-1.39, 0.28) ^b	
CD with knee osteoarthritis	LI, 78 wk	Static on force platform	DLS EO	Postural sway distance (in 10 sec)	4 first trial was practice, but not disclosed		-8.70 (-17.30, -0.10) ^b	-0.87 (-1.75, 0.01) ^b	
		Dynamic	SLS EO	Time up to 10 sec			-0.05 (-0.13, 0.03) ^b	-0.29 (-0.76, 0.18) ^b	
Functionally limited, CD	LI, 10 wk	Static	FICSIT 3	Time up to 10 sec, score 1-6	NR	26 (CI -17, 69)	-0.18 (-0.29, -0.07) ^b	-0.76 (-1.24, -0.27) ^b	
		Dynamic	SLS EO	Time up to 10 sec			1.00 (0.07, 1.93)	0.51 (0.03, 0.98)	

Continued next page

Table IV. Contd

Subjects	PRT intensity and duration	Balance outcome characteristics				Balance results			Reference	
		balance type	balance test	measure (max time, distance, or score)	trials allowed	% change \pm SD in PRT group ^a	significant between-group effect	WMD (95% CI)		effect size (95% CI)
Healthy, CD, moderately active	HI, 8 wk	Static	SLS EC	Time up to 30 sec	5 (NR whether best or mean time was recorded)	1		-0.12 (-3.01, 2.77) ^b	-0.04 (-0.87, 0.80) ^b	103
Healthy, CD	HI, 10 wk	Dynamic	Functional reach	Distance (cm)	NR	12.7 \pm 3.1	p < 0.003	3.60 (0.43, 6.77)	0.71 (0.07, 1.35)	21
Frail, mobility-limited	HI, 10 wk	Functional	Berg Balance Test	Score 0-56	NR	2 (CI -10, 14)	p = 0.001	3.10 (-4.77, 10.97)	0.22 (-0.34, 0.78)	30
Disabled women with CHD	HI, 26 wk	Functional	Balance and coordination domain of CS-PFP	Score 0-20	NR		p = 0.0001	14.5 (3.33, 25.67)	0.91 (0.19, 1.63)	20
Frail, institutionalized	LI, 52 wk	Functional	Berg Balance Test	Score 0-56	NR			-5.30 (-10.66, 21.26) ^b	0.30 (-0.59, 1.19) [authors report 0.32 [90% CI -0.09, 0.74)]	98
Frail, CD	MI, 10 wk	Functional	Berg Balance Test	Score 0-56	NR			-2.00 (-4.78, 0.78)	-0.19 (-0.45, 0.07)	92
Low level of physical function, CD	HI, 16 wk	Functional	Balance and coordination domain of CS-PFP	Score 0-20	NR			-0.40 (-12.25, 11.45)	-0.03 (-0.83, 0.78)	31
Mobility-limited, CD	LI (PT), 16 wk	Static	Balance and coordination domain of CS-PFP	Score 0-20	NR		Yes, p NR	10.1 (-0.92, 21.12)	0.75 (-0.08, 1.58)	
Healthy, CD	MI, 104 wk	Dynamic	Functional reach	Distance (cm)	NR	50		-1.98 (-6.04, 2.08) ^b	-0.44 (-1.33, 0.45) ^b	11
								0.40 (-3.63, 4.43)	0.06 (-0.53, 0.65)	99

Continued next page

Table IV. Contd

Subjects	PRT intensity and duration	Balance outcome characteristics				Balance results			Reference	
		balance type	balance test	measure (max time, distance, or score)	trials allowed	% change \pm SD in PRT group ^a	significant between-group effect	WMD (95% CI)		effect size (95% CI)
Healthy, CD with low bone mass	HI, 25 wk	Static	Postural sway (standing on foam)	Time up to 30 sec	1	-30.6	$p \leq 0.05$	-70.90 (-120.41, -21.39) ^b	-0.71 (-1.21, -0.20) ^b	25
Frail, functionally limited, CD	LI, 10 wk	Static	FICSIT 3 (HG)	Time up to 10 sec	NR			0.20 (-0.62, -1.02)	0.17 (-0.51, 0.85)	93
			FICSIT 3 (MG)					0.30 (-0.09, 0.69)	0.32 (-0.37, 1.00)	
			Tandem stand (HG)					-1.90 (-5.06, 1.26)	-0.42 (-1.11, 0.27)	
			Tandem stand (MG)					0.50 (-2.33, 3.33)	0.12 (-0.56, 0.81)	
Healthy, CD	MI, 12 wk	Functional	Balance and coordination subscale of ADAP	Score 1-5	1		$p = 0.05$	4.20 (-1.28, 9.68)	0.41 (-0.13, 0.95)	24
Healthy, active, CD	MI, 12 wk	Dynamic	Functional reach	Distance (cm)	NR	9.4	$p < 0.001$, $F = 39.23$	8.10 (2.19, 14.01)	1.23 (0.28, 2.19)	29
Healthy, CD	LI (PT), 10 wk	PG (Chattecx)	Balance index	Time up to 30 sec	Best of 3 if LOB occurred	-10.8 \pm 24.7	$p = 0.012$	-6.60 (-14.26, 1.06) ^b	-0.48 (-1.03, 0.08) ^b	32
			LOB			-13.0 \pm 24.7		-0.40 (-1.16, 0.36) ^b	-0.29 (-0.84, 0.26) ^b	
			Balance index			-2.1 \pm 10.4		2.20 (-4.51, 8.91)	0.18 (-0.37, 0.73)	
	MI (PT), 10 wk		LOB							
			Balance index			-2.1 \pm 22.4		0.30 (-0.40, 1.00)	0.24 (-0.31, 0.79)	
	HI (PT), 10 wk		LOB			-0.3 \pm 9.6		3.20 (-5.04, 11.44)	0.22 (-0.34, 0.77)	
			Balance Index							
Healthy, CD	HI, 8 wk	Static	LOB	Time over 6 m	NR	0.1 (33.1)%		0.30 (-0.56, 1.16)	0.19 (-0.36, 0.74)	101
			BTW					0.20 (-17.72, 18.12)	0.01 (-0.60, 0.61)	
		Dynamic	Functional reach	Distance (cm)	NR			7.10 (-10.90, 25.16)	0.24 (-0.37, 0.85)	

Continued next page

Table IV. Contd

Subjects	PRT intensity and duration	Balance outcome characteristics			Balance results		Reference		
		balance type	balance test	measure (max time, distance, or score)	trials allowed	% change ± SD in PRT group ^a		significant between-group effect	WMD (95% CI)
	MI (PT), 8 wk	Static	BTW	Time over 6 m	NR		0.30 (-18.42, 17.82)	-0.01 (-0.61, 0.59)	
		Dynamic	Functional reach	Distance (cm)	NR		5.70 (-12.43, 23.83)	0.19 (-0.41, 0.79)	

a Percentage change values are only presented if reported in studies.

b Although negative, sign indicates direction of change favours treatment.

ADAP = Assessment of Daily Activity Performance; **AP** = anterior-posterior; **BTW** = backward tandem walk; **CD** = community-dwelling; **CHD** = congestive heart disease; **CS-PFP** = Continuous Scale-Physical Functional Performance; **DLS** = double-leg stand; **EC** = eyes closed; **EO** = eyes open; **F** = female; **FB** = feedback; **FICSIT** = Frailty and Injuries: Cooperative Studies of Intervention Techniques; **HG** = high-attention group; **HI** = high intensity; **LI** = low intensity; **LOB** = loss of balance; **LOS** = loss of stability; **max** = maximum; **MG** = moderate-attention group; **MI** = moderate intensity; **NR** = not reported; **PG** = posturography; **PT** = power training; **SLS** = single-leg stand; **SOT** = Sensory Organisation Test; **WMD** = weighted mean difference (difference in mean of the change score in the intervention and control group).

test. In the 29 studies reviewed, a total of 68 balance tests with sufficient data points were analysed.

Fourteen studies reported that the PRT group performed from 2% to 98% better than the control group in a balance outcome. Studies that included multiple balance tests may have shown significant improvement in one or some, but not every, balance test. Fifteen of 68 tests (22%) showed significantly improved balance performance following PRT in: single-leg stance (SLS) eyes open,^[27,33] or closed,^[23,28] double-leg stance eyes open^[27] or closed,^[26] postural sway on a compliant surface,^[25] backward tandem walk,^[22] functional reach,^[21,29] Berg Balance Test,^[30] balance component of the Continuous Scale-Physical Functional Performance (CS-PFP),^[20,31] balance component of the Assessment of Daily Activity Performance (ADAP)^[24] and combined postural sway and static balance measured by posturography.^[32] Two of these studies conducted power training.^[31,32] Figure 2 depicts the percentage of balance tests showing statistically significant improvements compared with the total number of balance tests within each of the four balance types. Figure 3 shows the number of tests showing statistically significant improvements compared with the total number of balance tests for each specific balance outcome. Five balance tests in three studies^[26,27,95] showed nonsignificant results, but moderate to high ES, suggesting clinically relevant outcomes. The possibility of a type II error, indicating that the studies were underpowered, cannot be discounted. Three studies examined the relationship(s) between a change in strength^[100] and/or power^[31,32] and a change in balance. No such relationships were detected in any of these studies.

2.2.1 Static Balance

Static balance tests (n = 27) were the most widely used tests, comprising 40% of the total number of balance tests performed. Most studies assessing this outcome used low-intensity training; two used moderate intensity^[11,23] and four studies employed high intensity.^[25,33,94,101] Participants came from both healthy and frail/clinical cohorts. Six of 29 studies (seven tests, 26% of static tests) showed significantly improved static balance performance. The tests were conducted over 10–60 seconds or for as long as possible,^[28] and one to five trials were permitted. In




Fig. 2. Balance outcomes stratified by balance type. Studies showing significant balance improvement in: static balance,^[23,25-28,33] dynamic balance,^[21,22,29] functional balance^[20,24,30,31] and computerized dynamic posturography.^[32]

the eyes-open condition, SLS and tandem stance (see table I), [predominantly over 10 seconds] were overall not significantly different between control and intervention groups. Two of eight SLS^[27,33] and one of five tandem-stance^[27] tests showed significantly improved times. These studies employed maximum stance times of 30 and 60 seconds, respectively. No improvements in standing balance over 10 seconds were observed with the FICSIT tests^[93,96] or in postural sway on a stable platform.^[26]

In the eyes-closed condition, two of five studies showed better SLS time in 57 younger, healthy participants^[23] and 18 functionally limited women.^[28] Postural sway improved on a stable platform with eyes closed^[26] and on a compliant surface such as foam.^[25] Most studies used low-intensity training; two used moderate-intensity^[11,23] and four studies employed high-intensity training.^[25,33,94,103] Participants came from both healthy and frail/clinical cohorts.

2.2.2 Dynamic Balance

Dynamic balance tests (n = 22) comprised 32% of the total balance tests performed. Most studies of dynamic balance used moderate-^[29,99,101] or high-intensity training on resistance machines;^[21,22,94,95,101] three studies used low-intensity training with elastic tubing.^[28,102,105] Participants were generally healthy or frail/mobility-limited.^[28,100,102] Three of 29 studies (three tests, 14% of dynamic tests) showed significantly improved balance performance. The distance covered in the tests ranged from 1.8 to 9.1 m (6 to 30 ft) and outcome

measures were fastest time or number of incorrect steps for all tests. Of the 13 timed walk tests (three forward tandem, eight backward tandem and two beam balance walks), only one study^[22] demonstrated improved backward tandem walk in 39 healthy older adults. Functional reach, by contrast, improved by 9% and 13% in two of nine trials with healthy participants.^[21,29] Most studies used high-intensity training on resistance machines; three studies used low-intensity training with elastic tubing.^[28,102,105]

2.2.3 Functional Balance

Functional balance tests (n = 7) comprised 10% of the total balance tests performed. Four of 29 studies (four tests, 57% of functional tests) showed significantly improved balance performance. Frail, mobility-limited elderly demonstrated significantly improved balance in the Berg Balance Test after high-intensity PRT using resistance machines.^[30] In comparison, other studies of frail, elderly cohorts^[92,98] showed no change in the Berg Balance Test after low and moderate PRT using free weights and other weighted equipment. Two of three trials using CS-PFP testing improved after training on resistance machines.^[20,31] It was the power training group, in the study of Miszko et al.,^[31] and not strength training group that improved in this test. Healthy older adults training at moderate intensity




Fig. 3. Improvements in specific balance outcomes. **ADAP** = Assessment of Daily Activity Performance; **CS-PFP** = Continuous Scale-Physical Functional Performance; **EC** = eyes closed; **EO** = eyes open.

with elastic tubing and free weights improved in the balance component of the ADAP, although no statistical value was reported.^[24]

2.2.4 Computerized Dynamic Posturography

Balance testing using computerized dynamic posturography comprised 8% of the total balance tests (n = 12) performed.^[32,33,47] Training intensity ranged from low to high. Training equipment included resistance machines^[32,33] and Thera-bands™.^[47] All participants were healthy, and community dwelling. One of three studies (one test, 8% of posturography tests) showed significantly improved balance performance after power training at low intensity.^[32]

2.2.5 Subgroup Analysis

We looked for patterns and factors that might explain the heterogeneous findings by stratifying the balance tests showing significant improvements by cohort, intensity of training, training duration, training equipment, quality of the papers and use of blinded assessors. Chi-square testing was also conducted to determine statistical difference between strata. The number of balance tests (and percentage) showing a significant improvement following training compared with the total number of tests for most factors are presented in figures 4–9. In summary, no clear trends emerged. Balance improved most in longer duration studies (figure 4). Chi-square analysis showed a significant difference between studies lasting 40–104 weeks compared with those of shorter durations ($\chi^2 = 6.00$). Analysis of cohort




Fig. 4. Balance outcomes stratified by training duration. Studies showing significant balance improvement in training duration of: ≤ 12 weeks,^[21,23,24,28–30,32,33] 14–26 weeks^[20,25,31] and 40–104 weeks.^[22,26,27] Note that some studies have more than one balance outcome. * indicates significantly different from shorter duration training periods ($\chi^2 = 6.00$).




Fig. 5. Balance outcomes stratified by cohort. Studies showing significant balance improvement in: healthy cohorts^[21–25,27,29,31–33] and frail/clinical cohorts.^[20,26,28,30] Note that some studies have more than one balance outcome.

(figure 5), training intensity (figure 6), quality assessment (figure 7), assessor blinding (figure 8) and training equipment (figure 9) revealed no significant distinction between strata ($\chi^2 = 0.27–2.99$). Static balance improved after low-intensity training in three poorer quality studies,^[26–28] after moderate-intensity training,^[23] and after high-intensity training in two better quality studies.^[25,33] Dynamic balance improved with high-^[21,22] and moderate-intensity training.^[29] The better quality studies (PEDro score ≥ 6), showing significant improvements in balance performance, were predominantly studies of healthy cohorts undertaking high-intensity, short-term (10–12 weeks) training with resistance machines.

3. Discussion

This is the first systematic review to examine the efficacy of RCTs of PRT alone on balance performance in older adults. Although PRT is often stated to be beneficial for balance, this review provides evidence that PRT as an isolated intervention is not uniformly effective in improving balance performance. Only 22% of results from the balance tests examined offered support for the efficacy of PRT as a single modality to improve balance. One factor that must be considered is that balance was only one of a number of outcomes explored in all the studies reviewed, and therefore many studies may not have been optimally designed to investigate change in balance performance alone nor sufficiently powered to find such an effect.

The inconsistent effect of PRT on measures of balance may be due to: the heterogeneity of cohorts




Fig. 6. Balance outcomes stratified by intensity of resistance training. Studies showing significant balance improvement in: high-intensity,^[20-22,25,30,33] moderate-intensity^[23,24,29] and low-intensity^[26-28,31,32] resistance training. Note that some studies have more than one balance outcome.

and balance measures; a wide disparity in the conduct of balance tests; diversity of equipment used for PRT; inadequate or ineffective dose of the training programme; variability in sample size; lack of statistical power to detect between-group differences; inadequate compliance with the exercise programmes; and/or differences in overall quality of studies. The greater number of trials may have produced a learning effect in participants,^[38] further confounding comparisons across studies. However, it is also possible that PRT alone is not a robust balance-enhancing intervention.

Static balance tests were the most frequently used tests, with SLS accounting for 40% of all tests, supporting the view of Wolfson et al.^[33] that SLS is the most frequently used balance test in studies involving older adults. Generally, balance measured by SLS and tandem stance in the eyes-open condition, did not respond to training. The studies demonstrating significantly improved SLS and tandem stance employed stance times of 30–45 and 60 seconds, respectively. Most studies conducted the test only for 10 seconds. The authors using the FICSIT-3 tests^[93,96] suggested that the lack of significant finding was because the parallel and semi-tandem stances were too easy for their cohorts. Ferrucci et al.^[14] shared a similar view, finding that >45% of disabled women could maintain the tandem stance for 10 seconds. The static balance protocols using relatively short maximum test times may produce a ceiling effect because the test is too short to detect imbalance thus missing subtle balance impairments.

The ceiling effect may also explain results in a cohort without overt balance impairment that performs well initially, and cannot improve further without training. The result is, thus, a negative outcome^[94] and underestimation in balance adaptation.^[39] A recent meta-analysis of SLS times^[39] reported mean stance times in the eyes-open condition for three age groups (60–69, 70–79 and 80–99 years) as 27.0, 17.2 and 8.5 seconds, respectively suggesting that maximum times of ≤ 15 seconds for cohorts younger than 80 years may not detect any change in this test.

Training also provided no benefit to SLS with eyes closed. The two studies reporting improvements in this test,^[23,28] however, contained methodological problems. In one study,^[28] data were non-normally distributed and no indication that statistical normalization of data had been attempted prior to use of parametric analytic techniques. Nichols et al.,^[23] trained 57 younger, healthy participants at moderate intensity on resistance machines. The SLS test (two practice plus three trials) was conducted over 45 seconds on each leg and then the times summed. It is possible a learning effect could have occurred with the ten trials for this test. Furthermore, one group trained progressively at equal concentric/eccentric load, the other at greater eccentric load, the progression of which was equipment-limited. The balance results of the two groups were then combined for analysis, compromising the evaluation of the effect of PRT on balance.

There is considerable variation in the reported methodology of static and dynamic tests and a lack




Fig. 7. Balance outcomes stratified by quality assessment. Studies showing balance improvement in: poorer quality (1–5)^[20,22,23,26-29,31] and better quality (6–10)^[21,24,25,30,32,33] Physiotherapy Evidence Database (PEDro) scores. Note that some studies have more than one balance outcome.




Fig. 8. Balance tests stratified by assessor concealment. Significant balance improvement in: studies using blinded assessor,^[21,24,25,33] studies not reporting use of blinded assessor,^[20,22,23,26,28-30,101] and studies not using blinded assessor.^[27,31,32] Note that some studies have more than one balance outcome. **NR** = not reported.

of specific description in some studies. Standardization of methodology and better reporting of procedures will ensure greater comparability of results in future studies. Because of the multifactorial causes of balance impairment^[18] using several types of balance tests to develop a balance 'profile' in future studies may impart a more robust determination of effect following intervention.

We inspected the 15 (seven static, three dynamic, four functional and one posturography) balance tests demonstrating significant improvements for the presence of trends. No single balance test demonstrated prominence over the other tests. However, 57% of the functional tests exhibited significant improvements, greater than the other balance types. This could be explained by the Berg Balance, CS-PFP and ADAP tests having well defined test protocols, using quantitative scales and having strong intra- and inter-rater reliability.^[10,42] One factor to take into consideration is that in one study using the CS-PFP,^[31] a 39% withdrawal rate in the power training group compared with an overall 22% study withdrawal may compromise the validity of this significant finding.

When we stratified the balance tests by cohort, the balance tests performed by the frail/clinical group showed more improved outcomes (27%) than the healthy group. This is not an unexpected outcome as those individuals with some co-morbidities (depending on their nature) and/or mobility/functional limitations may be more likely to respond to PRT and show greater relative improvements than

healthy cohorts. It is plausible that fewer improvements were evident in tests conducted in the healthy populations because of a ceiling effect. The balance tests selected may not have been the most appropriate ones chosen to adequately challenge postural stability in minimally impaired individuals.

We also stratified dose-response characteristics to examine whether intensity and/or duration may affect balance outcomes. In studies of high-intensity training, significant balance gains were observed in six tests, equally distributed between static, dynamic and functional balance types. These improvements in balance performance were predominantly observed in higher quality studies (four of six) of healthy cohorts undertaking high-intensity, short-term (10–12 weeks) training, using resistance machines. A standardized and quantifiable dose of training can be achieved by training on resistance machines. High-intensity training may best deliver the stimulus required to increase muscle strength and to elicit neuromuscular benefits to enhance balance. Recent evidence has shown that muscle activation capacity and calf muscle-tendon mechanical properties relevant to contractile force production are highly correlated to more demanding balance tasks (SLS and tandem stance).^[106] Studies are needed to further examine whether high-intensity PRT can amplify the muscle-tendon properties, and determine which dose will provide optimal gains in balance.

In studies of low-intensity training, significant balance gains were observed in six tests, four of




Fig. 9. Balance test stratified by training equipment. Studies showing significant balance improvement using: resistance training machines,^[22-25,29,30,32] free weights^[21,26] and combined equipment.^[20,24,27,29,33] Note that some studies have more than one balance outcome.

which were static balance measures. These improvements in balance performance were predominantly observed in lower-quality studies (five of six) of various cohorts undertaking low-intensity training with free weights, Therabands™, and other equipment. The data suggest that if low-intensity training improves balance, it may preferentially improve static balance.

Balance tests were more likely to improve in longer-term than shorter-term studies (33% vs 22%), suggesting a threshold of total training duration or volume may be required to elicit gains. However, there were relatively few long-term studies in this review and statistical analysis of data just reached a level of significance.

The lack of a pronounced effect of PRT on balance (in 78% of tests) in this review may also imply that strength is not the major underlying mechanism for poor balance, and that other limiting factors influence postural stability. Balance control is determined by an integrated network of vestibular, visual, cognitive, somatosensory and motor systems. A 'unique combination of constraints' for each elderly individual will govern their postural orientation and equilibrium.^[107] In response to postural challenge, an individual will slow the centre of mass by generating muscle torque at the ankle or hip or by taking a step. The selection of one of these three strategies will depend on the base of support, location of centre of mass, speed of perturbation and surface characteristics.^[107] The key muscle groups used to effect these balance strategies are ankle dorsiflexors and plantarflexors, knee extensors and flexors and hip abductor and adductors. Aging dampens reaction time and muscle strength, impairing, in some people, the ability to control a fall. In older adults, lateral stability is the key contributor to maintaining balance control.^[108] The muscle torque required to maintain balance may be greater than the force that can be generated by older muscle. Specifically, weakness of hip abductors has been shown to compromise the ability to maintain lateral stability during stepping and thus maintain balance.^[109,110] This may have serious implications for those at risk, such as fallers who have demonstrated greater lateral sway than non-fallers.^[111] Furthermore, hip fractures, the most serious fractures resulting from falls, may be partly attributed to deficits in lateral stability,

resulting in a fall onto the greater trochanter specifically.^[112,113] Training of hip musculature has, for the most part, not been carried out. In the studies reviewed, eight studies^[21,24,26,28,94,100,102,104] (28%) trained hip abductor muscles, four of which showed improved balance.^[21,24,26,28] Rather than prescribing universal whole-body or lower-extremity strength training, it may be prudent to focus on specific muscles critical to balance such as hip abductors and adductors, knee flexors and extensors and ankle plantarflexors and dorsiflexors.

Muscle power generation may also be a limiting factor in the control of balance. The age-related decline in neural processing can diminish the ability for rapid force development necessary to respond to postural challenge.^[108,109,114] Healthy, community-dwelling, older women have recorded a 40% decline in the rate of force development in hip abduction and abduction compared with younger women, which may have a more marked effect on postural challenges than muscle strength.^[109] Chang and colleagues^[115] have documented the relationship between rate of force development of hip abductors and mediolateral stability in 30 community-dwelling older adults. Hip abductor rate of force development was a stronger predictor of compensatory stepping than centre of pressure displacement (sway). We have previously shown that muscle power is related to balance in non-frail older adults^[19] and obese, older adults with type 2 diabetes.^[116] In this review, 18% of studies (n = 4)^[11,31,32,101] used power training, two of which showed improved balance.^[31,32]

Because of the association between muscle weakness and falls or poor balance, most studies have focused on one adaptation to PRT, specifically the increase in muscle strength, as a mechanism for the improvement to balance control. That is, PRT increases muscle strength thereby increasing the force that muscles can generate in response to a loss of balance. Yet few studies in this review have tested this assumption by examining the relationship between the change in strength and the change in balance. In the two studies that have examined this relationship in healthy older adults, gains in strength were not associated with improved balance.^[32,33] Alternatively, other adaptations provided by PRT (e.g. increased neural drive to agonist muscles,^[114]

increased motor unit recruitment and activation,^[114] improved cognition^[117-119] decreased depression,^[120] reduced antagonist muscle co-contraction,^[114] enhanced stability of muscle co-ordination^[121] and improved force control^[114]) could also explain the efficacy of PRT on balance.

A potential limitation of this review is that we did not use abstracts, unpublished studies or thesis dissertations and a search/publication bias may have been introduced. We used only RCTs and did not include some well conducted dose-response studies that had fulfilled the criteria in all ways except that they lacked a randomized control group. This may have introduced a detection bias, where valid results have not been reported.

3.1 Implications for Research

Future directions to improve the empirical knowledge in this discipline would be to:

- employ robust RCT study designs;
- standardize the methodology of static and dynamic balance tests in order to make comparisons between studies more meaningful;
- assess a balance profile of the individual with a number of different tests to pinpoint weaknesses or deficits to individualize the intervention and optimize outcome;
- assess balance when delivering an unpredictable perturbation;
- conduct more studies using preclinical and frail populations and cohorts with multiple co-morbidities;
- investigate dose-response effects of PRT on balance performance;
- examine the effect of various intensities of training on static balance and to further define neuromuscular and other mechanisms that may underlie changes in balance with PRT.

More robust studies of longer duration are needed to confirm the preliminary evidence that balance improves when training takes place over longer periods. In addition, future studies should investigate the potential benefits of using resistance training to target muscle groups and to elicit neural adaptations specific to balance control. To this end, the merits of hip abductor and adductors strengthening as part of any exercise regimen aimed at improv-

ing lateral balance and reducing falls, and use of the less conventional power training, should be considered.

4. Conclusion

The results of this review suggest that PRT as an isolated intervention has not to date been consistently shown to improve balance performance in older adults. However, we found a wide disparity in the methodology of the static and dynamic outcomes assessed in these trials. Better reporting of the procedures in studies and regulation of the balance tests is recommended. Clinically relevant outcomes may have been overlooked due to the presence of type II errors in some studies. Nevertheless, before PRT is discounted as an efficacious intervention in the treatment of postural instability, future studies should examine possible relationships between accompanying adaptations to PRT and balance improvements that may provide further insights to the mechanism(s) by which PRT affects balance. At this time, recommendation to use PRT as an isolated intervention strategy for balance enhancement in an elderly cohort cannot be made based on the limited evidence presented in currently published data.

Acknowledgements

No sources of funding were used to assist in the preparation of this review. The authors have no conflicts of interest that are directly relevant to the content of this review.

References

1. Rubenstein LZ. Falls in older people: epidemiology, risk factors and strategies for prevention. *Age Ageing* 2006; 35 Suppl. 2: ii37-41
2. de Rekeneire N, Visser M, Peila R, et al. Is a fall just a fall: correlates of falling in healthy older persons – the Health, Aging and Body Composition study. *J Am Geriatr Soc* 2003; 51 (6): 841-6
3. Moreland JD, Richardson JA, Goldsmith CH, et al. Muscle weakness and falls in older adults: a systematic review and meta-analysis. *J Am Geriatr Soc* 2004; 52 (7): 1121-9
4. Lord SR, Ward JA, Williams P, et al. Physiological factors associated with falls in older community-dwelling women. *J Am Geriatr Soc* 1994; 42: 1110-7
5. Whipple RH, Wolfson LI, Amerman PM. The relationship of knee and ankle weakness to falls in nursing home residents: an isokinetic study. *J Am Geriatr Soc* 1987; 35 (1): 13-20
6. Lord SR, Clark RD, Webster IW. Postural stability and associated physiological factors in a population of aged persons. *J Gerontol A Biol Sci Med Sci* 1991; 46 (3): M69-76
7. Studenski S, Duncan PW, Chandler J. Postural responses and effector factors in persons with unexplained falls: results and methodologic issues. *J Am Geriatr Soc* 1991; 39 (3): 229-34

8. Shumway-Cook A, Gruber W, Baldwin M, et al. The effect of multidimensional exercises on balance, mobility, and fall risk in community-dwelling older adults. *Phys Ther* 1997; 77 (1): 46-57
9. Robbins AS, Rubenstein LZ, Josephson KR, et al. Predictors of falls among elderly people: results of two population-based studies. *Arch Intern Med* 1989; 149 (7): 1628-33
10. Berg KO, Wood-Dauphinee SL, Williams JI, et al. Measuring balance in the elderly: validation of an instrument. *Can J Pub Health* 1992; 83 Suppl. 2: S7-11
11. Bean JF, Herman S, Kiely DK, et al. Increased velocity exercise specific to task (InVest) training: a pilot study exploring effects on leg power, balance, and mobility in community-dwelling older women. *J Am Geriatr Soc* 2004; 52 (5): 799-804
12. Rantanen T, Guralnik JM, Ferrucci L, et al. Coimpairments as predictors of severe walking disability in older women. *J Am Geriatr Soc* 2001; 49 (1): 21-7
13. Reeves ND, Narici MV, Maganaris CN. Myotendinous plasticity to ageing and resistance exercise in humans. *Exp Physiol* 2006; 91 (3): 483-98
14. Ferrucci L, Guralnik JM, Buchner D, et al. Departures from linearity in the relationship between measures of muscular strength and physical performance of the lower extremities: the Women's Health and Aging study. *J Gerontol A Bio Sci Med Sci* 1997; 52A (5): M275
15. Wolfson L, Judge J, Whipple R, et al. Strength is a major factor in balance, gait, and the occurrence of falls. *J Gerontol A Bio Sci Med Sci* 1995; 50A (special issue): 64-7
16. Petrella JK, Miller LS, Cress ME. Leg extensor power, cognition, and functional performance in independent and marginally dependent older adults. *Age Ageing* 2004; 33 (4): 342-8
17. Skelton DA, Kennedy J, Rutherford OM. Explosive power and asymmetry in leg muscle function in frequent fallers and non-fallers aged over 65. *Age Ageing* 2002; 31: 119-25
18. Petrella JK, Kim J-S, Tuggle SC, et al. Age differences in knee extension power, contractile velocity, and fatigability. *J Appl Physiol* 2005; 98 (1): 211-20
19. Orr R, De Vos N, Singh N, et al. Novel relationships of balance to muscle power and mental health in healthy older adults [abstract]. *J Am Geriatr Soc* 2004; 52 Suppl. 4: S17
20. Ades PA, Savage PD, Cress ME, et al. Resistance training on physical performance in disabled older female cardiac patients. *Med Sci Sport Ex* 2003; 35 (8): 1265-70
21. Barrett CJ, Smerdely P. A comparison of community-based resistance exercise and flexibility exercise for seniors. *Aust J Physiother* 2002; 48 (3): 215-9
22. Nelson ME, Fiatarone MA, Morganti CM, et al. Effects of high-intensity strength training on multiple risk factors for osteoporotic fractures: a randomized controlled trial. *JAMA* 1994; 272 (24): 1909-14
23. Nichols JF, Hitzelberger LM, Sherman JG, et al. Effects of resistance training on muscular strength and functional abilities of community-dwelling older adults. *J Aging Phys Act* 1995; 3 (3): 238-50
24. de Vreede PL, Samson MM, van Meeteren NL, et al. Functional tasks exercise versus resistance exercise to improve daily function in older women: a feasibility study. *Arch Phys Med Rehabil* 2004; 85 (12): 1952-61
25. Liu-Ambrose T, Khan KM, Eng JJ, et al. Resistance and agility training reduce fall risk in women aged 75 to 85 with low bone mass: a 6-month randomized, controlled trial. *J Am Geriatr Soc* 2004; 52 (5): 657-65
26. Messier SP, Royer TD, Craven TE, et al. Long-term exercise and its effect on balance in older, osteoarthritic adults: results from the fitness, arthritis, and seniors trial (FAST). *J Am Geriatr Soc* 2000; 48 (2): 131-8
27. Rooks DS, Kiel DP, Parsons C, et al. Self-paced resistance training and walking exercise in community-dwelling older adults: effects on neuromotor performance. *J Gerontol A Biol Sci Med Sci* 1997; 52A (3): M161-8
28. Skelton DA, McLaughlin AW. Training functional ability in old age. *Physiother* 1996; 82 (3): 159-67
29. Sousa N, Sampaio J. Effects of progressive strength training on the performance of the functional reach test and the Timed Get-up-and-Go Test in an elderly population from the rural north of Portugal. *Am J Human Biol* 2005; 17 (6): 746-51
30. Timonen L, Rantanen T, Ryyanen O-P, et al. A randomized controlled trial of rehabilitation after hospitalization in frail older women: effects on strength, balance and mobility. *Scand J Med Sci Sports* 2002; 12 (3): 186-92
31. Miszko TA, Cress ME, Slade JM, et al. Effect of strength and power training on physical function in community-dwelling older adults. *J Gerontol A Biol Sci Med Sci* 2003; 58 (2): 171-5
32. Orr R, de Vos N, Singh N, et al. Power training improves balance in healthy older adults. *J Gerontol A Biol Sci Med Sci* 2006; 61A (1): 78-85
33. Wolfson L, Whipple R, Derby C, et al. Balance and strength training in older adults: intervention gains and tai chi maintenance. *J Am Geriatr Soc* 1996; 44 (5): 498-506
34. Latham NK, Bennett DA, Stretton CM, et al. Systematic review of progressive resistance strength training in older adults. *J Gerontol A Biol Sci Med Sci* 2004; 59 (1): 48-61
35. Province MA, Hadley EC, Hornbrook MC, et al. The effects of exercise on falls in elderly patients: a preplanned meta-analysis of the FICSIT trials – Frailty and Injuries: Cooperative Studies of Intervention Techniques. *JAMA* 1995; 273 (17): 1341-7
36. Rossiter-Fornoff JE, Wolf SL, Wolfson LI, et al. A cross-sectional validation study of the FICSIT common data base static balance measures – Frailty and Injuries: Cooperative Studies of Intervention Techniques. *J Gerontol A Biol Sci Med Sci* 1995; 50A (6): M291-7
37. Overstall PW. The use of balance training in elderly people with falls. *Rev Clin Gerontol* 2003; 13 (2): 153-61
38. Buchner DM, Cress ME, Wagner EH, et al. The Seattle FICSIT/MoveIt study: the effect of exercise on gait and balance in older adults. *J Am Geriatr Soc* 1993; 41 (3): 321-5
39. Bohannon RW. Single limb stance times: a descriptive meta-analysis of data from individuals at least 60 years of age. *Topics Geriatr Rehabil* 2006; 22 (1): 70-7
40. Duncan PW, Weiner DK, Chandler J, et al. Functional reach: a new clinical measure of balance. *J Gerontol A Biol Sci Med Sci* 1990; 45 (6): M192-7
41. Berg KO, Maki BE, Williams JI, et al. Clinical and laboratory measures of postural balance in an elderly population. *Arch Phys Med Rehabil* 1992; 73 (11): 1073-80
42. Cress ME, Buchner DM, Questad KA, et al. Continuous-scale physical functional performance in healthy older adults: a validation study. *Arch Phys Med Rehabil* 1996; 77 (12): 1243-50
43. Baloh RW, Fife TD, Zwergling L, et al. Comparison of static and dynamic posturography in young and older normal people. *J Am Geriatr Soc* 1994; 42: 405-12
44. Mattacola CG, Lebsack DA, Perrin DH. Intertester reliability of assessing postural sway using the Chattecx balance system. *J Athl Train* 1995; 30 (3): 237-42
45. Wolfson L, Whipple R, Derby CA, et al. A dynamic posturography study of balance in healthy elderly. *Neurology* 1992; 42: 2069-75
46. Ford-Smith CD, Wyman JF, Elswick Jr RK, et al. Test-retest reliability of the sensory organisation test in noninstitutionalised older adults. *Arch Phys Med Rehabil* 1995; 76: 77-81
47. Topp R, Mikesky A, Dayhoff NE, et al. Effect of resistance training on strength, postural control, and gait velocity among older adults. *Clin Nurs Res* 1996; 5 (4): 407-27

48. Middel B, van Sonderen E. Statistical significant change versus relevant or important change in (quasi) experimental design: some conceptual and methodological problems in estimating magnitude of intervention-related change in health services research. *Int J Integr Care* 2002; 2 (17): 1-18
49. Hedges L, Olkin I. *Statistical Methods for meta-analysis*. San Diego (CA): Academic Press, 1985
50. Higgins JPT, Green S. *Analysing and presenting results. Cochrane Handbook for Systematic Reviews of Interventions 4.2.5* [updated May 2005]; section 8. Chichester: John Wiley & Sons, Ltd, 2005 [online]. Available from URL: <http://www.Cochrane.Org/resources/handbook/hbook.htm> [Accessed 2006 Jun 3]
51. Erdfelder E, Faul F, Buchner A. Gpower: a general power analysis program. *Behav Res Meth Instr Comp* 1996; 28: 1-11
52. Maher CG, Sherrington C, Herbert R, et al. Reliability of the Pedro scale for rating quality of randomized controlled trials. *Phys Ther* 2003; 83 (8): 713-21
53. Jadad AR, Moore RA, Carroll D, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials* 1996; 17 (1): 1-12
54. Verhagen A, de Vet H, de Bie R, et al. The Delphi list: a criteria list for quality assessment of randomized clinical trials for conducting systematic reviews developed by Delphi Consensus. *J Clin Epidemiol* 1998; 51 (12): 1235-41
55. Alexander NB, Galecki AT, Grenier ML, et al. Task-specific resistance training to improve the ability of activities of daily living-impaired older adults to rise from a bed and from a chair. *J Am Geriatr Soc* 2001; 49 (11): 1418-27
56. Binder EF, Brown M, Sinacore DR, et al. Effects of extended outpatient rehabilitation after hip fracture: a randomized controlled trial. *JAMA* 2004; 292 (7): 837-46
57. Carter ND, Khan KM, McKay HA, et al. Community-based exercise program reduces risk factors for falls in 65- to 75-year-old women with osteoporosis: randomized controlled trial [published erratum appears in *CMAJ* 2003 Jan 21; 168 (2): 152]. *CMAJ* 2002; 167 (9): 997-1004
58. Carter ND, Khan KM, Petit MA, et al. Results of a 10 week community based strength and balance training programme to reduce fall risk factors: a randomised controlled trial in 65-75 year old women with osteoporosis. *Br J Sports Med* 2001; 35 (5): 348-51
59. Day L, Fildes B, Gordon I, et al. Randomised factorial trial of falls prevention among older people living in their own homes. *BMJ* 2002; 325 (7356): 128
60. Donald IP, Pitt K, Armstrong E, et al. Preventing falls on an elderly care rehabilitation ward. *Clin Rehabil* 2000; 14 (2): 178-85
61. Latham NK, Stretton CM, Ronald M. Progressive resistance strength training in hospitalised older people: a preliminary investigation. *NZ J Physiother* 2001; 29 (2): 41-8
62. Lord SR, Castell S. Physical activity program for older persons: effect on balance, strength, neuromuscular control, and reaction time. *Arch Phys Med Rehabil* 1994; 75 (6): 648-52
63. Lord SR, Castell S, Corcoran J, et al. The effect of group exercise on physical functioning and falls in frail older people living in retirement villages: a randomized, controlled trial. *J Am Geriatr Soc* 2003; 51 (12): 1685-92
64. Lord SR, Ward JA, Williams P. Exercise effect of dynamic stability in older women: a randomized controlled trial. *Arch Phys Med Rehabil* 1996; 77 (3): 232-6
65. Lord SR, Ward JA, Williams P, et al. The effect of a 12-month exercise trial on balance, strength, and falls in older women: a randomized controlled trial. *J Am Geriatr Soc* 1995; 43 (11): 1198-206
66. Peterson MGE, Ganz SB, Allegrante JP, et al. High-intensity exercise training following hip fracture. *Topics Geriatr Rehabil* 2004; 20 (4): 273-84
67. Ramsbottom R, Ambler A, Potter J, et al. The effect of 6 months training on leg power, balance, and functional mobility of independently living adults over 70 years old. *J Aging Phys Act* 2004; 12 (4): 497-510
68. Siebens H, Aronow H, Edwards D, et al. A randomized controlled trial of exercise to improve outcomes of acute hospitalization in older adults. *J Am Geriatr Soc* 2000; 48 (12): 1545-52
69. Skelton D, Dinan S, Campbell M, et al. Tailored group exercise (falls management exercise - FaME) reduces falls in community-dwelling older frequent fallers (an RCT). *Age Ageing* 2005; 34 (6): 636-9
70. Toulotte C, Fabre C, Dangremont B, et al. Effects of physical training on the physical capacity of frail, demented patients with a history of falling: a randomised controlled trial. *Age Ageing* 2003; 32 (1): 67-73
71. Lichtenstein MJ, Shields SL, Shiavi RG, et al. Exercise and balance in aged women: a pilot controlled clinical trial. *Arch Phys Med Rehabil* 1989; 70 (2): 138-43
72. Lindemann U, Rupp K, Mueche R, et al. Improving balance by improving motor skills. *Z Gerontol Geriatr* 2004; 37 (1): 20-6
73. Morgan RO, Virnig BA, Duque M, et al. Low-intensity exercise and reduction of the risk for falls among at-risk elders. *J Gerontol A Biol Sci Med Sci* 2004; 59A (10): 1062-7
74. Ourania M, Yvoni H, Christos K, et al. Effects of a physical activity program: the study of selected physical abilities among elderly women. *J Gerontol Nurs* 2003; 29 (7): 50-5
75. Sherrington C, Lord SR, Herbert RD. A randomized controlled trial of weight-bearing versus non-weight-bearing exercise for improving physical ability after usual care for hip fracture. *Arch Phys Med Rehabil* 2004; 85 (5): 710-6
76. McMurdo MET, Johnstone R. A randomized controlled trial of a home exercise programme for elderly people with poor mobility. *Age Ageing* 1995; 24: 425-8
77. Capodaglio P, Capodaglio EM, Ferri A, et al. Muscle function and functional ability improves more in community-dwelling older women with a mixed-strength training programme. *Age Ageing* 2005; 34 (2): 141-7
78. Henwood TR, Taaffe DR. Improved physical performance in older adults undertaking a short-term programme of high-velocity resistance training. *Gerontology* 2005; 51 (2): 108-15
79. Hess JA, Woollacott M. Effect of high-intensity strength-training on functional measures of balance ability in balance-impaired older adults. *J Manipulative Physiol Ther* 2005; 28 (8): 582-90
80. LaStayo PC, Ewy GA, Pierotti DD, et al. The positive effects of negative work: increased muscle strength and decreased fall risk in a frail elderly population. *J Gerontol A Biol Sci Med Sci* 2003; 58A (5): 419-24
81. Robitaille Y, Laforest S, Fournier M, et al. Moving forward in fall prevention: an intervention to improve balance among older adults in real-world settings. *Am J Public Health* 2005; 95 (11): 2049-56
82. Ryushi T, Kumagai K, Hayase H, et al. Effect of resistive knee extension training on postural control measures in middle aged and elderly persons. *Physiol Anthropol Appl Human Sci* 2000; 19 (3): 143-9
83. Shaw JM, Snow CM. Weighted vest exercise improves indices of fall risk in older women. *J Gerontol A Biol Sci Med Sci* 1998; 53 (1): M53-8
84. Mobily KE, Mobily PR, Lane BK, et al. Using progressive resistance training as an intervention with older adults. *Ther Recreation J* 1998; 32: 42-53
85. Holviala JHS, Sallinen JM, Kraemer WJ, et al. Effect's of strength training on muscle strength characteristics, functional capabilities, and balance in middle-aged and older women. *J Strength Cond Res* 2006; 20 (2): 336-44
86. Mangione KK, Craik RL, Tomlinson SS, et al. Can elderly patients who have had a hip fracture perform moderate- to

- high-intensity exercise at home? *Phys Ther* 2005; 85 (8): 727-39
87. Simons R, Anel R. The effects of resistance training and walking on functional fitness in advanced old age. *J Aging Health* 2006; 18 (1): 91-105
 88. Bruhn S, Kullmann N, Gollhofer A. The effects of a sensorimotor training and a strength training on postural stabilisation, maximum isometric contraction and jump performance. *Int J Sport Med* 2004; 25 (1): 56-60
 89. Brochu M, Savage P, Lee M, et al. Effects of resistance training on physical function in older disabled women with coronary heart disease. *J Appl Physiol* 2002; 92 (2): 672-8
 90. Judge JO, Whipple RH, Wolfson LI. Effects of resistive and balance exercises on isokinetic strength in older persons. *J Am Geriatr Soc* 1994; 42 (9): 937-46
 91. Rooks DS, Ransil BJ, Hayes WC. Self-paced exercise and neuromotor performance in community dwelling older adults. *J Aging Phys Act* 1997; 5 (2): 135-49
 92. Latham NK, Anderson CS, Lee A, et al. A randomized, controlled trial of quadriceps resistance exercise and vitamin D in frail older people: the frailty interventions trial in elderly subjects (FITNESS). *J Am Geriatr Soc* 2003; 51 (3): 291-9
 93. Boshuizen HC, Stemmerik L, Westhoff MH, et al. The effects of physical therapists' guidance on improvement in a strength-training program for the frail elderly. *J Aging Phys Act* 2005; 13 (1): 5-22
 94. Buchner DM, Cress ME, de Lateur BJ, et al. The effect of strength and endurance training on gait, balance, fall risk, and health services use in community-living older adults. *J Gerontol A Biol Sci Med Sci* 1997; 52A (4): M218-24
 95. Taafe DR, Duret C, Wheeler S, et al. Once-weekly resistance exercise improves muscle strength and neuromuscular performance in older adults. *J Am Geriatr Soc* 1999; 47 (10): 1208-14
 96. Westhoff MH, Stemmerik L, Boshuizen HC. Effects of a low-intensity strength-training program on knee-extensor strength and functional ability of frail older people. *J Aging Phys Act* 2000; 8 (4): 325-42
 97. Ware JE, Sherbourne CD. The MOS 36-item Short-Form Health Survey (SF-36). *Med Care* 1992; 30 (6): 473-83
 98. Baum EE, Jarjoura D, Polen AE, et al. Effectiveness of a group exercise program in a long-term care facility: a randomized pilot trial. *J Am Med Dir Assoc* 2003; 4 (2): 74-80
 99. Brandon LJ, Boyette LW, Lloyd A, et al. Resistive training and long-term function in older adults. *J Aging Phys Activity* 2004; 12 (1): 10-28
 100. Chandler JM, Duncan PW, Kochersberger G, et al. Is lower extremity strength gain associated with improvement in physical performance and disability in frail, community-dwelling elders? *Arch Phys Med Rehabil* 1998; 79 (1): 24-30
 101. Henwood TR, Taafe DR. Short-term resistance training and the older adult: the effect of varied programmes for the enhancement of muscle strength and functional performance. *Clin Physiol Funct Imaging* 2006; 26 (5): 305-13
 102. Jette AM, Lachman M, Giorgetti MM, et al. Exercise – it's never too late: the strong-for-life program. *Am J Publ Health* 1999; 89 (1): 66-72
 103. Schlicht J, Camaione DN, Owen SV. Effect of intense strength training on standing balance, walking speed, and sit-to-stand performance in older adults. *J Gerontol A Biol Sci Med Sci* 2001; 56A (5): M281-6
 104. Skelton DA, Young A, Greig C, et al. Effects of resistance training on strength, power, and selected functional abilities of women aged 75 and older. *J Am Geriatr Soc* 1995; 43 (10): 1081-7
 105. Topp R, Mikesky A, Wigglesworth J, et al. The effect of a 12-week dynamic resistance strength training program on gait velocity and balance of older adults. *Gerontologist* 1993; 33 (4): 501-6
 106. Onambele GL, Narici MV, Maganaris CN. Calf muscle-tendon properties and postural balance in old age. *J Appl Physiol* 2006; 100 (6): 2048-56
 107. Horak FB. Postural orientation and equilibrium: what do we need to know about neural control of balance to prevent falls? *Age Ageing* 2006; 35 Suppl. 2: ii7-11
 108. Maki BE, McLroy WE. Control of rapid limb movements for balance recovery: age-related changes and implications for fall prevention. *Age Ageing* 2006; 35 Suppl. 2: ii12-8
 109. Johnson ME, Mille M-L, Martinez KM, et al. Age-related changes in hip abductor and adductor joint torques. *Arch Phys Med Rehabil* 2004; 85 (4): 593-7
 110. Rogers MW, Mille M-L. Lateral stability and falls in older people. *Exerc Sport Sci Rev* 2003; 31 (4): 182-7
 111. Lord SR, Rogers MW, Howland A, et al. Lateral stability, sensorimotor function and falls in older people. *J Am Geriatr Soc* 1999; 47 (9): 1077-81
 112. Nevitt MC, Cummings SR, Black D, et al. Type of fall and risk of hip and wrist fractures: the study of osteoporotic fractures. *J Am Geriatr Soc* 1993; 41 (11): 1226-34
 113. Campbell AJ, Borrie MJ, Spears GF. Risk factors for falls in a community-based prospective study of people 70 years and older. *J Gerontol A Biol Sci Med Sci* 1989; 44 (4): M112-7
 114. Barry BK, Carson RG. The consequences of resistance training for movement control in older adults. *J Gerontol A Bio Sci Med Sci* 2004; 59A (7): 730
 115. Chang S-HJ, Mercer VS, Giuliani CA, et al. Relationship between hip abductor rate of force development and mediolateral stability in older adults. *Arch Phys Med Rehabil* 2005; 86 (9): 1843-50
 116. Orr R, Tsang T, Lam P, et al. Mobility impairment in type 2 diabetes: association with muscle power and effect of tai chi intervention. *Diabetes Care* 2006; 29 (9): 2120-2
 117. Ozkaya GY, Aydin H, Toraman FN, et al. Effect of strength and endurance training on cognition in older people. *J Sport Sci Med* 2005; 4 (3): 300-13
 118. Heyn P, Abreu BC, Ottenbacher KJ. The effects of exercise training on elderly persons with cognitive impairment and dementia: a meta-analysis. *Arch Phys Med Rehabil* 2004; 85 (10): 1694-704
 119. Kramer AF, Erickson KI, Colcombe SJ. Exercise, cognition, and the aging brain. *J Appl Physiol* 2006; 101 (4): 1237-42
 120. Singh NA, Clements KM, Singh MAF. The efficacy of exercise as a long-term antidepressant in elderly subjects: a randomised, controlled trial. *J Gerontol A Biol Sci Med Sci* 2001; 56A (8): M497-504
 121. Carroll TJ, Barry B, Riek S, et al. Resistance training enhances the stability of sensorimotor coordination. *Proc Biol Sci* 2001; 268 (1464): 221-7

Correspondence: Rhonda Orr, PO Box 170, Lidcombe, NSW 2141, Australia.

E-mail: r.orr@usyd.edu.au