

Effort-Reward Imbalance, Work-Privacy Conflict, and Burnout Among Hospital Employees. — [Source link](#)

Nadine Häusler, Matthias Bopp, Oliver Hämmig

Institutions: University of Zurich

Published on: 01 Apr 2018 - Journal of Occupational and Environmental Medicine (Ovid Technologies (Wolters Kluwer Health))

Topics: Burnout and Educational attainment

Related papers:

- [Adverse health effects of high-effort/low-reward conditions.](#)
- [The measurement of effort-reward imbalance at work: European comparisons.](#)
- [Reviewing the effort-reward imbalance model: drawing up the balance of 45 empirical studies](#)
- [Balancing Effort and Rewards at University: Implications for Physical Health, Mental Health, and Academic Outcomes.](#)
- [Effects of effort-reward imbalance, job satisfaction, and work engagement on self-rated health among healthcare workers](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/effort-reward-imbalance-work-privacy-conflict-and-burnout-2am09rp26q>



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Effort-Reward Imbalance, Work-Privacy Conflict, and Burnout Among Hospital Employees

Häusler, Nadine ; Bopp, Matthias ; Hämmig, Oliver

Abstract: **OBJECTIVE** Studies investigating the relative importance of effort-reward imbalance and work-privacy conflict for burnout risk between professional groups in the health care sector are rare and analyses by educational attainment within professional groups are lacking. **METHODS** The study population consists of 1422 hospital employees in Switzerland. Multivariate linear regression analyses with standardized coefficients were performed for the overall study population and stratified for professional groups refined for educational attainment. **RESULTS** Work-privacy conflict is a strong predictor for burnout and more strongly associated with burnout than effort-reward imbalance in the overall study population and across all professional groups. Effort-reward imbalance only had a minor effect on burnout in tertiary-educated medical professionals. **CONCLUSION** Interventions aiming at increasing the compatibility of work and private life may substantially help to decrease burnout risk of professionals working in a health care setting.

DOI: <https://doi.org/10.1097/JOM.0000000000001287>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-165296>

Journal Article

Updated Version

Originally published at:

Häusler, Nadine; Bopp, Matthias; Hämmig, Oliver (2018). Effort-Reward Imbalance, Work-Privacy Conflict, and Burnout Among Hospital Employees. *Journal of Occupational and Environmental Medicine*, 60(4):e183-e187.

DOI: <https://doi.org/10.1097/JOM.0000000000001287>

Effort–Reward Imbalance, Work–Privacy Conflict, and Burnout Among Hospital Employees

Nadine Häusler, MSc, Matthias Bopp, PhD, MPH, and Oliver Hämmig, PhD, MPH

Objective: Studies investigating the relative importance of effort–reward imbalance and work–privacy conflict for burnout risk between professional groups in the health care sector are rare and analyses by educational attainment within professional groups are lacking. **Methods:** The study population consists of 1422 hospital employees in Switzerland. Multivariate linear regression analyses with standardized coefficients were performed for the overall study population and stratified for professional groups refined for educational attainment. **Results:** Work–privacy conflict is a strong predictor for burnout and more strongly associated with burnout than effort–reward imbalance in the overall study population and across all professional groups. Effort–reward imbalance only had a minor effect on burnout in tertiary-educated medical professionals. **Conclusion:** Interventions aiming at increasing the compatibility of work and private life may substantially help to decrease burnout risk of professionals working in a health care setting.

Keywords: burnout, effort–reward imbalance, health professionals, hospital employees, work–privacy conflict

Burnout is a psychological syndrome characterized by emotional exhaustion and fatigue.¹ It is a consequence of occupational stress and results from long-lasting adverse and emotionally demanding working conditions.¹ Risk factors for burnout in a work environment are high workload, low personal control, insufficient rewards, low social support, lack of reciprocity between efforts and rewards, conflict between individual and organizational values,² and shift work.³ Health care professionals face various occupational stressors^{3,4} that result in an elevated burnout risk compared with the general population.⁵ Several studies report high rates of burnout in nurses^{6–8} as well as physicians,^{9,10} whereas the burnout rate varies by occupational groups and specialty.^{7,10–12} Studies have identified different predictors of work stress and burnout between occupational groups^{13–15} and between educational levels within an occupational group.¹⁶ However, most studies on burnout in the health care setting focus on one occupational group and therefore miss the opportunity to directly compare health professionals despite inter-professional teamwork in a hospital. Moreover, literature investigating differences in burnout risk between occupational groups and taking into account educational attainment is widely lacking, despite uncontentious differences between and within occupational groups regarding working conditions, demands, and responsibilities.¹⁷ These in turn are important with respect to burnout risk. Future interventions could profit in the way that the programs can be better targeted to the needs and environments of these subgroups.

Two widely used work-related stress models, *effort–reward imbalance* (ERI) and *work–privacy conflict* (WPC), provide a theoretical framework to explain the development of occupational

stress and subsequently burnout. ERI postulates that occupational stress arises when there is a lack of reciprocity between effort and reward at the workplace, that is, high efforts spent and low reward received. This perceived imbalance generates stress and results in various adverse health effects.¹⁸ Whereas ERI focuses on the imbalance within the working situation, the WPC focuses on the inter-role conflict between work and private life.¹⁹ Stress resulting of these inter-role conflicts may spill over from work to private life or vice versa and can be understood as a result of the changing working conditions in Western societies over the past decades.²⁰ Both of these stress measures are predictive for burnout.^{2,8,12,20–22} Although ERI is a measure for work-specific stressors and hence is in theory expected to be stronger associated with burnout than WPC,² WPC was found to be a stronger predictor for burnout in studies directly comparing their effect on burnout risk among various occupations²⁰ as well as among health professionals.¹² Interestingly, WPC was not much stronger associated with burnout than ERI across all medical professions.¹²

Burnout among health care professionals does not only harm their health and well-being but is a threat to patients safety^{23,24} and the continuity of care²⁵ and diminishes quality of care.¹⁰ Hence, identifying whether the imbalance of effort and reward at work or the imbalance between work and private life predicts burnout for different medical occupations can lead to better targeted interventions to prevent burnout. We therefore seek to answer the following research questions:

1. Is ERI or WPC the stronger predictor for burnout risk in the health care setting?
2. Is the relative predictive importance of ERI and WPC with respect to burnout risk different between professional groups and educational levels in the health care setting?

METHODS

Data were gathered through an anonymous employee survey on working conditions and health among six hospitals and rehabilitation clinics in German-speaking Switzerland. The six hospitals and clinics varied by type (regional, cantonal and university hospital, rehabilitation clinic), size (workforces ranging from 473 to 2182 employees), degree of specialization, and urbanization of catchment area. All permanent employees ($n = 4497$) received a questionnaire containing 100 questions. They were asked to return it with the enclosed postage paid envelope. Three weeks after initial distribution, a reminder was sent to all employees. In total, 1844 questionnaires were returned resulting in a response rate of 41%. Compared with Switzerland as a whole, our sample has a larger proportion of women (85% compared with 75% in Switzerland), a similar share of physicians and nurses, whereas the share of tertiary educated nurses was lower in our sample.^{26,27} We restricted our study population to women ($n = 1570$) with complete information on age, sex, and professional group ($n = 1422$) due to the low number of men with complete information on all variables of interest ($n = 247$) and the expected effect modification by sex. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical

From the Epidemiology, Biostatistics and Prevention Institute, University of Zurich, Zurich, Switzerland.

Funding for this study was provided by Swiss National Accident Insurance Fund (SUVA), Federal Office of Public Health (BAG), State Secretariat for Economic Affairs (SECO).

The authors have no conflicts of interest.

Address correspondence to: Oliver Hämmig, PhD, MPH, University of Zurich, Hirschengraben 84, 8001 Zurich, Switzerland (oliver.haemmig@uzh.ch).

Copyright © 2018 American College of Occupational and Environmental Medicine

DOI: 10.1097/JOM.0000000000001287

standards. No approval by an ethics committee was required, as this study is based on survey data that were collected on a voluntary and anonymous basis.

Burnout

We measured burnout risk with the validated German version of the personal burnout scale¹⁹ from the Copenhagen Burnout Inventory.¹ This scale consists of six questions regarding the prolonged physical and psychological fatigue. Participants are required to indicate how frequently they experience these symptoms with five answer categories ranging from *always* (100), *often* (75), *sometimes* (50), *seldom* (25) to *never/almost never* (0). The burnout risk score is the average score of the items answered, whereas at least three items have to be answered in order to get a valid burnout risk score. A score of 50 and above points to being at an increased risk of having or developing a burnout. The average burnout score was 33 in the original Danish study, whereas the reference score of the German validation study is 42.²⁸

Effort–reward Imbalance

We assessed occupational stress with the long version of the validated German ERI scale. There are two subscales: *effort* contains six items and *reward* contains ten items, both on a 4-point Likert scale ranging from *I totally agree* to *I totally disagree*. Items were summed up for each subscale and the ratio was calculated by dividing the sum score of the effort subscale through the sum score of the reward subscale and multiplied by the correction factor 10/6 to adjust for the unequal number of items per subscale. A ratio over one indicates an unfavorable imbalance characterized by high efforts spent which are not met with corresponding rewards.

Work–privacy Conflict

We used the German translation of the WPC scale,¹⁹ which is an adapted version of the work–family conflict scale measuring inter-role conflict between work and private life²⁹ and is part of the Copenhagen Psychosocial Questionnaire. It measures solely the work to family spillover and consists of five items with five answer categories ranging from *I totally agree* (100), *I agree* (75), *neither agree nor disagree* (50), *I disagree* (25), to *I totally disagree* (0). The WPC score is the averaged sum score, whereas at least three items must have been answered in order to get a valid WPC score.

Social Support

The questions to assess social support stemmed from the Stress Study of the State Secretariat for Economic Affairs in 2000. It was asked whether one could rely on social support from partner, friends, relatives, supervisors, closest colleague at work, and other work colleagues when they face private problems and when they face problems at the workplace. The answer categories were “never,” “sometimes,” “often,” and “always.”

In consultation with the participating hospitals, we formed the following professional categories: *administrative staff*, *nurses*, *academic staff* including medical-therapeutic experts, and medical-technical experts as well as *physicians*. Educational level was assessed in three levels according to the Swiss Health Survey and the International Standard Classification of Education (ISCED). Whereas academic staff and physicians inherently have tertiary education, administrative staff and nurses can be split into non-tertiary (primary/secondary) and tertiary education. Hence, professional groups resulted in the following six categories: *nontertiary administrative staff*, *non-tertiary nurses*, *tertiary administrative staff*, *tertiary nurses*, *academic staff*, and *physicians* (See Table 1).

Descriptive statistics are used to describe the study population according to burnout risk. We used Pearson correlation to check for the multi-collinearity of ERI and WPC and performed multivariate linear regression analyses to assess the effects of ERI and

WPC on burnout risk while adjusting for age, marital status, and work volume. By using standardized coefficients (beta), we were able to compare the relative importance of the two predictors ERI and WPC. To check for the robustness of the effects, we included further explanatory variables such as hours of overtime per week, social support in case of private problems, and work problems from direct supervisor, closest colleague, and other colleagues in the sensitivity analyses. We performed all analyses using STATA 14.0 (StataCorp, College Station, TX).

RESULTS

In our female study population, tertiary educated staff is younger than nontertiary staff (Table 1). Most participants are married, whereas in tertiary-educated hospital employees, the share of married and single is about the same. Tertiary professions mostly work full-time, whereas in nontertiary professions, working part- and full time is more balanced. Most missings are found in ERI, especially in administrative staff regardless of educational level.

The unadjusted mean burnout risk score is lowest among the nontertiary professional groups and highest among physicians (Table 2). Across all professional groups, the burnout risk score is higher than the original Danish study, whereas only the burnout score for physicians is also higher than the German reference score. There is a clear positive association between ERI as well as WPC and burnout risk among all professional groups. Pearson correlation between ERI and WPC varied from 0.44 in academic staff to 0.58 in tertiary administrative staff. As the interaction term for ERI on one hand and WPC on the other hand was not significant, it was not included in multivariate linear regressions to keep the model simple and the interpretation straightforward.

In the overall study population, both ERI and WPC are significantly and positively associated with burnout risk, whereas the beta coefficient for WPC is much higher (Table 3). Likewise, across all stratified analyses for the professional groups, WPC is the stronger predictor for burnout risk. The largest effect is found for administrative staff with tertiary education followed by physicians and nontertiary nurses and the beta coefficients vary between 0.43 and 0.55. ERI only has a significant effect on burnout risk in tertiary nurses, academics, and physicians. The beta coefficient for ERI in the stratified analyses varies between 0.20 and 0.27 and is about one-fourth the size of the beta coefficient for WPC in the overall study population. The largest effect of ERI on burnout risk is found among physicians followed by tertiary administrative staff, academic staff, and tertiary nurses.

In the overall study sample, ERI and WPC as well as the control variables age, marital status, and work volume can explain 31% of the variance in burnout risk (Table 3). The highest adjusted R^2 was found for tertiary administrative staff and physicians. Results from the sensitivity analyses point to stable results. The additional explanatory variables for overtime and social support are not associated with burnout risk, with few exceptions: In administrative staff with tertiary education, ERI is significantly associated with burnout risk when adjusted for overtime as well as social support from direct supervisor and colleagues in case of work and private problems.

DISCUSSION

For both ERI and WPC, we found clear gradients according to burnout risk. Moreover, a strong and constant association between WPC and burnout risk among all professional groups is evident, whereas ERI only seems to be relevant with respect to burnout risk in tertiary-educated professional groups.

Despite the well-investigated relationships between WPC and burnout as well as ERI and burnout, to our knowledge, this is the first study to compare the relative importance of these measures for burnout in different professional groups taking into account educational attainment within a common health care

TABLE 1. Descriptive Statistics for Overall Study Population and Stratified for Professional Groups

	Overall Staff N (%)	Nontertiary Administrative Staff N (%)	Nontertiary Nurses N (%)	Tertiary Administrative Staff N (%)	Tertiary Nurses N (%)	Academic Staff N (%)	Physicians N (%)
Total	1,547 (100)	215 (100)	358 (100)	81 (100)	462 (100)	281 (100)	150 (100)
Age							
Under 25	107 (7)	23 (11)	42 (12)	0 (0)	33 (7)	8 (3)	1 (1)
25–34	429 (28)	26 (12)	61 (17)	15 (18)	170 (37)	101 (36)	56 (37)
35–44	385 (25)	44 (20)	84 (23)	22 (27)	105 (23)	75 (27)	55 (37)
45–54	394 (25)	73 (34)	97 (27)	28 (35)	112 (24)	59 (21)	25 (17)
55+	232 (15)	49 (23)	74 (21)	16 (20)	42 (9)	38 (13)	13 (9)
Marital status							
Married	749 (48)	111 (52)	181 (51)	38 (47)	214 (46)	129 (46)	73 (49)
Single	614 (40)	62 (29)	120 (34)	28 (35)	204 (44)	131 (47)	68 (45)
Divorced/widowed	182 (12)	42 (19)	54 (15)	15 (18)	42 (9)	20 (7)	9 (6)
Missing	2 (0)	0 (0)	0 (0)	0 (0)	2 (1)	0 (0)	0 (0)
Work volume							
100%	471 (31)	60 (28)	80 (22)	30 (37)	148 (32)	74 (26)	79 (53)
80–99%	400 (26)	49 (23)	76 (21)	24 (30)	139 (30)	86 (31)	27 (18)
51–79%	419 (27)	62 (29)	116 (33)	24 (30)	96 (21)	82 (29)	39 (26)
30–50%	206 (13)	36 (16)	65 (18)	3 (4)	68 (15)	31 (11)	3 (2)
29% and below	49 (3)	8 (4)	20 (6)	0 (0)	11 (2)	8 (3)	2 (1)
Missing	1 (0)	0 (0)	1 (0)	0 (0)	0 (0)	0 (0)	0 (0)
ERI							
Lowest quartile	364 (24)	48 (22)	87 (24)	18 (22)	115 (25)	67 (24)	37 (25)
2nd lowest quartile	356 (23)	47 (22)	78 (22)	19 (24)	112 (24)	71 (25)	36 (24)
2nd highest quartile	360 (23)	48 (22)	84 (24)	17 (21)	108 (23)	64 (23)	33 (22)
Highest quartile	356 (23)	47 (22)	79 (22)	17 (21)	106 (23)	65 (23)	33 (22)
Missing	111 (7)	25 (12)	30 (8)	10 (12)	21 (5)	14 (5)	11 (7)
WPC							
Lowest quartile	419 (27)	74 (34)	90 (25)	23 (28)	138 (30)	75 (27)	39 (26)
2nd lowest quartile	354 (23)	48 (22)	95 (27)	17 (21)	116 (25)	79 (28)	45 (30)
2nd highest quartile	452 (29)	47 (22)	98 (27)	23 (28)	95 (21)	58 (21)	32 (21)
Highest quartile	317 (21)	45 (21)	74 (21)	16 (20)	113 (24)	68 (24)	34 (23)
Missing	5 (0)	1 (1)	1 (0)	2 (2)	0 (0)	1 (0)	0 (0)

setting. The only previous study with the same focus¹² neglected to refine the professional groups for educational attainment and was based on very small numbers for most professional groups.

In contrast to a previous study reporting highest burnout risk for nurses,⁷ we also found the highest burnout risk among physicians.¹² Generally, we found higher burnout risk scores in professional groups with tertiary education. This could be a result of greater perceived work stress¹⁶ through higher demands and

expectations as well as more responsibilities imposed through their higher qualifications.^{30,31} However, only minor differences in burnout risk were found between nurses with different educational levels.³⁰ The high explained variance in the overall study population as well as across different professional groups strongly suggests burnout being a result of occupational stress¹ due to ERI¹⁸ and WPC.¹⁹ However, a previous study found that the association between ERI as well as WPC and burnout risk is not fully mediated

TABLE 2. Unadjusted Mean of Burnout Risk Score (0–100) incl. Standard Deviation by Levels of ERI and WPC, for Overall Study Population and Stratified for Professional Groups

	Overall Staff M (SD)	Nontertiary Administrative Staff M (SD)	Nontertiary Nurses M (SD)	Tertiary Administrative Staff M (SD)	Tertiary Nurses M (SD)	Academic Staff M (SD)	Physicians M (SD)
Total	39.8 (17)	37.7 (18)	38.6 (16)	39.1 (17)	40.2 (16)	40.4 (17)	43.0 (16)
ERI							
Lowest quartile	32.2 (14)	30.6 (15)	31.7 (15)	31.5 (11)	32.9 (12)	31.0 (14)	32.1 (13)
2nd lowest quartile	36.5 (14)	37.1 (16)	35.7 (14)	33.1 (15)	36.8 (14)	39.5 (15)	40.0 (14)
2nd highest quartile	42.0 (16)	35.1 (21)	41.3 (15)	42.4 (13)	41.7 (14)	44.7 (17)	45.9 (13)
Highest quartile	48.6 (17)	47.2 (19)	47.8 (16)	53.9 (21)	49.6 (17)	47.1 (17)	54.3 (16)
WPC							
Lowest quartile	29.7 (15)	31.2 (17)	27.6 (14)	25.2 (12)	31.2 (13)	29.2 (14)	31.5 (12)
2nd lowest quartile	36.0 (14)	30.9 (16)	36.9 (15)	35.5 (13)	36.8 (13)	39.1 (14)	40.0 (12)
2nd highest quartile	43.3 (14)	38.2 (14)	40.0 (14)	44.0 (12)	44.2 (13)	44.8 (16)	51.0 (15)
Highest quartile	51.8 (15)	54.2 (17)	52.4 (13)	52.4 (15)	51.0 (16)	50.7 (16)	52.3 (14)

TABLE 3. Standardized Effects of ERI and WPC (Beta Coefficient) on Burnout Risk for Overall Study Population and Stratified for Professional Groups

	Overall Staff	Nontertiary Administrative Staff	Nontertiary Nurses	Tertiary Administrative Staff	Tertiary Nurses	Academic Staff	Physicians
Age	0.10 ^c	0.26 ^b	0.06 NS	0.15 NS	0.02 NS	0.19 ^b	0.14 NS
Marital status							
Married (reference)	0	0	0	0	0	0	0
Single	0.07 ^b	-0.06 NS	0.05 NS	-0.07 NS	0.17 ^b	0.06 NS	0.05 NS
Divorced/widowed	0.03 NS	0.11 NS	-0.04 NS	-0.03 NS	-0.02 NS	0.05 NS	0.03 NS
Work volume	0.06 ^b	-0.01 NS	-0.06 NS	0.08 NS	0.14 ^b	0.14 ^a	0.15 NS
WPC	0.42 ^c	0.44 ^c	0.48 ^c	0.55 ^c	0.43 ^c	0.43 ^c	0.49 ^c
ERI	0.10 ^c	0.14 NS	0.09 NS	0.22 NS	0.20 ^c	0.20 ^b	0.27 ^c
Adjusted R ²	0.31	0.31	0.32	0.45	0.33	0.31	0.39
N	1,422	190	323	69	436	265	139

NS, not significant.

^aP < 0.05.^bP < 0.01.^cP < 0.001.

by general stress,¹² suggesting that ERI and WPC have direct effects on burnout that cannot be explained by stress in general.

Regarding our first research question, namely whether ERI or WPC is the stronger predictor in the health care setting, we found a much stronger association with WPC than with ERI. Hence, inter-role conflicts between work and private life, in particular strain spilling over from work to private life, seem to be more strongly associated with burnout risk than an imbalance between efforts spent and reward received. This is against theoretical expectations, as conceptually speaking burnout is a syndrome resulting of long-standing adverse working conditions.² However, our findings are in line with previous studies reporting a stronger association for WPC than ERI with burnout.^{12,20} Remarkably, a previous Swiss study among hospital employees focusing on relative importance of ERI and WPC on burnout risk reported a coefficient for WPC with respect to burnout is of the same strength ($\beta = 0.43$), whereas they reported that a stronger beta coefficient for ERI with respect to burnout risk ($\beta = 0.24$) was stronger than in our overall study population ($\beta = 0.10$).¹² However, when we compared these results with our findings in stratified analyses, we saw similar standardized coefficients for both WPC and ERI with respect to burnout risk among tertiary nurses and academic staff, whereas we find stronger associations for physicians.

As regards our second research question, whether the relative predictive importance of ERI and WPC with respect to burnout risk differs between professional groups and educational levels, we found that WPC is stronger associated with burnout than ERI across all professional groups. Whereas WPC is a strong predictor for burnout risk in all six groups, ERI is only weakly associated in three professional groups and not associated in the remaining three groups. Interestingly, ERI only plays a role for burnout risk in the tertiary medical professions, whereas in tertiary administrative staff, ERI as an explanatory factor was close to significant ($P = 0.05$). Our findings do not fully support the results of a previous Swiss study revealing different associations between these stress measures and burnout for the different medical professions: For therapists and medical-technical staff (which we had to merge into academic staff) as well as for physicians, ERI seemed to play a more important role than WPC with respect to burnout risk.¹² Although this is not the case in our study population, for tertiary nurses, academic staff, and physicians, ERI has a weak but significant impact on burnout risk, which is in line with previous studies.^{8,22} However, no associations between ERI and burnout risk were found in the other professional groups, which is at variance with previous

studies.^{12,22} Hence, imbalance of efforts spent and rewards received at work does not seem to play an important role with respect to burnout in primary and secondary educated hospital staff, whereas such an imbalance matters for tertiary educated staff. As higher educated professionals usually score higher on ERI and over-commitment scales,³² an imbalance of effort and reward might be more important for them with respect to burnout. Insufficient rewards such as having professional esteem/status were found to be important for burnout risk in physicians.¹¹

There are several limitations of this study. We cannot generalize our findings for health professionals, as our study population is a nonrepresentative sample of health care professionals in Switzerland and we excluded men from the sample due to their low number. Although we conceptualize ERI and WPC as antecedents or predictors of burnout, we cannot make causal inferences as our data are cross-sectional. Further, the response rate of 41% may result in nonresponse and selection bias. Due to the small number of participants in some professional groups, the standard errors are relatively large. Moreover, the low number in some professional groups required to merge some categories despite of different working conditions.

It can be concluded that interventions aiming at reducing work-related stress, especially increasing the compatibility of work and private life, are an effective approach to reduce the burnout risk of hospital employees and thus increase patients safety^{23,24} and continuity²⁵ and quality of care.¹⁰ The cause for ERI to matter only for burnout risk in tertiary-educated hospital employees remains unclear and should be further investigated. Hence, larger studies evaluating the different causes of burnout for hospital employees with different educational levels are needed.

REFERENCES

- Kristensen TS, Hannerz H, Høgh A, Borg V. The Copenhagen Psychosocial Questionnaire: a tool for the assessment and improvement of the psychosocial work environment. *Scand J Work Environ Health*. 2005;31:438–449.
- Maslach C, Leiter MP. Early predictors of job burnout and engagement. *J Appl Psychol*. 2008;93:498–512.
- McVicar A. Workplace stress in nursing: a literature review. *J Adv Nurs*. 2003;44:633–642.
- Marine A, Ruotsalainen J, Serra C, Verbeek J. Preventing occupational stress in healthcare workers (Review). *Cochrane Database Syst Rev*. 2006;CD002892.
- Shanafelt TD, Boone S, Tan L, et al. Burnout and satisfaction with work-life balance among US physicians relative to the general US population. *Arch Intern Med*. 2012;172:1377–1385.

6. Aiken LH, Sermeus W, Van den Heede K, et al. Patient safety, satisfaction, and quality of hospital care: cross sectional surveys of nurses and patients in 12 countries in Europe and the United States. *BMJ*. 2012;344:e1717.
7. Chou L-P, Li C-Y, Hu SC. Job stress and burnout in hospital employees: comparisons of different medical professions in a regional hospital in Taiwan. *BMJ Open*. 2014;4:e004185.
8. Bakker AB, Killmer CH, Siegrist J, Schaufeli WB. Effort-reward imbalance and burnout among nurses. *J Adv Nurs*. 2000;31:884–891.
9. Arigoni F, Bovier PA, Mermillod B, Waltz P, Sappino AP. Prevalence of burnout among Swiss cancer clinicians, paediatricians and general practitioners: who are most at risk? *Support Care Cancer*. 2009;17:75–81.
10. Shanafelt TD, Hasan O, Dyrbye LN, et al. Changes in burnout and satisfaction with work-life balance in physicians and the general US working population between 2011 and 2014. *Mayo Clin Proc*. 2015;90:1600–1613.
11. Ramirez AJ, Graham J, Richards MA, Cull A, Gregory WM. Mental health of hospital consultants: the effects of stress and satisfaction at work. *Lancet*. 1996;347:724–728.
12. Hämmig O, Brauchli R, Bauer GF. Effort-reward and work-life imbalance, general stress and burnout among employees of a large public hospital in Switzerland. *Swiss Med Wkly*. 2012;142. w13577.
13. Gray-Toft P, Anderson JG. Stress among hospital nursing staff: its causes and effects. *Soc Sci Med A*. 1981;15:639–647.
14. Shanafelt TD, Sloan JA, Habermann TM. The well-being of physicians. *Am J Med*. 2003;114:513–519.
15. Klein J, Grosse Frie K, Blum K, von dem Knesebeck O. Burnout and perceived quality of care among German clinicians in surgery. *Int J Qual Heal Care*. 2010;22:525–530.
16. Jenkins R, Elliott P. Stressors, burnout and social support: nurses in acute mental health settings. *J Adv Nurs*. 2004;48:622–631.
17. Lunau T, Siegrist J, Dragano N, Wahrendorf M. The association between education and work stress: does the policy context matter? *PLoS One*. 2015;10:e0121573.
18. Siegrist J. Adverse health effects of high-effort/low-reward conditions. *J Occup Health Psychol*. 1996;1:27–41.
19. Nübling M, Stöbel U, Hasselhorn H-M, Michaelis M, Hofmann F. Measuring psychological stress and strain at work: evaluation of the COPSOQ Questionnaire in Germany. *Psychosoc Med*. 2006;3:Doc05.
20. Nübling M, Seidler A, Garthus-Niegel S, et al. The Gutenberg Health Study: measuring psychosocial factors at work and predicting health and work-related outcomes with the ERI and the COPSOQ questionnaire. *BMC Public Health*. 2013;13:538.
21. Kozak A, Kersten M, Schillmöller Z, Nienhaus A. Psychosocial work-related predictors and consequences of personal burnout among staff working with people with intellectual disabilities. *Res Dev Disabil*. 2013;34:102–115.
22. Schulz M, Damkröger A, Heins C, et al. Effort-reward imbalance and burnout among German nurses in medical compared with psychiatric hospital settings. *J Psychiatr Ment Health Nurs*. 2009;16:225–233.
23. Welp A, Meier LL, Manser T. Emotional exhaustion and workload predict clinician-rated and objective patient safety. *Front Psychol*. 2015;5:1573.
24. Hall LH, Johnson J, Watt I, Tsipa A, O'Connor DB. Healthcare staff wellbeing, burnout, and patient safety: a systematic review. *PLoS One*. 2016;11:e0159015.
25. Heinen MM, van Achterberg T, Schwendimann R, et al. Nurses' intention to leave their profession: a cross sectional observational study in 10 European countries. *Int J Nurs Stud*. 2013;50:174–184.
26. Federal Statistical Office. Krankenhausstatistik 2015 - Standardtabellen. Neuchâtel; 2017.
27. Mercay C, Burla L, Widmer M. Gesundheitspersonal in der Schweiz - Bestandsaufnahme und Perspektiven bis 2030. Obsan Bericht 71. Neuchâtel; 2016.
28. Nübling M, Stöbel U, Hasselhorn H-M, Michaelis M, Hofmann F. Methods for assessing psychological strain - testing a measurement instrument [German]. Bremerhaven: Wirtschftsverlag NW; 2005:1–142.
29. Netemeyer RG, Boles JS, McMurrian R. Development and validation of work-family conflict and family-work conflict scales. *J Appl Psychol*. 1996;81:400–410.
30. Sorgaard KW, Ryan P, Dawson I. Qualified and Unqualified (N-RC) mental health nursing staff: minor differences in sources of stress and burnout. A European multi-centre study. *BMC Health Serv Res*. 2010;10:163.
31. Maslach C, Schaufeli WB, Leiter MP. Job burnout. *Annu Rev Psychol*. 2001;52:397–422.
32. Siegrist J, Starke D, Chandola T, et al. The measurement of effort-reward imbalance at work: European comparisons. *Soc Sci Med*. 2004;58:1483–1499.