

Electronic and vibrational contributions to first hyperpolarizability of donor–acceptor-substituted azobenzene

Robert Zalesny,^{1,a)} Ireneusz W. Bulik,^{1,b)} Wojciech Bartkowiak,¹ Josep M. Luis,² Aggelos Avramopoulos,³ Manthos G. Papadopoulos,³ and Przemysław Krawczyk⁴

¹Theoretical Chemistry Group, Institute of Physical and Theoretical Chemistry, Wrocław University of Technology, Wyb. Wyspiańskiego 27, 50–370 Wrocław, Poland

²Institute of Computational Chemistry and Department of Chemistry, University of Girona, Campus de Montilivi, 17071 Girona, Catalonia, Spain

³Institute of Organic and Pharmaceutical Chemistry, National Hellenic Research Foundation, Vas. Constantinou 48, GR–11635 Athens, Greece

⁴Department of Physical Chemistry, Collegium Medicum, Nicolaus Copernicus University, Kurpińskiego 5, 85–950 Bydgoszcz, Poland

(Received 1 July 2010; accepted 25 October 2010; published online 28 December 2010)

In this study we report on the electronic and vibrational (hyper)polarizabilities of donor–acceptor-substituted azobenzene. It is observed that both electronic and vibrational contributions to the electric dipole first hyperpolarizability of investigated photoactive molecule substantially depend on the conformation. The contributions to the nuclear relaxation first hyperpolarizability are found to be quite important in the case of two considered isomers (*cis* and *trans*). Although the double-harmonic term is found to be the largest in terms of magnitude, it is shown that the total value of the nuclear relaxation contribution to vibrational first hyperpolarizability is a result of subtle interplay of higher-order contributions. As a part of the study, we also assess the performance of long-range-corrected density functional theory in determining vibrational contributions to electric dipole (hyper)polarizabilities. In most cases, the applied long-range-corrected exchange-correlation potentials amend the drawbacks of their conventional counterparts. © 2010 American Institute of Physics. [doi:10.1063/1.3516209]

I. INTRODUCTION

Among various molecular properties, molecular hyperpolarizabilities (β , γ) have attracted special attention of scientific community for last three decades. As a result, new theoretical approaches and computational protocols have been proposed for evaluation of electronic and vibrational hyperpolarizabilities.¹ The interest in accurate determining of β and γ is certainly because of potential applications of compounds with large hyperpolarizabilities in photonics. For example a plethora of organic and organometallic systems have been analyzed as potential optical switches.^{2,3} Among them, azobenzene derivatives (see Fig. 1) are still the subject of intense research.^{4,5} Azobenzene molecule is known to undergo photoisomerization from *trans* to *cis* conformer. The metastable *cis* form can return to *trans* isomer either thermally or after irradiation.⁶ The quite large value of the barrier to *trans*–*cis* isomerization, which can be modulated with donor and acceptor groups,^{6,7} makes the azobenzene derivatives candidate systems for their use as optical switches. A large difference between the nonlinear optical properties of both conformers will increase their potential as optical switches. In this paper we will analyze the dependence of the first hyperpolarizability on the *trans*–*cis* isomerization of the azobenzene derivative. Both electronic and vibrational

contributions are computed in this work. The latter, although sometimes neglected, has an important weight in the value of the Pockels first hyperpolarizability.

Despite evident progress in the development of methods for evaluation vibrational (hyper)polarizabilities,^{8–20} their calculation for medium and large size molecules is much more computationally demanding than the electronic counterpart. The bottleneck of the calculation of the vibrational hyperpolarizabilities are the required high-order derivatives of energy and electronic properties with respect to the nuclear coordinates required for their evaluation. A significant step forward for the efficient evaluation of vibrational contributions to molecular hyperpolarizabilities was introduction of field-induced coordinates (FICs) method.^{21,22} In this technique, instead of $3N-6$ normal coordinates one introduces a relatively small number of FICs to describe vibrational hyperpolarizabilities. It is now well settled that the vibrational hyperpolarizabilities can be even higher in terms of magnitude than their electronic counterpart.²³ It has been shown only quite recently by Torrent-Sucarrat *et al.*^{16,24–26} and also by other authors^{27,28} that electron correlation effects have significant impact on the vibrational contributions to hyperpolarizabilities. In some cases, the $P^{\text{vib}}/P^{\text{e}}$ ($P = \alpha, \beta, \gamma$) ratio might depend crucially on the applied level of theory.^{29,30} Although an accurate determination of electronic and vibrational contributions to (hyper)polarizabilities of molecules in the gas phase is interesting in its own right, computational modeling of nonlinear vibrational spectroscopies is still very challenging task.^{14,31,32} Highly accurate post-Hartree–Fock

^{a)} Author to whom correspondence should be addressed. Electronic mail: robert.zalesny@pwr.wroc.pl.

^{b)} Permanent address: Department of Chemistry, Rice University, 6100 Main Street, Houston, Texas 77005-1892, USA.




FIG. 1. Schematic representation of the investigated system.

(HF) theories are in most cases too computationally expensive to be efficiently applied. At first glance, the density functional theory (DFT) might appear to be a reliable tool in the field of vibrational spectroscopies. Unfortunately, still little is known about the performance of DFT in determining vibrational hyperpolarizabilities.^{17,33–35} Hence, one of the goals of this contribution is the assessment of recently proposed long-range-corrected (LRC) functionals in computing vibrational properties. As far as electronic contributions to nonlinear optical properties are concerned, these functionals were proven to be successful^{36–38} in curing an overshoot problem. Too large values of electronic (hyper)polarizabilities, reported at first for conjugated chain-like molecules,^{39–41} have their roots in self-interaction error.⁴² In the case of LRC potentials, the attenuation of Coulomb interactions is achieved by introduction of Hartree–Fock exchange at long range. Although the testing of exchange–correlation potentials is essential step in process of development of efficient methods to determine electric dipole (hyper)polarizabilities, the central point of the present study is the analysis of the change in the electronic and vibrational second-order nonlinear response of substituted azobenzene upon photoisomerization. Also of interest is the analysis of the weight of the anharmonicity on the nuclear relaxation first hyperpolarizabilities of D/A-substituted azobenzene. For this purpose we shall employ the FICs approach combined with the second-order Møller–Plesset perturbation theory to account for electron correlation effects.

II. COMPUTATIONAL DETAILS

In this section we will present only brief description of the methods used to determine electronic and vibrational (hyper)polarizabilities. The reader unfamiliar with the applied methods may refer to the vast literature on the subject.⁴³ In the presence of an uniform electric field, the total energy of molecule can be expressed as a Taylor series:

$$E(F) = E(0) - \mu_i F_i - \frac{1}{2!} \alpha_{ij} F_i F_j - \frac{1}{3!} \beta_{ijk} F_i F_j F_k - \frac{1}{4!} \gamma_{ijkl} F_i F_j F_k F_l \dots, \quad (1)$$

where $E(0)$ denotes the field-free total energy and expansion coefficients μ , α , β , ... are dipole moment, polarizability, and first- to n -th order hyperpolarizability, respectively. Equation (1) conforms to the so-called T-convention⁴⁴ and shall be further used for calculations of electric dipole (hyper)polarizabilities. Within the Born–Oppenheimer approximation one may separate electronic and vibrational contributions to the property of interest P ($P = \alpha$, β , γ , ...).^{43,45}

$$P = P^e + P^{\text{vib}}. \quad (2)$$

Numerous approximations can be adopted to evaluate the two terms in the above equation. The details of the methods used to compute electronic and vibrational contributions in this work are given in the sections below.

A. Electronic (hyper)polarizabilities

In order to evaluate electronic contributions to (hyper)polarizabilities, we followed the procedure outlined by Kurtz and co-workers,^{46,47} and used Eq. (1) for numerical differentiation of the total energy of the molecular system with respect to the electric field. Many-body perturbation theory and coupled cluster calculations presented in this study were performed employing field-relaxed Hartree–Fock orbitals. In the case of orientationally invariant (hyper)polarizabilities:⁴³

$$\bar{\alpha} = \frac{1}{3} \sum_{i=x,y,z} \alpha_{ii}, \quad (3)$$

$$\bar{\beta} = \sum_{i=x,y,z} \frac{\mu_i \beta_i}{|\mu|},$$

$$\text{where } \beta_i = \frac{1}{5} \sum_{j=x,y,z} (\beta_{ijj} + \beta_{jij} + \beta_{jji}), \quad (4)$$

the constant value of the field amplitude (equal to 0.001 a.u.) was assumed. In the case of diagonal tensor elements the Romberg differentiation procedure was employed.⁴⁸

B. Vibrational (hyper)polarizabilities

The vibrational contributions to a property P (P^{vib}) as defined in Eq. (2) may be further divided into nuclear relaxation P^{NR} and curvature P^{curv} terms:⁴³

$$P^{\text{vib}} = P^{\text{NR}} + P^{\text{curv}}. \quad (5)$$

The P^{NR} is usually larger than the curvature contribution. For this reason, the latter term is not computed here. Using the Bishop–Kirtman (BK) square bracket nomenclature the nuclear relaxation contributions to static (hyper)polarizabilities read⁴⁹

$$\alpha^{\text{NR}}(0; 0) = [\mu^2]^{(0,0)}, \quad (6)$$

$$\beta^{\text{NR}}(0; 0, 0) = [\mu\alpha]^{(0,0)} + [\mu^3]^{(1,0)} + [\mu^3]^{(0,1)}, \quad (7)$$

where for $[P]^{(n,m)}$ expressions, n and m are the order of electrical and mechanical anharmonicity, respectively. The terms inside square brackets denote energy/property derivatives. For instance the $[\mu\alpha]^{(0,0)}$ square bracket term is given by^{8–10}

$$[\mu\alpha]^{(0,0)} = \frac{1}{2} P_{abc} \sum_{i=1}^{3N-6} \left(\frac{\partial \alpha_{ab}}{\partial Q_i} \right) \left(\frac{\partial \mu_c}{\partial Q_i} \right) \omega_i^{-2}, \quad (8)$$

where ω_i are the vibrational frequencies, a, b, c are the Cartesian coordinates x, y, z , and P_{abc} indicates a sum over all the perturbations of the indices abc . For dc-Pockels first hyperpolarizability, within the infinite frequency approximation, only the lowest order term is sufficient to calculate the complete NR contribution.⁴⁹

$$\beta^{\text{NR}}(-\omega; \omega, 0)_{\omega \rightarrow \infty} = \frac{1}{3} [\mu\alpha]^{(0,0)}. \quad (9)$$

TABLE I. Comparison of electronic and vibrational contributions to static first hyperpolarizability evaluated at various levels of theory.

	$\bar{\beta}^e$	$[\mu\alpha]^{(0,0)}$	$\bar{\beta}^e$	$[\mu\alpha]^{(0,0)}$
	<i>trans</i> isomer		<i>cis</i> isomer	
HF/6-31G(d)	1687	3843	494	8094
HF/6-31+G(d)	1848	3891	565	8107
MP2/6-31G(d)	4112	4412	1058	8516
MP2/6-31+G(d)	4414	4901	1115	9597

The $[P]^{(0,0)}$ terms (see Tables I, III and IV) were computed using standard BK formulae, i.e., by summation over $3N-6$ normal modes of vibration. In that event, the numerical differentiation of either the dipole moment or polarizability with respect to normal modes of vibration was performed to obtain relevant derivatives.

The vibrational contributions presented in Table V (see values in parentheses) were computed using the method proposed by Bishop *et al.*,⁹ and later implemented by Luis *et al.*¹² In this approach, the molecular geometry is first optimized in the presence of a static electric field while strictly maintaining the Eckart conditions.¹² Then the difference in the static electronic properties due to the change in geometry induced by the field is expanded as a power series in the field. Each

TABLE II. Purely electronic contributions to first-order hyperpolarizability calculated for the geometry optimized at the MP2/6-31G(d) level of theory.

	β_{zzz}^e	$\bar{\beta}^e$	β_{zzz}^e	$\bar{\beta}^e$
	<i>trans</i> isomer		<i>cis</i> isomer	
B3LYP/PC1	-6589	4007	-2443	1490
B3LYP/PC2	-6834	4159	-2254	1525
B3LYP/6-31G(d)	-6367	3845	-2340	1426
CAM-B3LYP/PC1	-5626	3404	-1610	1081
CAM-B3LYP/PC2	-5814	3520	-1642	1113
CAM-B3LYP/6-31G(d)	-5337	3229	-1525	1030
BLYP/PC1	-6787	4142	-2697	1851
BLYP/PC2	-7164	4372	-2783	1919
BLYP/6-31G(d)	-6651	4024	-2903	1781
LC-BLYP/PC1	-5182	3134	-1422	961
LC-BLYP/aug-PC1	-5266	3194	-1416	967
LC-BLYP/PC2	-5402	3274	-1468	1005
LC-BLYP/6-31G(d)	-4913	2959	-1504	913
BOP/PC1	-6832	4170	-2705	1854
BOP/PC2	-7189	4387	-2780	1917
BOP/6-31G(d)	-6641	4054	-2600	1786
LC-BOP/PC1	-5169	3126	-1415	957
LC-BOP/PC2	-5382	3252	-1641	995
LC-BOP/6-31G(d)	-4891	2958	-1339	911
HF/PC1	-3752	2262	-945	573
HF/PC2	-3858	2330	-872	589
HF/6-31G(d)	-3530	2123	-889	539
MP2/cc-pVDZ	-6668	4136	-1537	1038
MP2/cc-pVTZ	-6998	4346	-1599	1090
MP2/6-31G(d)	-6654	4112	-1565	1058
MP2/aug-cc-pVDZ	-7274	4503	-1659	1140
CCSD(T)/6-31G(d)	-6176	3881	-1542	1074

term in the expansion yields the sum of a static electronic (hyper)polarizability plus a nuclear relaxation vibrational term. For example, the change of the dipole moment is given by⁹

$$\Delta\mu_i(F, R_F) = a_{1,ij}F_j + \frac{1}{2}b_{1,ijk}F_jF_k + \dots, \quad (10)$$

where

$$a_{1,ij} = \alpha_{ij}^e(0;0) + \alpha_{ij}^{\text{NR}}(0;0), \quad (11)$$

$$b_{1,ijk} = \beta_{ijk}^e(0;0,0) + \beta_{ijk}^{\text{NR}}(0;0,0). \quad (12)$$

The argument R_F refers to the field-relaxed geometry of a molecule.

In order to evaluate explicitly the first-order electrical and mechanical contributions to the vibrational first-hyperpolarizability (see middle columns of Table V), an approach based on the FICs, which are linear combinations of field-free normal coordinates associated with the change in equilibrium geometry induced by a static electric field, has been employed.²¹ The displacement of the i -th field-free normal coordinate at the field-relaxed geometry is given by²¹

$$Q_i^F(F_x, F_y, F_z) = - \sum_a^{x,y,z} q_1^{i,a} F_a - \sum_{a,b}^{x,y,z} \left[q_2^{i,ab} - \sum_{j=1}^{3N-6} \frac{a_{21}^{ij,a}}{a_{ii}^{jj}} q_1^{j,b} + \sum_{j,k=1}^{3N-6} \frac{3a_{30}^{ijk}}{2a_{ii}^{jj}} q_1^{j,a} q_1^{k,b} \right] F_a F_b + \dots, \quad (13)$$

where

$$q_{nm}^{ij\dots ab\dots} = \frac{1}{n!m!} \times \left(\frac{\partial^{(n+m)} V(Q_1, \dots, Q_{3N-6}, F_x, F_y, F_z)}{\partial Q_i \partial Q_j \dots \partial F_a \partial F_b \dots} \right)_{Q=0, F=0}, \quad (14)$$

$$q_1^{i,a} = \frac{a_{11}^{i,a}}{2a_{20}^{ii}}, \quad q_2^{i,ab} = \frac{a_{12}^{i,ab}}{2a_{20}^{ii}}. \quad (15)$$

The indices i, j, k refer to normal coordinates whereas a, b, c label the Cartesian directions along molecular axes. a_{nm} involves the n -th and m -th derivatives of the potential energy $V(Q, F)$ with respect to normal coordinates and field components, respectively. The first-order FICs are sufficient to compute static nuclear relaxation contribution to the linear polarizability and first hyperpolarizability:²¹

$$\chi_1^a = - \sum_{i=1}^{3N-6} q_1^{i,a} Q_i. \quad (16)$$

Depending on the nonlinear optical property of interest, a small set of FICs, independent of the number of atoms in the

TABLE III. Nuclear relaxation and electronic contributions to molecular first hyperpolarizability calculated using the 6-31G(d) basis set. All values are given in atomic units.

	$\bar{\alpha}^{\text{NR}}(0;0)$	$\bar{\alpha}^e$	$\bar{\beta}^{\text{NR}}(-\omega;\omega,0)$	$\frac{1}{3}\bar{\beta}^e$	$\bar{\alpha}^{\text{NR}}(0;0)$	$\bar{\alpha}^e$	$\bar{\beta}^{\text{NR}}(-\omega;\omega,0)$	$\frac{1}{3}\bar{\beta}^e$
	<i>trans</i> isomer				<i>cis</i> isomer			
BLYP	73.6	238.4	2433	1364	91.4	195.7	4139	643
BOP	60.9	239.3	2122	1393	94.8	195.8	4381	658
B3LYP	58.7	217.1	2314	1357	87.3	178.5	3799	552
LC-BLYP	55.8	188.1	1860	963	81.0	161.2	3519	333
LC-BOP	57.1	188.4	1890	965	79.3	161.3	3500	330
CAM-B3LYP	55.9	195.6	2082	1092	81.5	165.0	3415	392
HF	46.6	172.3	1279	562	66.1	150.2	2697	165
MP2	49.2	188.4	1471	1371	62.7	157.8	2839	353

molecule, may be employed to calculate the total NR contribution instead of using all $3N-6$ normal coordinates. For example, from the three first-order FICs (one for each Cartesian direction) one can compute the nuclear relaxation contributions to the static polarizability and first hyperpolarizability, as

$$\alpha_{ab}^{\text{NR}}(0;0) = \frac{1}{2} \sum P_{ab} \frac{\partial \mu_a}{\partial \chi_1^b} \frac{\partial \chi_1^F}{\partial F_b} = [\mu^2]^{(0,0)}, \quad (17)$$

$$\begin{aligned} \beta_{abc}^{\text{NR}}(0;0,0) &= \sum P_{abc} \left[\frac{1}{2} \frac{\partial \alpha_{ab}}{\partial \chi_1^c} \frac{\partial \chi_1^F}{\partial F_c} \right. \\ &\quad + \frac{\partial^2 \mu_a}{\partial \chi_1^b \partial \chi_1^c} \frac{\partial \chi_1^F}{\partial F_b} \frac{\partial \chi_1^F}{\partial F_c} \\ &\quad \left. - \frac{1}{6} \frac{\partial^3 V}{\partial \chi_1^a \partial \chi_1^b \partial \chi_1^c} \frac{\partial \chi_1^F}{\partial F_a} \frac{\partial \chi_1^F}{\partial F_b} \frac{\partial \chi_1^F}{\partial F_c} \right] \\ &= [\mu\alpha]^{(0,0)} + [\mu^3]^{(1,0)} + [\mu^3]^{(0,1)}, \quad (18) \end{aligned}$$

where $\sum P_{ab\dots}$ indicates the sum over all permutations of the indices a, b, \dots . More details about this method can be found elsewhere.²¹

Derivatives with respect to the FIC coordinate were obtained numerically, at the field-free optimum geometry, by applying a number of steps along the FIC vector.

Computations with the aid of post-Hartree–Fock methods were done within the frozen-core approximation. Electronic and vibrational contributions to α and β were determined using the GAMESS US,⁵⁰ the GAUSSIAN 09,⁵¹ and the modified version of the DALTON program.⁵²

III. RESULTS AND DISCUSSION

As mentioned in Introduction, the investigated system undergoes a significant structural transformation upon elec-

tronic excitation. The geometries of the molecule optimized at the MP2/6-31G(d) level of theory are presented in electronic supplementary information (ESI).⁵³ The average difference in bond lengths presented in ESI in Fig. 1 is not significant and is equal to 0.012 Å. Thus, the main structural difference upon *trans*–*cis* transformation is the change of dihedral angles involving $N=N$ moiety. In a result, the electronic dipole moment decreases from 7.4 to 4.1 D. The changes are also reflected in the values of average electronic and vibrational static first hyperpolarizability (see Table I). The absolute values of $\Delta \bar{\beta}_{\text{trans} \rightarrow \text{cis}}^e$ (and $\Delta \bar{\beta}_{\text{trans} \rightarrow \text{cis}}^v$) might suggest that *trans*→*cis* isomerization leads to large changes in both vibrational and electronic structure. The reliability of the double-harmonic oscillator model, used for computing part of the data from Table I, in determining vibrational counterpart of molecular hyperpolarizabilities shall be discussed further. As it has already been shown, the ratio β^v/β^e might crucially depend on the level of theoretical approximation.^{25,30} Indeed, the values of electronic and vibrational hyperpolarizabilities seem to confirm previous findings. In the case of both conformers, the inclusion of electron correlation effects diminishes the β^v/β^e ratio approximately by a factor of 2. This is because the HF approximation significantly underestimates the values of β^e in comparison with the MP2 results. The latter method is not always a good reference point; however, as it will be shown later, it provides very reliable estimate of electronic contributions to β for investigated molecules. The other issue is the choice of the basis set. The 6-31+G(d) basis set was found by Torrent-Succarrat *et al.* to provide the results of “semiquantitative accuracy.”²⁵ The difference in vibrational first hyperpolarizability computed in this study using the two basis sets at the MP2 level of theory is not large and does not exceed 15%. Hence, we decided to employ the smaller 6-31G(d) basis set for computing the vibrational properties.

In order to elucidate the effect of the level of theory (including the influence of basis set) on the electronic contributions to β , the extensive computations were performed using both *ab initio* methods as well as several exchange-correlation potentials commonly used within the density functional theory framework. The results are presented in Table II. Contrary to the data discussed above, all results of computations from Table II correspond to fixed geometries optimized at the MP2/6-31G(d) level of theory. This allows to compare directly the values of purely electronic

TABLE IV. $[\mu\beta]^{(0,0)}$ contribution to the vibrational second-order hyperpolarizability (γ_{zzzz} , given in a.u.) calculated at various levels of theory.

	B3LYP/6-31G(d)	CAM-B3LYP/6-31G(d)	MP2/6-31G(d)
<i>trans</i> isomer	143×10^3	483×10^3	647×10^3
<i>cis</i> isomer	126×10^3	403×10^3	542×10^3

TABLE V. Harmonic and anharmonic contributions to the vibrational first-order hyperpolarizability calculated at the MP2/6-31G(d) level of theory. The values calculated using the HF/B3LYP/CAM-B3LYP methods with the 6-31G(d) basis set are given in parentheses. All values are given in atomic units and should be multiplied by 10^3 .

	$[\mu\alpha]_{zzz}^{(0,0)}$	$[\mu^3]_{zzz}^{(1,0)}$	$[\mu^3]_{zzz}^{(0,1)}$	β_{zzz}^{NR}
<i>trans</i> isomer	-5.7 (-5.2/-10.2/-8.8)	0.8	-2.1	-7.1 (-8.2/-21.8/-18.8)
<i>cis</i> isomer	-12.6 (-13.7/-18.2/-17.2)	-2.1	-4.2	-18.9 (-21.1/-22.9/-26.9)

(hyper)polarizabilities computed using various approaches. Even though the 6-31G(d) basis set is not easily comparable to the cc-pVXZ basis sets, the values of β determined using the MP2 method show an acceptable convergence with respect to the basis set size. As expected, augmentation of the cc-pVDZ basis set with diffuse functions further increases the values of first hyperpolarizability. It should be underlined that in the case of both conformers the differences in electronic contributions to β arising from the basis set used in calculations are not large and does not exceed 10%. Similarly to other organic compounds, the MP2 method tends to overestimate the absolute values of hyperpolarizabilities as compared to the CCSD(T) approach. Likewise, the differences are not significant. Based on the *ab initio* data presented in both tables, the MP2 method combined with 6-31G(d) basis set was selected as reliable and quite accurate level of approximation to study vibrational properties which are of primary interest for this study. Turning toward the results presented in Table II, let us comment on the performance of density functional theory in determining electronic contributions to β . Besides the Pople's 6-31G(d) basis set, we have also computed electronic hyperpolarizabilities using Jensen's pc-*n* basis sets as they are specifically designed for HF and DFT calculations.^{54,55} The pc-1 and pc-2 basis set in terms of size correspond to Dunning's correlation consistent cc-pVDZ and cc-pVTZ basis sets. The first conclusion is that *all* applied exchange-correlation potentials improve upon Hartree-Fock approximation. The latter method leads to significantly underestimated values of β in comparison with the MP2 or the CCSD(T) treatments. Secondly, one finds that the long-range corrected functionals do not improve systematically upon their conventional counterparts. It is quite striking that in the case of *trans* conformer they worsen the results both for diagonal and average hyperpolarizability. On the contrary, for *cis* isomer they provide smaller values of hyperpolarizability which are in better agreement with the MP2/CCSD(T) data.

In order to make an assessment of the density functional theory in determining nonlinear optical response, arising from nuclear motions, the nuclear relaxation contributions to α and β were determined for three commonly used exchange-correlation potentials as well as for their long-range-corrected counterparts. The results of computations are presented in Table III. In each case, the equilibrium geometry and the values of electronic and vibrational properties were determined at the same level of theory. First, the electronic and the NR contributions to polarizability are overestimated by conventional functionals and the LRC potentials partially cure the overshoot problem. This is true for both isomers. It is important to remark that for α^{NR} HF gives more accurate values than any of the studied DFT functionals. As far as the nuclear

relaxation first hyperpolarizability is concerned, the general trends are similar to α^{NR} . The nuclear relaxation contributions to β for dc-Pockels process are predicted significantly better by the LRC functionals than by their conventional counterparts. Again, conventional DFT functionals overshoot the MP2 result for β^{NR} whereas the LRC functionals partially cure this problem. And also for β^{NR} no functional improves the results of HF, which gives quite accurate values.

Table III shows that the nuclear relaxation contribution to Pockels hyperpolarizability for *cis* isomer is about two times larger than their counterpart for *trans* isomer. This vibrational contribution is given by the square bracket $[\mu\alpha]^{(0,0)}$ term, which involves summation over derivatives of electric dipole moment and polarizability with respect to nuclear displacements and vibrational frequencies [see Eq. (8)]. In ESI (cf. Table II), the contributions from individual normal modes of vibration to first hyperpolarizability for the two isomers are presented.⁵³ For the *cis* conformer the lowest frequency mode is found to determine solely vibrational contribution to β . Contrary, in the case of *trans* form the largest contribution comes from mode denoted as 26. A close look at dipole moment and polarizability derivatives with respect to given normal mode reveals that $\partial\mu_z^{\text{trans}}/\partial Q_{26} \gg \partial\mu_z^{\text{cis}}/\partial Q_1$ but $\partial\alpha_{zz}^{\text{trans}}/\partial Q_{26} \approx \partial\alpha_{zz}^{\text{cis}}/\partial Q_1$. On the other hand $\omega_{26}^{\text{trans}}$ is about 20 times larger than ω_1^{cis} . Thus, it is the vibrational frequency of the key normal mode that determines the nuclear relaxation contribution which causes that vibrational first hyperpolarizability is larger for *cis* rather than for *trans* conformer.

The working formulae for $\beta_{\text{dc-p}}^{\text{NR}}$ involve summation over derivatives of electric dipole moment and polarizability with respect to nuclear displacements. Since the values of electronic first hyperpolarizability seem to be much more sensitive to the XC potential used in calculations than the polarizability is, the nuclear derivatives of β should depend to a larger extent on the applied exchange-correlation potential. Indeed, the results of calculations presented in Table IV indicate significant discrepancy in $[\mu\beta]^{(0,0)}$ values obtained using conventional B3LYP functional and the MP2 method. The computed term makes a contribution to vibrational second hyperpolarizability. Contrary to the purely electronic hyperpolarizabilities of extended π -conjugated systems,⁴⁰ no overshoot is observed here, i.e., the values predicted by the B3LYP potential are significantly underestimated. The long-range correction improves the poor performance of the B3LYP functional. The ratio of the $[\mu\beta]^{(0,0)}$ values for both isomers is the same for the CAM-B3LYP functional and the MP2 method.

The values of various contributions to the vibrational first hyperpolarizability presented in Table V were performed for equilibrium geometries oriented in such a way that in each case the dipole moment vector was parallel to

Cartesian z axis. This geometry dependence of the results could increase the discrepancies of the different theoretical methods. Unfortunately, due to the complexity of the computational protocol used to determine the vibrational properties, we have not attempted to evaluate the full hyperpolarizability tensor. Nevertheless, some valuable conclusions regarding theoretical methods can be drawn based on the data presented in Table V. First, despite the small differences in geometries and in the orientation in Cartesian coordinate system, the nuclear relaxation contributions to β are predicted by both applied functionals to be similar in terms of magnitude. Secondly, the sum of first-order corrections (i.e., $\beta_{zzz}^{\text{NR}} - [\mu\alpha]_{zzz}^{(0,0)}$) is negative at all applied levels of theory. Thirdly, also for the anharmonic NR contribution to β the HF results are far more similar to the MP2, which based on Table II data we use as reference, than the two DFT functionals studied. However, the contribution through first order in electrical anharmonicity might be either positive (*trans* isomer) or negative (*cis* isomer). In the considered case, the lowest order of the BK perturbation theory seem to be only crude approximation to the vibrational static first hyperpolarizability. As pointed previously, the calculation of the higher order anharmonic terms for the full tensor have a high computational cost. However, at this point it is important to remark that first order $[\mu^3]$ term has no role in the calculation of nuclear relaxation contribution to Pockels first hyperpolarizability within the infinite frequency approximation [see Eq. (9)].

IV. SUMMARY

We have considered both electronic and vibrational (hyper)polarizabilities of donor–acceptor-substituted azobenzene. Nowadays, azobenzene derivatives are widely used as building blocks of hybrid systems of high nonlinear optical activity. As a part of a model study, we have assessed the performance of density functional theory in determining electric dipole (hyper)polarizabilities. In particular, recently proposed long-range-corrected exchange-correlation potentials were considered. As it turned out, in many cases they partially amend the drawbacks of their conventional counterparts, as for instance being successful in determining vibrational first and second hyperpolarizability. We have also noted that of the two contributions to first hyperpolarizability, namely electronic and vibrational, the former is influenced to a much larger extent by electron correlation effects. As a result, the Hartree–Fock method does not seem to be fortunate choice for determining relations between vibrational and electronic hyperpolarizabilities. Nevertheless, in this particular case, HF seems to be a good approach to calculate the nuclear-relaxation contribution values.

The nonlinear optical response of D/A-substituted azobenzene crucially depends on the conformation. Orientationally invariant electronic first hyperpolarizability is four times larger for *trans* than for *cis* isomer. The vibrational contributions behave differently, i.e., they are approximately two times larger for the latter isomer. We have also determined, with the inclusion of electron correlation effects by means of the MP2 method, the anharmonic contributions to vibrational first hyperpolarizability through the first order in

electrical and mechanical anharmonicity. We noticed that the value of the so-called nuclear relaxation contribution to vibrational first hyperpolarizability is a subtle balance between the three contributions among which the double-harmonic term is found to be the largest in terms of magnitude. However, the values of individual components, and particularly their signs, are warning not to neglect the first-order anharmonic corrections as they might be quite important and strongly depend on the molecular geometry.

ACKNOWLEDGMENTS

This work was supported by computational grants from Wrocław Center for Networking and Supercomputing (WCSS) and ACK Cyfronet. One of the authors (R.Z.) is the recipient of the fellowship cofinanced by European Union within European Social Fund. R.Z. would also like to acknowledge support from a grant from Iceland, Liechtenstein, and Norway through the EEA Financial Mechanism—Scholarship and Training Fund. The support from the European Union (MTKD-CT-2006-042488), the Barcelona Supercomputing Center, and grants to access the TeraGrid facility (TG-CHE080063N) and the DEISA infrastructure are gratefully acknowledged. R.Z. and W.B. are grateful for financial support from Wrocław University of Technology.

- ¹P. Sałek, O. Vahtras, T. Helgaker, and H. Ågren, *J. Chem. Phys.* **117**, 9630 (2002).
- ²B. J. Coe, in *Nonlinear Optical Properties of Matter*, edited by M. G. Papadopoulos, J. Leszczynski, and A. Sadlej (Springer-Verlag, Berlin, 2006).
- ³K. A. Green, M. P. Cifuentes, T. C. Corkery, M. Samoc, and M. G. Humphrey, *Angew. Chem., Int. Ed.* **48**, 7867 (2009).
- ⁴S. Kawata and Y. Kawata, *Chem. Rev.* **100**, 1777 (2000).
- ⁵S. Wagner, F. Leyssner, C. Koedel, S. Zarwell, R. Schmidt, M. Weinelt, K. Rueck-Braun, M. Wolf, and P. Tegeder, *Phys. Chem. Chem. Phys.* **11**, 6242 (2009).
- ⁶N. Nishimura, S. Kosako, and Y. Sueishi, *Bull. Chem. Soc. Jpn.* **57**, 1617 (1984).
- ⁷C. R. Crecca and A. E. Roitberg, *J. Phys. Chem. A* **110**, 8188 (2006).
- ⁸D. M. Bishop and B. Kirtman, *J. Chem. Phys.* **95**, 2646 (1991).
- ⁹D. M. Bishop, M. Hasan, and B. Kirtman, *J. Chem. Phys.* **103**, 4157 (1995).
- ¹⁰B. Kirtman, J. M. Luis, and D. M. Bishop, *J. Chem. Phys.* **108**, 10008 (1998).
- ¹¹J. M. L. J. Martí, M. Duran, J. L. Andrés, and B. Kirtman, *J. Chem. Phys.* **108**, 4123 (1998).
- ¹²J. M. Luis, M. Duran, J. L. Andrés, B. Champagne, and B. Kirtman, *J. Chem. Phys.* **111**, 875 (1999).
- ¹³O. Quinet, B. Kirtman, and B. Champagne, *J. Chem. Phys.* **118**, 505 (2003).
- ¹⁴R. Zaleśny, W. Bartkowiak, and B. Champagne, *Chem. Phys. Lett.* **380**, 549 (2003).
- ¹⁵D. M. Bishop, J. M. Luis, and B. Kirtman, *J. Chem. Phys.* **116**, 9729 (2002).
- ¹⁶M. Torrent-Sucarrat, J. M. Luis, and B. Kirtman, *J. Chem. Phys.* **122**, 204108 (2005).
- ¹⁷A. J. Thorvaldsen, K. Ruud, and M. Jaszuński, *J. Phys. Chem. A* **112**, 11942 (2008).
- ¹⁸J. Kongsted and O. Christiansen, *J. Chem. Phys.* **127**, 154315 (2007).
- ¹⁹O. Christiansen, *Phys. Chem. Chem. Phys.* **9**, 2942 (2007).
- ²⁰P. Seidler, J. Kongsted, and O. Christiansen, *J. Phys. Chem. A* **111**, 11205 (2007).
- ²¹J. M. Luis, M. Duran, B. Champagne, and B. Kirtman, *J. Chem. Phys.* **113**, 5203 (2000).
- ²²J. M. Luis, M. Duran, and B. Kirtman, *J. Chem. Phys.* **115**, 4473 (2001).
- ²³D. P. Shelton and L. Ulivi, *J. Chem. Phys.* **89**, 149 (1988).

- ²⁴M. Torrent-Sucarrat, M. Sola, M. Duran, J. M. Luis, and B. Kirtman, *J. Chem. Phys.* **116**, 5363 (2002).
- ²⁵M. Torrent-Sucarrat, M. Sola, M. Duran, J. M. Luis, and B. Kirtman, *J. Chem. Phys.* **118**, 711 (2003).
- ²⁶M. Torrent-Sucarrat, M. Sola, M. Duran, J. M. Luis, and B. Kirtman, *J. Chem. Phys.* **120**, 6346 (2004).
- ²⁷W. Bartkowiak, W. Niewodniczański, T. Misiaszek, and R. Zaleśny, *Chem. Phys. Lett.* **411**, 8 (2005).
- ²⁸R. Zaleśny, W. Bartkowiak, P. Toman, and J. Leszczynski, *Chem. Phys.* **337**, 77 (2007).
- ²⁹B. Champagne, T. Legrand, E. A. Perpète, O. Quinet, and J. M. André, *Collect. Czech. Chem. Commun.* **63**, 1295 (1998).
- ³⁰R. Zaleśny, M. G. Papadopoulos, W. Bartkowiak, and A. Kaczmarek, *J. Chem. Phys.* **129**, 134310 (2008).
- ³¹O. Quinet, *Int. J. Quantum Chem.* **104**, 727 (2005).
- ³²E. Botek and B. Champagne, *Appl. Phys. B* **74**, 627 (2002).
- ³³E. A. Perpète, B. Champagne, and B. Kirtman, *Phys. Rev. B* **61**, 13137 (2000).
- ³⁴M. R. Pederson, T. Baruah, P. B. Allen, and C. Schmidt, *J. Chem. Theory Comput.* **1**, 590 (2005).
- ³⁵R. Zaleśny, G. Wójcik, I. Mossakowska, W. Bartkowiak, A. Avramopoulos, and M. G. Papadopoulos, *J. Mol. Struct.: THEOCHEM* **907**, 46 (2009).
- ³⁶B. Kirtman, S. Bonness, A. Ramirez-Solis, B. Champagne, H. Matsumoto, and H. Sekino, *J. Chem. Phys.* **128**, 114108 (2008).
- ³⁷O. Loboda, R. Zaleśny, A. Avramopoulos, J. M. Luis, B. Kirtman, N. Tagmatarchis, H. Reis, and M. G. Papadopoulos, *J. Phys. Chem. A* **113**, 1159 (2009).
- ³⁸D. Jacquemin, E. A. Perpète, M. Medved, G. Scalmani, M. J. Frisch, R. Kobayashi, and C. Adamo, *J. Chem. Phys.* **126**, 191108 (2007).
- ³⁹B. Champagne, E. A. Perpète, S. J. A. van Gisbergen, E. J. Baerends, J. G. Snijders, C. Soubra-Ghaoui, K. A. Robins, and B. Kirtman, *J. Chem. Phys.* **109**, 10489 (1998b).
- ⁴⁰B. Champagne, E. A. Perpète, D. Jacquemin, S. J. A. van Gisbergen, E. J. Baerends, C. Soubra-Ghaoui, K. A. Robins, and B. Kirtman, *J. Phys. Chem. A* **104**, 4755 (2000).
- ⁴¹S. J. A. van Gisbergen, P. R. T. Schipper, O. V. Gritsenko, E. J. Baerends, J. G. Snijders, B. Champagne, and B. Kirtman, *Phys. Rev. Lett.* **83**, 694 (1999).
- ⁴²A. Dreuw and M. Head-Gordon, *Chem. Rev.* **105**, 4009 (2005).
- ⁴³D. Bishop and P. Norman, *Handbook of Advanced Electronic and Photonic Materials and Devices*, Vol. 9, edited by H. S. Nalwa (Academic, San Diego, 2001).
- ⁴⁴A. Willets, J. Rice, D. Burland, and D. Shelton, *J. Chem. Phys.* **97**, 7590 (1992).
- ⁴⁵B. Champagne and B. Kirtman, *Handbook of Advanced Electronic and Photonic Materials and Devices*, Vol. 9, edited by H. S. Nalwa (Academic, San Diego, 2001).
- ⁴⁶H. Kurtz, J. Stewart, and K. Dieter, *J. Comput. Chem.* **11**, 82 (1990).
- ⁴⁷H. A. Kurtz and D. S. Dudis, *Reviews in Computational Chemistry*, Vol. 12, edited by K. B. Lipkowitz and D. B. Boyd (Wiley-VCH, Weinheim, 1998), pp. 241–279.
- ⁴⁸M. Medved, M. Stachová, D. Jacquemin, J.-M. André, and E. A. Perpète, *J. Mol. Struct.: THEOCHEM* **847**, 39 (2007).
- ⁴⁹B. Kirtman, B. Champagne, and J. M. Luis, *J. Comput. Chem.* **21**, 1572 (2000).
- ⁵⁰M. W. Schmidt, K. K. Baldrige, J. A. Boatz, S. T. Elbert, M. S. Gordon, J. H. Jensen, S. Koseki, N. Matsunaga, K. A. Nguyen, S. Su, T. L. Windus, M. Dupuis, and J. A. Montgomery, *J. Comput. Chem.* **14**, 1347 (1993).
- ⁵¹M. J. Frisch, G. W. Trucks, H. B. Schlegel *et al.*, Gaussian 09 Revision A.1, Gaussian Inc., Wallingford, CT, 2009.
- ⁵²DALTON, a molecular electronic structure program, Release 2.0 (2005), see <http://www.kjemi.uio.no/software/dalton/dalton.html>.
- ⁵³See supplementary material at <http://dx.doi.org/10.1063/1.3516209> for vibrational analysis.
- ⁵⁴F. Jensen, *J. Chem. Phys.* **115**, 9113 (2001).
- ⁵⁵F. Jensen, *J. Chem. Phys.* **116**, 7372 (2002).